LETTERS TO THE EDITOR

Send your letters to the Editor, British Dental Journal, 64 Wimpole Street, London, W1G 8YS
Email bdj@bda.org. Priority will be given to letters less than 500 words long. Authors must sign the letter, which may be edited for reasons of space. Readers may now comment on letters via the *BDJ* website (www.bdj.co.uk). A 'Readers' Comments' section appears at the end of the full text of each letter online.

PHARMACEUTICALS

Timing of extraction

Sir, I read with great interest the recently published paper *National clinical guidelines for the extraction of first permanent molars in children* (*BDJ* 2014; 217: 643) where the authors stated that evidence of radiographic root bifurcation development in the second permanent molars (SPM) is the ideal clinical radiographic clue for enforced extraction of compromised first permanent molars (FPM) which produced the best occlusal position for SPM.¹ This radiographic clue corresponds to Demirjian Stage E early bifurcation and Stage F late bifurcation of SPM.² It should be mentioned that this radiographic clue does not apply to all SPM as there are some clinical situations where bifurcation development does not occur at all as in cases of pyramidal roots, or bifurcation development is delayed as in cases of taurodontism.

Taurodontism has a racial variation with a prevalence rate of 0.25–11.3%, and in particular the SPM are the most commonly affected teeth. It is classified into three categories depending on the degree of apical displacement of bifurcation area.^{3,4} The clinical dilemma is with mesotaurodont (moderate) and hypertaurodont (severe) types where bifurcation development is severely delayed and can affect the timing of enforced extraction of FPM. Furthermore, the behaviour of bodily movement of taurodontic SPM after extraction of FPM is not known and whether they succeed in ultimate space closure or not. The only clue we have for the possibility of development of taurodontic SPM is their possible association with other dental anomalies including oligodontia, amelogenesis imperfecta, supernumerary teeth, pulpal calcification and microdontia-dens invaginatus.^{3,4}

F. B. Rihani, by email

1. Cobourne M T, Williams A, Harrison M. National clinical guidelines for the extraction of first permanent molars in children. *Br Dent J* 2014; 217: 643–648.
2. Teo T K, Ashley P F, Parekh S, Noar J. The evaluation of spontaneous space closure after the extraction of first permanent molars. *Eur Arch Paediatr Dent* 2013; 14: 207–212.

NEWER ORAL ANTICOAGULANTS

Sir, your article¹ and other sources have noted the main advantages and concerns about the newer target-specific oral anticoagulants (NOACs) such as dabigatran (a direct thrombin inhibitor) and rivaroxaban, edoxaban, and apixaban (direct Factor X inhibitors).² Initially developed and marketed as not requiring routine laboratory monitoring, it is clear that in several situations, knowing the level of anticoagulant activity of the NOACs is imperative for patients' safety and optimal dosing.³ Such situations include before invasive emergency surgery, when there is spontaneous/traumatic haemorrhage, in the very elderly, extremes of body weight, in those with renal impairment or in overdose situations.⁴

To date, the inability to directly reverse the NOACs and to monitor effects have been seen as their main drawbacks. The preferred method for monitoring NOAC according to latest British guidelines⁴ include the ecarin clotting time and the dilute thrombin time for monitoring dabigatran, and the anti-factor Xa assays for monitoring the direct factor X inhibitors.

Now a recent phase I clinical trial has appeared⁵ in which the anticoagulant effects of the factor X inhibitor edoxaban has been successfully and safely reversed by a novel agent, PER977, which, within 10–30 minutes, restored baseline haemostasis. PER977 had been shown earlier, *in vitro* and in animal studies, to bind and reverse the anticoagulant effect of all the above NOACs. Phase II clinical trials are underway on what promises to be an advance in this important field of anticoagulation.

A. N. Robinson, Singapore
C. Scully, London

1. Syeed N, Ansell M, Sood V. Dabigatran (Pradaxa®): surgeon's friend or foe? *Br Dent J* 2014; 217: 623–626.
2. Scully C. *Scully's medical problems in dentistry*, 7th ed. Churchill Livingstone, Elsevier, 2014.
3. Cohen D. Dabigatran: how the drug company withheld important analyses. *BMJ* 2014; 349: g4670.
4. Kitchen S, Gray E, Mackie I, Baglin T, Makris M. Measurement of non-coumarin anticoagulants and their effects on tests of Haemostasis: Guidance from the British Committee for Standards in Haematology. *Br J Haematol* 2014; 166: 830–841.
5. Ansell J E, Bakhr S H, Laulicht B E *et al*. Use of PER977 to reverse the anticoagulant effect of edoxaban. *N Engl J Med* 2014; 371: 2141–2142.

DOI: 10.1038/sj.bdj.2015.151

3. Jafarzadeh H, Azarpazhooh A, Mayhall J T. Taurodontism: a review of the condition and endodontic treatment challenges. *Int Endod J* 2008; 41: 375–388.
4. Haskova J E, Gill D S, Figueiredo J A, Tredwin C J, Naini F B. Taurodontism – a review. *Dent Update* 2009; 36: 235–236, 239–240, 243.

DOI: 10.1038/sj.bdj.2015.150

ORAL SURGERY

Checklists and double checks

Sir, I read with interest the article by Saksena *et al.*¹ describing the development of a checklist applicable to outpatient settings to 'prevent wrong tooth extraction'. The medical community has learned valuable lessons in 'error avoidance' from the experience of aviation industry.² To err is human and the importance of checklists and a double check system by another competent person (eg a co-pilot) is well recognised

in the aviation industry. With regards to prevention of 'wrong tooth extractions', I would like to point out the desirability (or even necessity) of having a second appropriately trained person who can double check the correct tooth identification prior to extraction. There are certain instances in which mere counting of teeth cannot ensure the correct identification of the tooth to be extracted eg distinction between a first and second premolar in the absence of a canine tooth with space closed. A minimum knowledge of dental anatomy to enable correct tooth identification, eg by the assistant nurse, could be considered essential to perform a meaningful double check.

In the outpatient settings of United Kingdom hospital dental services, dental surgeons are sometimes assisted by registered general nurses, who might not be

competent (or adequately knowledgeable in dental anatomy) to perform this double check safety verification. Furthermore, when dental/oral surgeons operate under general anaesthesia, dental nurses are NOT allowed to assist in an operating theatre environment and the assistant nurses are always general nurses.

Current rules do not allow dental nurses to attend the operating theatres, as their training syllabus does not fulfill the requirements for operating room attendance.³ I believe this needs to be debated by educators and regulators if we are to eliminate the possibility of future never events.

B. A. Beyqi, by email

1. Sakseena A, Pemberton M N, Shaw A, Dickson S, Ashley M P. Preventing wrong tooth extraction: experience in development and implementation of an outpatient safety checklist. *Br Dent J* 2014; **217**: 357–362.
2. Gawande A. *The checklist manifesto: how to get things right*. Metropolitan Books/Henry Holt & Company, 2010.
3. AORN (Association of perioperative Registered Nurses) Position Statements Related to First Assisting. Available at: [http://www.aorn.org/Clinical_Practice/RNFA_Resources/First_Assisting_\(RNFA\).aspx](http://www.aorn.org/Clinical_Practice/RNFA_Resources/First_Assisting_(RNFA).aspx) (accessed 10 February 2015).

DOI: 10.1038/sj.bdj.2015.152

ORAL HEALTH

Reverse referral

Sir, the oral health of 102 patients (age 14–88) admitted for 24 hours of intravenous antibiotics for odontogenic infections was considered over 12 months in a Surrey district general hospital. Orthopantomograms were used to assess dental disease and restorative status. Three patients required ITU admission.

The average number of carious teeth (over two thirds into dentine) was 3.1; endodontically treated teeth was 0.9 and 2.3 teeth had radiographic evidence of apical pathology. The most common infected source tooth was the mandibular first molar and 10.7% of infected source teeth were root filled. No significant radiographic bone loss patterns were identified. The average number of restored (non-root filled) teeth was 5.0.

Whilst assessment of oral health using radiographic examination alone is insufficient, it nevertheless provides a general overview of the dentition in a secondary care setting. Full oral health screenings are seen as irrelevant, time consuming and often intolerable when patients are systemically unwell from odontogenic infection. The data show that patients admitted tend to have untreated decay in multiple teeth which may be a source of future odontogenic infection, and very few source teeth have had endodontic therapy.

Patients are usually treated solely for the infected source tooth and discharged with the hope that the rest of the dentition will be managed by primary care services. However, on questioning, the overwhelming majority of such patients have no primary care dentist due to financial, social and psychological reasons; these patients may thus return with similar episodes of infection and morbidity which presents a cost burden on public health care facilities.^{1,2} There has been a 62% increase in the number of patients who require admission for surgical treatment of spreading odontogenic infections.³ The number of admissions and bed days as a result of drainage of a dental abscess almost doubled between 1998–99 and 2005–06.⁴

Whilst referrals to maxillofacial departments for routine dentoalveolar services from primary care are common, a reverse pathway should also be firmly established to ensure patients presenting in secondary care are followed up by primary care or community dental services on discharge.

F. Jamil, London

1. Jundt J S, Gutta R. Characteristics and cost impact of severe odontogenic infections. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2012; **114**: 558–566.
2. Ahmad N, Abubaker A O, Laskin D M, Steffen D. The financial burden of hospitalization associated with odontogenic infections. *J Oral Maxillofac Surg* 2013; **71**: 656–658.
3. Burnham R, Bhandari R, Bridle C. Changes in admission rates for spreading odontogenic infection resulting from changes in government policy about the dental schedule and remunerations. *Br J Oral Maxillofac Surg* 2011; **49**: 26–28.
4. Thomas S J, Atkinson C, Hughes C, Revington P, Ness A R. Is there an epidemic of admissions for surgical treatment of dental abscesses in the UK? *BMJ* 2008; **336**: 1219–1220.

DOI: 10.1038/sj.bdj.2015.153

Turn off the tap

Sir, I would like to make further comments to J. Hartley's letter on *Spit don't rinse*.¹ One response to this letter cited the evidence from some well-designed studies,² while another quoted from a paper recommending the 'spit don't rinse' message as an integral part of oral hygiene instruction.³ However, an additional environmental benefit has not been mentioned.

Research by SaveWaterSaveMoney has revealed that 64% of 7–10-year-olds admitted to leaving the tap running while brushing their teeth. Turning the tap off while brushing and not rinsing the mouth out when finished are two simple yet effective ways of saving water in the bathroom. Not only could it save around 12 litres of water every time, it also saves money.⁴

The long-term effects of 'spit don't rinse' will not only preserve teeth, but also save water. That's why spitting is better for the environment too!

C. A. Yeung, Bothwell

1. Hartley J. Spit don't rinse. *Br Dent J* 2014; **217**: 206.
2. Wanless M. Spitting evidence. *Br Dent J* 2014; **217**: 612.
3. McCall D R. Integral to oral hygiene. *Br Dent J* 2014; **217**: 612.
4. British Dental Health Foundation. Why spitting is better for the environment! 10 July 2012. Available at: <http://www.dentalhealth.org/news/details/619> (accessed December 2014).

DOI: 10.1038/sj.bdj.2015.154

ORTHODONTICS

Fast and furious

Sir, P. Huntley (*BDJ* 2015; **218**: 2–3) sadly reflects how many confuse even basic differences between short-term orthodontics (STO) and Fastbraces, which is a comprehensive orthodontic system designed to be easy to use and with less apical resorption¹ than traditional systems, offering a significant time saving and innovative approach where a rectangular wire is used in a triangular bracket giving 3D control from day one.

Increased complications in traditional orthodontics are associated with their longer treatment times extending beyond a year.² Unfortunately, older traditional rectangular bracket mechanics tend to be high-force and simply do not recommend rectangular wire usage routinely from day one.³ Thus, traditionally it is loose sloppy round wires that are used initially with mainly tipping mechanics, the very same type of STO treatment some specialists have recently been declaring are *higher* risk, waving apices around etc.⁴ There are no round wire disadvantages in the FastBraces system.

The single reference quoted by Huntley relates to ligatures in round wire ortho systems, a study clearly nothing to do with FastBraces' unique triangular brackets and rectangular wire 3D torque system from day one. It is not often one has a system new to the UK that has a great track record and where some 80% of orthodontic cases can be completed well and safely by the humble GDP; this neatly rebalances orthodontics with other common dental disciplines where only maybe 20% of perio, endo, restorative, oral surgery etc complex cases need referring out to a higher trained and valued specialist.

When it comes to ethics, one must also wonder are some orthodontic specialists who imply all ortho cases should be referred to them, totally altruistic in their motives – a recent USA survey found