The eye limits the brain's learning potential

SUBJECT AREAS:

BEHAVIOUR

PLASTICITY

DEVELOPMENT

SENSORY SYSTEMS

Jiawei Zhou¹, Yudong Zhang^{2,3}, Yun Dai^{2,3}, Haoxin Zhao^{2,3}, Rong Liu¹, Fang Hou¹, Bo Liang^{2,3}, Robert F. Hess⁴ & Yifeng Zhou¹

¹CAS Key Laboratory of Brain Function and Disease, and School of Life Sciences, University of Science and Technology of China, Hefei 230027, ²Institute of Optics and Electronics, Chinese Academy of Sciences, Chengdu, China, ³The Key Laboratory on Adaptive Optics, Chinese Academy of Sciences, Chengdu 610209, China, ⁴McGill Vision Research, Department of Ophthalmology, McGill University, Montreal, Canada.

Received

2 January 2012

Accepted

23 March 2012

Published

16 April 2012

The concept of a critical period for visual development early in life during which sensory experience is essential to normal neural development is now well established. However recent evidence suggests that a limited degree of plasticity remains after this period and well into adulthood. Here, we ask the question, “what limits the degree of plasticity in adulthood?” Although this limit has been assumed to be due to neural factors, we show that the optical quality of the retinal image ultimately limits the brain potential for change. We correct the high-order aberrations (HOAs) normally present in the eye's optics using adaptive optics, and reveal a greater degree of neuronal plasticity than previously appreciated.

Correspondence and requests for materials should be addressed to Y.-D.Z. (ydzhang@ioe.ac.cn) or Y.-F.Z. (zhouy@ustc.edu.cn)

The pioneering work of Hubel and Wiesel¹ established the concept of a critical period for visual development early in life during which sensory experience is essential to normal neural development. Although this is a fundamental concept in neurobiology it is also now recognized that some limited plasticity remains after this period well into adulthood^{2–5}. During recent decades, numerous studies have shown that a range of visual functions in normal adult subjects can be improved as a result of intensive training (termed perceptual learning). These functions include contrast sensitivity^{6–10}, motion perception^{11–13} and object recognition^{14,15}. However, there is abundant evidence to indicate that this kind of learning-induced plasticity in adults while being possible is also very limited in extent^{8,9,16–18}. We wished to know what limits visual improvements that can occur as a consequence of perceptual learning in the normal adult. This refers directly to the mechanisms of brain plasticity that operate beyond the critical period.

In the adult visual system, uncorrectable optical aberrations limit the quality of the retinal image^{19,20}, even when defocus is corrected by sphero-cylindrical lenses^{21–23}. These aberrations, termed higher order aberrations (HOAs)¹⁹, exist throughout the life span^{24–27}. We wondered if HOAs set a fundamental limit to the benefits obtained from perceptual learning for the adult visual system.

The aim of the current study is to assess whether perceptual learning in normal adults is limited by the eye's optical quality. To address this we measure the effects of perceptual learning on visual sensitivity with and without HOAs-correction, using a real-time closed-loop adaptive optics visual stimulator system²⁸ (for details, see Supplementary online). We found larger and more robust contrast sensitivity improvements when the HOAs were corrected than when they were left uncorrected. We show that this is not due to the better optical quality per se or by the brain's ability to utilize this information but a consequence of improved perceptual learning using images of higher optical quality. This also transferred to a significant improvement of visual acuity in the HOAs-corrected perceptual learning group, compared with that of the HOAs-uncorrected group. We confirm previous reports of brain plasticity well beyond the critical period and show that its benefits are even larger if the eye's higher-order optical aberrations are corrected.

Results

Visually normal adults were separated into two training groups, in one (Group1 – 13 subjects) training was undertaken under the HOAs-corrected condition whereas in the other (Group2 – 8 subjects) training was undertaken under normal viewing condition (i.e. HOAs-uncorrected). Learning effects were then analyzed in accordance with individuals' pre-training baseline conditions. In both cases, 10 training sessions of 1 hour were undertaken with a high spatial frequency stimulus. For each subject, a high spatial frequency was selected for training that corresponded to a pre-training contrast threshold of 0.4. In particular, subjects in Group1 were

trained using spatial frequency when contrast threshold at HOAs-corrected condition is 0.4; subjects in Group2 were trained using spatial frequency when contrast threshold at HOAs-uncorrected condition is 0.4. Correction of HOAs has been shown to improve contrast sensitivity at all spatial frequencies but more so at higher spatial frequencies^{29,30}. The contrast sensitivity benefits that we found when the HOAs were corrected were similar to that previously reported by Yoon and Williams³⁰ for a 3 mm pupil (see Supplementary Table S1 online). After training, Group1, the HOAs-corrected group, showed a 5.39 dB (86.1%) significant (Paired Samples Test, $t(12) = -7.66$, $P = 0.0000059$, 2-tailed) improvement of contrast sensitivity. The slope of the average learning curve was 0.41 log units, and reached a plateau after 7.35 training sessions (Fig. 1a). Group2, the HOAs-uncorrected group, showed a smaller (3.42 dB or 48.2%) but also significant (Paired Samples Test, $t(7) = -2.62$, $P = 0.03$, 2-tailed) improvement. The slope of the average learning curve was 0.35 log unit, and reached a plateau after 4.12 training sessions (Fig. 1a- individual learning curves data are given in Supplementary Table S2 online).

Contrast sensitivity functions (CSFs), which measure sensitivity to stimuli of different spatial frequencies, were measured at pre- and post-training stages for both groups. There were significant improvements after training in both Group1 (Fig. 1b, post- vs. pre-training: $F(1, 12) = 75.43$, $P < 0.00001$) and Group2 (Fig. 1c, post- vs. pre-training: $F(1, 7) = 5.46$, $P = 0.05$). The average magnitude of the contrast sensitivity improvements across observers and spatial frequencies were 3.11 dB and 1.31 dB in Group1 and Group2, respectively (Fig. 1d). For the subjects in Group1 and Group2 who showed significant contrast sensitivity improvement (all 13 subjects in Group1 and only 4 subjects in Group2. For method, see Supplementary online), while the magnitude of improvements at the training spatial frequency were not significantly different (Independent Samples Test, $t(15) = -0.58$, $P = 0.57$, 2-tailed), a different spatial frequency dependency was evident (Fig. 1e). A Gaussian function was fitted to the average normalized improvement curve⁶ (for detail, see Supplementary online): there was a specific learning effect (full width at half height bandwidth of 1.11 octaves) together with a more general increase in sensitivity at all spatial frequencies tested in Group1. The magnitude of this general increase was half the magnitude of the peak increase; while only a specific learning effect (bandwidth 1.42 octaves) in Group2.

Another important finding was that training also significantly improved visual acuity in Group1 (Paired Samples Test, $t(12) = 9.16$, $P = 0.0000091$, 2-tailed), but not in Group2 (Paired Samples Test, $t(7) = 1.42$, $P = 0.20$, 2-tailed). The average improvement of visual acuity in Group1 was 2.32 dB (or 31%), and this was larger than what was found in Group2 (Independent Samples Test, $t(19) = 3.97$, $P = 0.00082$, 2-tailed) (Fig. 2a). All subjects in Group1 had visual acuity improvements after training (Fig. 2b) that could sustain for at least 5 months (4 subjects in Group1 had visual acuity retested 5 months after training).

It is important to stress that correction of HOAs results in improved optical quality and therefore improved contrast sensitivity and visual acuity, however these purely optical improvements are incorporated in the pre-training baseline measurements from which any training improvements are assessed. To confirm that the CSF improvements we found after training were neural in origin, we also assessed pre- and post-training optical modulation transfer functions (MTF), to quantify the quality of the optics²⁰, for subjects in Group1 and Group2. We found significant improvements in the MTFs as a result of the correction of HOAs but no significant change in the MTFs after training for both groups. On the other hand, the neural transfer function (NTF), calculated from subtracting MTF from the CSF³¹, showed significant improvements after training in both groups (for details, see Supplementary online). These results demonstrate that optical quality did not alter significantly as a result

Figure 1 | Different training effects between Group1 and Group2. (a) Average learning curves of Group1 (red) and Group2 (blue). The first and last points in each condition were derived from pre- and post-training contrast sensitivity function (CSF) measurements, respectively. Error bars, SEM. Subjects in Group1 were trained under the HOAs-corrected condition. Red solid line represents a piecewise linear model fit to the average learning curve of Group1; Subjects in Group2 were trained under the HOAs-uncorrected condition. Blue solid line represents a piecewise linear model fit to the average learning curve of Group2. (b) Average post- and pre-training CSFs of Group1. (c) Average post- and pre-training CSFs of Group2. (d) Average magnitude of contrast sensitivity improvements across observers and spatial frequencies in Group1 (3.11dB) and Group2 (1.31dB); (e) Average normalized improvement curves of Group1 (red) and Group2 (blue). Dashed line represents no improvement. Solid lines represent Gaussian fit to the average normalized improvement curves.

of training; the training-based improvement to the CSF reflected neural changes.

To confirm that the neural benefits were contingent on perceptual learning and not just a passive consequence of having a sharper retinal image, we undertook another training experiment under the HOAs-corrected condition (Group3, 6 adults) using a spatial frequency which, while being optimal in terms of HOAs-corrected sensitivity, was sub-optimal in terms of perceptual training⁹. Group3

Figure 2 | Improvements of visual acuity after training in Group1 (red) and Group2 (blue). Visual acuity associated with 75% correct identification was measured with the Chinese Tumbling E Chart at HOAs-uncorrected condition, and converted to MAR acuity. (a) There were significant improvements of visual acuity in Group1 (Paired Samples Test, $t(12)=9.16$, $P=0.00000091$, 2-tailed), but not in Group2 (Paired Samples Test, $t(7)=1.42$, $P=0.20$, 2-tailed). (b) Visual acuity of all subjects before and after training in Group1. Abscissa represents visual acuity after training; ordinate represents visual acuity before training. Each red ‘◇’ point represents one subject. The dashed line indicates a prediction of no improvement. The 4 yellow ‘▲’ points represent visual acuity retested 5 months after training for 4 subjects (the corresponding pre-/post-training results of these 4 subjects are shown by red ‘◇’ points that are marked by black ‘□’).

was same as Group1, except subjects were trained at a spatial frequency that corresponded to the peak of the CSF, a frequency lower than previously used (Figs 1 & 2) and one known to be non-optimal in terms of perceptual learning⁹. If the previous improvements were simply a passive adaptation³² to the new HOAs-corrected image, (and not specifically a result of perceptual learning) we would expect to find similar results for Group3 as we did previously for Group1. Interestingly, we found much less improvement in Group3 (Fig. 3): the average magnitude of improvements across observers and spatial frequencies was 1.12 dB, which was much less than previously found in Group1 ($F(1,17)=9.86$, $P=0.006$); improvements were not generalized across spatial frequency and did not transfer to letter acuity

Figure 3 | Training effect in Group3. Subjects in Group3 were trained at a low spatial frequency (spatial frequency that has peak contrast sensitivity under the HOAs-corrected condition), when HOAs were corrected. Error bars, SEM. (a) Average post- and pre-training CSFs of Group3; (b) Average magnitude of improvements across observers and spatial frequencies in Group3 was 1.12 dB, less than in Group1 ($F(1,17)=9.86$, $P=0.006$); improvements of visual acuity in Group3 were not significant (Paired Samples Test, $t(5)=0.32$, $P=0.76$, 2-tailed), and less than in Group1 (Independent Samples Test, $t(17)=3.27$, $P=0.004$, 2-tailed).

(Paired Samples Test, $t(5)=0.32$, $P=0.76$, 2-tailed). In these respects they were different to that previously found in Group1 for the high spatial frequency training target, suggesting that a HOAs-corrected environment in itself is necessary but not sufficient to explain the results previously described for Group1 of this study; training at a near-cutoff spatial frequency is also required, which suggests the neural improvement is specific to perceptual training.

The neural improvements in contrast sensitivity after training under the HOAs-corrected condition could in principle be due to increased signal efficiency or reduced internal noise. Either effect would result in improved signal-to-noise and thus, lower thresholds^{33–36}. To separate between these two potential explanations, we assessed the training improvement in contrast thresholds for stimuli with and without added spatial noise. Training improvements were measured at a spatial frequency of 16 c/d and 2 c/d for 4 subjects in Group1 within the HOAs-corrected environment. These results are shown in Fig. 4 where contrast thresholds are plotted against the amplitude of the added white spatial noise, before and after training in the HOAs-corrected environment. The solid lines are best fits to an equivalent noise model of the form:

$$d = \sqrt{\frac{E \times K}{N + N_i}} \quad (1)$$

Where K is related to sampling efficiency, E is the squared contrast energy, N is the squared rms noise contrast, and N_i is related to the internal noise required to account for the measured thresholds. This is referred to as the equivalent noise model^{33,34,37}. The values of the two derived parameters (K and N_i) are shown in Fig. 4. We found that contrast sensitivity improvements under the HOAs-corrected condition for noise-free stimuli were similar to those found with noisy stimuli (Fig. 4), indicating an improved neural efficiency (K) as a consequence of training in the HOAs-corrected condition training³⁸, not a reduction in the internal neural noise (N_i).

Discussion

We demonstrate that the optical quality of the eye limits visual improvements from perceptual learning in adults well beyond the critical period. This in turn means that a greater degree of adult plasticity exists than previously thought. When the HOAs are corrected, we show larger visual improvements that are a consequence of perceptual learning, not the better viewing condition per se. Using the conventional method where training occurred in a HOAs-uncorrected environment (Group2), we found only half of the subjects exhibited improvement in contrast sensitivity, and the average bandwidth of the improvement from perceptual learning was 1.42 octaves. These results are consistent with previous results in the literature. For example, Huang et al.⁹ found only 7 out of 14 normal adults had performance improvement, and the average bandwidth was 1.40 octaves.

Interestingly, we found that identical training within the HOAs-corrected environment (Group1) produced very different results: all subjects exhibited visual performance improvement; training not only produced the expected specific improvement⁹ but also produced a general benefit for all spatial frequencies tested. General benefits after training have been reported previously. Sowden et al.⁶ reported that training subjects at a parafovea site resulted in an improvement with a bandwidth of 1.30 octaves and a general learning effect of 0.05 log units. They claimed that the general learning effect was due to using naive subjects who were not pre-trained. However, in our study, the general learning effect is not amenable to this explanation. First of all, all the subjects in our study received 1.5 h of practice to make sure that they were familiar with the task requirements before the experiment commenced. Second, because Group1 and Group2 were measured using an identical procedure, the explanation for the general improvement displayed by Group1 cannot be in terms of any general procedural learning effect. It must be the consequence of

Figure 4 | Averaged sensitivity (squared contrast) of four subjects in Group1 (trained with HOAs-corrected) as functions of external noise levels (squared rms) for spatial frequencies of 16 c/d and 2 c/d. Contrast thresholds corresponded to 79.3% correct. '○', pre-training; '●', post-training; Error bars, SEM. Solid lines are curve fits to the equivalent noise (see text) and demonstrate that perceptual training improvements are the result of improved neural efficiency rather than diminished neural internal noise.

learning within a HOAs-corrected environment, as this is the only difference in the experimental manipulations between Group1 and Group2.

The general improvement we report at all spatial frequencies was only found in the HOAs-corrected group (Group1). It is not caused by the improved quality of the retinal image per se because this improvement, which is shown in Supplementary Table S3 online, is instantaneous³⁹, easily verified by the improved MTFs and factored out since it is incorporated in our pre-training baseline. Furthermore it cannot be the results of slow term optical changes over the duration of training because it was not reflected in the optical transfer functions we obtained (see Supplementary Fig. S2 online). Therefore the improvement that occurred over the training period must be neural in origin. This neural effect is not a passive consequence of extended viewing under the HOAs-corrected condition³² but rather a consequence of perceptual learning per se under the HOAs-corrected condition. The hallmarks of perceptual learning^{9,10} are its magnitude³⁹, its generalization to lower spatial frequencies and its transfer to letter acuity. These three conditions are met only when perceptual training is undertaken at a high spatial frequency (Group1) not at a peak spatial frequency (Group3). Passive neural adaptation cannot provide an explanation because 1. No similar effects were seen for Group3 who had similar HOAs-corrected experience, 2. For each subject, exposure to the HOAs-corrected condition was lasting only 1 hr a day with at least 15 hrs of normal viewing and 3. The improvements were sustained in that visual acuity improvements under normal viewing condition were still present 5 months later. Therefore, we are left to conclude that the visual improvement (i.e. contrast sensitivity and letter acuity) exhibited by Group1 are a consequence of perceptual training when undertaken under the HOAs-corrected viewing condition.

We also show that the neural improvements reported here are due to improved transduction efficiencies rather than reductions in internal neural noise. Using an equivalent noise model³⁶ to separate these different components of perceptual learning, we show, by evaluating the effects of additive spatial noise, that the benefits are multiplicative rather than additive in nature, consistent with the notion of improved signaling efficiency. This is in agreement with a number of studies that have investigated this distinction for other perceptual learning tasks, such as position discrimination^{40,41}, letter identification⁴², faces and textures³⁸ and contrast detection⁸. The general finding is that training improves efficiency (position discrimination, letter identification, faces & textures) but that there are also, in some cases, reductions in internal noise (contrast detection). Thus the mechanisms underlying the effects we report may not be different in principle to those reported by others; simply our effects are

larger due to the improved optical quality of our participants. Parenthetically, the fact that the improvements are due to efficiency (or sampling efficiency), rules out a purely optical explanation, as this would have been manifested as a reduction in the equivalent noise measure^{36,43}.

It has been argued that the typical perceptual learning that exhibits a selectivity for spatial frequency, contrast and field position is more to do with changes in the higher-level decision stage than it is to do with improved efficiency of neural responses at lower level of visual processing⁴⁴. The improvements we report using adaptive optics and trained at a high spatial frequency involve two components, one is the typical leaning effect that is spatial frequency specific and does not transfer to other functions such as letter acuity^{8,9} whereas the other extends to lower spatial frequencies and represents not only a more generalized improvement at all spatial frequencies but a more generalized improvement in visual function in general, extending to letter acuity (Fig. 2). This latter component is due to perceptual learning because it is not present when training is undertaken at a low spatial frequency even when the HOAs have been corrected (i.e. Group3). It is possible that the sites of these two components are different and that the more generalized benefit occurs at an earlier site of processing.

Visual function rapidly improves after birth reaching asymptotic values for acuity and contrast sensitivity by 9 months of age. While it is true that infants do suffer from refractive errors in early life^{45–47}, there is good evidence^{48,49} that their depth of field, due to their relatively large pupils, is sufficiently large to make them resistant to the HOAs that limit adult vision. We surmise that initial neural development reaches asymptotic levels that are eventually matched to and limited by the optical quality of the adult eye. Improvements in the eye's optical quality during adulthood by adaptive optics can facilitate concurrent gains in the efficiency of neural learning in visual areas of the brain. We reveal two components to this plasticity, one tuned for spatial frequency and limited to high spatial frequencies and another one that is untuned for spatial frequency. This latter component is only revealed when the optical quality is improved and a high spatial frequency stimulus is used to engage perceptual learning. Importantly, this second component allows benefits to extend to the detection of stimuli that are more spatially broadband such as letters. In turn this implies the visual benefit will transfer to the detection of real world objects. Our finding in normals with improved optics may help explain one of the mysteries of perceptual learning in amblyopes where, unlike normals, the visual benefits do transfer from the trained spatial frequency to letters⁵⁰. In amblyopes, the optics do not limit function due to the presence of neural loss and to this extent amblyopes are like normal subjects whose HOAs have

been corrected and for which more generalized spatial frequency improvements are obtainable. Our finding of enhanced visual plasticity in the adult, in both magnitude at the trained spatial frequency and extent of transfer to other spatial frequencies, is directly relevant to the development of new therapies applied in later life to redress brain dysfunction resulting from anomalous visual development earlier in life^{9,10,51,52}.

Methods

Observers. Twenty seven adults (Age: 19–26) with normal or corrected to normal vision (slightly myopic, ≤ -3.0 D) were randomly assigned into three groups. There was no substantial difference between these groups in terms of mean age or mean refractive error. There were 13 observers, 8 observers and 6 observers in Group1, Group2 and Group3 respectively. Group1 was trained at the HOAs-corrected cut-off spatial frequency (spatial frequency when contrast threshold at HOAs-corrected condition is 0.4 which was 29.53 ± 1.68 c/d (s.e.m.)), with HOAs corrected. Group2 was trained at the HOAs-uncorrected cut-off spatial frequency (spatial frequency when contrast threshold at HOAs-uncorrected condition is 0.4 which was 28.75 ± 1.88 c/d (s.e.m.)), with HOAs uncorrected. Group3 was the same as Group1, except subjects were trained at a lower spatial frequency (spatial frequency corresponding to peak contrast sensitivity at HOAs-corrected condition which was 6.00 ± 0.89 c/d (s.e.m.)).

The study was approved by the Institutional Review Board of University of Science and Technology of China. All subjects were naive to the purpose of the experiment and informed consent was obtained from each of them.

Apparatus. All experiments were conducted on a real-time closed-loop adaptive optics visual stimulator system (AOVS)²⁸ in a dark room. It consists of a Hartmann-Shack wavefront sensor (WFS) with 97 lenslets, operating at 25 Hz, and a 37-actuator PZT deformable mirror with a stroke of about 2 microns. The control bandwidth of the system is about 1 Hz. See Supplementary online for diagram (Fig. S1) and more detail. The aberrations are measured for a 4 mm artificial pupil up to 35 Zernike polynomials (7th order) according to OSA wavefront standards¹⁹.

Design. The experiment in each group consisted of four consecutive stages: a pre-training practice stage, a pre-training test stage, a training stage and a post-training test stage. For each subject, only one eye was used in the experiment, the other eye was covered by an opaque fabric. The tested eye having normal or corrected to normal vision was selected randomly for each observer. At the pre- and post-training test stages, visual acuity was measured under conditions where the higher order aberrations (HOAs) were uncorrected. Contrast sensitivity functions (CSFs) were measured under both the HOAs- corrected and uncorrected conditions. Visual acuity corresponding to 75% correct identification was measured with the Chinese Tumbling E Chart, and converted to MAR acuity. Contrast sensitivity, defined as the reciprocal of contrast threshold for detecting a sine-wave grating with 79.3% accuracy, was measured at spatial frequencies 0.6, 1, 2, 4, 8, 16, 24, 36 c/deg on the AOVS. See Supplementary online for more detail.

Procedure. A two-interval forced-choice procedure was used for both training and threshold measurements. In training and CSF measurements, the presentation sequence in each trial was as follows: a 267-ms fixation cross signalled by a brief tone in the beginning, a 117-ms interval, a 500-ms inter-stimulus interval blank, a 267-ms fixation signalled by a brief tone in the beginning, a 117-ms second interval, and blank until response. See Supplementary online for more detail.

- Hubel, D. H. & Wiesel, T. N. The period of susceptibility to the physiological effects of unilateral eye closure in kittens. *J Physiol* **206**, 419–436 (1970).
- Goldstone, R. L. Perceptual learning. *Annu Rev Psychol* **49**, 585–612 (1998).
- Gilbert, C. D., Sigman, M. & Crist, R. E. The neural basis of perceptual learning. *Neuron* **31**, 681–697 (2001).
- Sasaki, Y., Nanez, J. E. & Watanabe, T. Advances in visual perceptual learning and plasticity. *Nat Rev Neurosci* **11**, 53–60 (2010).
- Lu, Z. L., Hua, T., Huang, C. B., Zhou, Y. & Doshier, B. A. Visual perceptual learning. *Neurobiol Learn Mem* (2010).
- Sowden, P. T., Rose, D. & Davies, I. R. L. Perceptual learning of luminance contrast detection: specific for spatial frequency and retinal location but not orientation. *Vision Research* **42**, 1249–1258 (2002).
- Hua, T. *et al.* Perceptual learning improves contrast sensitivity of V1 neurons in cats. *Curr Biol* **20**, 887–894 (2010).
- Huang, C. B., Lu, Z. L. & Zhou, Y. Mechanisms underlying perceptual learning of contrast detection in adults with anisometropic amblyopia. *J Vis* **9**, 24 21–14 (2009).
- Huang, C. B., Zhou, Y. & Lu, Z. L. Broad bandwidth of perceptual learning in the visual system of adults with anisometropic amblyopia. *Proc Natl Acad Sci U S A* **105**, 4068–4073 (2008).
- Zhou, Y. *et al.* Perceptual learning improves contrast sensitivity and visual acuity in adults with anisometropic amblyopia. *Vision Research* **46**, 739–750 (2006).
- Liu, Z. L. Perceptual learning in motion discrimination that generalizes across motion directions. *P Natl Acad Sci USA* **96**, 14085–14087 (1999).
- Ball, K. & Sekuler, R. A specific and enduring improvement in visual motion discrimination. *Science* **218**, 697–698 (1982).
- Stewart, L., Battelli, L., Walsh, V. & Cowey, A. Motion perception and perceptual learning studied by magnetic stimulation. *Electroencephalogr Clin Neurophysiol Suppl* **51**, 334–350 (1999).
- Furmanski, C. S. & Engel, S. A. Perceptual learning in object recognition: object specificity and size invariance. *Vision Res* **40**, 473–484 (2000).
- Golcu, D. & Gilbert, C. D. Perceptual learning of object shape. *J Neurosci* **29**, 13621–13629 (2009).
- Fiorentini, A. & Berardi, N. Perceptual-Learning Specific for Orientation and Spatial-Frequency. *Nature* **287**, 43–44 (1980).
- Ahissar, M. & Hochstein, S. Task difficulty and the specificity of perceptual learning. *Nature* **387**, 401–406 (1997).
- Fahle, M. Perceptual learning: specificity versus generalization. *Curr Opin Neurobiol* **15**, 154–160 (2005).
- Thibos, L. N., Applegate, R. A., Schwiegerling, J. T., Webb, R. & Taskforce, V. S. in *Vision Science and its Applications (VSIA)* (Santa Fe, New Mexico, 2000).
- Williams, D. R. *et al.* (eds Ronald R. Krueger, Raymond A. Applegate & Macrae, S. M.) 19–38 (Slack, 2004).
- Smirnov, M. S. Measurement of the wave aberration of the human eye. *Biofizika* **6**, 776–795 (1961).
- Howland, B. & Howland, H. C. Subjective measurement of high-order aberrations of the eye. *Science* **193**, 580–582 (1976).
- Liang, J. Z., Grimm, B., Goelz, S. & Bille, J. F. Objective Measurement of Wave Aberrations of the Human Eye with the Use of a Hartmann-Shack Wave-Front Sensor. *Journal of the Optical Society of America a-Optics Image Science and Vision* **11**, 1949–1957 (1994).
- Brunette, I., Bueno, J. M., Parent, M., Hamam, H. & Simonet, P. Monochromatic aberrations as a function of age, from childhood to advanced age. *Invest Ophthalmol Vis Sci* **44**, 5438–5446 (2003).
- Porter, J., Guirao, A., Cox, I. G. & Williams, D. R. Monochromatic aberrations of the human eye in a large population. *J Opt Soc Am A Opt Image Sci Vis* **18**, 1793–1803 (2001).
- McLellan, J. S., Marcos, S. & Burns, S. A. Age-related changes in monochromatic wave aberrations of the human eye. *Invest Ophthalmol Vis Sci* **42**, 1390–1395 (2001).
- Kirwan, C., O’Keefe, M. & Soeldner, H. Higher-order aberrations in children. *American Journal of Ophthalmology* **141**, 67–70 (2006).
- Li, S. *et al.* Effects of Monochromatic Aberration on Visual Acuity Using Adaptive Optics. *Optometry and Vision Science* **86**, 1–7 (2009).
- Williams, D. *et al.* Visual benefit of correcting higher order aberrations of the eye. *J Refract Surg* **16**, S554–S559 (2000).
- Yoon, G. Y. & Williams, D. R. Visual performance after correcting the monochromatic and chromatic aberrations of the eye. *J Opt Soc Am A Opt Image Sci Vis* **19**, 266–275 (2002).
- Campbell, F. W. Probing the Human Visual-System - a Citation-Classic Commentary on Optical and Retinal Factors Affecting Visual Resolution by Campbell, F.W., and Green, D.G. *Cc/Life Sci* 16–16 (1990).
- Yehezkel, O., Sagi, D., Sterkin, A., Belkin, M. & Polat, U. Learning to adapt: Dynamics of readaptation to geometrical distortions. *Vision Res* **50**, 1550–1558 (2010).
- Burgess, A. E., Wagner, R. F., Jennings, R. J. & Barlow, H. B. Efficiency of human visual signal discrimination. *Science* **214**, 93–94 (1981).
- Pelli, D. *Effects of visual noise*. (University of Cambridge, 1981).
- Barlow, H. Retinal and central factors in human vision limited by noise. *Vertebrate photoreception*, 337–C358 (1977).
- Kersten, D., Hess, R. & Plant, G. Assessing contrast sensitivity behind cloudy media. *Clinical Vision Science* **2**, 143–C158 (1988).
- Legge, G. & Rubin, G. Contrast sensitivity function as a screening test: a critique. *American journal of optometry and physiological optics* **63**, 265 (1986).
- Gold, J., Bennett, P. J. & Sekuler, A. B. Signal but not noise changes with perceptual learning. *Nature* **402**, 176–178 (1999).
- Rossi, E. A. & Roorda, A. Is visual resolution after adaptive optics correction susceptible to perceptual learning? *J Vis* **10**, 11 (2010).
- Li, R. W. & Levi, D. M. Characterizing the mechanisms of improvement for position discrimination in adult amblyopia. *J Vis* **4**, 476–487 (2004).
- Li, R. W., Klein, S. A. & Levi, D. M. Prolonged perceptual learning of positional acuity in adult amblyopia: perceptual template retuning dynamics. *J Neurosci* **28**, 14223–14229 (2008).
- Chung, S. T., Levi, D. M. & Tjan, B. S. Learning letter identification in peripheral vision. *Vision Res* **45**, 1399–1412 (2005).
- Legge, G. E., Kersten, D. & Burgess, A. E. Contrast discrimination in noise. *J Opt Soc Am A* **4**, 391–404 (1987).
- Yu, C., Klein, S. A. & Levi, D. M. Perceptual learning in contrast discrimination and the (minimal) role of context. *Journal of Vision* **4** (2004).
- Atkinson, J. & French, J. Astigmatism and orientation preference in human infants. *Vision Res* **19**, 1315–1317 (1979).
- Gwiazda, J., Scheiman, M., Mohindra, I. & Held, R. Astigmatism in children: changes in axis and amount from birth to six years. *Invest Ophthalmol Vis Sci* **25**, 88–92 (1984).
- Gwiazda, J., Thorn, F., Bauer, J. & Held, R. Emmetropization and the Progression of Manifest Refraction in Children Followed from Infancy to Puberty. *Clin Vision Sci* **8**, 337–344 (1993).

48. Green, D. G., Powers, M. K. & Banks, M. S. Depth of focus, eye size and visual acuity. *Vision Res* **20**, 827–835 (1980).
49. Powers, M. K. & Dobson, V. Effect of focus on visual acuity of human infants. *Vision Res* **22**, 521–528 (1982).
50. Astle, A. T., Webb, B. S. & McGraw, P. V. The pattern of learned visual improvements in adult amblyopia. *Invest Ophthalmol Vis Sci* **52**, 7195–7204 (2011).
51. Thompson, B., Mansouri, B., Koski, L. & Hess, R. F. Brain plasticity in the adult: modulation of function in amblyopia with rTMS. *Curr Biol* **18**, 1067–1071 (2008).
52. Polat, U., Ma-Naim, T., Belkin, M. & Sagi, D. Improving vision in adult amblyopia by perceptual learning. *Proc Natl Acad Sci U S A* **101**, 6692–6697 (2004).

Acknowledgement

This research was supported by the National Basic Research Program of China (973 Programs, 2011CBA00405 and 2009CB941303), National Science Foundation of China (NSFC 30630027 to YZ and 60808031 to YD), Frontier Research Foundation of Institute of Optics and Electronics: C09K006 (YD), and the CIHR (# MT108-18) to RFH.

Author Contributions

J.-W.Z., Y.-D.Z., R.-F.H. and Y.-F.Z. conceived the experiments. J.-W.Z., Y.D., H.-X.Z., R.L., F.H. and B.L. performed the experiments. J.-W.Z., Y.D., H.-X.Z. and R.-F.H. analyzed the data. J.-W.Z., Y.-D.Z., R.-F.H. and Y.-F.Z. interpreted the data and wrote the manuscript. J.-W.Z. and Y.-D.Z. are co-first authors of this work. All authors reviewed the manuscript.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: There is NO Competing Interest. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

License: This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>

How to cite this article: Zhou, J. *et al.* The eye limits the brain's learning potential. *Sci. Rep.* **2**, 364; DOI:10.1038/srep00364 (2012).