SCIENTIFIC REPORTS



SUBJECT AREAS: NANOBIOTECHNOLOGY NANOPARTICLES CELL SIGNALLING CHEMICAL BIOLOGY

> Received 29 May 2012

Accepted 23 July 2012

Published 7 August 2012

Correspondence and requests for materials should be addressed to C.F. (fchh@sinap.ac. cn) or H.S. (hysong@ sibs.ac.cn)

> * These authors contribute equally to this work.

Akt signaling-associated metabolic effects of dietary gold nanoparticles in *Drosophila*

Bin Wang^{1,3*}, Nan Chen^{2*}, Yingliang Wei^{1*}, Jiang Li², Li Sun¹, Jiarui Wu¹, Qing Huang², Chang Liu³, Chunhai Fan² & Haiyun Song¹

¹Laboratory of Systems Biology, Shanghai Institute of Biochemistry and Cell Biology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai, 200031, China, ²Laboratory of Physical Biology, Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800, China, ³Jiangsu Key Laboratory for Molecular and Medical Biotechnology and College of Life Sciences, Nanjing Normal University, Nanjing, Jiangsu, China.

Gold nanoparticles (AuNPs) are often used as vehicles to deliver drugs or biomolecules, due to their mild effect on cell survival and proliferation. However, little is known about their effect on cellular metabolism. Here we examine the *in vivo* effect of AuNPs on metabolism using *Drosophila* as a model. *Drosophila* and vertebrates possess similar basic metabolic functions, and a highly conserved PI3K/Akt/mTOR signaling pathway plays a central role in the regulation of energy metabolism in both organisms. We show that dietary AuNPs enter the fat body, a key metabolic tissue in *Drosophila* larvae. Significantly, larvae fed with AuNP show increased lipid levels without triggering stress responses. In addition, activities of the PI3K/Akt/mTOR signaling pathway and fatty acids synthesis are increased in these larvae. This study thus reveals a novel function of AuNPs in influencing animal metabolism and suggests its potential therapeutic applications for metabolic disorders.

old nanoparticles (AuNPs) possess many attractive optical and electronic properties that have proven to be of high utility in biomedical applications¹. Biomolecular and cellular interactions of AuNPs are particularly interesting and versatile, which have led to numerous AuNPs-based diagnostic, imaging and therapeutic tools²⁻⁷. More recently, AuNPs were found to be cell permeable^{8,9} in addition to their well known low cytotoxicity and high biocompatibility^{10,11}. Consequently, several groups including ours have shown that AuNPs can serve as nanoscale vehicles for drug delivery^{2,8,12}. Despite the well-documented progress, direct dietary use of AuNPs in living organisms and possible consequences on cellular metabolism are little explored.

Living organisms can sense nutritional conditions to coordinate cellular metabolism. Interestingly, *Drosophila* and vertebrates are very similar in their basic metabolic functions, i.e., sugar, lipid and amino acid homeostasis^{13,14}. In both organisms, a highly conserved phosphatidylinositol 3-kinase (PI3K) pathway has a pivotal role in nutrition uptake and storage^{15–18}. Aberrant PI3K signaling in human results in metabolic disorders and often occurs in type-2 diabetes or cancers^{15,17}. As a response to dietary nutrient, the upstream signaling molecule PI3K is activated, leading to re-localization and activation of Akt¹⁵. The Akt kinase has two major metabolic target molecules, FOXO and mTOR^{19,20}. Phosphorylation by Akt inactivates FOXO²¹, a key transcription factor involved in catabolism, while activates the mTOR complex and its downstream targets. Activation of the mTOR complex in turn stimulates the activity of sterol regulatory element binding protein (SREBP)^{22,23}, a transcription factor that controls expression of key enzymes of lipogenesis including Fatty Acid Synthase (FAS) and Acetyl-CoA Carboxylase (ACC)²⁴.

In this work, we aimed to explore whether dietary AuNPs affect cellular metabolism and used *Drosophila* larvae as an *in vivo* model. We fed larvae with several concentrations of AuNPs and found that they were absorbed by the fat body, a key metabolic tissue in *Drosophila* larvae. In these larvae, stress responses including expression of heat shock genes, activation of the JNK pathway or induction of autophagy were not induced compared to those in control larvae. Despite their little effects on growth or stress responses, lipid levels were increased in larvae fed with AuNPs. By examining metabolic pathways that control lipid anabolism, we found that dietary AuNPs increased activities of the PI3K/Akt/mTOR signaling pathway and fatty acids synthesis in these larvae. In addition, inhibition of mTOR abolished these effects of AuNPs, suggesting that dietary AuNPs could affect *Drosophila* lipid metabolism by promoting the PI3K/Akt/mTOR signaling pathway.



Figure 1 | Uptake of AuNPs by fly larva tissues. (A) Hydrated diameter of AuNPs (23 nm) were measured by dynamic light scattering (DLS). (B) 15 nm AuNPs characterized by transmission electron microscopy (TEM). (C) Uptake of AuNPs by fly larva tissues, as measured by quantitative inductively coupled plasma mass spectrometry (ICP-MS). This bar graph shows representative data of four individual sample analyses.

Results

AuNPs of 15 nm (Fig. 1A,B, supplementary Fig. S1) were synthesized and formulated in the diet. *Drosophila* larvae fed with AuNPs were dissected to determine the distribution of AuNPs among different tissues. Quantitative analysis with inductively coupled plasma mass spectrometry (ICP-MS) detected localization of AuNPs in the fat body (equivalent to mammalian white adipose tissue and liver), but not in brain, salivary gland or imaginal discs (Fig. 1C). Further analysis by transmission electron microscopy (TEM) showed that AuNPs localized to lipid droplets of fat body cells (supplementary Fig. S4C, D).

We first examined whether cellular stress was up-regulated by dietary AuNPs. Cellular stress could increase expression of heat shock proteins²⁵, activate the c-Jun N-terminal Kinase (JNK) pathway²⁶, and trigger autophagy^{27,28}. In larvae fed with 0.5 or 2 nM AuNPs, expression of *hsp83* or *hsp70* was unchanged (Fig. 2A, B, supplementary Fig. S2A, B); activity of the JNK pathway was unchanged or slightly reduced, as measured by expression of its downstream gene *puckered* (Fig. 2C, supplementary Fig. S2C)²⁹; autophagy was also unchanged or slightly reduced, as measured by lysotracker staining (Fig. 2D, supplementary Fig. S2D). However, 5 nM dietary AuNPs did slightly increase expression of *hsp70* and *puckered*, but not that of *hsp83* (supplementary Fig. S2A-C). As a

conclusion, up to 2 nM dietary AuNPs did not trigger cellular stress responses in larvae, although higher concentration of AuNPs might have a mild effect. For this reason we used 2 nM AuNPs for the rest of experiments.

The localization of AuNPs to lipid droplets of fat body cells triggered our interest to examine its possible consequences on lipid metabolism. *Drosophila* and mammals regulate lipid metabolism via the highly conserved SREBP pathway³⁰, which activates expression of key enzymes of fatty acid synthesis including FAS and ACC²⁴. In larvae fed with AuNPs, triacylglycerol (TAG, the main stored form of lipid) levels were slightly higher (~10%, Fig. 3A). In addition, expression of *FAS* and *ACC* were increased, suggesting that enhanced fatty acid synthesis was responsible for the change of lipid levels (Fig. 3B,C). The fly food provided excessive nutrition, and lipid anabolism in larvae already reached a high level. Thus we also fed larvae in the condition of calorie restriction (CR), where lipid anabolism was limited. Under this background, larvae fed with AuNPs showed even higher TAG levels (~20%) compared to the CR control (Fig. 4A).

The SREBP pathway is a downstream effector of the PI3K/Akt/ mTOR pathway^{22,23}. To test whether this signaling pathway was activated by dietary AuNPs, we examine the activity of PI3K, Akt, and expression of their downstream genes. In standard diet, the activity of



Figure 2 | **Dietary AuNPs do not trigger stress responses in fly larvae.** In larvae fed with control food or 2 nM AuNPs, stress responses were measured by quantitative RT-PCR (A–C) or lysotracker staining (D). (A), (B) Dietary AuNPs did not affect expression of heat shock protein genes *hsp83* (A) or *hsp70* (B). ns, not significant. (C) Dietary AuNPs slightly reduced activation of JNK pathway downstream gene *puckered.* *, P<0.05. (D) Dietary AuNP slightly reduced autophagy in fat body.





Figure 3 | Dietary AuNPs increases lipid levels and fatty acids synthesis in larvae. (A) Normalized TAG levels in larvae fed with AuNPs were 10% higher than that in control. *, P < 0.05. (B), (C) Expression of key enzymes for fatty acids synthesis FAS (B) and ACC (C) were increased by dietary AuNPs. **, P < 0.01.

Akt in larvae fed with AuNPs was comparable to that in control, consistent with the result that AuNPs only had a mild effect on lipid metabolism under this condition. In the presence of CR, dietary AuNPs significantly enhanced Akt activity (Fig. 4B). We also measured membrane bound tGPH in the fat body, whose levels reflected the activity of PI3K16. Larvae fed with AuNPs showed higher activity of PI3K in the fat body than the control (Fig. 4C,D). These results confirmed that the upstream signaling molecules (PI3K and Akt) were activated by dietary AuNPs. Activated Akt can enhance expression of FAS and ACC via the mTOR/SREBP cascade²², and repress expression of InR and 4EBP via FOXO (supplementary Fig. S3A)²¹. In both standard diet and CR diet, larvae fed with AuNPs showed increased expression of FAS and ACC, and reduced expression of InR and 4EBP (supplementary Fig. S3B-E), confirming that dietary AuNPs activated the Akt pathway and affected Akt target gene expression.

Dietary AuNPs may affect other metabolic pathways besides Akt signaling. To test whether the effect of AuNPs on *Drosophila* fat metabolism was dependent on the PI3K/Akt/mTOR pathway, we

blocked Akt signal transduction by rapamycin, an inhibitor of mTOR³¹, and examined the effect of AuNPs on fat levels and fatty acid synthesis. In the presence of rapamycin, larvae fed with AuNPs had similar fat levels compared to the control, either in standard diet or CR diet (Fig. 5A,B). In addition, dietary AuNPs no longer increased expression of *FAS* or *ACC* (Fig. 5C,D), suggesting that rapamycin blocked the stimulation of fatty acid synthesis by AuNPs. Therefore, dietary AuNPs possibly affect *Drosophila* fat metabolism via the PI3K/Akt/mTOR pathway (Fig 5E).

Discussion

There are several ways that AuNPs can affect PI3K/Akt signaling. PI3K/Akt signaling generally senses nutritional conditions to coordinate metabolism. Although larvae fed with or without AuNPs consume the same amount of food (supplementary Fig. S4A), it is possible that AuNPs may promote nutrient uptake at cellular levels. To test this possibility, we examined total sugar levels as sugar catabolism provided acetyl-CoA precursors (pyruvate and citrate) for fatty acid synthesis. We found that larvae fed with AuNPs had higher





Figure 4 | Calorie restriction enhances effects of AuNPs on fat metabolism and Akt signaling. (A) Dietary AuNPs increased larva TAG levels by 10% in standard food, and by 20% under the condition of calorie restriction (CR). *, P < 0.05. (B) Under CR, larvae fed with AuNPs showed increased Akt activity compared to the control (lane 4 vs. lane 3). (C) (D) Under CR, larvae fed with AuNPs (D) had higher PI3K activity in the fat body compared to the control (C).

sugar levels than the control in CR food, while this difference was insignificant in standard food (supplementary Fig. S4B). These data suggest that AuNPs enhance nutrient uptake in CR food and may at least partially explain our observations. Alternatively, AuNPs may affect activities of signaling molecules of the PI3K/Akt pathway. Activations of both PI3K and Akt occur on the cell membrane. It is also likely that endocytosis of AuNPs facilitates recruitment of PI3K and Akt on the cell membrane and promotes their activations. Consistent with this model, higher levels of tGPH are detected at the cell membrane (Fig. 4C,D), reflecting that higher levels of PI3K are localized to the cell membrane. Therefore this mechanism may also contribute to the effect of AuNPs.

AuNPs have shown great promise in biomedicine, e.g. they may serve as vehicles to deliver chemical drugs or biomolecules. While it is generally considered that AuNPs are biocompatible and non-toxic for cell growth, other biochemical activities of AuNPs are little





standard diet

*

A 0.5

0.4

Figure 5 | The effect of AuNPs on larva fat metabolism is dependent on Akt/mTOR signaling. (A) (B) In standard diet (A) and CR (B), the increase in larva TAG levels by dietary AuNPs was abolished when the Akt/mTOR signaling pathway was blocked by rapamycin. *, P<0.05. ns, not significant. (C) (D) Increased expression of FAS (C) and ACC (D) by dietary AuNPs was abolished when the Akt/mTOR signaling pathway was blocked. **, P<0.01. ns, not significant. (E) A diagram for the effect of dietary AuNPs on larva fat metabolism.

studied. In this work, we focus on the metabolic effects of dietary AuNPs in *Drosophila* larvae. We find that dietary AuNPs promote the Akt signaling pathway and alter lipid metabolism, an unexpected *in vivo* effect of nanomaterials on cellular metabolism. Given that metabolic diseases such as diabetes have become worldwide epidemic, our new observations shed new light on using AuNPs, either by themselves or coupled with other functional drugs, as nanomedicine against metabolic disorders.

Methods

Preparation and characterization of AuNPs. AuNPs of 15 nm were synthesized by reduction of HAuCl₄ as previously reported³², and their sizes were characterized with transmission electron microscopy and dynamic light scattering.

Fly strain and culture. *yw D. melanogaster* was used in this study. Flies were raised in cages. 0–1 hour embryos were collected from fruit juice agar plates, counted and distributed on SY food (10% sucrose, 10% yeast, 1.5% agar)³³ with or without AuNPs. AuNPs were coated with 2 ug/ul BSA for 10 minutes and mixed with the food before solidification. Same amount of BSA is also included in the control food. Larvae were harvested 108 hours later for analysis. For CR, the fly food contained 25% nutrients of standard food, and larvae were harvested 156 hours after egg laying. To prepare larval tissues for detection of AuNPs distribution, larvae fed with AuNPs were transferred to control food for 8 hours, allowing evacuation of AuNP food. Larvae were then transferred to 6 cm dishes and washed 10 times in 1×PBST before dissection. Dissected tissues were transferred to microcentrifuge tubes and washed 10 times in 1×PBST.

Measurement of mRNA, protein and lipid levels. Total RNAs were extracted from larvae with TRIzol reagent (Invitrogen). The first-strand cDNAs were generated by iScriptTMcDNA Synthesis Kit (Bio-Rad). Quantitative RT-PCR assays were performed with SYBR Green PCR Master Mix and the Applied Biosystems ABI Prism 7000 Sequence Detection System (Applied Biosystems; Foster City, CA). The average expression levels of *actin*, *tubulin* and *TBP* were used as internal controls for normalization of mRNA levels. For western blotting, anti-phospho-Akt and anti-Tubulin antibodies (Cell Signaling) were used. For measurement of TAG, larvae were suspended in 400 ul chilled lysis buffer (0.01 mM KH₂PO₄, 1 mM EDTA, protease inhibitors cocktail, 0.05% Tween 20), lysed by Tissuelyser II (Qiagen, set at 30 Hz for 3 min), and heated at 70°C for 5 min. Cleared supernatant was used to measure TAG with TAG determination kits (Sigma,TR0100). Lipid levels were normalized to total protein levels.

- Daniel, M. C. & Astruc, D. Gold nanoparticles: assembly, supramolecular chemistry, quantum-size-related properties, and applications toward biology, catalysis, and nanotechnology. *Chem. Rev.* 104, 293–346 (2004).
- Ghosh, P., Han, G., De, M., Kim, C. K. & Rotello, V. M. Gold nanoparticles in delivery applications. *Adv. Drug Deliv. Rev.* 60, 1307–1315 (2008).
- Zhang, J., Song, S., Wang, L., Pan, D. & Fan, C. A gold nanoparticle-based chronocoulometric DNA sensor for amplified detection of DNA. *Nat. Protoc.* 2, 2888–2895 (2007).
- Liu, J. & Lu, Y. Fast colorimetric sensing of adenosine and cocaine based on a general sensor design involving aptamers and nanoparticles. *Angew. Chem. Int. Ed.* 45, 90–94 (2005).
- Xiao, Y., Patolsky, F., Katz, E., Hainfeld, J. F. & Willner, I. "Plugging into Enzymes": nanowiring of redox enzymes by a gold nanoparticle. *Science* 299, 1877–1881 (2003).
- Zhang, J., Liu, Y., Ke, Y. & Yan, H. Periodic square-like gold nanoparticle arrays templated by self-assembled 2D DNA Nanogrids on a surface. *Nano Lett.* 6, 248–251 (2006).
- Chen, P. et al. Gold nanoparticles for high-throughput genotyping of long-range haplotypes. Nature Nanotechnol. 6, 639–644 (2011).
- Dhar, S., Daniel, W. L., Giljohann, D. A., Mirkin, C. A. & Lippard, S. J. Polyvalent Oligonucleotide Gold Nanoparticle Conjugates as Delivery Vehicles for Platinum(IV) Warheads. J. Am. Chem. Soc. (2010).
- Rosi, N. L., Giljohann, D. A., Thaxton, C. S., Lytton-Jean, A. K., Han, M. S. & Mirkin, C. A. Oligonucleotide-modified gold nanoparticles for intracellular gene regulation. *Science* 312, 1027–1030 (2006).
- Rosi, N. L. & Mirkin, C. A. Nanostructures in biodiagnostics. *Chem. Rev.* 105, 1547–1562 (2005).
- Connor, E. E., Mwamuka, J., Gole, A., Murphy, C. J. & Wyatt, M. D. Gold nanoparticles are taken up by human cells but do not cause acute cytotoxicity. *Small* 1, 325–327 (2005).
- Wei, M. C., H., Li, J., Yin, M., Liang, L., He, Y., Song, H., Fan, C. & Huang, Q. Polyvalent immunostimulatory nanoagents with self-assembled CpG oligonucleotide-conjugated gold nanoparticles. *Angew. Chem. Int. Ed.* 51, 1202–1206 (2012).

- Baker, K. D. & Thummel, C. S. Diabetic larvae and obese flies-emerging studies of metabolism in Drosophila. *Cell Metabolism* 6, 257–266 (2007).
- Leopold, P. & Perrimon, N. Drosophila and the genetics of the internal milieu. Nature 450, 186–188 (2007).
- 15. Cantley, L. C. The phosphoinositide 3-kinase pathway. *Science* **296**, 1655–1657 (2002).
- Britton, J. S., Lockwood, W. K., Li, L., Cohen, S. M. & Edgar, B. A. Drosophila's insulin/PI3-kinase pathway coordinates cellular metabolism with nutritional conditions. *Dev. Cell* 2, 239–249 (2002).
- 17. Saltiel, A. R. & Kahn, C. R. Insulin signalling and the regulation of glucose and lipid metabolism. *Nature* **414**, 799–806 (2001).
- Engelman, J. A., Luo, J. & Cantley, L. C. The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat. Rev. Genet.* 7, 606–619 (2006).
- Huang, H. & Tindall, D. J. Dynamic FoxO transcription factors. J. Cell. Sci. 120, 2479–2487 (2007).
- 20. Hay, N. The Akt-mTOR tango and its relevance to cancer. *Cancer Cell* **8**, 179–183 (2005).
- Puig, O., Marr, M. T., Ruhf, M. L. & Tjian, R. Control of cell number by Drosophila FOXO: downstream and feedback regulation of the insulin receptor pathway. *Genes Dev.* 17, 2006–2020 (2003).
- 22. Porstmann, T., Santos, C. R., Griffiths, B., Cully, M., Wu, M., Leevers, S., Griffiths, J. R., Chung, Y. L. & Schulze, A. SREBP activity is regulated by mTORC1 and contributes to Akt-dependent cell growth. *Cell Metab.* 8, 224–236 (2008).
- Peterson, T. R. *et al.* mTOR complex 1 regulates lipin 1 localization to control the SREBP pathway. *Cell* 146, 408–420 (2011).
- Magana, M. M., Lin, S. S., Dooley, K. A. & Osborne, T. F. Sterol regulation of acetyl coenzyme A carboxylase promoter requires two interdependent binding sites for sterol regulatory element binding proteins. *J. Lipid Res.* 38, 1630–1638 (1997).
- Singh, M. P., Reddy, M. M., Mathur, N., Saxena, D. K. & Chowdhuri, D. K. Induction of hsp70, hsp60, hsp83 and hsp26 and oxidative stress markers in benzene, toluene and xylene exposed Drosophila melanogaster: role of ROS generation. *Toxicol. Appl. Pharmacol.* 235, 226–243 (2009).
- Stronach, B. E. & Perrimon, N. Stress signaling in Drosophila. Oncogene 18, 6172–6182 (1999).
- Simonsen, A., Cumming, R. C. & Finley, K. D. Linking lysosomal trafficking defects with changes in aging and stress response in Drosophila. *Autophagy* 3, 499–501 (2007).
- Scott, R. C., Schuldiner, O. & Neufeld, T. P. Role and regulation of starvationinduced autophagy in the Drosophila fat body. *Dev. Cell* 7, 167–178 (2004).
- McEwen, D. G. & Peifer, M. Puckered, a Drosophila MAPK phosphatase, ensures cell viability by antagonizing JNK-induced apoptosis. *Development* 132, 3935–3946 (2005).
- 30. Rawson, R. B. The SREBP pathway--insights from Insigs and insects. *Nat. Rev. Mol. Cell. Biol.* 4, 631-640 (2003).
- Brown, E. J., Albers, M. W., Shin, T. B., Ichikawa, K., Keith, C. T., Lane, W. S. & Schreiber, S. L. A mammalian protein targeted by G1-arresting rapamycinreceptor complex. *Nature* 369, 756–758 (1994).
- 32. Hayat, M. A. Colloidal gold : principles, methods, and applications (Academic Press, 1989).
- Magwere, T., Chapman, T. & Partridge, L. Sex differences in the effect of dietary restriction on life span and mortality rates in female and male Drosophila melanogaster. J. Gerontol. A-Biol. Sci. Med. Sci. 59, 3–9 (2004).

Acknowledgements

This work was supported in part by grants from the Chinese Ministry of Science and Technology (MOST, 2011CB943900 and 2012CB932600), the National Natural Science Foundation of China (NSFC, 31071145, 90913014). H.S. is a scholar of the "Hundred Talent Program."

Author contributions

CF and HS designed the project, BW, NC and YW performed experiments, JL, LS, JW, QH and CL provided new materials, BW, NC, CF and HS wrote the main manuscript text and BW prepared figures 1–5. All authors reviewed the manuscript.

Additional information

Supplementary information accompanies this paper at http://www.nature.com/ scientificreports

Competing financial interests: The authors declare no competing financial interests.

License: This work is licensed under a Creative Commons

Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit http://creativecommons.org/licenses/by-nc-sa/3.0/

How to cite this article: Wang, B. et al. Akt signaling-associated metabolic effects of dietary gold nanoparticles in *Drosophila*. Sci. Rep. 2, 563; DOI:10.1038/srep00563 (2012).