



OPEN

SUBJECT AREAS:

CELL LINEAGE

DIFFERENTIATION

EMBRYONIC INDUCTION

BODY PATTERNING

Single blastomere expression profiling of *Xenopus laevis* embryos of 8 to 32-cells reveals developmental asymmetry

Monika Flachsova^{1,2}, Radek Sindelka¹ & Mikael Kubista^{1,3}

¹Laboratory of Gene Expression, Institute of Biotechnology, Academy of Sciences of the Czech Republic, Prague, Czech Republic, ²Charles University in Prague, Faculty of Science, Department of Cell Biology, Prague, Czech Republic, ³TATAA Biocenter AB, Göteborg, Sweden.

Received
13 May 2013Accepted
5 July 2013Published
24 July 2013

Correspondence and requests for materials should be addressed to M.K. (mikael.kubista@img.cas.cz)

We have measured the expression of 41 maternal mRNAs in individual blastomeres collected from the 8 to 32-cell *Xenopus laevis* embryos to determine when and how asymmetry in the body plan is introduced. We demonstrate that the asymmetry along the animal-vegetal axis in the oocyte is transferred to the daughter cells during early cell divisions. All studied mRNAs are distributed evenly among the set of animal as well as vegetal blastomeres. We find no asymmetry in mRNA levels that might be ascribed to the dorso-ventral specification or the left-right axis formation. We hypothesize that while the animal-vegetal asymmetry is a consequence of mRNA gradients, the dorso-ventral and left-right axes specifications are induced by asymmetric distribution of other biomolecules, probably proteins.

Understanding the formation of the embryonic body plan is essential in developmental biology. It is generally accepted that mechanisms of cellular differentiation driving development are based on differential transcription and translation of key regulatory elements. Recent development of highly sensitive and reliable molecular diagnostic methods allows for precise measurements of genes' transcriptional activities in very small samples including single cells¹. In an excellent study by Guo *et al.*, mouse embryos from 8 to 64 cells were dissociated into individual blastomeres and gene expression profiles were measured using the high-throughput microfluidic quantitative real-time PCR (qPCR) system BioMark². They found early mouse embryos are composed of three types of cells representing the trophoectoderm, epiblast, and primitive endoderm. The three cell types were readily distinguished by the expression profile of 48 genes. Pluripotency of mouse single blastomeres from preimplantation embryos (1 to 16-cell stages) was also demonstrated with multiplex RT-PCR expression analysis by May *et al.*³. Expression profiles of trophoectoderm markers, inner cell mass markers and stemness markers have also been measured on human single blastomeres from 5 to 8-cell embryos by Galan *et al.*, using microarray⁴ and in single putative stem cells from human adult ovarian surface epithelium⁵.

The *Xenopus* oocyte has two differentially colored hemispheres known as animal and vegetal. The separation into the hemispheres creates the first developmental axis of the embryo referred to as animal-vegetal. The darker color of the animal hemisphere is due to accumulation of pigmented granules called melanosomes. The vegetal hemisphere stores yolk and has light color⁶. *De novo* transcription in the developing *Xenopus* embryo is silenced until the mid-blastula transition stage (MBT), which takes place after twelve cell divisions post fertilization. All mRNA molecules required for the development into MBT must be present in the oocyte. These were transcribed during oogenesis from solely maternal chromosomes. The maternal mRNAs are asymmetrically distributed along the animal-vegetal axis of the oocyte and direct the specialization of the animal and vegetal parts⁷⁻⁹. The second embryonic body axis is formed after fertilization and separates the embryo into a dorsal and a ventral part. In *Drosophila* embryos the mechanism determining the dorsal-ventral axis is triggered by a concentration gradient of *gurken* maternal mRNA, which is accumulated at the future dorsal site. Absence of *gurken* mRNA at the ventral site leads to translocation of transcription factor Dorsal from the cytoplasm into the nuclei, where the Dorsal activates genes that specify the ventral part¹⁰. The mechanistic details of the dorso-ventral axis formation in zebrafish are not known, but differences between the dorsal and ventral parts appear during the gastrula stage, when the ventral side of the embryo gets thinner than the dorsal side¹¹. The formation of the dorso-ventral axis in *Xenopus laevis* embryo can be discerned already at the 4-cell stage¹². The sperm enters the oocyte through the animal hemisphere and in about 25 minutes the cortical cytoplasm of the oocyte rotates some 30 degrees relative to the inner cytoplasmic mass in opposite direction to the sperm's entrance^{13,14}. Dorsal maternal factors, such as



the Dishevelled protein (Dvl), move to the future dorsal side in the vegetal hemisphere of the oocyte¹⁵. Dvl inactivates gsk3 β protein and protects β -catenin from degradation. The accumulation of β -catenin in the dorsal site of the embryo promotes local Wnt signaling, which regulates transcription of zygotic *xnr3* and *siamois*, leading to dorsalization of the embryo^{16,17}. The first cell division cuts the oocyte through the point of sperm entrance in the animal pole and the groove elongates to the vegetal pole. The second cell division is perpendicular to the first and separates the embryo into the future dorsal and ventral halves. All four blastomeres contain cytoplasm from both the animal and vegetal hemispheres, but vary in size and pigmentation¹². The third and last important body axis of *Xenopus laevis* embryos creates the right-left asymmetry, which manifests at the gastrula stage when the production of embryonic mRNAs has been initiated. Maternal *vg1* protein is more active in the left side of the embryo and activates the expression of *xnr1* (*Xenopus* nodal-related 1). *Xnr1* induces *pitx2* production, which is required for the left side formation. Injection of *vg1* protein to the future right side of the embryo causes left-right deformation manifested by random localization of the heart and of the digestive tube¹⁸.

We have previously used quantitative real-time PCR (qPCR) expression profiling of cryostat sections (qPCR tomography) of a single oocyte to identify transcripts that form gradients along the animal-vegetal axis^{8,9}. These transcripts will become differentially distributed among the animal and the vegetal blastomeres formed at the third cell division. The purpose of the present study is to elucidate if there are expression gradients throughout the embryo that also reflect the dorsal-ventral axis specification and the left-right axis formation by measuring expression of selected 41 maternal genes in individual blastomeres collected from *Xenopus* embryos between 8 to 32-cell stages.

Results

First we extensively optimized our experimental procedure for high extraction and reverse transcription yields, and high qPCR efficiencies. We standardized the protocol for minimum technical variation for the expression profiling of single blastomeres. The vegetal blastomeres store yolk, which is strong inhibitor of biochemical reactions and can induce high variation in the reverse transcription yield and compromise qPCR efficiencies. An RNA spike (*in vitro* transcribed

artificial RNA with 3' polyA tail and 5' cap, TATAA Biocenter) was used to validate the reverse transcription reaction and the qPCR. The spike was added to the RNA extracted from the samples together with the random sequence hexamers and oligo-dT primers for reverse transcription. The amount of reverse transcribed spike was then measured along with the endogenous transcripts. For the optimized protocol the standard deviation of the RNA spike Cq values across all samples was below 0.20 cycles, which evidences excellent reproducibility between sample preparations. Embryos from the same mother also showed high concordance. Our protocol for expression profiling is based on collecting the individual blastomeres in random order and we can only keep track from which hemisphere (animal or vegetal) the different blastomeres originated. This was necessary to avoid introducing bias during collection. With this protocol any dorso-ventral (or possibly left-right) asymmetry would be reflected by heterogeneity in expression profiles among the blastomeres collected from the same hemisphere. As markers to probe the formation of the three developmental axes we chose ten maternal transcripts that have been implicated in the dorso-ventral patterning: *dvl2*, *dvl3*, *lrp6*, *wnt11*, *tcf3*, *gsk3b*, *ctnnb1* (= β -catenin), *foxh1*, *trim36* and *axin1*^{16,19–21}, one important marker for left-right specification: *vg1*¹⁸, and nine genes implicated in animal-vegetal orientation: *dazl*, *cdx1* (= *xcad2*), *wnt11*, *vg1*, *vegt*, *trim36*, *ddx25* (= *deadsouth*), *otx1*, and *maml1*^{9,19,22,23}. We also included 25 mRNAs that have previously been observed in the mature oocyte: *fzd7*, *bmp2*, *pias1*, *foxr1*, *frat1*, *mapk8*, *odc1*, 18S rRNA, 5S rRNA, *cycl1*, *acta*, *tubb*, *gapdh*, *eef1a1*, RNA polymerase II, U3 snoRNA, *par1*, *oct60*, *est1*, *apc*, *zpc*, *mos*, *stat3*, and *pax6*^{8,9}.

Expression data were collected for a total of 224 single blastomeres. The data were analyzed with principal component analysis (PCA) and hierarchical clustering, which are the most powerful multivariate methods to classify samples based on the collective expression of multiple genes²⁴. PCA clearly separated cells originating from the animal and from the vegetal hemispheres into clusters (Fig. 1), and indicated that the vegetal blastomeres are more heterogeneous than the animal blastomeres. Neither of the clusters, however, divided into subgroups that would indicate additional asymmetry that could be ascribed to the formation of the other body axes (e.g. dorsal-ventral and left-right asymmetry). The PCA results were confirmed by classifications using SOM (not shown) and hierarchical clustering. The hierarchical clustering is presented in a

Principal component analysis




Figure 1 | PCA of profiled blastomeres collected from early developmental stages of *Xenopus laevis*. Triangles indicate blastomeres from the 8-cell stage; squares from the 16-cell stage, and circles from the 32-cell stage. Blastomeres originating from the animal hemisphere are shown in red and those from the vegetal hemisphere are shown in blue. Left and right graphs are data from two different females (left: four embryos from 8-cell stage, four embryos from 16-cell stage, and one 32-cell stage embryo, in total 128 blastomeres; right: two embryos from 8-cell stage, three from 16-cell stage, and one from 32-cell stage, in total 96 blastomeres).



heatmap (Fig. 2), which is a graphical illustration of all the genes' expressions in all the blastomeres with the level indicated by color, and dendrograms clustering the blastomeres and the genes, respectively, shown on the top and left side. Separate heatmaps were calculated for each embryo from 8-cell (six embryos), 16-cell (seven embryos), and 32-cell (two embryos) stages. Blastomeres of 8 and 16-cell stages separated into well-defined animal and vegetal clusters, but no subclusters that could be ascribed to the formation of dorsal-ventral or left-right body axes can be discerned. Blastomeres from the 32-cell stage also divided into an animal and a vegetal cluster. The 32-cell stage vegetal cluster was diffuse, reflecting heterogeneity among the vegetal blastomeres possibly caused by the blastomeres being arranged into two ventral layers and some transcripts, such as germ plasm determinants, being concentrated in the extreme ventral cortical layer. The genes cluster into four well separated groups at all the developmental stages studied. The first group contains ubiquitously expressed genes represented by *18S rRNA*, *5S rRNA*, and *cyc1*. These genes are expressed at high level in all the blastomeres at all stages. A second group is composed of *vg1*, *cdx1*, *vegt*, *dazl*, *wnt11*, *otx1*, *trim36*, and *ddx25*, which are most abundant in the vegetal blastomeres. Several of these transcripts have previously been found in the vegetal part of the mature oocyte⁹. Third and fourth groups of transcripts are more abundant in the animal blastomeres, but subtle difference in expression patterns separate them into two clusters. *Fzd7*, *bmp2*, *pias1*, *dvl2*, *dvl3*, *lrp6*, *frat1*, *mapk8*, *axin1*, *est1*, and *U3 snoRNA* are expressed at slightly lower level (indicated by the green shade in the heatmap in Fig. 2). Than the fourth set of genes; *ctnnb1*, *foxh1*, *odc1*, *zpc*, *mos*, *maml1*, *eef1a1*, *RNA polymerase II*, *foxr1*, *tcf3*, *gsk3b*, *oct60*, *stat3*, *par1*, *acta*, *tubb*, *gapdh*, *apc*, and *pax6* (indicated by the green shade turning to light red in the heatmap in Fig. 2). The separation of genes into these two groups shows some variability across embryos (data not shown). The separation of the gene transcripts into clusters is even more evident in the principal component analysis (PCA) (Fig. 3). One cluster characterizing genes expressed in animal blastomeres: *fzd7*, *bmp2*, *pias1*, *dvl2*, *dvl3*, *lrp6*, *frat1*, *mapk8*, *axin1*, *est1*, *U3 snoRNA*, *ctnnb1*, *foxh1*, *odc1*, *zpc*, *mos*, *maml1*, *eef1a1*, *RNA polymerase II*, *foxr1*, *tcf3*, *gsk3b*, *oct60*, *stat3*, *par1*, *acta*, *tubb*, *gapdh*, *apc*, and *pax6*; a second cluster of genes expressed in vegetal blastomeres: *vg1*, *cdx1*, *vegt*, *dazl*, *wnt11*, *otx1*, *trim36*, and *ddx25*; a third cluster of high expressed genes: *18S rRNA*, *5S rRNA*, and *cyc1*.

Discussion

Maternal mRNAs are synthesized during oogenesis and translated during the early development of *Xenopus laevis*. It is generally accepted that the temporal regulation and the spatial distribution of these maternal mRNAs are important for the establishment of the body axes and subsequently the whole body pattern. The animal-vegetal asymmetry is the first body axis formed during oogenesis. The germ layers, endoderm, mesoderm, and ectoderm, are formed along the animal-vegetal axis such that the vegetal part of the oocyte becomes the endoderm and the most distal animal part of the oocyte gives rise to the ectoderm^{25,26}. Transcripts of genes coding specific endodermal and mesodermal factors, such as *vg1*, *wnt11*, and *vegt*, are localized in the vegetal hemisphere^{8,9,22}. Genes responsible for primordial germ cell formation, such as *dazl* and *cdx1*, are localized towards the extreme vegetal pole⁹. Previously we described the new method qPCR tomography to quantify maternal mRNA distribution along the animal-vegetal axis formed during oogenesis, and found two dominant profiles: one with animal and one with vegetal mRNA localization⁸. Subsequent high-resolution qPCR tomography allowed us to discern two vegetal profiles; one encompassing germ plasm determinants (*dazl*, *cdx1*, *ddx25*) and one encompassing the other vegetal genes investigated (*vg1*, *vegt*, *wnt11*, *otx1*, *eg6*)⁹. In this study we show that the animal-vegetal polarizations observed in the oocyte remain in the 8, 16, and 32-cell stage embryos. The vegetal

blastomeres had higher levels of *dazl*, *cdx1*, *ddx25*, *vg1*, *vegt*, *wnt11*, *otx1*, as well as *trim36*, which was not included in the oocyte study. At the 8 and 16-cell stages the vegetal mRNAs were distributed evenly among the vegetal blastomeres; the determinants of the germ plasm being present in all the vegetal cells. At the 32-cell stage, when the vegetal blastomeres form two layers, heterogeneity among the vegetal blastomeres was introduced. *18S rRNA*, *5S rRNA*, *cyc1*, *acta*, *tubb*, *gapdh*, *eef1a1*, *RNA polymerase II*, *U3 snoRNA*, *oct60*, *axin1*, *est1*, *apc*, *tcf3*, *zpc*, *gsk3b*, *maml1*, *ctnnb1*, *mos*, *foxh1*, *stat3*, *fzd7*, *par7*, *bmp2*, *pias1*, *dvl2*, *dvl3*, *lrp6*, *foxr1*, *frat1*, *mapk8*, *odc1* and *pax6* transcripts were more abundant in the animal blastomeres, and were evenly distributed among the individual cells. Multivariate statistical analysis revealed higher variability among the vegetal blastomeres than among the animal. The difference was particularly pronounced at the 32-cell stage. The heterogeneity seems random, possibly caused by yolk platelets that may interfere with PCR inducing technical noise or introduced by the formation of two ventral layers; but we find no evidence for systematic distribution of transcripts neither along the dorsal-ventral axis nor the left-right axis.

An essential question in developmental biology is how amphibians establish the dorsal-ventral axis. The site of sperm entry determines the future ventral half of the embryo. The dorsal part forms on the opposite side and is determined by the accumulation of maternal dorsal determinants during the cortical rotation^{27–29}. Consequently, blastomeres of the early *Xenopus* embryos can be separated into a dorsal and a ventral group with equal number of animal and vegetal cells in each. In *Drosophila* the dorsalizing determinants have been shown to be maternal mRNA molecules that are asymmetrically distributed within the oocyte and regulate translation that leads to the polarization of the embryo^{30,31}. In our study we tested if the dorsalizing factors in *Xenopus* embryogenesis are also mRNAs coding for dorsal specific proteins within the Wnt pathway. We found that among the maternal transcripts that have been implicated in the induction of dorso-ventral asymmetry *ctnnb1*, *dvl2*, *dvl3*, *lrp6*, *gsk3b*, *tcf3*, *foxh1*, and *axin1*, are present predominantly in the animal blastomeres, while *trim36* and *xwnt11* are more abundant in the vegetal blastomeres. This supports the idea that the majority of the dorsal-ventral inducing molecules is localized in the dorsal animal blastomeres of the 16-cell stage embryo³², although we found no evidence of asymmetric distribution of maternal mRNAs among the dorsal or ventral blastomeres. In contrast to Tao *et al.*³³, we found *wnt11* distributed evenly between the dorsal and ventral cells. Our results rather support the finding of Schroeder *et al.*³⁴, who found maternal *wnt11* mRNA symmetrically distributed along the dorsal-ventral axis, and proposed that spatially regulated translation leads to asymmetric distribution of the Wnt11 protein. None of the 41 maternal mRNAs included in this study shows dorso-ventral polarization, although dorso-ventral asymmetry is observed starting from the 4-cell stage¹². Of course, the dorso-ventral asymmetry could be induced by entirely different genes, but since our panel includes ten genes (*dvl2*, *dvl3*, *lrp6*, *wnt11*, *tcf3*, *gsk3b*, *ctnnb1*, *foxh1*, *trim36*, and *axin1*) previously implicated in dorsal-ventral patterning^{16,19–21}, this would be surprising. Rather, we suggest the dorsal-ventral asymmetry in *Xenopus laevis* is due to early accumulation of dorsal proteins that regulate signaling pathways, which after the midblastula transition activate the expression of downstream dorsalizing factors at the dorsal site of the embryo. We speculate these proteins are present already in the oocyte and accumulate in the future dorsal site during the cortical rotation. Our set includes only one, gene (*vg1*) previously shown to be involved in the left-right specification¹⁸. We find no evidence of asymmetric distribution of *vg1* or of any other of the studied transcripts that could be implicated in left-right specification.

Methods

Ethics statement. This study was carried out in accordance with the Act No 246/1992 Coll., on the protection of animals against cruelty. Official permission was issued to Faculty of Science, Charles University in Prague by the Central Commission for



Heatmaps




Figure 2 | Hierarchical clustering of blastomeres and mRNAs of *Xenopus laevis* early embryos presented as heatmaps. One embryo from each developmental stage was arbitrarily chosen for the cluster analysis. (A) 8-cell, (B) 16-cell and (C) 32-cell embryo. Green color indicates low expression and red high expression. The dendrograms clustering blastomeres and genes are shown in the top and left side of the heatmap, respectively. In the dendrograms similarity between blastomeres/genes is indicated by the height at which they are joined.




Figure 3 | Principal component analysis of maternal genes. (A) 8-cell stage (four embryos/32 cells analyzed), (B) 16-cell stage (four embryos/64 cells analyzed), and (C) 32-cell stage (32 cells analyzed) embryos. Three clusters are seen in all the developmental stages. First cluster comprises *18S rRNA*, *5S rRNA*, and *cyc1* (mitochondrial cytochrome *c*); second cluster *vg1*, *cdx1* (*xcad2*), *wnt11*, *dazl*, *vegt*, *ddx25* (*deadsouth*), *otx1*, and *trim36*; third cluster (red squares) *fzd7*, *par1*, *bmp2*, *pias1*, *dvl2*, *dvl3*, *lrp6*, *foxr1*, *fart1*, *mapk8*, *odc1*, *axin1*, *est1*, *U3 snoRNA*, *apc*, *gapdh*, *acta*, *eef1a1*, *tcf3*, *zpc*, *gsk3b*, *ctnnb1*, *mos*, *foxh1*, *stat3*, *pax6*) were designed with Primer3Plus (<http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi>). Annealing temperature was set to 60°C and the length of qPCR products was around 120bp. Specificity of the assays was tested in silico using blast (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Primer sequences are available upon request. qPCR and melting curve validations were measured on the real-time CFX384 cyclers system (BioRad). qPCR mix contained 2 µl of cDNA, 0.5 µl of forward and reverse primers (mixture 1:1, 10 µM each), 5 µl of SYBR Green JumpStart™ Taq Ready Mix™ (Sigma), and water in a total volume of 10 µl. The cycling program was: 2 min at 95°C for activation of the polymerase, followed by 40 cycles of denaturation at 95°C for 15 sec, annealing at 60°C for 20 sec and elongation at 72°C for 30 sec. Post PCR melting curves were measured from 65 to 95°C in 0.5°C intervals to validate the formation of expected PCR products.

The animal-vegetal distinction is found along the y-axis (PC2), while differences in expression level of the maternal transcripts is reflected along the x-axis (PC1).

Animal Welfare under the Ministry of Agriculture of Czech Republic (accreditation No.24773/2008-10001, date of expiry 10.12.2013). The experiment was also approved by the institutional ethics committee.

In vitro fertilization and collection of single blastomeres. Two *Xenopus laevis* females were injected with 450 U of human chorionic gonadotrophin hormone (hCG) and kept overnight at room temperature. Ovulated oocytes were obtained 12 hours after stimulation by manual squeezing. Testes of *Xenopus laevis* males were removed and homogenized in L-15 Leibowitzs medium with 15% of fetal bovine serum. Oocytes were *in vitro* fertilized by sperm suspension and 1× MMR medium (Marcs Modified Ringers medium: 0.1 M NaCl, 2.0 mM KCl, 1 mM MgCl₂, 2 mM CaCl₂, 5 mM HEPES, pH 7.7) was added to cover the fertilized oocytes. Jelly coats were removed by cysteine treatment (2.2% in MMR, for 8 minutes) at 20 minutes after fertilization. During cell division the embryos were transferred to Ca²⁺ and Mg²⁺ free medium (88 mM NaCl, 1 mM KCl, 2.4 mM NaHCO₃, 7.5 mM Tris, pH 7.7) and kept there for 60 minutes in order to separate single blastomeres. Vitelline membranes of the embryos containing 8, 16, and 32 cells were manually torn and individual blastomeres were collected. The blastomeres were collected separately from animal and vegetal hemispheres, but in random order to avoid introducing any technical bias and subjective dorso-ventral specification before measurement.

Isolation of total RNA and reverse transcription. The individual blastomeres collected from the 8 and 16-cell stages were homogenized in 200 µl and blastomeres from the 32-cell stage in 100 µl of TRIzol reagent (Invitrogen), and immediately frozen at -80°C. After thawing, the samples were carefully vortexed for 2 minutes to pulp yolk. Manufacturer's instructions were followed during isolation of total RNA, and glycogen was added to enhance RNA precipitation yield. RNA from individual cells from the 8 and 16-cell stages was dissolved in 20 µl of RNase-DNase free water. Samples from the 32-cell stage were dissolved in 12 µl of water. RNA concentration was measured using the Nanodrop ND1000 quantification system. cDNA was produced with SuperScript™ III Reverse transcriptase kit (Invitrogen). 10 ng of total RNA, 0.5 µl of a mixture of oligo-dT and random hexamers (mixture 1:1, 50 µM each), 0.5 µl of dNTPs (10 mM each), and 0.5 µl of RNA spike (*in vitro* transcribed artificial RNA with 3' polyA tail and 5' cap, TATAA Biocenter) were mixed with sterile water to a total volume of 6.5 µl. Samples were incubated for 5 min at 70°C, followed by 20 sec at 25°C, and then cooled at 4°C for 1 min. 100 U of SuperScript III reverse transcriptase, 20 U of RNaseOUT™ (recombinant ribonuclease inhibitor, Invitrogen), 0.5 µl of 0.1 M DTT and 2 µl of 5× first strand synthesis buffer was added to a final volume of 10 µl. The following temperature gradient profile was used to synthesize cDNA: 5 min at 25°C, 60 min at 50°C, 15 min at 55°C, and 15 min at 75°C. cDNA samples were diluted 8 times to a final volume of 80 µl and were stored at -20°C.

Primer design and quantitative PCR. Primer assays for the selected 41 maternal genes (*fzd7*, *otx1*, *bmp2*, *pias1*, *dvl2*, *dvl3*, *lrp6*, *trim36*, *foxr1*, *frat1*, *mapk8*, *odc1*, *18S rRNA*, *5S rRNA*, *cyc1*, *acta*, *tubb*, *gapdh*, *maml1*, *eef1a1*, RNA polymerase II, *U3 snoRNA*, *par1*, *oct60*, *ddx25*, *dazl*, *cdx1*, *wnt11*, *vg1*, *vegt*, *axin1*, *est1*, *apc*, *tcf3*, *zpc*, *gsk3b*, *ctnnb1*, *mos*, *foxh1*, *stat3*, *pax6*) were designed with Primer3Plus (<http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi>). Annealing temperature was set to 60°C and the length of qPCR products was around 120bp. Specificity of the assays was tested in silico using blast (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Primer sequences are available upon request. qPCR and melting curve validations were measured on the real-time CFX384 cyclers system (BioRad). qPCR mix contained 2 µl of cDNA, 0.5 µl of forward and reverse primers (mixture 1:1, 10 µM each), 5 µl of SYBR Green JumpStart™ Taq Ready Mix™ (Sigma), and water in a total volume of 10 µl. The cycling program was: 2 min at 95°C for activation of the polymerase, followed by 40 cycles of denaturation at 95°C for 15 sec, annealing at 60°C for 20 sec and elongation at 72°C for 30 sec. Post PCR melting curves were measured from 65 to 95°C in 0.5°C intervals to validate the formation of expected PCR products.

Data analysis. Data were analyzed with GenEx (MultiD, version 5.3.). Off scale data were removed during pre-processing using a cut off at 36 cycles and outliers were identified with Grubb's test. Missing data were imputed using the average Cq of the other blastomeres from the same hemisphere in the same embryo (roughly 6% of the data was imputed). All data were normalized to the spike and converted to relative quantities (relative to the highest Cq for each gene, corresponding to arbitrary assigning an expression of 1 to the least expressed sample). Last step of the pre-processing was to transform the data to log₂ scale. 8-cell, 16-cell and 32-cell embryos were analyzed separately. The combined expression of the individual blastomeres adds up to the expression of the original oocyte, which is expected, since *de novo* expression does not start until MBT. Three multivariate statistical methods were used for classification: principal component analysis (PCA), Kohonen self-organizing map (SOM), and hierarchical clustering presented as heatmaps. PCA of individual blastomeres (Fig. 1) was performed on data autoscaled along genes, while heatmaps (Fig. 2) and PCA of genes (Fig. 3) were performed on data autoscaled along samples³⁵.

- Bengtsson, M., Ståhlberg, A., Rorsman, P. & Kubista, M. Gene expression profiling in single cells from the pancreatic islets of Langerhans reveals lognormal distribution of mRNA levels. *Genome Res.* **15**, 1388–1392 (2005).



2. Guo, G. *et al.* Resolution of cell fate decisions revealed by single-cell gene expression analysis from zygote to blastocyst. *Dev. Cell* **18**, 675–685 (2010).
3. May, A. *et al.* Multiplex RT-PCR expression analysis of developmentally important genes in individual mouse preimplantation embryos and blastomeres. *Biol. Reprod.* **80**, 194–202 (2009).
4. Galan, A. *et al.* Functional genomics of 5- to 8-cell stage human embryos by blastomere single-cell cDNA analysis. *PLoS One* **5**, e13615 (2010).
5. Virant-Klun, I. *et al.* Expression of pluripotency and oocyte-related genes in single putative stem cells from human adult ovarian surface epithelium cultured in vitro in the presence of follicular fluid. *BioMed Res. Int.* **2013**, 1–18 (2012).
6. Danilchik, M. V. & Gerhart, J. C. Differentiation of the animal-vegetal axis in *Xenopus laevis* oocytes. I. Polarized intracellular translocation of platelets establishes the yolk gradient. *Dev. Biol.* **122**, 101–112 (1987).
7. King, M. L., Messitt, T. J. & Mowry, K. L. Putting RNAs in the right place at the right time: RNA localization in the frog oocyte. *Biol. Cell* **97**, 19–33 (2005).
8. Sindelka, R., Jonak, J., Hands, N., Bustin, S. A. & Kubista, M. Intracellular expression profiles measured by real-time PCR tomography in the *Xenopus laevis* oocyte. *Nucleic Acids Res.* **36**, 387–392 (2008).
9. Sindelka, R., Sidova, M., Svec, D. & Kubista, M. Spatial expression profiles in the *Xenopus laevis* oocytes measured with qPCR tomography. *Methods* **51**, 87–91 (2010).
10. Roth, S. The origin of dorsoventral polarity in *Drosophila*. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **358**, 1317–1329 (2003).
11. Aanstad, P. & Whitaker, M. Predictability of dorso-ventral asymmetry in the cleavage stage zebrafish embryo: an analysis using lithium sensitivity as a dorso-ventral marker. *Mech. Dev.* **88**, 33–41 (1999).
12. Sive, L. H., Grainger, M. R. & Harland, M. R. Early development of *Xenopus laevis* – A laboratory manual. Cold Spring Harbor Laboratory Press 2000, chapter 2, figure 2.1.
13. Vincent, J. P. & Gerhart, J. C. Subcortical rotation in *Xenopus* eggs: an early step in embryonic axis specification. *Dev. Biol.* **123**, 526–539 (1987).
14. Denegre, J. M. & Danilchik, M. V. Deep cytoplasmic rearrangements in axis-respecified *Xenopus* embryos. *Dev. Biol.* **160**, 157–164 (1993).
15. Miller, J. R. *et al.* Establishment of the dorsal-ventral axis in *Xenopus* embryos coincides with dorsal enrichment of dishevelled that is dependent on cortical rotation. *J. Cell Biol.* **146**, 427–437 (1999).
16. Marikawa, Y., Li, Y. & Elinson, R. P. Dorsal determinants in the *Xenopus* egg are firmly associated with the vegetal cortex and behave like activators of the Wnt pathway. *Dev. Biol.* **191**, 69–79 (1997).
17. Darras, S., Maikawa, Y., Elinson, R. P. & Lemaire, P. Animal and vegetal pole cells of early *Xenopus* embryos respond differently to maternal dorsal determinants: implications for the patterning of the organiser. *Development* **124**, 4275–4286 (1997).
18. Hyatt, B. A., Lohr, J. L. & Yost, H. J. Initiation of vertebrate left-right axis formation by maternal Vg1. *Nature* **384**, 62–65 (1996).
19. King, M. L., Messitt, T. J. & Mowry, K. L. Putting RNAs in the right place at the right time: RNA localization in the frog oocyte. *Biol. Cell* **97**, 19–33 (2005).
20. Kofron, M. *et al.* New roles for FoxH1 in patterning the early embryo. *Development* **131**, 5065–5078 (2004).
21. Kofron, M. *et al.* Wnt11/ β -catenin signaling in both oocytes and early embryos acts through LRP6-mediated regulation of axin. *Development* **134**, 503–513 (2007).
22. Cuykendall, T. N. & Houston, D. W. Identification of germ plasm-associated transcripts by microarray analysis of *Xenopus* vegetal cortex. *Dev. Dyn.* **239**, 1838–1848 (2010).
23. Cuykendall, T. N. & Houston, D. W. Vegetally localized *Xenopus* trim36 regulates cortical rotation and dorsal axis formation. *Development* **136**, 3057–3065 (2009).
24. Bergkvist, A. *et al.* Gene expression profiling – Clusters of possibilities. *Methods* **50**, 323–335 (2010).
25. Keller, R. E. Vital dye mapping of the gastrula and neurula of *Xenopus laevis*: I. Prospective areas and morphogenetic movements of the superficial layer. *Dev. Biol.* **42**, 222–241 (1975).
26. Keller, R. E. Vital dye mapping of the gastrula and neurula of *Xenopus laevis*: II. Prospective areas and morphogenetic movements of the deep layer. *Dev. Biol.* **51**, 118–137 (1976).
27. Kikkawa, M., Takano, K. & Shinagawa, A. Location and behavior of dorsal determinants during first cell cycle in *Xenopus* eggs. *Development* **122**, 3687–3696 (1996).
28. Weaver, C. & Kimelman, D. Move it or lose it: axis specification in *Xenopus*. *Development* **131**, 3491–3499 (2004).
29. Medina, A., Wendler, S. R. & Steinbeisser, H. Cortical rotation is required for the correct spatial expression of nr3, sia and gsc in *Xenopus* embryos. *Int. J. Dev. Biol.* **41**, 741–745 (1997).
30. Kugler, J. M. & Lasko, P. Localization, anchoring and translational control of oskar, gurken, bicoid and nanos mRNA during *Drosophila* oogenesis. *Fly (Austin)* **3**, 15–28 (2009).
31. Steinhauer, J. & Kalderon, D. Microtubule polarity and axis formation in the *Drosophila* oocyte. *Dev. Dyn.* **235**, 1455–1468 (2006).
32. Hainski, A. M. & Moody, S. A. *Xenopus* maternal mRNAs from dorsal animal blastomere induce a secondary axis in host embryo. *Development* **116**, 347–355 (1992).
33. Tao, Q. *et al.* Maternal Wnt11 activates the canonical Wnt signaling pathway required for axis formation in *Xenopus* embryos. *Cell* **120**, 857–871 (2005).
34. Schroeder, K. E., Condic, M. L., Eisenberg, L. M. & Yost, H. J. Spatially regulated translation in embryos: asymmetric expression of maternal wnt-11 along the dorso-ventral axis in *Xenopus*. *Dev. Biol.* **214**, 288–297 (1999).
35. Ståhlberg, A., Rusnakova, V., Forootan, A., Anderova, M. & Kubista, M. RT-qPCR work-flow for single-cell data analysis. *Methods* **59**, 80–88 (2013).

Acknowledgements

This study was supported by Grant agency of Czech Republic 301/09/1752; Ministry of Youth, Education and Sports of the Czech Republic AV0Z50520701; BIOCEV CZ.1.05/1.1.00/02.0109 from the ERDF and by the grant SVV–2013-267212.

Author contributions

M.F. performed the experiments and together with M.K. analyzed the obtained data. R.S. and M.K. conceived and designed the experiments. All authors co-wrote the manuscript and approved the final paper.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Flachsova, M., Sindelka, R. & Kubista, M. Single blastomere expression profiling of *Xenopus laevis* embryos of 8 to 32-cells reveals developmental asymmetry. *Sci. Rep.* **3**, 2278; DOI:10.1038/srep02278 (2013).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported license. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0>