



OPEN

SUBJECT AREAS:

BIOTECHNOLOGY

MICROBIOLOGY

BIOCATALYSIS

MICROBIAL ECOLOGY

Diversity and functional significance of cellulolytic microbes living in termite, pill-bug and stem-borer guts

Zeenat Bashir^{1*}, Vamsi Krishna Kondapalli^{1*}, Nidhi Adlakha¹, Anil Sharma², Raj K. Bhatnagar², Girish Chandel³ & Syed Shams Yazdani¹Received
22 April 2013Accepted
16 August 2013Published
30 August 2013

Correspondence and requests for materials should be addressed to S.S.Y. (shams@icgeb.res.in)

* These authors contributed equally to this work.

¹Synthetic Biology and Biofuels Group, ²Insect Resistance Group, International Centre for Genetic Engineering and Biotechnology, Aruna Asaf Ali Marg, New Delhi, ³Department of Plant Molecular Biology and Biotechnology, College of Agriculture, Indira Gandhi Agricultural University, Raipur (C.G.).

Arthropods living on plants are able to digest plant biomass with the help of microbial flora in their guts. This study considered three arthropods from different niches - termites, pill-bugs and yellow stem-borers - and screened their guts for cellulase producing microbes. Among 42 unique cellulase-producing strains, 50% belonged to Bacillaceae, 26% belonged to Enterobacteriaceae, 17% belonged to Microbacteriaceae, 5% belonged to Paenibacillaceae and 2% belonged to Promicromonosporaceae. The distribution of microbial families in the three arthropod guts reflected differences in their food consumption habits. Most of the carboxymethylcellulase positive strains also hydrolysed other amorphous substrates such as xylan, locust bean gum and β -D-glucan. Two strains, A11 and A21, demonstrated significant activity towards Avicel and p-nitrophenyl- β -D-cellobiose, indicating that they express cellobiohydrolase. These results provide insight into the co-existence of symbionts in the guts of arthropods and their possible exploitation for the production of fuels and chemicals derived from plant biomass.

Rising concern over the cost and sustained availability of fossil fuels has increased demand for the development of alternative and preferably renewable fuel resources, with lignocellulosic biomass as one promising resource^{1,2}. Lignocellulose represents ~90% of the dry weight of all plant materials³, is primarily composed of the sugar polymers cellulose (35–50%) and hemicellulose (20–35%) together with lignin (5–30%) that provides structural support for the plant⁴. Cellulose and hemicellulose represent abundant and inexpensive sources of fermentable sugar for the production of various chemicals and biofuels, but finding efficient catalysts for their saccharification to monomeric sugar and producing those catalysts in a cost-effective manner are major hurdles⁵.

Biological catalysts such as cellulase/hemicellulase enzymes play significant roles in the production of ethanol using plant biomass⁶. The enzymatic hydrolysis of cellulose polymers is achieved by the synergistic activity of three different groups of cellulase enzymes¹: endoglucanase², exoglucanase or cellobiohydrolase, and³ β -glucosidase^{7,8}. Arabinoxylanases and xylanases are groups of enzymes that hydrolyse hemicellulose components. A large number of these hydrolytic enzymes have been discovered in fungi⁹, but they are typically associated with certain limitations, such as stability to low pH and thermal stresses and low specific activities¹⁰. As bacteria are able to tolerate environmental extremes, they represent an ideal source for the screening and isolation of novel cellulolytic enzymes to help overcome these challenges¹⁰. Therefore, researchers are investigating natural systems that have evolved to decompose plant cell wall polymers as sources for enzyme discovery.

Arthropods are the most abundant and successful species on earth that can survive in diverse ecological niches^{11,12}. The digestive tracts of arthropods contain microbiota members from the bacterial, fungal, protozoan and archaeal groups, which assist with various physiological functions including food digestion, nitrogen fixation, nutrition and pheromone production¹³. Comprehensive analyses of the transcriptomes and metagenomes of termites and symbiotic microbes have generated huge amounts of genomic information for the screening of cellulase encoding genes^{14,15}. However, enzymes screened through metagenomic and metatranscriptomic studies could have issues with heterologous overexpression in terms of folding and yield^{16,17} because little is known about the host organism. Therefore, natural cellulolytic bacteria that can be cultured under laboratory conditions are excellent platforms for the study of the properties of hydrolytic enzymes.

In the present work, we have selected three arthropods with different dietary habits, i.e., two insects (termites and yellow stem-borers) and a crustacean (pill-bugs), and explored their guts as sources for the isolation and



characterisation of cellulose-degrading bacteria. The ability of these arthropods to feed on wood, foliage and detritus is likely to involve catalysis by different types of cellulases/hemicellulases that are secreted by gut microbiota to digest the structural and recalcitrant lignocellulosic residues in their foods. Thus, the main goals of this study were to explore the guts of these arthropods that feed on potential biofuel feedstocks and to compare their hydrolytic activities. We successfully isolated 42 cellulose-degrading bacterial strains from these arthropods and characterised their abilities to produce hydrolytic enzymes.

Results

Screening and characterisation of cellulolytic bacteria from the guts of arthropods. The guts of arthropods feeding on woody and agricultural biomass are an ideal place for symbiotic cellulolytic microbes to thrive. We selected two insects, termites and yellow stem borers, and a crustacean, pill-bugs, because of their ability to live in diverse niches. Termites were collected from the trunks and roots of dead trees; pill-bugs were collected from dampened soil containing decomposed leaf litters; and yellow stem borers living in rice stems were collected from an agricultural rice field (Figure 1). Arthropod families were identified by performing BLAST with the nucleotide sequences of the mitochondrial cytochrome oxidase I (CO-I) genes from each organism and further confirmed by morphological characterisation. The nearest possible hits for termites, pill-bugs and yellow stem borers were *Odontotermes hiananensis*, *Armadillidium* sp. and *Scirpophaga incertulas*, respectively. The guts of these arthropods were dissected as described in the Methods section, and the microbes living in the guts were grown on agar plates containing CMC and trypan blue at various dilutions under both aerobic and anaerobic conditions. Unique isolated colonies showing clearance zones on a blue background were purified by re-streaking, and their cellulolytic capabilities were further confirmed by Congo red staining of agar plates¹⁸. Approximately 42 bacterial colonies were selected by the above screening method for phylogenetic analysis and enzymatic activity analysis. Among these, 26 bacteria were isolated by screening under aerobic conditions, and 16 were isolated under anaerobic conditions (Table 1).

Phylogenetic analysis of gut bacteria. Among the 42 bacterial colonies selected for further characterisation, 17 were isolated from termite guts, 8 were isolated from pill-bug guts and 17 were isolated from yellow stem borer guts (Table 1). Phylogenetic analysis of the 16S rDNA sequences revealed that all identified bacterial species belonged to five families - 50% belonged to Bacillaceae, 26% belonged to Enterobacteriaceae, 17% belonged to Microbacteriaceae, 5% belonged to Paenibacillaceae and 2% belonged to Promicromonosporaceae (Figure 2A). Between 60 and 70% of the bacteria from the guts of termites and pill-bugs were from the Bacillaceae family, while bacteria from the Microbacteriaceae and Enterobacteriaceae families were more dominant in the guts of yellow stem borers (Figure 2B).

Characterisation of bacterial enzymes hydrolysing plant biomass.

Microbes residing in the guts of biomass feeding arthropods are likely to produce a variety of hydrolytic enzymes. In previous studies we showed that *Paenibacillus* ICGB2008 isolated from the gut of the cotton bollworm produced several biomass-degrading enzymes, including cellulases and hemicellulases^{18,19}. Therefore, we tested the expression levels in the gut microbes of major glycosyl hydrolase enzymes belonging to the categories of cellulase, hemicellulase, mannanase and glucanase. The expression levels of endoglucanase, β -glucosidase, xylanase, β -xylosidase, mannanase and β -D-glucanase were determined using the substrates CMC, pNPG, Xylan, pNPX, locust bean gum and barley β -D-glucan, respectively. To test exocellulase or cellobiohydrolase expression, both Avicel and pNPC (in the presence of the β -glucosidase inhibitor glucono- δ -lactone) were used as substrates. A potential biofuel feedstock was also used, namely alkali treated rice straw, and it mainly contained crystalline cellulose as well as some amorphous forms of cellulose. These assays were used to test both extracellular and cell associated enzymes. Except for A13, A16 and T3, all natural isolates produced at least one type of glycosyl hydrolase (Figure 3A and 3B). The reason that these three strains are negative in the liquid assay despite being positive for CMC activity on the agar plates could be related to the stringency criteria set as described in the Methods section. The majority of microbes produced more than one kind of hydrolytic enzyme (Figure 3C). There were 23 microbes that produced at least one hydrolytic enzyme from each category of biomass degrading enzymes. Several strains produced only one category of enzyme, such as cellulases (5 strains) and hemicellulases (1 strain), which hydrolysed pNPG and pNPX, respectively, but did not produce significant amounts of any other type of hydrolase (Figure 3). Extracellular fractions were found to be rich in enzymes specific for amorphous polymers (Figure 3A). Only a few microbes secreted enzymes that were able to hydrolyse dimeric or oligomeric substrates, such as pNPG, pNPC and pNPX, or complex structures, such as rice straw. On the other hand, the majority of cell associated fractions hydrolysed dimeric substrates such as pNPG and pNPX (Figure 3B). However, cell associated fractions of none of the microbes were able to hydrolyse pNPC. Some strains also produced enzymes in the extracellular or cell-associated fractions for the hydrolysis of complex structures such as rice straw and Avicel.

Quantitative assessments of the hydrolytic activities in both the extracellular and cell-associated fractions enabled us to analyse their relative contributions to biomass degradation (Figure 4, 5 and 6). The majority of strains secreted high quantities of enzymes specific for amorphous polymers such as CMC, xylan, locust bean gum and barley β -D-glucan (Figure 4). Relatively fewer strains produced enzymes that were able to hydrolyse significant amounts of crystalline polymers such as Avicel and rice straw (Figure 5D, 6A and 6D). The strains hydrolysing the greatest amounts of rice straw were A11, A18, A21, A22, S2 and U8 (Figure 5D and 6D), while strains A11, A17, A21, A24, U5 and U10 hydrolysed Avicel (Figure 6A). Enzymes



Termite

Pill bug

Rice Stem Borer

Figure 1 | Pictures of termites, pill-bugs and rice stem-borers collected from various ecological niches.



Table 1 | Coding of cellulolytic microbes isolated from insect guts

Insect	Aerobic screening				Anaerobic screening			
	Strain ID	GenBank Accession	Nearest relative	Strain ID	GenBank Accession	Nearest relative		
Termite	A1	KC434960	Bacillus sp. 6063	T1	KC434986	Bacillus subtilis M50		
	A4	KC434961	Klebsiella sp. clone F7	T2	KC434987	Bacillus licheniformis ACO1		
	A5	KC434962	Trabulsirella guamensis GTC1379	T3	KC434988	Trabulsirella guamensis GTC1379		
	A6	KC434963	Bacillus pumilus BSH4	T4	KC434989	Bacillus sp. DY9-35		
	A7	KC434964	Bacillus sp. SCS510					
	A9	KC434965	Pantoea agglomerans WAB1927					
	A10	KC434966	Bacillus licheniformis EdyKolBI23					
	A11	KC434967	Bacillus licheniformis BCRC 15413					
	A12	KC434968	Bacillus licheniformis SubatMucBI16					
	A13	KC434969	Bacillus cereus TAUC5					
	A16	KC434970	Bacillus cereus F837/76					
	A17	KC434971	Bacillus subtilis K21					
	A18	KC434972	Paenibacillus polymyxa DSM 36T					
	PillBug	A19	KC434973	Paenibacillus polymyxa YRL13	W1	KC434990	Bacillus subtilis KL-073	
		A21	KC434974	Bacillus subtilis M50	W6	KC434991	Bacillus thuringiensis 2PR56-10	
		A22	KC434975	Bacillus subtilis M16K	W7	KC434992	Bacillus tequilensis VITJAM2	
		A23	KC434976	Enterobacter aerogenes KCTC 2190				
		A24	KC434977	Cellulosimicrobium sp. TUT1242				
Yellow Stem Borer	S1	KC434978	Bacillus species BB2_1A	U2	KC434993	Bacillus sp. PR 1.7		
	S2	KC434979	Bacillus subtilis AQ1	U3	KC434994	Pantoea species NCCP116		
	S3	KC434980	Microbacteriaceae bacterium HIB-6	U4	KC434995	Pantoea agglomerans WAB1927		
	S4	KC434981	Microbacteriaceae bacterium BMC-3	U5	KC434996	Klebsiella pneumoniae L-13		
	S5	KC434982	Microbacterium oleivorans CCGE2277	U6	KC434997	Klebsiella pneumoniae L-13		
	S6	KC434983	Microbacterium arborescens DSM 20754	U7	KC434998	Microbacteriaceae bacterium BMC-3		
	S7	KC434984	Microbacterium arborescens DSM 20754	U8	KC434999	Bacillus subtilis C1Y001		
	S8	KC434985	Microbacterium arborescens JB8_2B	U9	KC43500	Enterobacter species DHM1T		
			U10	KC43501	Serratia marcescens PSI			




Figure 2 | Classification and phylogenetic analysis of natural isolates from the guts of termites, pill-bugs and yellow stem-borers. (A) 16s rDNA sequences of 42 isolates screened for their ability to produce cellulolytic enzymes obtained from the guts of termites (●), pill bugs (○) and yellow stem borers (■) were used to perform blast searches to identify their nearest neighbors, and MEGA5 was used to construct the phylogenetic tree. The natural isolates were found to belong to five families. (B) Relative abundance of various bacterial families in the guts of these three arthropods.




Figure 4 | Quantitative assessment of the enzymatic activities of extracellular fractions against amorphous polysaccharides. All bacterial strains that were found to be cellulase positive by the agar plate assay were grown in liquid medium, and their extracellular fractions were tested for the hydrolysis of various substrates. Each graph indicates the substrate. Black bars represent strains that exhibited significantly higher activities compared to the negative controls, while white bars represent negative strains. The data represent the average and standard deviation of two different assays.




Figure 5 | Quantitative assessment of the enzymatic activities of extracellular fractions against chromomeric substrates and rice straw. All bacterial strains that were found to be cellulase positive by the agar plate assay were grown in liquid medium, and their extracellular fractions were tested for the hydrolysis of various substrates. Each graph indicates the substrate. Black bars represent strains that exhibited significantly higher activities compared to the negative controls, while white bars represent negative strains. The data represent the average and standard deviation of two different assays.





Figure 6 | Quantitative assessment of the enzymatic activities of the cell-bound fractions. All bacterial strains that were found to be cellulase positive by the agar plate assay were grown in liquid medium, the cells were lysed by sonication, and the lysed cell samples were tested for the hydrolysis of various substrates. Each graph indicates the substrate. Black bars represent strains that exhibited significantly higher activities compared to the negative controls, while white bars represent negative strains. The data represent the average and standard deviation of two different assays.



A.



B.




Figure 7 | Zymograms to determine the enzymatic activities of the extracellular fractions of natural isolates against (A) CMC and (B) xylan.

production. Nature has created reservoirs of catalysts that perform this function in a variety of niches. We have attempted to mine these biological reservoirs by selecting arthropods from three diverse niches and screening for microbes capable of biomass hydrolysis in their guts. While other methodologies such as metagenomics and metatranscriptomics are becoming popular approaches to screen for biomass hydrolysing enzymes, they are often associated with issues of low heterologous expression yield and protein misfolding¹⁶. Our approach of identifying culturable microbes with biomass hydrolysing capabilities has the advantage of studying the enzyme characteristics in both homologous and heterologous hosts.

This study has revealed significant knowledge regarding the diversity of microbes existing in the guts of arthropods and their roles in biomass degradation. Among the arthropods we selected for this study, termites are the best-studied arthropods in terms of cellulolytic microbes^{13–15}, although the majority of the cellulolytic microbes that have been identified are difficult to culture in the lab. No previous studies have reported on the nature of cellulolytic microbes existing in the guts of pill-bugs and yellow stem borers. We found that greater than 60% of the cellulolytic microbes isolated from the guts of termites and pill-bugs belonged to the Bacillaceae family, while greater than 75% of the cellulolytic microbes isolated from the guts of yellow stem borers belonged to the Microbacteriaceae and Enterobacteriaceae families. These differences in the microbial flora reflected differences in the food habits of the arthropods. While termites and pill-bugs were collected from decomposed trees and litter, yellow stem borer larvae were collected from rice plants at the time of seeding. The pill-bug gut also contained *Cellulosimicrobium* sp., a known natural producer of cellulases and xylanases.

Hydrolytic enzymes have been reported in both the extracellular and cell-bound fractions^{18–21}. More often, the enzymes hydrolysing amorphous polysaccharides are secreted, while those hydrolysing crystalline substrates are found in the membrane-bound fraction of the bacteria^{18,20}. Enzymes hydrolysing dimeric sugars into monomeric forms can be found in both fractions^{19,22}. Therefore, we selected substrates to analyse the hydrolysing enzymes present in both the extracellular and cell-bound fractions. The majority of the natural isolates were expected to produce CMCase as they were selected on CMC containing agar plates. Among a total of 42 selected strains, 17 secreted significant amounts of endocellulase. The remaining strains did not pass the stringency criteria of the liquid assay. Most of the CMCase producing strains also secreted hydrolytic enzymes for other amorphous substrates. We found a large number of strains producing cell-bound enzymes to hydrolyse dimeric substrates, although few strains also secreted these enzymes into the extracellular medium. Few strains demonstrated significant Avicelase activity in the cell-bound fraction or pNPC activity in the extracellular fraction, indicating the presence of hydrolytic enzymes against crystalline substrates.

Enzyme production by the major CMCase and xylanase producers was further confirmed by performing zymograms. A number of enzymes have been reported in the literature with CMCase activity in the ranges of 40 kDa and 90 kDa and xylanase activity in the range of 60 kDa^{18,23,24}. The xylanase activity with a much higher molecular mass is intriguing and could represent oligomers.

In conclusion, the results of this study provide insight into the presence of diverse microbial flora in the guts of three arthropods differing in their food habits and into the roles of these floras in the




Figure 8 | Kinetic properties of enzyme preparations from strains A11 and A21. Substrate (para-nitrophenyl cellobiose (pNPC)) saturation kinetics of the (A) A11 and (B) A21 strains. (C) Hanes-Woolf graph for the calculation of K_m and V_{max} . The data represent the average of two different assays.

degradation of plant biomass. The knowledge gained from these studies could be exploited for the identification of novel enzymes and the development of enzyme systems for the effective hydrolysis of plant biomass.

Methods

Media composition and chemical reagents. Tryptic Soya broth (1.7% Tryptone, 0.3% Soya peptone, 0.25% K_2HPO_4 and 0.5% NaCl, pH 7.0) (Sigma, USA) was used as the medium for the cultivation of microbes in liquid culture and on agar Petri-plates. Reagents for analytical assays such as cellobiose, Avicel, birchwood xylan, carboxymethylcellulose (CMC), p-nitrophenyl β -D-cellobioside (pNPC), p-nitrophenyl β -D-glucopyranoside (pNPG) and p-nitrophenyl β -D-xylopyranoside (pNPX) and other chemicals were purchased from Sigma (Saint Louis, MO, USA). Alkali-treated rice straw was obtained from ICT, Mumbai²⁵.

Collection of host organisms. Approximately 25 wood-feeding worker termites and litter-feeding pill bugs were collected from different locations of ICGEB, New Delhi, India, and 25 yellow stem-borer insect larvae were collected from the paddy fields of the Biotechnological Research Experiments field, Raipur University, Chhattisgarh, India during the month of October, 2011. The guts of the insects were processed to isolate microbes, and host DNA was isolated from the carcasses for species identification.

Screening for cellulase producing bacteria. To isolate gut microbes, the insects were first pre-chilled on ice for 2–3 min, followed by surface sterilisation with 70% ethanol for 1 min. All the dissections were performed in the laminar flow cabinet. Guts were carefully removed with the help of aseptic needles and placed in a 1.5 ml sterile microfuge tube containing 100 μl of ice-cold buffered saline solution (pH 7.2)¹⁸. The guts were then homogenised using a micropestle, and the suspension was serially diluted from 10^{-1} to 10^{-7} , plated on Tryptic Soya agar (TSA) plates containing 0.5% CMC and 0.1% trypan blue, and incubated for 72 h at 30 °C. Isolated bacterial colonies showing clearance zones on the plates were further confirmed to produce cellulase by spreading on a TSA-CMC plate and observing the clearance zone via the Congo-red method¹⁸, and the results of this experiment were then used to select colonies for further characterisation. To isolate anaerobic bacteria, all dissections and platings were performed under anaerobic conditions in an anaerobic chamber (Shel Lab).

Identification of microbial and arthropod families. All the bacterial strains showing clearance zones were streaked onto TSA plates and sent to Macrogen for 16S rDNA sequencing. Phylogenetic analysis was performed based on the blast results of the 16S rDNA sequence. The evolutionary history was inferred using the Neighbor-Joining method²⁶. The evolutionary distances were computed using the Kimura 2-parameter method²⁷, and evolutionary analyses were conducted in MEGA5²⁸.

The host species were identified using the molecular marker gene mitochondrial CO-I based on the previously described method²⁹. Genomic DNA from the host organisms was isolated by the conventional hexadecyl-trimethyl-ammonium bromide (CTAB) method³⁰, and the universal DNA primers LCO1490 (5'-gtgcaacaaatcataagattgg-3') and HCO2198 (5'-taaactcagggtgacaaaatac-3') were used to amplify the 710-bp region of the mitochondrial CO-I gene. The PCR product was analysed on a 1% agarose gel in the presence of ethidium bromide, purified using a gel extraction kit (Genetix) and sent to Macrogen for sequencing. The nucleotide sequences were then used in a Blast search to identify the arthropod families. The arthropod families were further confirmed by morphological identification^{31,32}.

Cultivation of microbes under laboratory conditions. Microbial strains isolated from the guts of insects were grown in Tryptic soya broth medium supplemented with 1% alkali pretreated rice straw at 37 °C at 150 rpm for 48 h. To grow natural isolates under anaerobic conditions, the cultures were grown at 37 °C for 72 h in anaerobic chambers. After the cell growth reached the stationary phase, the culture broth was centrifuged at $8000 \times g$ for 20 min, and then the supernatant was used as the source of extracellular enzymes. The cells were re-suspended in sodium phosphate buffer (pH 7.0) and lysed either by sonication in the case of aerobically grown cultures or by glass beads inside the anaerobic chamber in the case of anaerobically grown cultures. The resulting crude lysate was used as the cell-bound fraction.

Assays for enzymatic activity. The enzymatic activities of endocellulase (β -1,4-endoglucanase), cellobiohydrolase (β -1,4-exoglucanase), glucosidase (β -1,4-glucosidase), xylanase (β -1,4-endoxylanase), xylosidase (β -1,4-xylosidase), glucanase (β -D-glucanase) and mannanase (β -1,4-mannanase) were assayed using CMC, Avicel and pNPC, pNPG, birchwood xylan, pNPX, barley β -D-glucan and locust bean gum, respectively. Alkali pre-treated rice straw was also used as a substrate, as it was a potential feedstock for biofuel generation. The enzymatic activities in the supernatant were tested for their ability to hydrolyse CMC, pNPC, pNPG, xylan, pNPX, glucan, locust bean gum and rice straw. The enzymatic activities in the cell lysate were measured based on their activity against Avicel, pNPC, pNPG, pNPX and rice straw.

The reducing sugar released upon the hydrolysis of sugar polymers was measured by the dinitrosalicylic acid (DNSA) method, and the para-nitrophenol released upon the hydrolysis of the chromogenic substrate was measured by monitoring the absorbance at 405 nm^{19,27}. Briefly, a crude enzyme solution (0.125 ml) was mixed with 0.125 ml of a 1% sugar polymer solution in 0.05 M sodium phosphate buffer (pH 7.0) and incubated at 50 °C for 30 min. Enzymatic reactions containing Avicel and rice straw as substrates were incubated for 60 min. The reducing sugar produced in these experiments was measured by the DNS reagent at 540 nm¹⁸. One unit of enzymatic activity was defined as the amount of enzyme that released 1 μmol of reducing sugar from the substrate per minute under the above conditions.

For enzymatic assays with chromogenic substrates, 0.125 ml of enzyme sample was mixed with 0.55 ml of 5 mM substrate in 50 mM citrate phosphate buffer (pH 6.0) and incubated at 50 °C for 30 min. The enzyme assay with pNPC substrate was set up in the presence of 100 mM glucono- δ -lactone, an inhibitor of β -glucosidase activity, in a 0.2 ml volume. The reaction was stopped after incubation by the addition of 1 ml of 2 M Na_2CO_3 , and the absorbance of the released p-nitrophenol was measured at 405 nm. One unit of enzymatic activity was defined as the amount of enzyme catalysing the release of 1 μmol of p-nitrophenol per minute under the above conditions¹⁹.



All assays were performed in duplicate, and the data presented here represent the average of the two readings. *E. coli* DH5 α grown under similar conditions was taken as a negative control for all of the enzymatic assays. The data obtained for the experimental conditions and negative controls were compared in multiple comparisons using Student's *t*-test. The statistical significance of differences between various test samples and the control sample were determined using two-tailed unpaired Student's *t*-tests. *P* values ≤ 0.05 were considered statistically significant.

The kinetic parameters of enzymes produced by strains A11 and A21 against pNPC were determined as follows. The strains were grown in 1 L of TSB medium and harvested to obtain the extracellular fractions. The extracellular fractions were concentrated 50-fold using a 10 kDa cut-off poly-ether-sulphone membrane in a LabScale XL system (Merck Millipore) and buffer exchanged. This partially purified enzyme preparation was used for enzyme assays with different concentrations of pNPC ranging from 100 μ M to 5120 μ M. The substrate affinity (K_m) and catalytic efficiency (V_{max}) of these enzyme preparations were calculated using the Hanes-Woolf linear transformation³⁵.

Analytical methods. To assess the CMCase or xylanase activity of the protein by zymograms, the samples were resolved on a 12% SDS-PAGE gel containing either 0.5% (wt/vol) CMC or 0.5% (wt/vol) xylan. After electrophoresis, the gel was washed once with 20% (vol/vol) isopropanol in PBS for 1 min followed by three washes of 20 min each in PBS. The gel was incubated in PBS at 37°C for 1 h, stained with 0.1% (wt/vol) Congo red for 30 min, and destained with 1 M NaCl. Clear bands against the red background indicated CMCase or xylanase activity. Protein concentrations were estimated with the bicinchoninic acid (BCA) protein assay kit (Pierce) using bovine serum albumin as a standard.

Nucleotide sequence accession numbers. The nucleotide sequences of 16S rDNA obtained in this study have been deposited in the NCBI gene bank, and the GenBank accession numbers of the bacterial isolates are listed in Table 1.

- Himmel, M. E. & Bayer, E. A. Lignocellulose conversion to biofuels: current challenges, global perspectives. *Curr. Opin. Biotechnol.* **20**, 316–317 (2009).
- Jager, G. & Buchs, J. Biocatalytic conversion of lignocellulose to platform chemicals. *Biotechnol. J.* **7**, 1122–1136 (2012).
- Howard, R. L., Abotsi, E., Jansen van Rensburg, E. L. & Howard, S. Lignocellulose biotechnology: issues of bioconversion and enzyme production. *Afr. J. Biotechnol.* **2**, 602–619 (2003).
- Lange, J. P. Lignocellulose conversion: an introduction to chemistry, process and economics. *Biofuels Bioprod. Bioref.* **1**, 39–48 (2007).
- Stephanopoulos, G. Challenges in engineering microbes for biofuels production. *Science* **315**, 801–804 (2007).
- Beguín, P. & Aubert, J. P. The biological degradation of cellulose. *FEMS Microbiol. Rev.* **13**, 25–58 (1994).
- Bhat, M. K. & Bhat, S. Cellulose degrading enzymes and their potential industrial applications. *Biotechnol. Adv.* **15**, 583–620 (1997).
- Dashtban, M., Maki, M., Leung, K. T., Mao, C. & Qin, W. Cellulase activities in biomass conversion: measurement methods and comparison. *Crit. Rev. Biotechnol.* **30**, 302–309 (2010).
- Dashtban, M., Schraft, H. & Qin, W. Fungal bioconversion of lignocellulosic residues; opportunities & perspectives. *Int. J. Biol. Sci.* **5**, 578–595 (2009).
- Maki, M., Leung, K. T. & Qin, W. The prospects of cellulase-producing bacteria for the bioconversion of lignocellulosic biomass. *Int. J. Biol. Sci.* **5**, 500–516 (2009).
- May, R. M. The dimensions of life on earth. *Nature and Human Society: the Quest for a Sustainable World* (ed. by Raven, P. H. & Williams, T.), pp. 30–45 (2000).
- Levine, M. How insects lose their limbs. *Nature* **415**, 848–849 (2002).
- Hongoh, Y. Diversity and genomes of uncultured microbial symbionts in the termite gut. *Biosci. Biotechnol. Biochem.* **74**, 1145–1151 (2010).
- Warnecke, F. *et al.* Metagenomic and functional analysis of hindgut microbiota of a wood-feeding higher termite. *Nature* **450**, 560–565 (2007).
- Tartar, A. *et al.* Parallel metatranscriptome analyses of host and symbiont gene expression in the gut of the termite *Reticulitermes flavipes*. *Biotechnol. Biofuels* **2**, 25 (2009).
- de Lorenzo, V. Problems with metagenomic screening. *Nat. Biotechnol.* **23**, 1045 (2005).
- Uchiyama, T. & Miyazaki, K. Functional metagenomics for enzyme discovery: challenges to efficient screening. *Curr. Opin. Biotechnol.* **20**, 616–622 (2009).
- Adlakha, N., Rajagopal, R., Kumar, S., Reddy, V. S. & Yazdani, S. S. Synthesis and characterization of chimeric proteins based on cellulase and xylanase from an insect gut bacterium. *Appl. Environ. Microbiol.* **77**, 4859–4866 (2011).
- Adlakha, N., Sawant, S., Anil, A., Lali, A. & Yazdani, S. S. Specific fusion of β -1,4-endoglucanase and β -1,4-glucosidase enhances cellulolytic activity and helps in channeling of intermediates. *Appl. Environ. Microbiol.* **78**, 7447–7454 (2012).
- Fontes, C. M. & Gilbert, H. J. Cellulosomes: highly efficient nanomachines designed to deconstruct plant cell wall complex carbohydrates. *Annu. Rev. Biochem.* **79**, 655–681 (2010).
- Wood, T. M., Wilson, C. A. & Stewart, C. S. Preparation of the cellulase from the cellulolytic anaerobic bacterium *Ruminococcus albus* and its release from the bacterial cell wall. *Biochem. J.* **205**, 129–137 (1982).
- Ma, L., Zhang, J., Zou, G., Wang, C. & Zhou, Z. Improvement of cellulase activity in *Trichoderma reesei* by heterologous expression of a beta-glucosidase gene from *Penicillium decumbens*. *Enzyme Microb. Technol.* **49**, 366–371 (2011).
- Kleman-Leyer, K. M., Siika-Aho, M., Teeri, T. T. & Kirk, T. K. The Cellulases Endoglucanase I and Cellobiohydrolase II of *Trichoderma reesei* Act Synergistically To Solubilize Native Cotton Cellulose but Not To Decrease Its Molecular Size. *Appl. Environ. Microbiol.* **62**, 2883–2887 (1996).
- Ng, T. K. & Zeikus, J. G. Purification and characterization of an endoglucanase (1,4-beta-D-glucan glucanohydrolase) from *Clostridium thermocellum*. *Biochem. J.* **199**, 341–350 (1981).
- Lali, A. M. *et al.* Method for preparation of fermentable sugars from biomass. patent application PCT/IN2010/000355 (2010).
- Saitou, N. & Nei, M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**, 406–425 (1987).
- Kimura, M. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* **16**, 111–20 (1980).
- Tamura, K. *et al.* MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28**, 2731–2739 (2011).
- Folmer, O., Black, M., Hoeh, W., Lutz, R. & Vrijenhoek, R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol. Mar. Biol. Biotechnol.* **3**, 294–299 (1994).
- Ausubel, F. M. *et al.* *Current Protocols in Molecular Biology*. New York City, NY: John Wiley & Sons Inc (1994).
- Brusca, R. C., Coelho, V. & Taiti, S. A guide to the coastal Isopods of California. Internet address: http://tolweb.org/notes/?note_id=3004.x (2001).
- Roonwal, M. L. & Chhotani, O. B. Termite fauna of Assam region, Eastern India. *Proc. Natn. Inst. Sci. India* **28**, 282–406 (1962).
- Segel, I. H. *Biochemical Calculations: How to Solve Mathematical Problems in General Biochemistry*, 2nd edn. New York: John Wiley (1976).

Acknowledgements

We are thankful to Dr. Arvind Lali for providing us alkali pretreated rice straw. We acknowledge financial support from the Department of Biotechnology of the Government of India.

Author contributions

S.S.Y., R.B. and N.A. designed the experiments; Z.B., V.K.K., N.A. and G.C. collected the arthropods; Z.B., V.K.K., N.A. and A.S. conducted the experiments; Z.B. and S.S.Y. performed the computational analysis and prepared the figures; Z.B., V.K.K. and S.S.Y. wrote the manuscript. All authors reviewed the manuscript.

Additional information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Bashir, Z. *et al.* Diversity and functional significance of cellulolytic microbes living in termite, pill-bug and stem-borer guts. *Sci. Rep.* **3**, 2558; DOI:10.1038/srep02558 (2013).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported license. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0>