OPEN

Evolution of NO₂ levels in Spain from 1996 to 2012

SUBJECT AREAS:

ATMOSPHERIC
CHEMISTRY

ENVIRONMENTAL MONITORING

Carlos A. Cuevas¹, Alberto Notario², José Antonio Adame³, Andreas Hilboll⁴, Andreas Richter⁴, John P. Burrows⁴ & Alfonso Saiz-Lopez¹

¹Atmospheric Chemistry and Climate Group, Institute of Physical Chemistry Rocasolano, CSIC, Madrid, Spain, ²Departamento de Química Física, Facultad de Ciencias Químicas, Universidad de Castilla-La Mancha, Ciudad Real, Spain, ³Atmospheric Sounding Station-El Arenosillo, National Institute for Aerospace Technology (INTA), Atmospheric Research and Instrumentation Branch, Mazagón-Huelva, Spain, ⁴Institute of Environmental Physics and Remote Sensing, University of Bremen, Bremen, Germany.

Received
16 April 2014Accepted
14 July 2014Published
30 July 2014

Correspondence and requests for materials should be addressed to C.A.C. (ccuevas@iqfr.csic.es)

We report on the evolution of tropospheric nitrogen dioxide (NO₂) over Spain, focusing on the densely populated cities of Barcelona, Bilbao, Madrid, Sevilla and Valencia, during 17 years, from 1996 to 2012. This data series combines observations from in-situ air quality monitoring networks and the satellite-based instruments GOME and SCIAMACHY. The results in these five cities show a smooth decrease in the NO₂ concentrations of ~2% per year in the period 1996–2008, due to the implementation of emissions control environmental legislation, and a more abrupt descend of ~7% per year from 2008 to 2012 as a consequence of the economic recession. In the whole Spanish territory the NO₂ levels have decreased by ~22% from 1996 to 2012. Statistical analysis of several economic indicators is used to investigate the different factors driving the NO₂ concentration trends over Spain during the last two decades.

Ozone, fine particulate matter (PM), nitrogen oxides (NO_x = NO + NO₂), and sulfur dioxide (SO₂) are the most important air pollutants in the troposphere, which represent a serious threat for ecosystems and human health¹. Although the sources of these pollutants in the European Union (EU) have considerably been reduced in the last ten years, the air quality is still far from the levels required by the EU legislation. The EU environmental policies requirements are currently being reviewed to be adapted to safer levels for human health².

NO_x are important in tropospheric chemistry for several reasons, including i) their role in the photochemical production of ozone³, ii) their atmospheric processing and subsequent deposition which leads to acidification and eutrophication in ecosystems through the formation of nitric acid (HNO₃)^{4,5}, iii) their role as precursors of the nitrate radical (NO₃) (which controls the oxidizing capacity of the nocturnal atmosphere^{6,7}), and iv) their effect on the radiative forcing of the atmosphere, directly through the absorption of radiation by NO₂, or indirectly through ozone formation and altering the lifetimes of several greenhouse gases^{8,9}. Although NO_x are also naturally produced by soil emissions, biomass burning and lightning, their main source is anthropogenic through the combustion of fossil fuels such as coal, oil (mainly in urban traffic) and gas. Hence, NO_x emissions in a country or region are directly related to its industrial and economic development, and are consequently affected by changes in the economic and industrial activity, in addition to the implementation of environmental policies^{10–12}.

Most of the NO_x emissions from combustion systems are in the form of NO, although in diesel vehicles NO₂ emissions can reach up to 70% of the emitted NO_x¹³. From 1996 to 2011, the gasoline/gas-oil percentage ratio in Spain has changed from 75/25 to 45/54 (1% corresponds to electric vehicles)¹⁴, therefore a relatively higher amount of NO_x emissions is currently produced in the form of NO₂. As sunlight produces an equilibrium between NO and NO₂ through a photochemical cycle involving ozone⁹, space-based observations of NO₂ can be used to locate sources of NO_x emissions due to the short lifetime of NO₂¹⁵. The first studies about the temporal evolution of tropospheric NO₂ were reported by Richter et al.¹⁶ for the 1996–2004 time period in the main industrialized and populated areas of the world, and Irie et al.¹⁷ for 1996–2002 focusing on East Asia. The first study reporting the effect of the economic downturn in the NO₂ levels was published by Lin and McElroy¹², showing a clear decrease in China from late 2008 to late 2009. Since these studies there have been other reports about the evolution of tropospheric NO₂ as seen from space^{18–20}. More recently, the global study on tropospheric NO₂ trends by Richter et al.¹⁶ has been updated for the 1996–2011 period by Hilboll et al.¹⁵. Previous reports of NO₂ trends have also highlighted a clear reduction in the NO₂ levels over urban areas due to both the application of anthropogenic NO_x emission control policies and the effects of the global economic crisis^{10,11,21}.

In this work we report on the NO₂ geographical distribution and concentration trends over Spain using combined multiple satellite datasets and ground-based observations from five metropolitan areas in the country: Madrid (3,233,527 inhabitants), Barcelona (1,620,943 inhabitants), Valencia (797,028 inhabitants), Sevilla (702,355 inhabitants) and Bilbao (351,629 inhabitants)²². Ground-based observations in León and Ponferrada are also reported, as these relatively small cities are located in the region where satellite measurements detect the higher decrease. We assess the contribution of environmental policies and the recent severe economic recession to the observed reduction in NO₂ levels over Spain in the last two decades. Finally, the variation of the main economic indicators of the country is compared with the measured NO₂ levels, to infer the possible relationship between the NO_x emissions and the economic evolution of Spain.

Results

Satellite-based NO₂ observations and trends. Figure 1 shows averaged tropospheric NO₂ Vertical Column Density (VCD) maps for 1996 and 2011 over the Iberian Peninsula, with data from the GOME and SCIAMACHY instruments, respectively. Due to the different ground pixel size between these different sensors, a resolution correction method is applied to the low spatial resolution data from GOME¹⁵ (see Methods). This procedure allows building a time-consistent NO₂ dataset from 1996 to 2012. The figure includes annual, winter (Dec-Jan-Feb), and summer (Jun-Jul-Aug) NO₂ column averages for the first and last years of the time series. Data from the GOME resolution corrected data product have been used for 1996. The plot for the last year of the series is made from 2011 SCIAMACHY measurements, as the operative cycle of this instrument ended in April 2012. Figure 1 clearly illustrates the difference between concentrations and geographical distribution of tropospheric NO₂ in 1996 and 2011. The seasonality of NO₂ values is also evident for both years, showing higher concentrations in winter that contrast with summer values. As expected in mid-latitude developed countries, the lower solar radiation levels and the higher energy consumption, due to building heating systems, produce higher values of tropospheric NO₂ in the winter season²³.

The highest levels of NO₂ are observed in the regions of Madrid and Cataluña, the two largest urban areas of Spain. Madrid's economy is based on the service, construction and industry sectors. Its location in the centre of the country also makes Madrid the main transport knot within the Iberian Peninsula. Cataluña is located in the north-east coast of Spain, and its economy is mainly based on service, tourism and industry. Most of the economic activity in these two regions is located close to the metropolitan areas of the cities of Barcelona and Madrid.

The north and east coastal regions of the peninsula are also areas with considerable industrial activity, and is therefore reflected in high NO₂ levels (Fig. 1). The north-west part of the peninsula (Asturias, León and Palencia provinces, see Figure S1 for their location) is an important region of mining and power generation industries. Also, considerable maritime transport, mining and steel industries activities take place in the north coast area of the Gulf of Vizcaya, close to the city of Bilbao. At the east coast, most of the economic activity is located in the Valencia region and the Ebro basin, whereas the Guadalquivir basin concentrates most of the economic and industrial activity in the south-west part of Spain, where cities such as Cordoba, Sevilla and Huelva are located.

The reduction in the NO₂ levels over recent times is evident when comparing the first and last years of the satellite dataset (Fig. 1). The decrease is clear in the annual and summer averages, although the change in winter is less pronounced. Both annual and summer averages show a marked reduction in Madrid, Cataluña, and the north-west part of the peninsula. In winter the drop is less apparent, but still appreciable in Madrid, east coast and north-west areas. Note

that the 2011 winter was particularly cold²⁴ and thus could be responsible for a higher use of heating systems, incrementing NO_x emissions during that winter. The more widespread NO₂ distributions over the regions of Madrid and Cataluña in 1996 are reduced and limited by 2011 to the metropolitan and localised industrial areas around the cities of Madrid and Barcelona. The decrease in NO₂ levels and its geographical spread is also clear along the north and east coasts, and Ebro and Guadalquivir basins (Fig. 1).

To study the time evolution of NO₂ levels over the Iberian peninsula, the trend model described in Hilboll et al.¹⁵ has been applied here to the NO₂ measurements from the GOME and SCIAMACHY instruments. Figure 2 shows the integrated evolution of tropospheric NO₂ levels during the 1996–2011 time period, including absolute and relative changes. Both plots illustrate the decrease in NO₂ concentrations over a large part of the peninsula, and particularly in the main economic and industrial areas. NO₂ levels in the area of Madrid have decreased on an annual average by up to 1.9×10^{14} molecules cm⁻², which adds up to 53% of the levels measured in 1996. This reduction is much higher than the 20% previously reported for a different period (2004–2010) using OMI measurements by Castellanos and Boersma, 2012¹⁰. This difference is due to the different time period considered and also note the difference in local time at which different instruments overpass Spain. The measurement times for GOME and SCIAMACHY are ~10:30 and ~10:00 local time, while OMI is ~13:45. GOME and SCIAMACHY measurements are closer to the diurnal NO₂ peak in the early morning, when NO_x emissions are more affected by transport sources at the rush hour. Emissions from industry and power plants have a less pronounced diurnal variation; consequently lower NO₂ VCDs are expected from the OMI measurements resulting in lower relative decreases. These differences in the SCIAMACHY product compared to OMI have been previously studied^{25–27}, reporting higher NO₂ levels from SCIAMACHY in the northern midlatitudes and regions where the NO_x emissions come mainly from fossil fuel sources. Other reasons are the different pixel size of OMI and the differences in the instruments and retrieval algorithm. Considerable reductions of ~40, ~36, and ~51% are measured in the areas of Barcelona, Bilbao and Valencia, respectively. In Sevilla, however, a smaller decrease of 27% is observed.

The most widespread and remarkable reduction in NO₂ levels is detected in the northwest industrial area and the north coast, with the highest decreases in the Asturias region and the provinces of León and Palencia (Fig. 2). These regions have experienced a large reduction in the industrial, mining and coal power plants activities, which together with the application of new emission control technologies have resulted in a drastic reduction in NO₂ concentrations. The decline in mining and industrial activities in these regions from 2004 to 2009, have led to a fall in the NO_x emissions of ~52% from the main power plant stations of the region²⁸. A summary of the satellite NO₂ trends by geographical areas is given in Table 1, including absolute and relative changes.

Recently, Castellanos and Boersma¹⁰ for Europe and Vrekoussis et al.²¹ for Greece, have also reported a general negative trend in NO₂ concentrations accelerated since the beginning of the economic recession (i.e., 2008). The trend model can also be applied to the pre-recession and recession time periods, although the nonlinearities in the general trend and the short length of this individual periods yield few significant data. Nevertheless the purpose of this work is to show the evolution of the NO₂ levels rather than specifically analyse the short time period trends. Overall, from 1996 to 2011, the integrated NO₂ VCD as observed from space over the entire Spanish territory has been reduced by ~22%.

NO₂ trends as measured by *in-situ* air quality networks. To quantify the variation of surface NO₂ levels in the cities of Barcelona, Bilbao, Madrid, Sevilla and Valencia, data from their ground air quality monitoring networks (AQMN) are used. All the *in-situ*

Figure 1 | Averaged tropospheric NO_2 VCD for different seasons during 1996, using the GOME resolution-corrected dataset (a, c and e), and 2011/2012 using SCIAMACHY data (b, d, f). Top: annual averages. Middle: summer (Jun-Jul-Aug) averages. Bottom: winter (Dec-Jan-Feb) averages. Figure created by the authors using Matlab R213a.

monitoring stations employ the chemiluminescence method described in the Methods section. Figure 3 shows the annually-averaged NO_2 levels from 1996 to 2012 measured by the available AQMN. All cities show a general decrease in the NO_2 levels during the last two decades. Regression analysis yields considerable absolute

reductions of 67% for Valencia and 52% in Sevilla, while Barcelona exhibits a shallower 18% decrease. Madrid and Bilbao show intermediate decreases of 37% and 35%, respectively. Overall, the average of all cities considered here shows a decrease of 44% in the NO_2 concentrations.

Figure 2 | Change in the NO_2 tropospheric VCD over the Spanish territory derived from the shift trend model for the years 1996–2011 (data with significant trends at the 95% level). Figure created by the authors using Matlab R213a.

The results of the statistical regression analysis performed for different cities and time periods are shown in Table 2, including absolute and relative changes. This summary also includes the combined average of all studied cities. Barcelona, Madrid, and Valencia show two well differentiated slopes for the pre-recession and recession periods (-0.5 , -0.7 , and $-3.1/\mu\text{g m}^{-3} \text{ yr}^{-1}$ in Barcelona, Madrid and Valencia, respectively, during the pre-recession period, and -0.9 , -4.3 , and $-3.9/\mu\text{g m}^{-3} \text{ yr}^{-1}$, respectively, during the recession period). In these three cases the slopes are higher during the recession period. These slopes correspond to annual decreases of 0.9%, 1.1% and 4.3 for Barcelona, Madrid and Valencia, respectively, during the pre-recession period, and 1.9%, 7.8% and 9.1%, respectively, during the economic recession. Sevilla shows similar slopes for both periods, being slightly higher during the pre-recession period ($-1.7/\mu\text{g m}^{-3} \text{ yr}^{-1}$) than during the economic recession ($-1.5/\mu\text{g m}^{-3} \text{ yr}^{-1}$). By contrast with the other cities, Bilbao shows a higher slope during the pre-recession period than during the economic recession (-1.4 and $-0.8/\mu\text{g m}^{-3} \text{ yr}^{-1}$ respectively). Overall, all cities except Bilbao, show higher reductions during the recession period than during the pre-recession. This result suggests an initial reduction driven by the progressive implementation of environmental policies and technology, whereas the second slope can be attributed to the rapid decrease of the economic and industrial activity during the economic downfall between 2008 and 2012.

In order to confirm the pronounced decline in NO_2 levels reported by the satellite measurements in the North-West region, AQMN data in the cities of Ponferrada and León (province of León) were also

investigated. These data show a decrease of 80% in Ponferrada and 66% in León since 1997. Both cities also show a higher decrease during the economic recession period, 72% and 75%, compared to 63% and 41% during the pre-recession period for Ponferrada and León, respectively.

The general annually-averaged NO_2 trend from both satellite and ground stations is shown in Figure 4. The satellite averages have been made integrating over the entire Spanish territory in the Iberian Peninsula, while the ground stations data correspond to yearly average in the studied cities. Hence the smoother shape in the AQMN dataset, which is more affected by local effects than the satellite measurements. Note that both datasets are not directly comparable, although their combination provides a clear picture about the evolution of NO_2 levels in Spain during the last two decades.

Both datasets show two distinct slopes corresponding to the pre-recession and recession periods, the slope for the last period being higher. The purpose of this work is to show how the NO_2 levels have changed in absolute terms during the last two decades, rather than to perform a statistical trend study. Therefore we refer here to ‘general trends’, despite the evident variable nonlinear trends observed in the time series. The nonlinearity in the NO_x emission trends have already been reported by Konovalov et al.²⁹ for the case of Madrid.

Discussion

In order to gain insights about how the economic activity affects NO_x emissions in Spain, the levels of NO_2 are compared to the main economic indicators, including Gross Domestic Product (GDP),

Table 1 | Absolute and relative changes in the NO_2 tropospheric VCD from 1996–2011, derived from the trend model for different geographic regions. (See Table S1 for latitude-longitude coordinates of the studied regions)

Geographic region	Absolute significant changes/ molecule cm^{-2}	Error/ $\pm \text{molecule cm}^{-2}$	Relative significant changes/%	Error/ $\pm \%$
East coast of Valencia	$-1,27\text{E} + 15$	$3,60\text{E} + 14$	-33	9
Ebro Basin and Cataluña	$-1,34\text{E} + 15$	$4,97\text{E} + 14$	-23	9
Area of Madrid	$-3,10\text{E} + 15$	$9,48\text{E} + 14$	-53	16
Area of Sevilla	$-1,16\text{E} + 15$	$5,33\text{E} + 14$	-27	12
Area of Barcelona	$-2,98\text{E} + 15$	$9,89\text{E} + 14$	-40	13
Area of Bilbao	$-2,25\text{E} + 15$	$7,29\text{E} + 14$	-36	12
North-West	$-2,87\text{E} + 15$	$9,22\text{E} + 14$	-46	15
Gulf of Vizcaya	$-1,34\text{E} + 15$	$6,24\text{E} + 14$	-24	11
Guadalquivir basin	$-6,36\text{E} + 14$	$3,86\text{E} + 14$	-17	10
Area of Valencia	$-2,21\text{E} + 15$	$5,89\text{E} + 14$	-51	13
Integrated over Spain	$-9,47\text{E} + 14$	$3,55\text{E} + 14$	-22	8

Figure 3 | NO₂ annual averages from air quality monitoring stations.

Industrial Production Index (IPI, base 2005) and Oil Consumption. The GDP is the market value of all officially recognized final goods and services produced within a country in a given period of time. The IPI measures the real production output of manufacturing, mining and utilities of a country. The Oil Consumption refers to the number of barrels consumed in a country per day. NO_x natural emissions account for 2.8–3.8% of total emissions in Spain (National Emission Inventory³⁰) during the studied period. Therefore the trends in economy and industrial activities can be considered the main source of variation for the NO_x emissions in Spain. Bottom-up emissions data from the National Emission Inventory³⁰ and the Centre on Emissions Inventories and Projections³¹ are shown in Figures S2 and S3. Note however that the trends in these emission inventories, which follow the trends in the economic parameters of the country, show a discrepancy with both on-ground and satellite data (see Figs. 5, S2 and S3). This highlights potential uncertainties in the trends of emission inventories and suggests that further research is needed to reconcile differences in the trends of observed NO₂ and the emission inventories during the pre-recession period in Spain.

Although Spain has failed to achieve the Gothenburg protocol targets, it has reduced the NO_x emissions by about 20% from 1990 to 2011, as reported by the European Environment Agency³². This fact is reflected in the NO₂ measurements through a reduction by 22% in the NO₂ levels integrated over Spain as observed from space, and by an average of 45% in the AQMN observations in the five cities studied here. This reduction is considerably higher in the main industrial and economic cores of the country, such as Madrid and the north-west part of the Iberian Peninsula, with reductions of up to 50% in the areas of Madrid and Valencia. In general, there is a correspondence between the tendencies of on-ground and satellite

data (Figure 4), having two well differentiated slopes in the trends (i.e. pre-recession and economic recession periods).

On the other hand, the main economic indicators were gradually increasing since 1996 until the beginning of the economic crisis in 2008, when all indicators suddenly fell. These economic metrics refer to the whole country, so satellite NO₂ VCDs integrated over the whole Spanish territory in the Iberian Peninsula are used to assess the correlation between economic fall and NO₂ concentrations. In general we find a good correspondence between the economic indicators and the evolution of NO₂, as shown in Figure 5, despite the 2001–2007 period in which the trends are different. The NO₂ increase from 1996 to 2000 coincides with a period of economic growth marked by a huge increase of the IPI and a small, but appreciable, increase in the GDP. During this first period it can be considered that the economic rise is directly reflected in the NO₂ levels. From 2000 onwards the IPI slows down its rising while the NO₂ VCDs fall from 4.3×10^{15} to 3.6×10^{15} molecules cm⁻², and the GDP undergoes a huge and constant growth until the start of the recession in 2008. Consequently, during the period in which the economic activity of the country is rising, the application of environmental policies and the optimization in the combustion processes are likely responsible for the reduction in the emissions of NO_x measured from 2000 onwards. Finally during the recession (2008–2012) both indicators and the NO₂ levels decrease considerably, showing a clear effect of the economy on the levels of anthropogenic pollutants.

The same information can be extracted from the oil consumption evolution in Spain (Figure 5). From 1996 to 2008 it has been growing at constant rate, starting a deep decrease in 2008 with the recession. Note that the 33% increase in the oil consumption from 1996 to 2008 should have led to an increase in the NO₂ levels during this time period. This indicates the positive effect of emissions legislation upon combustion technologies.

Our results show a lack of correlation between NO₂ levels and economic indicators from 2001 to 2007, however Vrekousis et al.²¹ reported good agreement between NO₂ and economic markers in Greece from 2000 to 2011. This difference may be due to the fact that Greece has not reduced its NO_x emissions, as a result of environmental legislation, at the same rate as Spain³³, and hence the better correspondence between economic evolution and NO_x emissions in Greece. This reduction in the NO₂ levels in Spain during the 2001–2007 period, when all the economic factors increase, is a clear signal that the application of environmental policies are efficiently reducing the NO_x emissions. This has also been reported for other cities of the European Union as shown by Castellanos and Boersma¹⁰.

To summarize, the datasets presented here can be divided into two differentiated parts: i) from 1996 to 2008 the application of environmental policies reduced the NO_x emissions, with a slight increase from 1996 until 2000 before the effects of environmental policies start to pay off; ii) during the 2008–2012 recession period the eco-

Table 2 | Regression analysis of NO₂ trends as measured by air quality monitoring stations. Slopes in Table S2

City	1996–2012		1996–2008		2008–2012	
	Decrease/% yr ⁻¹	Absolute decrease/%	Decrease/% yr ⁻¹	Absolute decrease/%	Decrease % yr ⁻¹	Absolute decrease/%
Barcelona	−1.16*	−18.60	−0.92	−11.99	−1.89*	−7.56
Bilbao	−2.19*	−35.02	−2.89	−37.59	−2.27*	−9.08
Madrid	−2.20	−37.45	−1.11	−14.42	−7.84	−39.18
Sevilla	−3.28**	−52.43	−3.64**	−43.66	−4.77	−23.87
Valencia	−4.47	−67.09	−4.32	−47.50	−9.13	−45.66
Average***	−2.64	−44.93	−2.23	−29.02	−6.86	−34.29

*No data available for 2012.

**No data available for 1996.

***Average of the five cities

Figure 4 | NO_2 trends in satellite- and ground-based observations during the pre-recession and economic recession periods.

economic crisis remarkably reduced the emission even more than the environmental policies. These trends are observable through both datasets, although the absolute changes are higher in the ground-based measurements, showing decreases up to 67% in the city of Valencia. These differences between the two datasets are, as explained in the Methods section, due to the different type of measurements considered. The ground stations represent surface observations within the cities, while the satellite measurements result from integrating the tropospheric column over the whole ground pixel of the instrument, which is larger than the largest of the studied cities. Thus we regard the ground measurements as representative of the local NO_2 levels and trends in the cities, while the satellite-based observations represent column-integrated NO_2 levels over larger areas around the cities. The 20% emissions reduction reported by the European Union agree better with the satellite dataset, as these emissions reductions refers to the whole country rather than to specific cities or regions.

The lack of correlation between the economic parameters of Spain and the levels of NO_2 during the 2001–2007 period, with NO_2 levels decreasing as the GDP, IP and Oil Consumption grow until 2008, shows that the improvement in the combustion technologies and environmental policies implemented to reduce the nitrogen oxides emissions are paying off. Therefore the air quality is improving, although there is still margin for further improvement as demonstrated by the fact that the daily thresholds of NO_2 and O_3 levels are often exceeded in Spanish cities. The steeper slope in the NO_2 reduction during the recession, when the economy falls, is clear in both datasets. Hence for the case of Spain, still on recession during 2012, it can be expected that the slope of the decrease will recover to the pre-recession values once the economy starts growing again. Finally, this decline in the NO_2 levels during the last two decades has led to an overall improvement of the air quality in the Iberian Peninsula, and is likely affecting the chemistry of the troposphere in three possible ways: 1) with the reduction in the NO_x levels, a variation in the

Figure 5 | Satellite tropospheric NO_2 VCD averaged over the Spanish territory, along with the main economic metrics of Spain, Gross Domestic Product (GDP), Industrial Production Index (IPI) and Oil Consumption.

tropospheric ozone levels is also expected, 2) this denoxification of the atmosphere has probably led to lower nitrogen inputs to ecosystems, and 3) the levels and seasonality of HNO_3 formation as influenced by traffic emissions³⁴ have likely been affected.

Finally, it is worth mentioning that future monitoring of air pollution in megacities and industrial regions will greatly benefit from geostationary satellite missions, which will considerably enhance the temporal and spatial resolution of the observations. Therefore it will provide a more accurate tool to verify the effect of implementing environmental policies and to follow the atmospheric impacts of changing emissions.

Methods

Description of datasets. Two main datasets including space-based observations and air quality monitoring networks (AQMN) measurements are used to study the evolution of tropospheric NO_2 over Spain. The satellite observations provide an overview of the NO_2 geographical distribution and variation not only over major urban areas, but also over the entire Iberian Peninsula, while *in-situ* ground monitoring stations provide insightful information of the pollutant levels evolution within cities. Note that the spatial resolution of current satellite-based spectrometers is still too low to properly assess detailed NO_x emissions at a sub-urban scale or from medium-size cities. On the other hand AQMN provide information of pollutants at surface level, while satellite based measurements integrate the NO_2 vertical column throughout the troposphere (once the stratospheric contribution has been removed). Therefore, both types of measurements are not directly comparable but complementary.

The satellite dataset is based on measurements from the GOME and SCIAMACHY instruments, using the trend model developed by Hilboll et al.¹⁵. This model accounts for the ground pixel size and instrument differences between GOME and SCIAMACHY, as described below in this section. The *in-situ* dataset has been built using data from the AQMN in each of the cities included in this exercise. Data from a total number of ~55 monitoring stations of urban and sub-urban type have been used during the period of study, with different typology and locations representative of the major emission points within each individual city. These AQMN are operated and maintained by the regional governments of the country, according to current European legislation.

Surface measurements. The method used to detect NO_2 in the monitoring station is described in EN 14211:2005 'Ambient air quality – Standard method for the measurement of the concentration of nitrogen dioxide and nitrogen monoxide by chemiluminescence, defined by the European Directive (2008/50/CE) as reference method. In this method a pre-reactor separates the NO - NO_2 reaction from the background chemiluminescence, allowing accurate auto-zeroing of the analyzer. The equipment works under a rigid maintenance program being periodically tested and calibrated.

Satellite measurements. In this work measurements performed by the satellite based instruments GOME³⁵ on board ERS-2, and SCIAMACHY^{36,37} on board ENVISAT are employed. The ground pixel of these instruments is $40 \times 320 \text{ km}^2$ and $30 \times 60 \text{ km}^2$ for GOME and SCIAMACHY, respectively. The raw data have been analysed as described in Hilboll et al., 2013a¹⁵. In short, differential optical absorption spectroscopy³⁸ in the 425–450 nm wavelength window³⁹ is applied to the measured radiances. The influence of stratospheric NO_2 has been corrected for using the algorithm detailed in Hilboll et al. 2013⁴⁰, using stratospheric NO_2 fields from the B3dCTM model^{41–43}. Tropospheric air mass factors have been calculated with the radiative transfer model SCIATRAN⁴⁴.

Correction for the difference in ground pixel size between GOME and SCIAMACHY. To show the spatial distribution of tropospheric NO_2 columns (see Fig. 1) from GOME measurements, the retrieved data have been post-processed to account for the very low spatial resolution of the GOME instrument. The method is based on spatial averaging of earthshine spectra measured by SCIAMACHY and the extraction of a spatial pattern of the effect of ground pixel size. This effect of ground pixel size is subsequently super-imposed on the original GOME measurements, yielding a data set "GOME res.corr". In this data set, the magnitude of the tropospheric NO_2 columns is retrieved from GOME measurements, while the spatial structure is derived from the full SCIAMACHY data set 2003–2011. The method is described in Section 4 of Hilboll et al., 2013¹⁵.

Trend model accounting for the difference between GOME and SCIAMACHY measurements. The rate of linear change of tropospheric NO_2 columns has been calculated using the algorithm described in Section 5 of Hilboll et al., 2013¹⁵. In short, the trend model explicitly accounts for (a) an offset between the two instruments and (b) for a change in the amplitude of the seasonal variation. The time series of monthly averages of NO_2 measurements $Y(t)$ is described by the equation

$$Y(t) = \mu + \omega t + \delta U(t) + S(t) + N(t) \quad (1)$$

where μ is the VCD_{trop} measurement at time $t = 0$, ω is the monthly trend

component, and t is the time in months since January 1996. δ is the level shift between GOME and SCIAMACHY measurements occurring at time $t = T_0$ (which we set to January 2003), and $U(t)$ is the step function

$$U(t) = \begin{cases} 0, & t < T_0 \\ 1, & t \geq T_0 \end{cases} \quad (2)$$

The seasonal component $S(t)$ has the same shape (represented by harmonic functions) and varying amplitude for the two instruments, and is modelled by four sine and cosine components, each. The noise process $N(t)$ is assumed to be autoregressive with lag 1, and only trends which are significant at a 95% level are considered.

- Solomon, P. A., Wexler, A. S. & Sioutas, C. Special Issue of Atmospheric Environment for Air Pollution and Health: Bridging the Gap from Sources-to-Health Outcomes. *Atmos. Environ.* **45**, 7537–7539 (2011).
- World Health Organization. Review of evidence on health aspects of air pollution – REVIHAAP Project. (<http://www.euro.who.int/data/assets/pdf/0020/182432/e96762-final.pdf>). (2013, Date of Access: 16/04/2014).
- Neuman, J. A. et al. Relationship between photochemical ozone production and NO_x oxidation in Houston, Texas. *J. Geophys. Res., [Atmos.]* **114**, D00F08 (2009).
- Vrekoussis, M. et al. Formation of HNO_3 and NO_3 – in the anthropogenically-influenced eastern Mediterranean marine boundary layer. *Geophys. Res. Lett.* **33**, L05811 (2006).
- Liu, X. et al. Enhanced nitrogen deposition over China. *Nature* **494**, 459–462 (2013).
- Platt, U., Perner, D., Schröder, J., Kessler, C. & Toennissen, A. The diurnal variation of NO_3 . *J. Geophys. Res., [Oceans]* **86**, 11965–11970 (1981).
- Heintz, F., Platt, U., Flentje, H. & Dubois, R. Long-term observation of nitrate radicals at the Tor Station, Kap Arkona (Rügen). *J. Geophys. Res., [Atmos.]* **101**, 22891–22910 (1996).
- Seinfeld, J. H. & Pandis, S. N. Atmospheric chemistry and Physics - From Air Pollution to Climate Change -2nd Edition. *John Wiley & Sons* (2006).
- Solomon, S. et al. On the role of nitrogen dioxide in the absorption of solar radiation. *J. Geophys. Res., [Atmos.]* **104**, 12047–12058 (1999).
- Castellanos, P. & Boersma, K. F. Reduction in nitrogen oxides over Europe driven by environmental policy and economic recession. *Sci. Rep.* **2**, 265, doi:10.1038/srep00265 (2012).
- Russell, A. R., Valin, L. C. & Cohen, C. Trends in OMI NO_2 observations over the United States: effect of emission control technology and the economic recession. *Atmos. Chem. and Phys.* **12**, 12197–12209 (2012).
- Lin, J. T. & McElroy, M. B. Detection from space of a reduction in anthropogenic emissions of nitrogen oxides during the Chinese economic downturn. *Atmos. Chem. Phys.* **11**, 8171–8188, doi:10.5194/acp-11-8171-2011 (2011).
- Grice, S. et al. Recent trends and projections of primary NO_2 emissions in Europe. *Atmos. Environ.* **43**, 2154–2167 (2009).
- Dirección General de Tráfico DGT. Series históricas parque de vehículos 1997–2011. (www.dgt.es). (2011, Date of access: 16/04/2014).
- Hilboll, A., Richter, A. & Burrows, J. P. Long-term changes of tropospheric NO_2 over megacities derived from multiple satellite instruments. *Atmos. Chem. Phys.* **13**, 4145–4169 (2013).
- Richter, A., Burrows, J. P., Nuss, H., Granier, C. & Niemeier, U. Increase in tropospheric nitrogen dioxide over China observed from space. *Nature* **437**, 129–132 (2005).
- Irie, H. et al. Evaluation of long-term tropospheric NO_2 data obtained by GOME over East Asia in 1996–2002. *Geophys. Res. Lett.* **32**, L11810 (2005).
- Konovalov, I. B., Beekmann, M., Richter, A. & Burrows, J. P. Inverse modelling of the spatial distribution of NO_x emissions on a continental scale using satellite data. *Atmos. Chem. Phys.* **6**, 1747–1770 (2006).
- van der A, R. J. et al. Trends, seasonal variability and dominant NO_x source derived from a ten year record of NO_2 measured from space. *J. Geophys. Res., [Atmos.]* **113** (2008).
- Konovalov, I. B., Beekmann, M., Burrows, J. P. & Richter, A. Satellite measurement based estimates of decadal changes in European nitrogen oxides emission. *Atmos. Chem. Phys.* **8**, 2623–2641 (2008).
- Vrekoussis, M., Richter, A., Hilboll, A., Burrows, J. P. & Gerasopoulos, E. Economic crisis detected from space: Air quality observations over Athens/Greece. *Geophys. Res. Lett.* **40**, 458–463 (2013).
- Instituto Nacional de Estadística de España (www.ine.es). (2012, Date of access: 16/04/2014).
- Beirle, S., Platt, U., Wenig, M. & Wagner, T. Weekly cycle of NO_2 by GOME measurements: a signature of anthropogenic sources. *Atmos. Chem. Phys.* **3**, 2225–2232 (2003).
- AEMET. Resúmenes climatológicos de España. <http://www.aemet.es/es/serviciosclimaticos/vigilanciaclima/resumenes> (2012, Date of access: 16/04/2014).
- Boersma, K. F. et al. Validation of urban NO_2 concentrations and their diurnal and seasonal variations observed from the SCIAMACHY and OMI sensors using in situ surface measurements in Israeli cities. *Atmos. Chem. Phys.* **9**, 3867–3879 (2009).
- Boersma, K. F. et al. Intercomparison of SCIAMACHY and OMI tropospheric NO_2 columns: Observing the diurnal evolution of chemistry and emissions from

- space. *Journal of Geophysical Research: Atmospheres* **113**, D16S26, doi:10.1029/2007jd008816 (2008).
27. Lin, J. T., McElroy, M. B. & Boersma, K. F. Constraint of anthropogenic NO_x emissions in China from different sectors: a new methodology using multiple satellite retrievals. *Atmos. Chem. Phys.* **10**, 63–78, doi:10.5194/acp-10-63-2010 (2010).
 28. Ministerio de Agricultura, Alimentación y Medio Ambiente de España. Registro Estatal de Emisiones y Fuentes Contaminantes (www.prtr-es.es/). (2009, Date of access: 16/04/2014).
 29. Konovalov, I. B., Beekmann, M., Richter, A., Burrows, J. P. & Hilboll, A. Multi-annual changes of NO_x emissions in megacity regions: nonlinear trend analysis of satellite measurement based estimates. *Atmos. Chem. Phys.* **10**, 8481–8498 (2010).
 30. MAGRAMA. Inventario Nacional de Emisiones de Contaminantes a la Atmósfera. Ministerio de Agricultura, Alimentación y Medio Ambiente del Gobierno de España (<http://www.magrama.gob.es/es/calidad-y-evaluacion-ambiental/temas/sistema-espanol-de-inventario-sei/>). (2014, Date of access: 6/06/2014).
 31. CEIP. Centre on Emission Inventories and Projections. (<http://www.ceip.at/>). (2014, Date of access: 8/07/2014).
 32. European Environment Agency - Nitrogen oxides (NO_x) emissions (APE 002) - Assessment published Jan 2014 (<http://www.eea.europa.eu/data-and-maps/indicators/eea-32-nitrogen-oxides-nox-emissions-1/assessment.2010-08-19.0140149032-3>). (2014, Date of access: 16/04/2014).
 33. European Environment Agency. Air quality in Europe - 2013 report (<http://www.eea.europa.eu/publications/air-quality-in-europe-2013>). (2013, Date of access: 16/04/2014).
 34. Saiz-Lopez, A., Notario, A., Albaladejo, J. & McFiggans, G. Seasonal Variation of NO_x Loss Processes Coupled to the HNO₃ Formation in a Daytime Urban Atmosphere: A Model Study. *Water Air Soil Poll.* **182**, 197–206 (2007).
 35. Burrows, J. P. *et al.* The Global Ozone Monitoring Experiment (GOME): mission concept and first scientific results. *J. Atmos. Sci.* **56**, 151–175 (1999).
 36. Bovensmann, H. *et al.* SCIAMACHY: mission objectives and measurements modes. *J. Atmos. Sci.* **56**, 127–150 (1999).
 37. Burrows, J. P., Hölzle, E., Goede, A., Visser, H. & Fricke, W. SCIAMACHY - scanning imaging absorption spectrometer for atmospheric chartography. *Acta Astronaut.* **35**, 445–451 (1995).
 38. Platt, U. & Stutz, J. *Differential Optical Absorption Spectroscopy: Principles and Applications*. (Springer, Berlin, 2008).
 39. Richter, A. & Burrows, J. P. Tropospheric NO₂ from GOME measurements. *Adv. Space Res.* **29**, 1673–1683 (2002).
 40. Hilboll, A. *et al.* Improvements to the retrieval of tropospheric NO₂ from satellite - stratospheric correction using SCIAMACHY limb/nadir matching and comparison to Oslo CTM2 simulations. *Atmos. Meas. Tech.* **6**, 565–584 (2013).
 41. Sinnhuber, B.-M., Weber, M., Amankwah, A. & Burrows, J. P. Total ozone during the unusual Antarctic winter of 2002. *Geophys. Res. Lett.* **30**, 1580 (2003).
 42. Sinnhuber, M. *et al.* A model study of the impact of magnetic field structure on atmospheric composition during solar proton events. *Geophys. Res. Lett.* **30**, 1818 (2003).
 43. Winkler, H. *et al.* Modeling impacts of geomagnetic field variations on middle atmospheric ozone responses to solar proton events on long timescales. *J. Geophys. Res., [Atmos.]* **113**, D02302 (2008).
 44. Rozanov, A., Rozanov, V., Buchwitz, M., Kokhanovsky, A. & Burrows, J. P. SCIATRAN 2.0 - A new radiative transfer model for geophysical applications in the 175–2400 nm spectral region. *Adv. Space Res.* **36**, 1015–1019 (2005).

Acknowledgments

This work was funded by the Consejo Superior de Investigaciones Científicas (CSIC). The authors are grateful to the Department of Environment of the Regional Government of Madrid, Sevilla and Valencia, and the Spanish Ministry of Agriculture, Food and Environment for providing data from the air quality monitoring networks. A. Hilboll gratefully acknowledges funding by the “Earth System Science Research School (ESSReS)”, an initiative of the Helmholtz Association of German research centres (HGF) at the Alfred Wegener Institute for Polar and Marine Research. A. Hilboll, A. Richter, and J. P. Burrows would like to acknowledge the European Union Seventh Framework Programme (FP7/2007–2013) project CityZen (Grant Agreement no. 212095).

Author contributions

C.A.C. and A.S.-L. designed research; C.A.C., A.S.-L., A.H. and A.R. performed research; C.A.C., A.N., J.A.A., A.H., A.R. and J.P.B. analyzed data; C.A.C., A.S.-L. and A.H. wrote the manuscripts and supplementary information, and all authors reviewed the paper.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Cuevas, C.A. *et al.* Evolution of NO₂ levels in Spain from 1996 to 2012. *Sci. Rep.* **4**, 5887; DOI:10.1038/srep05887 (2014).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>