



## OPEN

## Activity of the novel fungicide SYP-Z048 against plant pathogens

SUBJECT AREAS:  
CHEMICAL BIOLOGY  
MICROBIOLOGYFengping Chen<sup>1,2</sup>, Ping Han<sup>2,3</sup>, Pengfei Liu<sup>2</sup>, Naiguo Si<sup>4</sup>, Junli Liu<sup>4</sup> & Xili Liu<sup>2</sup>

<sup>1</sup>Key Laboratory of Plant Virology of Fujian Province, Institute of Plant Virology, Fujian Agriculture and Forestry University, Fuzhou, 350002, China, <sup>2</sup>Department of Plant Pathology, China Agricultural University, Beijing 100193, China, <sup>3</sup>Beijing Research Center for Agricultural Standards and Testing, Beijing 100097, China, <sup>4</sup>State Key Laboratory of the Discovery and Development of Novel Pesticide, China ShenYang Research Institute of Chemical Industry, Shenyang 110021, China.

Received  
2 April 2014Accepted  
19 August 2014Published  
25 September 2014Correspondence and  
requests for materials  
should be addressed to  
X.L.L. (seedling@cau.  
edu.cn)

**In *in vitro* tests with 18 plant pathogens, the fungicide 3-[5-(4-chlorophenyl)-2,3-dimethyl-3-isoxazolidinyl]pyridine (SYP-Z048) was highly effective on inhibiting mycelial growth of various ascomycota and basidiomycota, with EC<sub>50</sub> values ranging from 0.008 to 1.140 µg/ml. SYP-Z048 had much weaker activity against growth of oomycota with EC<sub>50</sub> values > 100 µg/ml. In a second *in vitro* test with *Monilinia fructicola* isolates, SYP-Z048 inhibited mycelial growth (EC<sub>50</sub> = 0.013 µg/ml), germ tube elongation (EC<sub>50</sub> = 0.007 µg/ml), and sporulation but did not affect spore germination. In a detached pear fruit assay inoculated with *M. fructicola* isolates, SYP-Z048 showed protective and curative activity. Field tests indicated that SYP-Z048 was an efficacious fungicide for control of brown rot disease in two peach orchards. When applied to a single spot on a tomato leaflet in a compound leaf, SYP-Z048 suppressed the growth of *Botrytis cinerea* isolates on the rest 4 leaflets, indicating that the fungicide has systemic movement in plant tissues. These results indicate that SYP-Z048 has potential for management of brown rot causing by *M. fructicola* and other diseases caused by ascomycota and basidiomycota.**

Fungi are among the crop pests that reduce crop yield and food quality, and they also can produce toxic compounds that make the produce harmful for consumers. Damage caused by plant-pathogenic fungi is often controlled by the application of fungicides. Currently more than 80% of the fruit and vegetable crops grown in the U.S. receive at least one fungicide application every season; if the crops were left untreated, the yields of most fruit and vegetables would reduce 50 to 95 percent<sup>1</sup>. Fungicides are estimated to increase farm income in the U.S. by nearly \$13 billion annually<sup>1</sup>. In China, pesticide application has been increasing annually. Fungicides have become an integral part of efficient food production in China and other countries.


Many fungicides that target different pathogens and diseases have been reported<sup>2–5</sup>, but because of problems relating to toxic residues, disease resurgence and pathogen resistance<sup>6–10</sup>, the use of many fungicides is restricted. Therefore, researchers have been searching for fungicides with low toxicity, high selectivity, and high activity against fungal strains that are resistant to other fungicides.

SYP-Z048, 3-[5-(4-chlorophenyl)-2,3-dimethyl-3-isoxazolidinyl] pyridine, a mixture of two diastereomers (Fig. 1), is a novel fungicide developed for control of plant-pathogenic fungi by China Shenyang Research Institute of Chemical Industry in 1997. SYP-Z048 have been registered for control of grey mold caused by *Botrytis cinerea* on tomato in China as Junsiqi (25% EC, China Shenyang Research Institute of Chemical Industry, Shenyang, China)<sup>11</sup>. SYP-Z048 inhibits ergosterol synthesis in *B. cinerea*<sup>12</sup> and has been demonstrated to be a sterol 14 $\alpha$ -demethylase inhibitor (DMI)<sup>13</sup>.

DMI fungicides have been playing an important role in chemical control of many diseases since they were developed in 1970s, providing high efficacy against a broad spectrum of fungal pathogens. DMIs consist of many structurally heterogeneous compounds, belonging to different subclasses. The subclasses of imidazole and triazole are among the most widely used fungicides, as well as pyridines<sup>14</sup> and buthiobate<sup>15</sup>. SYP-Z048 is a pyridine derivative that includes a isoxazolidine ring.

It has been documented SYP-Z048 has highly activity on *M. fructicola*, the causal agent of brown rot. The mean EC<sub>50</sub> values for baseline population of *M. fructicola* is 0.017 µg/ml<sup>13</sup>, which is similar to the values for propiconazole (0.03 µg/ml)<sup>16</sup> and tebuconazole (0.016 µg/ml)<sup>17</sup>. The antifungal activity of SYP-Z048 against other plant-pathogenic fungi or on *M. fructicola* causing brown rot in field, however, is unknown.

The objectives of this study were to determine: (a) the *in vitro* antifungal activity of SYP-Z048 against 18 plant-pathogenic fungi; (b) the fungicidal activities of SYP-Z048 on different developmental stages of *M. fructicola*; (c)



**Figure 1** | Structure of SYP-Z048 used in the experiments, a mixture of two stereoisomers: 3-[(3R, 5S)-5-(4-chloro-phenyl)-2,3-dimethyl-3-isoxazolidinyl] pyridine (1) and its (3R, 5R)-isomer (2), drawn by Fengping Chen.

the preventative and curative activities of SYP-Z048 by detached fruits assay and control efficacy in field on brown rot and (d) systemic activity in tomato leaves.

## Results

**In vitro activity of SYP-Z048 against 18 plant pathogens.** SYP-Z048 inhibited the *in vitro* mycelial growth of many plant-pathogenic fungi, including ascomycota and basidiomycotac (Table 1) with  $EC_{50}$  values of 0.008–1.140  $\mu\text{g/ml}$ . In contrast, SYP-Z048 was much less effective against oomycota with  $EC_{50}$  values  $> 100 \mu\text{g/ml}$  (Table 1).

**Effect of SYP-Z048 on the different developmental stages of *M. fructicola*.** SYP-Z048 substantially inhibited mycelial growth, germ tube elongation, and sporulation of *M. fructicola* (Table 2). The  $EC_{50}$  values were 0.013 and 0.001  $\mu\text{g/ml}$  for mycelial growth and germ tube elongation, respectively. The inhibition activity of SYP-Z048 was greater for initial germ tube elongation than for mycelial growth. Spore production was significantly inhibited by SYP-Z048 when the concentration was  $> 0.02 \mu\text{g/ml}$ . Spore germination was not affected by SYP-Z048 (Table 2).

**Protective and curative activity of SYP-Z048 on detached pear fruit.** Whether applied 24 h before inoculation (protective activity) or 24 h after inoculation (curative activity), SYP-Z048 provided substantial control of *M. fructicola* on pear fruit (Table 3). The efficacy for protective and curative activity was 89.3% and 85.2%, respectively (Table 3). Comparison of disease incidence and lesion

**Table 2** | The *in vitro* effect of SYP-Z048 on different developmental stages of *M. fructicola*

SYP-Z048 concentration ( $\mu\text{g/ml}$ )	Inhibition of mycelial growth (%)	Inhibition of germ tube elongation (%)	Spore production $10^4$ spores/ $\text{cm}^2$	Spore germination (%)
0.00	-	-	15.94a*	85.6a
0.005	27.7	65.5	14.91ab	85.6a
0.01	48.9	68.7	12.99ab	85.5a
0.02	56.8	73.1	7.12bc	83.0a
0.03	71.7	74.3	1.15c	85.0a
0.06	81.6	81.1	0.33c	84.8a
$EC_{50}$	0.013	0.001	-	-

\*Means in a column followed by different letters are significantly different ( $P < 0.05$ ).

area after inoculation of *M. fructicola* for 3 days indicated that the fungicide was slowing down disease progression.

**Control efficacy of SYP-Z048 on brown rot in field.** Significant differences among fungicides treatments were observed in both peach orchards, (Fig. 2). All fungicides treatments were statistically superior to water control ( $P < 0.05$ ). SYP-Z048 treatment controlled brown rot significantly better than thiophonate-methyl and propineb when the concentration was  $> 100 \text{ g/L}$  in both locations. Compared to fenbuconazole at same dose, lower efficacy was observed with SYP-Z048 application at the Liaoning location whereas no difference among these treatments was observed at the Shandong location.

**Systemic activity of SYP-Z048 in tomato leaves.** In the first experiment concerning systemic activity of SYP-Z048, the fungicide SYP-Z048 inhibited *B. cinerea* on the leaflet where the fungicide was applied (the fourth leaflet) (Fig. 3) and suppressed lesion developing on the other four leaflets, although the fungicide was less effective on the untreated leaflets than on the treated leaflet (Fig. 4).

In the second experiment concerning movement of SYP-Z048 in tomato leaves, inhibition of *B. cinerea* on the upper epidermis side of the leaflet was similar to inhibition on the lower epidermis side of the leaflet when the fungicide was added on the opposite side of the leaflet (Fig. 5).

## Discussion

SYP-Z048 is a novel fungicide, being both pyridine derivative and also carrying a isoxazolidine ring. Both groups are effective on some pathogens. The combination of two active groups in SYP-Z048 would make this compound behave specially. Isoxazolidine or oxazole derivatives have been synthesized for antitumor, antimicrobial, herbicidal, and fungicidal activities<sup>18–23</sup>; many are effective against human pathogens. Activity against plant pathogens by other oxazolidinones and oxazolidinone-diones involves inhibition of RNA polymerase I and cytochrome b<sup>24,25</sup>. Pyridine fungicides possess different modes of action according to the specific fungicide.

**Table 1** | *In vitro* inhibition of plant pathogens by SYP-Z048


Organism	Species	Correlation* coefficient	$EC_{50}$ ( $\mu\text{g/ml}$ )
Ascomycota	<i>Gibberella zeae</i>	0.976	0.047
	<i>G. fujikuroi</i>	0.991	0.044
	<i>Fusarium oxysporum</i>	0.994	0.315
	<i>Valsa mali</i>	0.978	0.067
	<i>Physalospora piricola</i>	0.996	0.033
	<i>Monilinia fructicola</i>	0.989	0.013
	<i>Sclerotinia sclerotiorum</i>	0.995	0.009
	<i>Gaeumannomyces graminis</i>	0.995	0.081
	<i>Cladosporium cucumerium</i>	0.988	0.046
	<i>Cochliobolus lunatus</i>	0.986	0.008
	<i>Botrytis cinerea</i>	0.976	0.306
	<i>Magnaporthe oryzae</i>	0.989	1.140
	<i>Passalora fulva</i>	0.986	0.126
<i>Alternaria solani</i>	0.992	0.418	
Basidiomycota	<i>Ceratobasidium cerealis</i>	0.988	0.573
	<i>Sphaelotheca reiliana</i>	0.969	0.079
Oomycota	<i>Pythium aphanidermatum</i>	-	$> 100$
	<i>Phytophthora capsici</i>	-	$> 100$

\*Correlation between the probability value transformed by percentage of inhibition relative to the control (without fungicide) and the concentration (log transformed) of the fungicide.

**Table 3** | Protective and curative effect of SYP-Z048 on brown rot in detached pears after inoculation of *M. fructicola* for 3 days

Treatment	Disease Incidence (%)	Lesion area ( $\text{mm}^2$ )	Control Efficacy (%)
Water spray with inoculation	100a*	193.7a	-
Curative treatment	65.1b	28.7b	85.2
Protective treatment	66.3b	20.8b	89.3
Water spray without inoculation	0c	0c	-

\*Means in a column followed by different letters are significantly different ( $P < 0.05$ ).



**Figure 2** | Efficacy of SYP-Z048 and other fungicides for brown rot disease control in peach orchards in Liaoning and Shandong province in China in 2013.


Most of pyridines are respiration inhibitors, for example succinate dehydrate inhibitors boscalid and fluopyram<sup>5</sup> and succinate-cytochrome c reductase inhibitor pyribencarb<sup>26</sup>; some pyridines are 14 $\alpha$ -demethylase inhibitors such as buthiobate<sup>15</sup> and pyrifeno<sup>14</sup>; dipyrithione is response to lipopolysaccharide<sup>27</sup> and fluopicolide is involve in modification of the cellular localization of a spectrin-like protein<sup>28</sup>. SYP-Z048 has typical 14 $\alpha$ -demethylase inhibitor activity<sup>13</sup>, which should be a novel member of the demethylation inhibitors of ergosterol biosynthesis, but whether the activity of SYP-Z048 also results from other mechanism is not clear yet.

The lack of inhibition on *M. fructicola* spore germination supports the modes of action of SYP-Z048 which was implied as a DMI fun-


gicide<sup>13</sup>. In general, DMI fungicides are known to inhibit hyphal growth strongly rather than spore germination<sup>29</sup>. For example, DMIs doesn't affect the urediniospore germination of *Puceinia hemerocallidis*<sup>30</sup> and conidium germination of *Spilocaea oleagina*<sup>31</sup>. But in some pathogens, it is effective on spore germination<sup>32,33</sup>. The behavior observed for DMIs in pathogens differs due to the stage of starting sterol biosynthesis. Conidia of some species are able to synthesize sterols in the early stage of germination, whereas other initiate sterol synthesis at the very end of the germ tube phase<sup>33</sup>.

SYP-Z048 has potential for control of diseases caused by 16 plant-pathogenic ascomycota and basidiomycota fungi but not the oomycota. According to our *in vitro* experiment, the EC<sub>50</sub> values for SYP-Z048 < 1  $\mu$ g/ml to all experimental fungi except *M. oryzae*. The data collected under controlled conditions might not be the same as field study, but it reflects the real efficacy to some extent. For example, *M. fructicola* was inhibited strongly by SYP-Z048 *in vitro* (EC<sub>50</sub> = 0.013  $\mu$ g/ml), and in consistent, brown rot caused by *M. fructicola* was controlled efficaciously in the field with the application dose of 100  $\mu$ g/ml (The efficacy > 80%).


SYP-Z048 exhibits substantial activity on brown rot of stone fruit, caused by *M. fructicola*. *M. fructicola* is a world widespread pathogen, including Asia, Europe, Africa, North America, Central America, as well as South America and Oceania<sup>34</sup>. The fungus can cause not only fruit rot but also blossom blight and twig canker, especially causing damage in postharvest fruit<sup>35,36</sup>. The most effective management



**Figure 3** | Schematic of a tomato leaf (with five numbered leaflets, created by Fengping Chen) used in the experiments to determine systemic activity of SYP-Z048. A drop of fungicide was added to the fourth leaflet (arrow) before leaflets were inoculated with mycelium plug (circles) colonized by *B. cinerea*. See Methods section for additional details.



**Figure 4** | The efficacy of SYP-Z048 against *B. cinerea* after 2 days inoculation when the fungicide was applied to the fourth leaflet and the fungus was inoculated to all five leaflets on a tomato leaf. See Fig. 3 and the text for more details.



**Figure 5** | The efficacy of SYP-Z048 against *B. cinerea* after 2 days inoculation when the fungicide was applied to the upper epidermis (leaf-front) or lower epidermis (leaf-back) of a tomato leaflet and the fungus was inoculated to the other side of the same leaflet.

approaches are pre-harvest fungicide applications. Although resistance to benzimidazoles<sup>37</sup>, dicarboxamides<sup>38</sup>, succinate dehydrogenase inhibitors (SDHIs) and quinone outside inhibitor fungicides (QoIs)<sup>39</sup> has been reported in *M. fructicola*, SYP-Z048 is not cross-resistant to compounds belonging to those classes<sup>13</sup>. The comparison of efficacy in two peach orchards showed that SYP-Z048 was more efficacious than thiophonate-methyl (a representative of benzimidazoles) and propineb (a protective fungicide) in this study, even the application concentration was 8 times and 15 times less, respectively. All the facts taken together suggest that SYP-Z048 is a fungicide with practical significance in brown rot management.

SYP-Z048 possesses both protective and curative fungicidal activity. Such fungicides are key components of modern agriculture due to their ability of preventing disease expanding. The detached fruit assay showed SYP-Z048 controlled brown rot well when applied before inoculation, but also showed high efficacy when applied after inoculation which is not significantly different from the pre-treatment. The movement of SYP-Z048 in plant tissue was confirmed using the compound leaf of tomato. The data we obtained indicated that SYP-Z048 could penetrate into leaflet and move acropetally, horizontally, and basipetally in plant, and also moved through the leaflet tissues, after application.

In conclusion, SYP-Z048 has substantial *in vitro* activity against a very broad spectrum of plant pathogenic fungi. The fungicide is a systemic fungicide with both protective activity and curative activity against brown rot disease. The novel structure fungicide SYP-Z048 may be expected to be an effective agrochemical agent in plant disease management.

## Methods

**Media, fungicides, and pathogens.** Potato dextrose agar (PDA medium) was used for the routine culture of the fungi and contained 200 g of potato, 15 g of glucose, and 12 g of agar in 1 L of sterile deionized water. The medium was autoclaved for 20 min at 121°C. Water agar was used for the spore germination and tube elongation tests and contained 12 g of agar in 1 L of sterile deionized water, without sterilization. The chemicals were purchased from Syngenta China Ltd. (Beijing). The same batch of medium was used for all replicates within each experiment to reduce variability.

The fungicide SYP-Z048 (84%) was kindly provided by China ShenYang Research Institute of Chemical Industry. The chemical was dissolved in methanol and diluted to 10 mg/ml as a stock solution, which was stored at 4°C in the dark to preserve fungicide activity. For each assay on PDA plate, the stock solution was diluted to working concentrations with sterile deionized water. In all cases, the final amount of solvent never exceeded 0.1% in treated and control samples. The formulations of SYP-Z048 (25% Junsiqi EC) was provided by Shenyang Sciencreat Chemicals Co., Ltd; fenbuconazole (24% Yingde SC), thiophonate-methyl (70% Jiajiliujunling WP) and propineb (70% bingsenxin WG) were purchased from Dow AgroScience, Jiangsu Rotam Chemistry Co., Ltd., and NantongBaoye Chemical Co., Ltd., respectively.

The activity of SYP-Z048 was tested against the following plant-pathogenic fungi: *Gibberella zeae*, *G. fujikuroi*, *Fusarium oxysporum*, *Valsa mali*, *Physalospora piricola*, *M. fructicola*, *Sclerotinia sclerotiorum*, *Gaeumannomyces graminis*, *Cladosporium*

*cucumerium*, *Cochliobolus lunatus*, *Botrytis cinerea*, *Magnaporthe oryzae*, *Passalora fulva*, *Alternaria solani*, *Ceratobasidium cerealis*, *Sphacelotheca reiliana*, *Pythium aphanidermatum* and *Phytophthora capsici*. *S. reiliana* was cultured in Richard's liquid medium (10.0 g of KNO<sub>3</sub>, 5.0 g of KH<sub>2</sub>PO<sub>4</sub>, 2.5 g of MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.02 g of FeCl<sub>3</sub>, 30.0 g of glucose and 1 L of sterile deionized water) at 20°C. The other fungi were cultured on PDA at 25°C.

**Inhibitory activity of SYP-Z048 *in vitro*.** The stock solution of SYP-Z048 in methanol was diluted serially in autoclaved PDA (cooled to around 50°C) to achieve final concentrations according to preliminary test (data not show). Before it solidified, the PDA plus fungicide was added to 9-cm diameter Petri plates, with three replicate plates per concentration. Control plates contained PDA without fungicide but with corresponding concentrations of methanol (≤0.1%). For each of the 18 test fungi listed in Table 1, except *S. reiliana*, a mycelium plug (5 mm) was transferred from the leading edge of an actively growing colony on PDA to the dishes amended with 0 to different concentrations fungicide, with percentage inhibition from 10% to 90%. The dishes were incubated at 25°C in darkness. When the colony in control plates (without fungicide) had covered 75% of the agar surface, the diameter of all colonies of that microorganism were measured. The percentage of inhibition of radial growth was calculated. The tests were replicated three times for each microorganism.

The effect of SYP-Z048 on growth of *S. reiliana*, which does not grow on PDA, was determined with the ELISA plate turbidity method. *S. reiliana* teliospores (10 mg) collected from diseased corn were suspended in 10 ml of 0.03% Tween 20 for 4 h. The suspension was centrifuged at 4000 g for 10 min. The pelleted teliospores were added to a 0.25% KMnO<sub>4</sub> solution. After 25 min, sterilized water was added and then centrifuged again. The pelleted teliospores were immersed in 0.05 M HCl for 6 h before they were mixed into Richard's medium adjusted to approximately 10<sup>5</sup> teliospores/ml. Liquid containing 75 µl teliospores suspension and 75 µl SYP-Z048 of each concentration were added to each well of ELISA plates. The plates were incubated at 28°C on a rotary shaker (120 r/min) in darkness for 84 h. There were three wells per fungicide concentration, and the experiment was performed twice. The transmittance and optical density at 650 nm (OD<sub>650</sub>) of the suspension in each well was measured with a microplate reader (Bio-rad model550) and inhibition based on differences in optical densities between fungicide treatments and controls. The median effective concentration value (EC<sub>50</sub>) was calculated by regressing the percentage inhibition (probability value transformed) against the logarithm value of fungicide concentration (log transformed). Regression analysis was performed with Microsoft Excel 2007.

**Effect of SYP-Z048 on different developmental stages of *M. fructicola*.** Effect of SYP-Z048 on mycelial growth of *M. fructicola* isolate was measured as described above. The fungicide concentrations were 0, 0.005, 0.01, 0.02, 0.03, and 0.06 mg/l. After colony diameters were measured, the cultures were used to determine the effect of SYP-Z048 on spore production. In cultures with < 0.02 mg/l, six colonized agar plugs, 5 mm in diameter, were excised from each replicate plate; the selected plugs were taken along the radius from the center of the culture to the margin of the colony. In plates with higher concentrations of fungicide, the colony was too small to collect plugs along the radius, therefore a single plug was collected. The plugs from each plate were put into a 15-ml tube with 2.0 ml of sterile deionized H<sub>2</sub>O, agitated with a sterile needle, and shaken for 5 min with a laboratory vortex mixer. Spores were counted using a haemocytometer. Spore production was expressed as the number of spores per mm<sup>2</sup> of colony area.

To determine the effect of SYP-Z048 on spore germination and germ tube elongation, spores were harvested from a colony of *M. fructicola* isolate growing on PDA without fungicide by adding sterile deionized water to the plate and agitating. The spore suspension was adjusted to approximately 10<sup>6</sup> spores/ml. One hundred µl of the spore suspension was spread on water agar in 9 cm diam petri dishes containing 0, 0.005, 0.01, 0.02, 0.03 and 0.06 µg/ml SYP-Z048. Germination was measured after 6 h at room temperature in the dark by examining 100 spores in three locations on the petri dish at 100× magnification. A spore was scored as germinated if the germ tube had reached at least the full length of the spore. The dishes were then incubated for an additional 4 h under the same conditions and germ tube length was measured on 60 germinated spores per dish at 400× magnification using Image-Pro Plus 5.0 software. Three replicate plates were used for each concentration of SYP-Z048.

**Preventive and curative activity of SYP-Z048 on pear fruit.** The methods for determining the preventive and curative activity of *M. fructicola* on pear fruits were similar to those described previously<sup>40,41</sup>. The pears of "Fengshui" variety were purchased from a supermarket. Before inoculation, pears were kept at 23°C overnight and were rinsed with tap water and air-dried. After the pears were peeled and cut in half, they were placed in a plastic box with a wet filter paper at the bottom, cut side facing down, and were then wounded with a sterile needle on the peeled but uncut surface (six wounds per fruit half). Each wounded site was inoculated with 10 µl of an *M. fructicola* spore suspension (5 × 10<sup>6</sup> conidia/ml) obtained from other pears that had been previously inoculated and infected with *M. fructicola*. SYP-Z048 (200 µg/ml) was applied either 24 h before the fruit was inoculated (protective treatment) or 24 h after the fruit was inoculated (curative treatment); the fungicide was applied to runoff with a fine-mist sprayer. Two control treatments were included. In one control (positive control), fruit were inoculated with *M. fructicola* 24 h after spraying with sterile water. In the other control (negative control), wounded fruit were sprayed with water but not inoculated. The moisture boxes were sealed with clear plastic and



arranged randomly in climate control chamber maintained at 23°C, >80% RH, and a 12-h photoperiod. After 4 days, the number of infected wounds and the diameter of the lesions around the wounds were recorded. The experiment was performed twice with 3 replicates per experiment. Each replicate consisted of 12 inoculations.

**Control efficacy of SYP-Z048 on brown rot in field.** To assess the efficacy of SYP-Z048 for control of brown rot in the field, the fungicides of SYP-Z048 (25% Junsiqi EC), fenbuconazole (24% Yingde SC), thiophonate-methyl (70% Jiajiliujunling WP) and propineb (70% bingsenxin WG) were applied to trees in two orchards. Three concentrations of SYP-Z048 were tested (50, 100 and 200 g/L) and compared to fenbuconazole, thiophonate-methyl and propineb applied at 100, 800, and 1500 g/L, respectively. Control treatments were trees sprayed with water. Treatments were replicated four times in a randomized block with four trees per replication. All fungicides were applied by sprayer to runoff four times before harvest. 'Jinlu' peach trees were treated on 6 May, 20 May, 4 June and 18 June 2013 in Liaoning province in China. 'Baifeng' peach trees were treated on 29 April, 16 May, 30 May, and 15 June 2013 in Shandong province in China. When fruit were at commercial shipping maturity, a sample consisting of 150 fruit, was randomly harvested from each tree and the total number of diseased fruit was counted. Disease incidence was expressed as percentage of diseased fruit and control efficacy was expressed as: Control efficacy (%) =  $(1 - A) \times 100$ , where A = disease incidence with fungicide treatment divided by mean disease incidence with water application.

**Systemic activity of SYP-Z048 against *B. cinerea* on tomato leaves.** To investigate potential movement and activity of the fungicide in acropetal, horizontal, and basipetal directions, the fungicide was applied in one spot on a tomato leaf with inoculation in the same spot and other locations in two experiments. In the first experiment, pinnately compound leaves (with five leaflets) were collected at the 3<sup>rd</sup> leaf position from tomato (*Lycopersicon esculentum* Mill cv. Jiabao) plants growing in the greenhouse. The leaves were rinsed with tap water, air dried, and placed in a plastic box with wet filter paper at the bottom. In the first experiment 10- $\mu$ L containing 0, 500, 1000, or 2000  $\mu$ g ml<sup>-1</sup> SYP-Z048 was applied as a drop to the adaxial side of the fourth leaflet (Fig. 3). The boxes were sealed with clear plastic and arranged randomly in a climate control chamber at 23°C, >80% RH, and a 12-h photoperiod. One day after application, a 3-mm-diam mycelium plug from the margin of an actively growing colony of *B. cinerea* on PDA was placed on the adaxial side of each of the five leaflets. After 2 days the diameter of the *B. cinerea* colony formed around each plug was recorded. The experiment was performed twice with three replicates per treatment. Each replicate consisted of 10 leaves.

In a second experiment, 10  $\mu$ L containing 0, 250, 750, or 1500  $\mu$ g ml<sup>-1</sup> SYP-Z048 was applied to the upper epidermis side of the leaflet or lower epidermis side. After 24 h, the leaflet was incubated as above with a 3-mm-diam mycelium plug placed on the opposite side of the leaflet; i.e., one plug was on the upper epidermis, when the fungicide was treated on the lower epidermis. Colonies were measured and efficacy was calculated as described in the previous paragraph. The experiment was performed twice with 30 replicate leaves per treatment.

**Statistical analysis.** All data were analyzed using SAS software ver. 8.0. Difference among means was determined with the Duncan's multiple range test at P = 0.05.

- Gianessi, L. & Reigner, N. The importance of fungicides in U.S. crop production. *Outlook Pest Manag.* **10**, 209–213 (2006).
- Bartlett, D. W. *et al.* The strobilurin fungicides. *Pest Manag. Sci.* **58**, 649–662 (2002).
- Becher, R. & Wirsal, S. G. R. Fungal cytochrome P450 sterol 14 $\alpha$ -demethylase (CYP51) and azole resistance in plant and human pathogens. *Appl. Microbiol. Biotechnol.* **95**, 825–840 (2012).
- Kumar, L. *et al.* Imidazole derivatives as possible microbicides with dual protection. *Eur. J. Med. Chem.* **45**, 817–824 (2010).
- Avenot, H. F. & Michailides, T. J. Progress in understanding molecular mechanisms and evolution of resistance to succinate dehydrogenase inhibiting (SDHI) fungicides in phytopathogenic fungi. *Crop Prot.* **29**, 643–651 (2010).
- Bridges, B. A. Mutagenicity of captan and related fungicides. *Mutat. Res.* **32**, 3–34 (1975).
- Hurley, P. M., Hill, R. N. & Whiting, R. J. Mode of carcinogenic action of pesticides inducing thyroid follicular cell tumors in rodents. *Environ. Health Perspect.* **106**, 437–445 (1998).
- Clary, T. & Ritz, B. Pancreatic cancer mortality and organochlorine pesticide exposure in California, 1989–1996. *Am. J. Ind. Med.* **43**, 306–313 (2003).
- Gordon, E. Captan: Transition from 'B2' to 'not likely'. How pesticide registrants affected the EPA Cancer Classification Update. *J. Appl. Toxicol.* **27**, 519–526 (2007).
- FRAC, List of plant pathogenic organisms resistant to disease control agents. (2013) Available from: [http://www.frac.info/publication/anhang/List%20of%20resistant%20plant%20pathogenic%20organisms\\_February%202013%20updated.pdf](http://www.frac.info/publication/anhang/List%20of%20resistant%20plant%20pathogenic%20organisms_February%202013%20updated.pdf) (Date of access: 10/7/2014).
- Si, N. G. *et al.* Biological activity and application of a novel fungicide SYP-Z048 (I). *Chin. J. Pestic.* **43**, 16–18 (2004).
- Han, P., Liu, X., Liu, P. & Si, N. Effect of novel fungicide 5-(4-chloro phenyl)-2,3-dimethyl-3-(pyridine-3)-oxazoline on ergosterol biosynthesis in *Botrytis cinerea* by high performance liquid chromatography. *Chin. J. Anal. Chem.* **34**, 1467–1470 (2006).
- Chen, F. P. *et al.* Baseline sensitivity of *Monilinia fructicola* from China to the DMI fungicide SYP-Z048 and analysis of DMI-resistant mutants. *Plant Dis.* **96**, 416–422 (2012).
- Masner, P. & Kerkenaar, A. Effect of the fungicide pyrifenoxy on sterol biosynthesis in *Ustilago maydis*. *Pestic. Sci.* **22**, 61–69 (1988).
- Aoyama, Y., Yoshida, Y., Hata, S., Nishino, T. & Katsuki, H. Buthiobate: A potent inhibitor for yeast cytochrome P-450 catalyzing 14 $\alpha$ -demethylation of lanosterol. *Biochem. Biophys. Res. Commun.* **115**, 642–647 (1983).
- Zehr, E. I., Luszcz, L. A., Olien, W. C., Newall, W. C. & Toler, J. E. Reduced sensitivity in *Monilinia fructicola* to propiconazole following prolonged exposure in peach orchards. *Plant Dis.* **83**, 913–916 (1999).
- Yoshimura, M. A., Luo, Y., Ma, Z. H. & Michailides, T. J. Sensitivity of *Monilinia fructicola* from stone fruit to thiophanate-methyl, iprodione, and tebuconazole. *Plant Dis.* **88**, 373–378 (2004).
- Shaber, S. H., Zhang, L., Szapacs, E. M. & Quinn, J. A. Heterocyclic substituted isoxazolidines and their use as fungicides. U.S. Patent 6313147.B1 (2001).
- Damodiran, M. *et al.* Antibacterial Activity, Quantitative structure-activity relationship and diastereoselective synthesis of isoxazolidine derivatives via 1,3-dipolar cycloaddition of d-glucose derived nitron with olefin. *Chem. Biol. Drug Des.* **74**, 494–506 (2009).
- Liu, X.-H., Lv, P.-C., Xue, J.-Y., Song, B.-A. & Zhu, H.-L. Novel 2,4,5-trisubstituted oxazole derivatives: Synthesis and antiproliferative activity. *Eur. J. Med. Chem.* **44**, 3930–3935 (2009).
- Turan-Zitouni, G., Kaplancikli, Z. A., Ozdemir, A., Reviel, G. & Guven, K. Synthesis and antimicrobial activity of some 2-(benzodioxol/benzodimidazol-2-ylthio)-N-(9H-fluoren-9-yl)acetamide derivatives. *Phosphorus Sulfur Silicon Relat. Elem.* **182**, 639–646 (2007).
- Vicentini, C. B. *et al.* Synthesis and biological evaluation of a series of substituted pyrazolo 3,4-d -1,2,3-triazoles and pyrazolo 3,4-d oxazoles. *Arch. Pharm.* **331**, 269–272 (1998).
- Bhat, A. R., Athar, F. & Azam, A. New derivatives of 3,5-substituted-1,4,2-dioxazoles: Synthesis and activity against *Entamoeba histolytica*. *Eur. J. Med. Chem.* **44**, 926–936 (2009).
- Xiao, K. H. *et al.* Effect of famoxadone on photoinduced electron transfer between the iron-sulfur center and cytochrome c1 in the cytochrome bc1 complex. *J. Biol. Chem.* **278**, 11419–11426 (2003).
- Jiang, Y. Effects of oxadixyl treatment on the ultrastructure of *Phytophthora infestans* infecting tomato leaves. *Pestic. Sci.* **35**, 153–159 (1992).
- Kataoka, S., Takagaki, M., Kaku, K. & Shimizu, T. Mechanism of action and selectivity of a novel fungicide, pyribencarb. *J. Pestic. Sci.* **35**, 99–106 (2010).
- Liu, Z. *et al.* Dipyrithione inhibits lipopolysaccharide-induced iNOS and COX-2 up-regulation in macrophages and protects against endotoxic shock in mice. *FEBS Lett.* **582**, 1643–1650 (2008).
- Toquin, V. *et al.* A new mode of action for fluopicolide: modification of the cellular localization of a spectrin-like protein. *Pflanzenschutz-Nachrichten Bayer* **59**, 171–184 (2006).
- Sisler, H. D. & Ragsdale, N. N. [Biochemical and cellular aspects of the antifungal action of ergosterol biosynthesis inhibitors] *Mode of action of antifungal agents* [257–282] (Cambridge, UK, 1984).
- Buck, J. W. & Williams-Woodward, J. L. The effect of fungicides on urediniospore germination and disease development of daylily rust. *Crop Prot.* **22**, 135–140 (2003).
- Obanor, F. O., Walter, M., Jones, E. E. & Jaspers, M. V. *In vitro* effects of fungicides on conidium germination of *Spilocaea oleagina*, the cause of olive leaf spot. *N. Z. Plant Prot.* **58**, 278–282 (2005).
- Clarkson, J. P., Kennedy, R., Phelps, K., Davies, J. & Bowtell, J. Quantifying the effect of reduced doses of propiconazole (Tilt) and initial disease incidence on leek rust development. *Plant Pathol.* **46**, 952–963 (1997).
- Pontzen, R. & Scheinpflug, H. Effects of triazole fungicides on sterol biosynthesis during spore germination of *Botrytis cinerea*, *Venturia inaequalis* and *Puccinia graminis* f. sp. *tritici*. *Eur. J. Plant Pathol.* **95**, 151–160 (1989).
- EPPO/CABI, Data sheets on quarantine pests: *Monilinia fructicola*. (2010) Available from: [https://www.eppo.int/QUARANTINE/fungi/Monilinia\\_fructicola/MONIFC\\_ds.pdf](https://www.eppo.int/QUARANTINE/fungi/Monilinia_fructicola/MONIFC_ds.pdf) (Date of access: 11/7/2014).
- Hong, C., Holtz, B. A., Morgan, D. P. & Michailides, T. J. Significance of thinned fruit as a source of the secondary inoculum of *Monilinia fructicola* in California nectarine orchards. *Plant Dis.* **81**, 519–524 (1997).
- Luo, Y. & Michailides, T. J. Risk analysis for latent infection of prune by *Monilinia fructicola* in California. *Phytopathology* **91**, 1197–1208 (2001).
- Jones, A. L. & Ehret, G. R. Isolation and characterization of benomyl-tolerant strains of *Monilinia fructicola*. *Plant Dis.* **60**, 765–769 (1976).
- Elmer, P. A. G. & Gaunt, R. E. The biological characteristics of dicarboximide-resistant isolates of *Monilinia fructicola* from New Zealand stone-fruit orchards. *Plant Pathol.* **43**, 130–137 (1994).
- Amiri, A., Brannen, P. M. & Schnabel, G. Reduced sensitivity in *Monilinia fructicola* field isolates from South Carolina and Georgia to respiration inhibitor fungicides. *Plant Dis.* **94**, 737–743 (2010).



40. Harada, Y. *et al.* *Monilia mumeicola*, a new brown rot fungus on *Prunus mume* in Japan. *J. Gen. Plant Pathol.* **70**, 297–307 (2004).
41. Valencia-Chamorro, S. A., Perez-Gago, M. B., Del Rio, M. A. & Palou, L. Curative and preventive activity of hydroxypropyl methylcellulose-lipid edible composite coatings containing antifungal food additives to control citrus postharvest green and blue molds. *J. Agric. Food Chem.* **57**, 2770–2777 (2009).

## Acknowledgments

This work was supported by the Special Fund for Agroscientific Research in the Public Interest of China (201303023). This work was also partially supported by National High Technology Research and Development Program of China (2012CB111401). We thank J. L. Starr for his editorial efforts and critique on the manuscript.

## Author contributions

L.X. and C.F. designed the experiments. C.F. and H.P. performed the experiments *in vitro*. S.N. and L.J. performed the experiments in field. C.F. and L.P. interpreted the results. L.X. and C.F. wrote the paper. All authors reviewed the manuscript.

## Additional information

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Chen, F. *et al.* Activity of the novel fungicide SYP-Z048 against plant pathogens. *Sci. Rep.* **4**, 6473; DOI:10.1038/srep06473 (2014).



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>