SCIENTIFIC REPORTS

OPEN

SUBJECT AREAS: PROTEIN ANALYSIS KIDNEY MOLECULAR EVOLUTION

> Received 29 April 2013

Accepted 4 February 2015

Published 12 March 2015

Correspondence and requests for materials should be addressed to W.L. (whliu@stu.edu. cn) or D.W. (wangd@ ihb.ac.cn)

* These authors contributed equally to this work.

Physicochemical Evolution and Molecular Adaptation of the Cetacean Osmoregulation-related Gene UT-A2 and Implications for Functional Studies

Jingzhen Wang^{1,2*}, Xueying Yu^{1,3*}, Bo Hu¹, Jinsong Zheng¹, Wuhan Xiao¹, Yujiang Hao¹, Wenhua Liu² & Ding Wang¹

¹Key Laboratory of Aquatic Biodiversity and Conservation of the Chinese Academy of Sciences; Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, Hubei 430072, China, ²Marine Biology Institute, Shantou University, Shantou, Guangdong 515063, China, ³University of Chinese Academy of Sciences, Beijing 100039, China.

Cetaceans have an enigmatic evolutionary history of re-invading aquatic habitats. One of their essential adaptabilities that has enabled this process is their homeostatic strategy adjustment. Here, we investigated the physicochemical evolution and molecular adaptation of the cetacean urea transporter UT-A2, which plays an important role in urine concentration and water homeostasis. First, we cloned UT-A2 from the freshwater Yangtze finless porpoise, after which bioinformatics analyses were conducted based on available datasets (including freshwater baiji and marine toothed and baleen whales) using MEGA, PAML, DataMonkey, TreeSAAP and Consurf. Our findings suggest that the UT-A2 protein shows folding similar to that of dvUT and UT-B, whereas some variations occurred in the functional S₀ and S₁ regions of the selectivity filter. Additionally, several regions of the cetacean UT-A2 protein have experienced molecular adaptations. We suggest that positive-destabilizing selection could contribute to adaptations by influencing its biochemical and conformational character. The conservation of amino acid residues within the selectivity filter of the urea conduction pore is likely to be necessary for urea conduction, whereas the non-conserved amino acid replacements around the entrance and exit of the conduction pore could potentially affect the activity, which could be interesting target sites for future mutagenesis studies.

t is well known that cetaceans have a unique evolutionary history. They originated from terrestrial tetrapod mammals¹⁻⁴, and were eventually distributed into aquatic environments after experiencing a period of semiaquatic living⁵⁻⁹. Therefore, they have developed unique adaptive osmoregulatory strategies to maintain internal homeostasis. Osmoregulation in marine mammals, including cetaceans, has been of interest to physiologists for over a century¹⁰⁻¹⁴. Previous research indicates that three main factors facilitate this osmoregulation: 1) reducing skin and respiratory system water loss¹⁵⁻¹⁹; 2) obtaining water from food, seawater and their body fat^{10,11,14,20-22}; and 3) concentrating urine and reabsorbing water^{10,11,23-25}. All of these strategies are physiological mechanisms for water and electrolyte regulation that allow cetaceans to maintain a relatively stable and constant internal state in the face of environmental changes.

One of the most fundamental and efficient osmoregulation mechanisms is the capability of the kidneys to produce concentrated or diluted urine to match the intake and elimination of water and electrolytes²⁶. In regard to the urine-concentrating mechanism, any hypothesis regarding the mechanism by which the inner medulla concentrates urine needs to include a urea effect^{26,27}. As a crucial factor in osmolytic balance of the blood and urine^{28–32}, urea can benefit water re-absorption in mammals^{26,33–35}. Cloning of the protein urea transporter (UT, also known as solute carrier family 14, SLC14), which mediates urea conduction, has provided additional insight into the urine-concentrating mechanism²⁸. The UT, which is located in the descending limbs of the loop of Henle and the inner medullary collecting ducts, facilitates urea accumulation in the renal medulla and equilibrates the urea concentrations between the urine and the interstitium, thereby influencing water re-absorption in the kidney^{29,31,35–38}. Previous animal experiments have also verified the essential roles of UTs in animal homeostasis regulation. For example, mice lacking UTs cannot efficiently conserve water^{39–41} and mutant UTs in humans generate bladder cancer^{42,43} and create blood pressure problems⁴⁴. Cetaceans usually consume lots of protein,

which will generate large quantities of urea by catabolism. Therefore, cetaceans also face the challenge of osmotic diuresis; if urea in the collecting ducts is highly concentrated and UT regulation is lacking, it will draw water from the kidney interstitium and lead to undesirably increased water excretion. This means that UT is also essential to cetacean osmoregulation. Recently, a few studies have been conducted to investigate UTs in cetaceans. Janech and colleagues sequenced the first cetacean UT-A2 from the kidney of a shortfinned pilot whale (whUT-A2)45. WhUT-A2 shows a high level of sequence identity to terrestrial mammals, and greatly enhanced the urea uptake of Xenopus oocvtes in a heterologous expression experiment, indicating that the whale UT-A2 conducts urea. Subsequently, several UT-A2 genes from the kidney of baleen and toothed whales were cloned. It is proposed that there are possible regulation differences between these genes (different protein phosphorylation sites) leading to their high urinary concentration capability⁴⁶; however, other studies indicate that UT-A2 is not stimulated by cAMP analog-dependent protein phosphorylation^{28,47-50}. In 2009 and 2012, Levin and colleagues used X-ray crystallography to study urea transporter structures in detail, and they elucidated the permeation mechanisms of UT-B and dvUT in bovine and bacteria^{51,52}. They indicated that both dvUT and UT-B are homotrimers and each subunit contains a urea conduction pore with a narrow selectivity filter. The binding sites in the filter and the well-placed α -helix dipoles accommodate multiple dehydrated urea molecules in single file that pass through the pore, acting like a molecular version of a coin-operated machine²⁹. However, previous studies have not yet experimentally determined the structure and mechanisms of UT-A2, neither have they resolved the effects of molecular adaptation and physicochemical evolution of UTs. These pieces of information will be important for understanding the evolutionary urine-concentrating strategies adjustments that occurred in organisms to counteract external osmotic changes and providing a treasure trove of information on UT protein structure and function.

Marine cetaceans that inhabit a hyperosmotic environment (with little to no access to fresh water) appear to conserve water better and produce more highly concentrated urine (1353-2129 mOsm/kg) than humans and cattle, to which cetaceans are closely and can access fresh water from time to time^{10,12,14}. Our recent studies showed that the Yangtze finless porpoise (Neophocaena asiaeorientalis asiaeorientalis, YFP), which resides in hypo-osmotic freshwater (Yangtze river) likely excretes relatively less concentrated urine (934 mOsm/ kg) compared to its marine counterparts that include Bohai finless porpoises (Neophocaena asiaeorientalis sunameri)²⁵. These observations regarding urine concentration lead to two questions. 1) Have the UT-A2 genes (which are correlated with urine concentration abilities in mammals) evolved adaptively during cladogenesis, when cetaceans transitioned from land to the marine environments? And 2) are there any molecular adaptations in UT-A2 between the extant seawater and freshwater cetaceans that could provide an adaptive advantage in facing different osmoregulatory challenges?

To address these questions, we sequenced the urea transporter UT-A2 from the kidney of freshwater YFP. Bioinformatics analyses of phylogenetic, molecular evolution and radical amino acid property changes were conducted on UT-A2s among cetaceans (both marine and freshwater cetaceans) and several terrestrial mammals to investigate the molecular evolution and adaptation of UT-A2s. Additionally, the newly revealed dvUT and UT-B 3D structure and mechanisms were introduced to explore the structural and functional conservation and variation of mammalian UT-A2. Furthermore, the molecular differences in the UT-A2 protein that may lead to function divergence between cetaceans and terrestrial mammals were discussed. These analyses may provide new insight in understanding the molecular mechanism of urea permeation and the molecular basis of osmoregulation in cetaceans and provide instructive

information for understanding human kidney diseases correlated with urea regulation.

Results

UT-A2 sequences. As shown in Fig. 1B, the fragments and entire length of UT-A2 were amplified from mRNAs of two YFP specimens, using RT-PCR with the degenerate primers F1-R1, F2-R2, F3-R3 and F1-R3 (Table S1). After sequencing, splicing, and manually checking the amplicons, the full-length contiguous coding sequence of 1194 bp, encoding a 397-amino acid protein (the sequences of UT-A2 from the two YFP were absolutely concordant) was obtained. The alignments of coding sequences (CDS) and amino acids from UT-A2 of cetaceans and terrestrial mammals are shown in Fig. S1. The CDS of the YFP UT-A2 gene was submitted to the NCBI Data Bank (Gene Bank accession no.: KP318829).

Phylogenetic analysis. In MrBayes analysis, the best-fit HKY + G model (-lnL = 3452.8572, K = 24, BIC = 7075.7559) was selected by Modeltest 3.7 using the Bayesian Information Criterion (BIC)⁵³. The base frequencies of A, T, C, G were 0.2118, 0.3185, 0.2423 and 0.2275, respectively. The transition/transversion (Ti/Tv) ratio was 2.9435, and the putative variable site gamma distribution shape parameter was 0.2360. After the phylogenetic analyses, very similar dendrograms, with high bootstrap values, were constructed in all cases (Fig. 2).

Molecular evolutionary analysis. The average ω ratio for overall sequence pairs was 0.085, 0.114 and 0.086, as calculated from MEGA4, DataMonkey and PAML codeml (M0), respectively, indicating purifying selection on UT-A2. In PAML tests, the estimated parameters of different models and the LRT results are provided in Table 1. Rejection of the M0 model by the free-ratios model (p < 0.01) presented the variable ω values among branches. In a further analysis, an *a-priori* hypothesis was tested by specifying the cetacean-specific branch with a different ω rate; the one-ratio model was rejected by an unrestricted two-ratios model (p < 0.05), and the alternative hypothesis (estimated ω) eliminated the constrained null hypothesis ($\omega 1 = 1$) in the LRT tests (p < 0.001). The ω value estimated for cetaceans was 0.14, about twice the value estimated for non-cetacean mammals ($\omega = 0.07$). We also tested the freshwater cetacean lineages for the baiji and YFP, but these were not significant relative to the null models. To evaluate ω ratio variation among codon positions, an LRT test based on sitespecific models M3 (K = 3) versus M0 and M3 (K = 2) versus M0 was conducted. Both results (p < 0.001) provided evidence for the existence of rate-heterogeneous selection pressure at amino acid sites. In the analyses, M2a had the same likelihood values and identical parameter estimates as M1a, indicating that approximately 93.3% of the sites are under purifying selection ($\omega =$ 0.05). M7 is not rejected in favor of M8, suggesting an "L" shape of the beta distribution (0.198, 1.738). In LRT tests, the branchsite Model A performed significantly better than model M1a (p <0.01), but was not significant against the null model A (p = 0.14) in the cetacean branch. However, the branch-site Model B was significantly better (p < 0.05) than model M3 (K = 2), and identified 3 positive sites, which is consistent with Model A-M1a. In the DataMonkey tests, the SLAC tests inferred substitutions by site using the shortest evolutionary paths (Fig. 3A), and the branch-site model MEME showed evidence that 2 sites are under episodic diversifying selection.

Detection of positive selection at the amino acid level. To extrapolate the potential structure and mechanism of the UT-A2 protein, Consurf was introduced to compute the position-specific conservation scores. Then, graded conservation scores were projected onto the space-filled 3D structure of UT-A2's homolog dvUT and UT-B, and finally visualized (Fig. 4A, B): identical



Figure 1 | RT-PCR amplification of Yangtze finless porpoise UT-A2. (A) Schematic diagram of the UT-A2 coding sequence. The primers used for RT-PCR are indicated by the green bar and labeled with letters F and R. The amplified fragments for the following sequencing and splicing are indicated by the dark brown bars. (B) Products of RT-PCR amplification with a set of primers. Lane a: a 454-bp fragment amplified using the primers F1 and R1; lane b: a 652-bp fragment amplified using the primers F2 and R2; lane c: a 534-bp products amplified using the primers F3 and R3; lane d, e and f: the full length 1194-bp fragment amplified using the primers F1 and R3 from two Yangtze finless porpoise individuals.

amino acids are shown in dark purple and variable ones in light blue. The results revealed that the functionally important regions on both the DNA and the protein were conserved. From the superimposed 3D structure, we can see that the individual protomers of UT-A2 have the same overall folding structure as do dvUT and UT-B, which contain two homologous halves with opposite orientations in the membrane (Fig. 4D, structural Motif 1 and Motif 2). A membrane-spanning pore is formed at the interface of the two halves that each contains six helices (Pa, T1a-5a and Pb, T1b-5b, Fig. 4B, C and D).

For its important role in urea conduction, the differences between the UT selectivity filters in the mammalian UT-A2, UT-B and the bacterial dvUT were further investigated. The results indicated that the UT-A2 residues of the S_m regions (L127, T176, L291 and T338) were completely consistent with their corresponding sites in dvUT (L84, T130, L247, T294) and UT-B (L123, T172, L287, T334). However, the sites in the S_o and S_i regions showed some variation between dvUT, UT-B and UT-A2 (Fig. 5). The S_o region sites E187, F190, L293 of dvUT shifted to glutamine (Q231), glycine (G234) and



Figure 2 | Phylogram resulting from NJ, MP and Bayesian analyses on UT-A2 genes from several cetaceans and terrestrial mammals. The numbers on each branch denote the bootstrap proportion inferred by each method. The cetacean drawings were modified from the book of Chinese cetaceans¹⁰⁷.

Table 1 Log likelihood values a	ind parameter estimates of d	ifferent models by PAML. $2\Delta\ell$: d	ouble likelihood difference			
Model	h L	Estimates of parameters	Models compared	2∆ℓ	P value	Sites positively selected
Branch models (A) M0: oneratio (B) free ratio	-3268.299851 -3249.845099	(1) = 0.08625 variable (1)	A vs B	88.318236	2.9×10 ⁻¹¹ **	
		by branch: (Human0.0509, (Cow0.1447, ([((Bottlenose_ di()butlenose_ dolphin0.3848, Pilot_whde 0.1284)0.5365, Finles_porpoise 0.000110.0001.				
		Baiji(0, 3906) 0,0001 (Minke_ whale0.7312, (Sei_whale0.0001, Brydes_whale 0.000110.0991) 0.234110.09111 0.0506 (Mouse 0.1079 Rati0.09881				
(C) all branches	-3499.431148	0.0558 $\omega = 1$	A vs C	462.262594	$1.5 \times 10^{-102**}$	
have same $\omega = 1$ (D) Two ratio	-3266.219683	0.0 = 0.07497	A vs D	4.160336	0.041*	
(branch cetacean) (E) neutral model	-3294.004217	0.1 = 0.1390/0.00 = 0.0705	D vs E	55.569068	9.0×10 ^{-14**}	
(F) Two ratio	-3265.523929	0.1 = 1 0.0 = 0.08178 0.0 = 0.08273	A vs F	5.551844	0.018*	
(G) neutral model (branch baiii)	-3266.393028	0.0 = 0.37322 0.0 = 0.08171 0.1 = 1	F vs G	1.738198	0.187	
(B) Two ratio	-3267.143786	00 = 0.08754 00 = 0.00010	A vs H	2.31213	0.128	
(I) neutral model (branch YFP)	-3273.394515	$\omega 0 = 0.08749$ $\omega 0 = 1.00000$	H vs I	12.501458	0.0004	
Sife models M3(k = 3)	-3247.386350	p0 = 0.63514 p1 = 0.33045 p2 = 0.03441	MO vs M3 (k = 3)	41.827002	1.8×10 ⁻⁸ **	
M3(K = 2)	-3247.617742	00 = 0.00000 01 = 0.21227 02 = 0.82386 p0 = 0.79369 p1 = 0.20631 00 = 0.01869 01 = 0.39995	M0 vs M3 (k = 2)	41.364218	1.0×10 ^{-9**}	

Table 1 Continued						
Model	h L	Estimates of parameters	Models compared	$2\Delta\ell$	P value	Sites positively selected
M7: beta	-3247.510906	p = 0.19794 2 - 173773				
M8: beta & w	-3247.505476	p0 = 0.29513 p0 = 0.29684 p = 0.20684 p = 1.89985 p1 = 0.00487	M7: beta vs M8: beta &	0.01086	0.995	
Mla	-3251.362169	p0 = 0.93299 p1 = 0.06701 m0 = 0.05006				
M2a	-3251.362169	p0 = 0.93299 p1 = 0.05530 p2 = 0.01172 m0 = 0.05006 m1 = 1.02 = 1	Mla vs M2a	0	-	
Branch- site models (Cetrecon)						
Model A	-3246.391824	0.0 = 0.04693 0.1 = 1.00000 0.2 = 3.14931 p0 = 0.92209 p1 = 0.05334 p2a = 0.02322 p2b = 0.001322	M1a vs Model A	9.94069	0.007**	154, 264, 328
Model A null	-3247.455370	0 = 0.04365 0 = 0.04365 0 = 1 0 = 1 0 = 0.88482 0 = 0.88482 0 = 0.05560 0 = 0.05560	Null model A vs model A	2.127092	0.144	
Model B	3243.468657	$p_{20} = 0.00352$ 00 = 0.02018 01 = 0.39328 02 = 2.86277 $p_{0} = 0.80077$ $p_{1} = 0.17119$ $p_{20} = 0.02310$ $p_{20} = 0.002310$	M3 (K = 2) v_s Model B	8.29817	0.016*	154, 264, 328
Branch- site models (Baiji)						



Table 1 Continued						
Model	hL	Estimates of parameters	Models compared	2Δ <i>€</i>	P value	Sites positively selected
Model A	-3247.955988	$ \begin{array}{l} \begin{array}{l} \begin{array}{l} \begin{array}{l} \begin{array}{l} \begin{array}{l} \begin{array}{l} \begin{array}{l}$	M1a vs Model A	6.812362	0.033* 7	, 149, 331
Model A null	-3247.956900	p2b = 0.01881 c00 = 0.04425 c01 = 1 c02 = 1 p0 = 0.63832 p1 = 0.04833 p2a = 0.29129	Null model A vs model A	0.001824	0.966	
Model B	-3243.587111	$\begin{array}{l} \begin{array}{l} \begin{array}{l} \begin{array}{l} \begin{array}{l} \begin{array}{l} \begin{array}{l} \begin{array}{l} $	M3 (K = 2) vs Model B	8.061262	0.018*	
Branch- site models //ED)						
Model A	-3251.362172	0.0 = 0.05006 0.1 = 1.00000 0.2 = 1.000 0.2 = 1.000 0.2 = 0.000 0.1 = 0.06701 0.2 = 0.000	M1a vs Model A	6×10 ⁻⁶	0.9999	
Model A null	-3251.362169	$p_{20} = 0.05006$ $p_{10} = 1$ $p_{20} = 0.05006$ $p_{20} = 0.03832$ $p_{1} = 0.06701$ $p_{20} = 0.000$	Null model A vs model A	6×10 ⁻ °	8 66.0	
Model B		pzb = 0.000 mol = 0.40446 mol = 0.40446 mol = 0 pol = 0 pol = 0 p2a = 0.79236 p2b = 0.20764	M3 (K = 2) vs Model B	2.307064	0.315	
(**': p < 0.01 , highly significant;	05, significant).					





Figure 3 | SLAC-inferred substitutions by site and detection of the positive destabilizing selection of the amino acids that were picked up by the sliding windows (windows size = 20) using TreeSAAP3.2. (A) The number of synonymous and non-synonymous substitutions on each codon were counted. For counting substitutions, the shortest evolutionary paths are assumed, and when multiples exist, the average over all is taken. Fractional counts are due to averaging over multiple pathways when multiple nucleotides are substituted along a single branch. (B) Regions above the z-score of 3.09 were significantly different than neutrality. Five types of significant physicochemical amino acids changes were identified and noted in different colors. (C) The clustered radical amino acid shift regions were correlated with aspects of protein 3D structure.





Figure 4 | Topology and structure of the urea transporter superimposed with the multiple sequence alignment (MSA). (A) The *Bos taurus* UT-B trimer chain A was superimposed with MSA: (*a*) viewed from the extracellular side; (*b*) viewed from within the plane of the membrane. The amino acids of the monomer are colored by their conservation grades using the color-coding bar, with turquoise through maroon, indicating variable through conserved amino acid positions. (*a1*) cartoon representation of mapped UT; (*a2*) 15 variable residues were detected and presented using a space-filled model, the letter abbreviations ahead of the site-numbers are the corresponding *Bos taurus* UT-B amino acid, the letter abbreviations behind the site-numbers are the variable amino acids of the MSA; (*a3*) the variable and conserved sites were both presented using a space-filled model. (B) The superimposed dvUT viewed from the extracellular side (a) and from within the plane of the membrane (b). (C) The multiple amino acid sequence alignment of UT-A2. The colored bars indicated the location of intramembrane helices in the UT-A2 structure according to the superimposed dvUT and UT-B. The red-boxed residues correspond to the signature sequences of UT. The black-, petunia-, and pink-boxed residues correspond to the signature sequences of UT. The cyan- and orange-boxed residues are the selected sites tested by Consurf and other evolutional assessment tools (treeSAAP, PAML and DataMonkey). (D) Annotated topology of the urea transporter with the evolutionary selective amino acid sites. The red dots represent the sites selected by PAML, DataMonkey, ConSurf and TreeSAAP.





Figure 5 | **Specific amino acid changes of the UT-A2 protein.** (A) Stereo view of residues in the selectivity filter S_o and S_i regions of UT-A2. (B) Specific amino acid changes among UT-A2s of the mammals. (C) Stereoviews of the mutation sites 264, 154 and the possible correlation sites.

$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	CodonBranch or clade3Baleen whale7Baiji7Baiji12Marine toothed w59Minke whale65Minke whale86Cow100Cetartiodactyla100Baleen whale149Baiji151Bottlenose dolphir151Bottlenose dolphir151Bottlenose dolphir200Cetacean	AA change $G u \rightarrow lys$ $G u \rightarrow Asp$ $I e \rightarrow Val$							higherit
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	 3 Baleen whale 7 Cow 12 Marine toothed w 59 Minke whale 65 Minke whale 65 Minke whale 86 Cow 100 Cetartiodactyla 108 Baleen whale 149 Baiji 151 Cetacean 151 Cetacean 200 Cetacean 	$\begin{array}{l} G \upsilon \rightarrow lys\\ G \upsilon \rightarrow Asp\\ Ile \rightarrow Val \end{array}$	ConSurf (Bayesian and LG)	Datamonkey (MEME)	PAML (MB-M3 and MA-M1a)	TreeSAAP	Domain	property	Category(+/-shift)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	7 Cow 59 Marine toothed w 57 Marine toothed w 65 Minke whale 86 Cow 100 Cetartiodactyla 108 Baleen whale 149 Baiji 151 Bottlenose dolphir 151 Cetacean 154 Cetacean 200 Cetacean	$Glu \rightarrow Asp$ Ile $\rightarrow Val$				+	N-terminus	Hu	8(+6.52)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	7 Baiji 12 Marine toothed w 59 Minke whale 65 Minke whale 86 Cow 100 Cetartiodactyla 108 Baleen whale 1149 Baiji 151 Bottlenose dolphir 154 Cetacean 154 Cetacean 150 Cetacean	lle → Val				• +		<u>؟</u> کمـ	6 (+0.72)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	 12 Marine toothed w 59 Minke whale 65 Minke whale 86 Cow 100 Cetartiodactyla 108 Baleen whale 149 Baiji 151 Bottlenose dolphir 154 Cetacean 200 Cetacean 					+	N-terminus	pK,	8 (+0.96)
	 59 Minke whale 65 Minke whale 86 Cow 100 Cetartiodactyla 108 Baleen whale 149 Baiji 151 Bottlenose dolphir 154 Cetacean 200 Cetacean 	hale Gly→Asp		+		+	N-terminus	.uŤ	6(-0.38)
$ \begin{array}{cccccc} 6.5 & \text{Minke whole} & \overline{\mathrm{Min}} \lor \mathrm{Volc} & \mathrm{Hin} \lor \mathrm{Vol} & \mathrm{Hin} \lor \mathrm{Volc} & \mathrm{Hin} \lor \mathrm{Volc} & \mathrm{Hin} \lor \mathrm{Volc} & \mathrm{Hin} & Hin$	 A5 Minke whale 86 Cow 100 Cetartiodactyla 108 Baleen whale 149 Baiji 151 Bottlenose dolphir 154 Cetacean 200 Cetacean 	Asp → Gly				+	Pa	шŤ	6(+0.38)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	86 Cow 100 Cetartiodactyla 108 Baleen whale 1149 Baiji 151 Bottlenose dolphir 154 Cetacean 200 Cetacean	Thr → Val	+	+		+	Pa	N_{s}, R_{α}	6(+1.81), 7(+4.54)
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	 100 Cetartiodactyla 108 Baleen whale 149 Baiji 151 Bottlenose dolphit 154 Cetacean 200 Cetacean 	Val → Ile	+			+	Tla	pK	8(-0.96)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	 Baleen whale Baiji Baiji Bottlenose dolphit Cetacean Bottlenose dolphit Cetacean Cetacean 	$IIe \rightarrow VaI$	+			+	T2a	pK,	8(+0.96)
$ \begin{array}{ccccccc} 149 & \text{Baiji} & \text{Val} \rightarrow \text{Abical loop} & \text{Pai, E} & \text{Paical loop} & \text{Pai, Paical loop} & \text{Paical loop}$	149 Baiji 151 Bottlenose dolphir 154 Cetacean Bottlenose dolphir 200 Cetacean	lle → Ser				+	T2a	N _s , B _r , R _F , P _c ,	6(-1.98), 6(-0.33),
$ \begin{array}{ccccccc} 149 & \text{Boili} \\ 151 & \text{Boillenose dolphin} & \text{Val - Ile} \\ 154 & Cencean control of the set of the set$	 149 Baiji 151 Bottlenose dolphir 154 Cetacean 200 Cetacean 							pK', F, R $_{\alpha}$, H _t , P _t	6(-10.2), 6(+0.68), 71+0 851 71+0 371
	149 Baiji 151 Bottlenose dolphir 154 Cetacean 200 Cetacean								8(-5.26), 7(-3.08), 8(-5.26), 7(-3.08),
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	 149 Baiji 151 Bottlenose dolphir 154 Cetacean 200 Cetacean 								8(+0.96)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	 151 Bottlenose dolphir 154 Cetacean Bottlenose dolphir 200 Cetacean 	Val → Ile				+	T4a	pK'	8(-0.96)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	154 Cetacean Bottlenose dolphir 200 Cetacean	Val → Ile	+			+	T4a	pK'	8(-0.96)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Bottlenose dolphir 200 Cetacean	Met → Val	+		+ +		T4a		
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	200 Cetacean	Val → Met					:	:	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$:	$Thr \rightarrow Lys$	+			+	Apical loop	рН <u>.</u> 	6(+4.08), 6 (-0.4)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	baili C .	Lys → Ihr	-				-	рН _і , Е,	6(-4.08), 6(+0.4)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	200 Ceracean	Ald → Ser	÷			+	Apical loop	rα, rc, rt	0(-0.03), 0(+0.03), 6(±0.77)
$ \begin{array}{ccccccc} \mbox{Mouse} & \mbox{Matrix} & \$	206 Rodentia	$Ser \to Ala$	+			+	Apical loop	P_{∞} , P_{c} , P_{t}	6(+0.65), 6(-0.63),
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Mouse	$Ala \to Thr$				+		P ₂₀	6(-0.59)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Rat	Ala → Val							
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	207 Cetartiodactyla	Ala \rightarrow Ser	+			+	Apical loop	P_{∞} P_{c} , P_{t}	6(-0.65), 6(+0.63),
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	208 Cow	Met → Val	+				Apical loop		
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	216 Cow	Val → Ile	+			+	Apical loop	pK'	8(-0.96)
$ \begin{array}{cccc} \mbox{Marine toothed whale} & \mbox{Thr} \rightarrow \mbox{Ala} & \mbox{Ala} & \mbox{Thr} \rightarrow \mbox{Ala} & \mbox{Ala} & \mbox{Ala} & \mbox{Met} \rightarrow \mbox{Ile} & \mbox{H} & \mbox{H} & \mbox{H} & \mbox{T2b} & \mbox{pK'} & \mbox{Ala} & \mbox{Ala} & \mbox{Ala} & \mbox{Ala} & \mbox{Ala} & \mbox{H} & \mbox{H} & \mbox{H} & \mbox{Ala} & \mbox{H} & \mbox{H} & \mbox{H} & \mbox{Ala} & \mbox{H} & \mbox{H} & \mbox{Ala} &$	264 Baleen whale	$Thr \rightarrow Ala$	+		+ +	+	TŹb .	.ط	6(+0.59)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Marine toothed w	hale Thr→Ala				+			
312CetaceanIle \rightarrow Val+T4bpK'328CetaceanMet \rightarrow Val++T4b28Pilot whaleVal \rightarrow Ile++pK'	265 Cow	Met → Ile	+			+	T2b	pK'	8(-0.92)
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	312 Cetacean	$ e \rightarrow Va $				+	T4b	pK'	8(+0.96)
Pilot whale Val → Ile + pK'	328 Cetacean	Met → Val	÷		+ +	+	T4b		
	Pilot whale	Val → Ile				+		pK'	8(-0.96)
376 Cow and Rodentia Tyr → His + C-terminus F	376 Cow and Rodenti	ז Tyr → His	+			+	C-terminus	щ	4(-0.21), 6(-0.32)



Property	Description
Ρ _α	Decreases in α-helical tendency broadens the outer surface area of a helix, and allows a more flexible surface area contact with neighboring protein constituents, whereas increases have the opposite effect ⁷⁹ .
Ns	N _s is number of residues surrounding a residue within the effective distance of influence ⁷³ .
Br	Buried nature of residues within a globular protein ⁷³ .
R _F	R_F is characteristic migration rate in a solvent-absorbent system ⁷³ .
Pc	The probability of a residue being in a coil ⁷³ .
рК′	pK' is a biochemical amino acid property that is a measure of the net charge in the overall charge resulting from the ionization of carboxyl groups ⁷³ .
рН _і	"pHithe isoionic point of the free amino acidincludes the ionizable character of either the sidechain or amino group plus the carboxyl group of the molecule" ¹⁰⁵ .
F	This parameter relates to the amount of displacement of any element with its distance from the centroid of the protein. Each protein is characterized by its specific internal residue distribution and associated fluctuational situation ¹⁰⁶ .
V°	Partial specific volume is the reciprocal of density ⁷³ .
Esm	Sum of all non-bonded energy between atoms within ten residues along a polypeptide chain ⁷³ .
R _α	Ratio of accessibility of water "from a hypothetical extended state to the native fold state", the increase in this value suggest that proteins could have become bulkier and allowing more space for active site formation ⁷³ .
H,	''-intrinsic hydrophobic character of the (amino acid) side chain- ^{''73} .
E,	Sum of short-, medium-, and long-range non-bonded energies ^{73,74} .
Pt	P_t refers to the ability of a specific amino acid to contribute to or initiate a turn ⁷³ .

cysteine (C337) in UT-A2, respectively (Fig. 5A). I228 in UT-B was replaced by value in UT-A2 (V232), which is identical to V188 in dvUT. In the S_i region, F80 (G115 in UT-B) and L129 in dvUT shifted to tyrosine (Y123) and phenylalanine (F175) in UT-A2, respectively (Fig. 5 A).

Although these ligand binding sites in the selectivity filter changed among the UT-A2, dvUT and UT-B proteins, residues in the UT-A2 in all of these mammals were in full accord (Fig. 4C). To determine the molecular adaptations in UT-A2, TreeSAAP was further used for identifying and characterizing adaptation in terms of shifts in the physicochemical properties of amino acid replacements. Eight amino acid properties (alpha-helical tendencies (P_{α}), bulkiness (B_{l}), equilibrium constant-ionization of COOH (pK'), partial specific volume (V°), power to be at the C-terminus (α_c), solvent accessible reduction ratio (R_{α}) , total non-bonded energy (E_t) and turn tendency (P_t)) were found to be undergoing positive-destabilizing selection (p < 0.001). To maximize accuracy, the three properties B_{l} , V° and α_{c} that tend to generate false-positive results were eliminated from the results⁵⁴. The sliding window graphs illustrate the spatial dynamics of selection across the primary sequence of the UT-A2 gene (Fig. 3B). Regions clustering around the radical physiochemical shifts (categories 6, 7 and 8, p < 0.001) from natural amino acid replacements were analyzed and visualized. The correlated 3D structure of the urea transporter showed the influenced protein regions that have been historically affected by selection on specific amino acid properties (Fig. 3C). The five amino acid property changes involve the protein secondary structure (P_{α} and P_t), ionization constants (pK'), nonbonded energy (E_t) and solvent accessibility (R_{α}). P_{α} and P_t are prone to be clustered at the apices of T5a, Pb and the loop between these helices. Et tends to be clustered at the apices of T5a and the loop affecting the non-bonded energies. Both of these radical changes are in the entrance region of the urea conduit. The shifts in pK' clustered at the surface of the protein and the entrance of the UT pore, whereas R_{α} is located at the exit region of the UT pore.

Among the non-synonymous substitutions, codons under positive-destabilizing selection were identified from 20 amino acid properties (the 11 exchanges that performed the worst were eliminated) in TreeSAAP and were summarized in Table 2. The major physicochemical amino acid properties that yielded positive analytical results are summarized in Table 3. The cetacean clade was detected by treeSAAP to have substitutions on sites 200, 205 and 312. Site 200 is located on the loop between T5a and Pb. The substitution (Thr \rightarrow Lys) on this site could increase the protein pH_i and decrease the E_t property. The amino acid shift from Ala to Ser at site 205 and the substitutions at sites 206 and 207 are located on the apical loop of the periplasmic side between T5a and Pb, which could influence the conformation of the protein (decrease P_{α} and increase P_{c} and P_{t}). The substitution at site 312 (Ile \rightarrow Val), which is located on the T4b helix cytoplasmic side, could increase the pK'. Baleen whales underwent positive-destabilizing selection at sites 3 and 108. Site 3 in the N-terminus (Glu \rightarrow Lys); pHi was increased, whereas site 108 (Ile \rightarrow Ser), which is located on the cytoplasm side of T2a helix, could cause many radical amino acid property changes (N_s, B_r, R_F, P_c, pK', F, R₇, H_t and P_t). We found that freshwater YFP and baiji shared two identical amino acids at sites 264 and 12, whereas marine toothed whales have substitutions of Thr to Ala and Gly to Asp. These changes could increase the P_{α} of T2b and decrease the E_t of the Nterminus. There are some substitutions pertaining to baiji (sites 7, 149 and 200) and minke whale (sites 59 and 65), which are different from their counterparts. The properties of the amino acid at site 200 in the baiji displayed a decrease in pH_i and an increase in E_t. Substitution at site 7 (Ile \rightarrow Val), which is in the N-terminal region, could increase the pK', whereas substitution at site 149 (Val \rightarrow Ile), which is in the T4a helix on the periplasmic side, could also decrease the pK'. In this study, we also detected some selection sites pertaining to baleen whale UT-A2 (sites 3 and 108). At site 59, substitution (Asp \rightarrow Gly) increased the E_t. At site 65, substitution (Thr \rightarrow Val) increased the N_{s} and $R_{\alpha}\!.$ The radical amino acid changes were marked on the multiple sequences alignment (MSA) (Fig. 4C), the topology (Fig. 4D) and the 3D structure of the urea transporter (Fig. 5B).

Discussion

The UT is a type of integral membrane protein that facilitates the diffusion of the osmolytic urea and plays important roles in *in vivo* water equilibrium regulation^{26,28,35,51,52,55,56}. In the present study, we looked for evidence of adaptive molecular evolution of the osmoregulation-correlated gene UT-A2 in the cetacean lineage (marine and freshwater), relative to their terrestrial counterparts. We cloned the UT-A2 gene from kidneys of two freshwater YFP and bioinformatics analyses were performed. The gene was found to be highly conserved (87–99% identity) and undergoing purifying selection. However, the cetacean branch has a significantly higher ω ratio relative to terrestrial mammals. Different selection sites in the cetacean branches were picked up with several complementary approaches, including d_N/d_s (PAML, Datamonkey), MM01 (TreeSAAP) and Rate4site (ConSurf) (Table 2). Furthermore, we

found that the UT-A2 individual protomer has an overall fold similarity with dvUT and UT-B (Fig. 4), although several variants occurred in the functional S_o and S_i regions of the UT selectivity filter (Fig. 5A). Finally, the radical amino acid changes (particular substitutions and cluster regions properties) were superimposed on the 3D structure (Fig. 3C and 5B, C), and then the molecular adaptations that may lead to the potential functional differences were discussed based on the urea permeation mechanisms of dvUT, UT-B and the urea channel of *Helicobacter pylori*^{51,52,57}.

In the evolutionary process, molecular adaptations that optimize the organism's survival in a given habitat can be accumulated and inherited. Effectively neutral nucleotide substitutions, and either purifying selection (maintaining functions) or positive selection (favoring new functions), play major roles in evolution at the molecular level. To understand what evolutionary forces drive protein evolution and why protein sequences differ in different species, several methods have been proposed to detect selection or adaptation since the genetic code was discovered in the 1960s. The most common or broadly accepted approach is the family of methods that contrast the rates of nonsynonymous and synonymous substitutions (d_N/d_S) , also known as Ka/Ks or ω); these methods have enjoyed a great deal of success^{58,59}. However, because most substitutions are expected to be neutral or deleterious⁶⁰, adaptive changes will be difficult to find using ω ratios, particularly if we only look for ratios greater than one⁶¹. In fact, relatively few mean ω estimates of genes in cetaceans are significantly greater than 1 (e.g., only 228/10,000 were found in a bottlenose dolphin and nine other genome comparisons⁶² and only 24/10,423 were found in the common branch of the baiji and dolphin versus the other four genomes⁶³). Additionally, for a conserved gene, sometimes even a single amino acid change can be adaptive if it provides a biochemically advantage relative to contemporaneous alternatives, which seems not sensitive enough for the d_N/d_S method^{64–68}. In the analyses of nucleotide substitution patterns in this study, the ω ratios tested as significantly less than 1 (approximately 0.1), showing that a strong purifying selection plays a central role in the evolution of UT to keep its important functions in urea conduction. Furthermore, to find the drivers of adaptation to the different levels of osmoregulatory challenges (different urea levels generated by protein catabolism and infrequent access to water), branch-specific, site and branch-site models were conducted to explore the variation of evolution rates in specific sites in cetaceanspecific, baiji and YFP lineages. However, the results showed weak evidence of positive selection using this pattern (Table 1). The tworatio model detected that the baiji and cetacean-specific branches have a significantly higher evolutionary rate than the background, indicating that the constraint on functional change of UT-A2 in the cetacean and baiji branches has been relaxed. Although LRT tests of Model B-M3 (K = 2) (p < 0.05) or MA-M1a (p < 0.01) picked up several positively selected sites in the cetacean and baiji clades, the result of the stringent testing (test 2, MA vs MA null) is not significant⁶⁹.

As a reaction to potential bias in d_N/d_S patterns^{64,66-68}, we applied a physicochemical property model (MM01 methods) to the possible molecular adaptation of UT-A2 among cetaceans, which has been proven to be a sensitive way to detect the subtle effects of positive selection^{64,66,68,70}. Because the 3D structure of a protein determines its function, the molecular adaptations resulting from changes in physicochemical amino acid properties were discussed based on the revealed 3D structure and mechanisms of the UT analogs. In the analysis of this pattern, five radical amino acid properties changes that involve protein secondary structure (P_{α} and P_t), ionization constants (pK'), non-bonded energy (E_t) and solvent accessibility (R_{α}) were detected using the sliding window mode (Fig. 3B, C). Using the analytical software TreeSAAP, we also characterized several positive destabilizing selections (selection that results in radical structural or functional shifts in local regions of the protein) that correlated with 12 biochemical shifts (Table 2), which represented the unambiguous signature of molecular adaptation⁶⁴.

In the 3D structure characterization, we found that the individual protomer of UT-A2 has the same overall folding pattern as do dvUT and UT-B (Fig. 4), containing 6 pairs of well-placed α -helices, which form the urea conduction pore. The conserved amino acids of the S_o region (Q231, V232, G234 and C337), S_m region (L127, T176, L291 and T338) and S_i region (V68, Q 67, Y123 and F175) are brought together to form the selectivity filter that helps urea to travel through the protein; these are stick-presented in Fig. 5B. When we compared the UT-A2 selectivity filter with that of dvUT, several residue changes in the S_o and S_i regions (Fig. 5A) were observed, indicating the potential differences in urea permeation regulation of these two proteins. The shift from F (phenylalanine) to G (glycine) on site 234, minus the planar, aromatic side chains, which were theorized to fix the orientation of the urea molecules entering the pore⁵¹, as in coin access to "coin slots"29, may indicate a structural relaxation of the UT-A2 entrance and supply enough space for urea passing through when there is a higher urea concentration in the periplasm. The shift from E (glutamic acid) to Q (glutamine) on site 231, along with a change from the carbonyl to amide group, which retains significant polarity and is a rich source of hydrogen bonds, would make it easier to bind urea and urea-water compounds. These two site shifts may functionally increase the rate and efficiency of urea conduction through enlargement of the urea acceptability of the entrance. The hypothesis could be verified by structural and functional studies of UT-B⁵¹, which displays nearly the same amino acids shifts. In the S_i region, F80 and L129 shifted to Y119 and F175, respectively, adding a hydroxyl and phenyl group. The hydroxyl of Y119 could be a binding site for urea and the hydrophobic phenyl group likely helps the coinshaped urea molecule to exit in good order, which may be important for osmotic changes in the cytoplasm. These two sites likely also increase the efficiency of urea discharge. To conclude, compared to bacterial dvUT, the urea conduction pore of UT-A2 seems to have a structurally relaxed entrance and a much more orderly exit, as well as enhanced urea binding capacity; therefore, it could promote the rate of urea conduction, which might be a demand caused by the higher efficiency of protein catabolism in mammals.

After superimposing the molecular adaptation sites onto the topology and 3D structure (Fig. 4, 5) we found that the positive-destabilizing selection in cetaceans mainly occurred at the N-terminus, the periplasmic side of the helices and loops, and the outside helices (T2a, b and T4a, b) of the UT-A2 protein (Table 2). We found that most of the radical amino acid changes are located on the regions that were exposed to the solvent, versus those buried in the interior of the folded structure. This may indicate that the interior residues are subject to stronger constraints against the exposed residues, which is consistent with the sliding-window amino acid properties cluster analyses (Fig. 3C). Conant suggested that the phenomenon of "neutral evolution on mammalian protein surfaces" was due to the selection to Ne (the variation in effective population sizes)⁷¹. However, TreeSAAP, the method that we used for detecting the amino acid changes, is not based on mutation frequencies within populations, but upon the sequences themselves⁷². Therefore, in this study, we assume that a few residues (e.g., the interacting surface) were under selective constraint or directional selection. The sliding-window analyses indicated that the entrance region of UT-A2 has undergone P_{α} , Et and Pt property changes (Fig. 3C), which may influence the accessibility of protein interfaces and interactions with other molecules^{73,74}. Previous studies suggested that residues in the entrance region of urea channel are responsible for gathering urea-water compounds and gating urea entrance through the pore⁷⁵. In the absence of an experimental loop structure of a urea transporter or urea channel, the function of the loop was not yet well illustrated^{51,52,75}. Mutation studies of the Helicobacter pylori urea channel have shown that the histidine residues on the periplasmic loops are essential for pH gating⁷⁶,

suggesting that the opening and closing of the channel might be controlled by reversible folding of one or both loops into the periplasmic vestibule of the channel⁷⁵. In this study, several radical amino acid substitutions occurred on sites 200, 205, 206 and 207, which are located on the periplasmic loop (Fig. 5B). Residue 200 (Thr \rightarrow Lys, hydroxyl amino acid, lower side chain flexibility and polar to methylene groups crowned with a amine nitrogen, higher side chain flexibility, ionic interaction mode, lower hydrophobicity and positive property⁷⁷) could increase the protein pH_i (Tables 2, 3) and make the isoelectric point more basic, decreasing its tendency to aggregate at or below neutral pH due to electrostatic interactions⁷⁸. This may have a role similar to histidine residues on the loops of the Helicobacter pylori urea channel, affecting the gating of the UT-A2 protein. Furthermore, the codon substitution decreases the Et property, affecting the short-, medium- and long-range interactions with atoms in the same and other molecules^{73,74}. Site 205 (Ala \rightarrow Ser, to 3 potential side chain H-bonds and polar) is also located on the apical loop of the periplasm side between T5a and Pb, which could decrease P_{α} and increase P_{c} and P_{t} (Tables 2, 3), resulting in increased accessibility at protein interfaces^{73,79}. Together with the Ser at site 206, 207, it provides a relatively larger polar surface binding the water-urea compounds⁸⁰, and could potentially gather more urea to enter the urea conduction pore. Site 328 (Met \rightarrow Val, shift to lower side chain flexibility and slight higher hydrophobicity) is also located on the region adjacent to the entrance to the pore (Fig. 5B). Met and Val have different restrictive capacities ($\Delta S^{contact}$) from other hydrophobic residues, which could affect the conformation and function of a protein⁸¹. In the α -helix, aliphatic residues (Leu, Val, Ile) more strongly affect each other, whereas Met residues are less affected by other residues⁸¹. Therefore, the apical side of the T4b helix of cetaceans seems to be more restricted, and may help to provide a wider entrance space for the urea transporter, affecting the protein functional activity. In previous studies, shifts in the Met and Val residues have been found to affect some genes' activity and function^{82,83}. Other types of amino acid changes in the cetacean occurred on the surface of UT-A2, such as sites 154 and 264. Site 154 also had a shift of Met to Val in the cetacean branch, along with a decrease of side chain flexibility and an increase in hydrophobicity⁷⁷, which has different restrictive capacities ($\Delta S^{contact}$) and could affect the conformation and function of a protein⁸¹. When we mapped the site on the 3D structure of UT-A2, we found that the region, where the site occupied, was across from the S_m region of the urea conduction pore (Fig. 5C), which may indirectly affect the urea conduction through the changes in $\Delta S^{\text{contact}}$ or through conformation changes. Site 264, which is located on the T2b helix, has an amino acid shift of Thr to Ala (change to limited side chain flexibility, no H-bonds, higher hydrophobicity and nonpolar⁷⁷) in cetaceans except for the freshwater YFP and baiji. These changes could increase the P_{α} of T2b (Table 2), leading to a longer, more rigid alpha helix with a more stable lipid raft composition, which, in turn, may alter its interactions with amino acid motifs and its efficiency of combination with other substances^{84,85}; thus, it may raise the efficiency of UT-A2 of marine cetaceans compared with the freshwater cetaceans. The region was across from the S_m region of the urea selectivity filter regulating the rate of urea conduction, which may increase the hydrophobicity of the S_m region, and raise the energy of dehydration, affecting urea conduction (Fig. 5C). The third type of amino acid changes occurred on the N-terminus of UT-A2. On site 3 (Glu \rightarrow Lys, containing a carboxylic acid group and negative charge to containing amine nitrogen and positive charge⁷⁷), the pH_i was increased, thus altering the interaction with electrostatic particles and the surrounding residues or subunits^{73,78} (Tables 2, 3). On site 12, marine toothed whales have the substitution of Gly to Asp (a change to moderate side chain flexibility, ionic interaction mode, four potential side chain H-bonds, lower hydrophobicity and polar properties⁷⁷), which can decrease the Et, affecting the interaction of the amino acid with the surrounding

atoms, residues, or subunits73,86. However, in the absence of functional studies of the N-terminus of the urea transporter or urea channel, we cannot deduce the role of amino acid changes within this region. Only when mutagenesis studies have been accomplished can this issue be illustrated. However, in this study, we predict some potential mutations that may influence protein function, which may help narrow down where scientists should concentrate future research. Collectively, we found conservation of the urea selectivity filter of UT-A2, despite some changes in the S_o and S_i regions, versus bacterial dvUT, which may correlate with the adaptation of highlevel protein catabolism in mammals. Also the radical amino acid changes tend to occur on the surface of the protein and may generate mildly advantageous effects for adapting to a changed environment. The selection may provide cetaceans a relaxed or wider urea channel entrance or increased S_m hydrophobicity to enhance the efficiency of urea conduction. However, in the absence of mutagenesis experiments, the conclusions drawn from the simulations remain speculative.

Based on these selection-pattern analyses and the potential functional characterization, we suspect that the period of adaptive evolution and rapid amino acid substitution may only have occurred for a short time period, followed by a long period of selective constraint (purifying selection) on the improved protein, which would obscure the selective processes and make it difficult to detect positive selection using the d_N/d_S pattern⁸⁷. The similar evolutionary rates in the branches of freshwater and marine whales suggest that baiji and YFPs may have similar molecular UT adaptations to their marine counterparts, although they live in freshwater and generate relatively diluted urine²⁵. In fact, the higher level urea transporting capacity that they inherited from their marine ancestor is not a burden to their physiological functions, as they can easily obtain sufficient fresh water from the environment to dilute the urine and eliminate the redundant urea. Physical and biochemical analyses of plasma and urine showed that both baiji and YFP excrete urine and that the urea concentration in the animals is still higher than the concentration in plasma^{25,88}, indicating active transport of urea in the kidney. The physical experiment (fresh and seawater ingestion) on bottlenose dolphins has shown that if a marine toothed whale were provided with sufficient freshwater, it would also excrete relatively diluted urine12. When a certain amount of deionized water was fed to the bottlenose dolphin, the urine osmolality could decrease to 813 ± 30 mOsm/kg, and the urea concentration in urine could decrease to 11-38 mEq/L, similar levels to those in YFP and baiji^{12,25,88}. The real challenge that the baiji and YFP face may be to produce more diluted urine (e.g. adaptation of water regulation through aquaporin). We believe that this hypothesis can be further confirmed, as UT-A2 sequences from marine finless porpoises are sequenced and the water channels are studied in detail. The other question is: why is the UT-A2 of YFP much more conserved with marine toothed whales than baiji, although baiji and YFP reside in similarly adapted osmotic conditions (similar osmosis, food and similar freshwater surroundings)? We attribute the dissimilarity to relaxation of purifying selection of UT-A2 in baiji. There are two aspects to purifying selection: (1) past selection works against highly deleterious mutations, which are eliminated quickly; and (2) ongoing selection works against slightly deleterious mutations, which may be relatively inefficient if the effective population size is small^{89,90}. Previous studies suggest that river dolphins, including baiji, escaped extinction and adapted to their riverine habitats during continental flooding in the Middle Miocene caused by the rise in global sea levels^{91–93}. Thus, we argue that baiji must have undergone reduced effective population sizes (N_e) that were influenced by oceanic events in their long-term fresh-water living conditions (approximately 21.5 Ma)⁹², and have experienced less efficient purifying selection. They are expected to accumulate more slightly deleterious mutations and thus to show a greater dissimilarity to their ancestors. The genome study of baiji also suggested that a bottleneck occurred near the end of the last deglaciation⁶³, which is consistent with our hypothesis. From another viewpoint, a change in effective population size can be distinguished from a change in selection intensity because a change in population size is expected to affect all genes along a lineage, producing a genome-wide effect on the ω ratio^{94,95}. In this study, a variance no greater than expected was observed between the d_N/d_S of the UT-A2 of the baiji lineage (0.39) and the baiji genome d_N/d_S $(0.32)^{63}$, also indicating that the relatively increased ω value was due to the confounding effects of population demographic history. However, YFP, the sole species of freshwater finless porpoise, entered the Yangtze River approximately 2.00-0.72 Ma along with the Quaternary glaciations⁹⁶⁻⁹⁸, and are considered to have diverged from their marine origin in the Yellow Sea. Until now, YFP was classified as a narrow-ridged finless porpoise species (N. asiaeorientalis), with the other marine subspecies of East Asian finless porpoise (N. a. sunameri)^{99,100}. If that were the case, it means that the subdivision and freshwater adaptation of the YFP was a recent historical event. Therefore the baiji have likely lived for a much longer period of time in fresh water and may have undergone slightly deleterious mutations that are reflected in their differential base substitutions, and even in the molecular divergence of their UT-A2.

It happens that there is a similar case, marine osmosis habitats, where the driving forces of UT-A2 are different between the baleen whale and toothed whale. One hypothesis is that differential feeding led to different urea concentration levels of their internal milieu along with different selective pressures on the UT gene. Marine toothed whales such as bottlenose dolphins and pilot whales feed mainly on marine fish, whereas baleen whales, such as Bryde's whale and sei whales, feed on invertebrates²⁰. Most invertebrates are osmoconformers, such that whales ingest seawater passively. Consequently, through food ingestion, relatively higher levels of urea concentration could be observed in the internal milieu of toothed whales^{10,14,25}. The physiological and biochemical analyses of cetacean plasma and urine also indicate the importance of feeding for osmoregulation¹⁴. This hypothesis is limited to explaining the different relaxed selection constraints that occurred on UT-A2s between minke whale and the sei/Bryde's whale, and between the baiji lineage and YFP lineage, as they feed on nearly the same osmotic foods and reside in the same aquatic surroundings²⁰. In this instance, a new or modified function is not necessary for the already adapted gene. The MEME results further indicate that some sites of marine toothed whale (site 12, empirical Bayes factor > 100) and minke whale (site 65, empirical Bayes factor > 1000) are under episodic diversification, indicating that the selective forces on UT-A2 are not constant across taxa. Therefore, there could be some transient periods of adaptive evolution masked by the prevalence of purifying or neutral selection on other branches. We argue that UT-A2 of marine cetaceans must have undergone different transitional periods of adaptation, influenced by these major oceanographic events, and that they are expected to have accumulated different mutations and thus to show a greater dissimilarity. In terms of potential positive selection, relaxation of purifying selection and episodic diversification in independent lineages of cetaceans in similar environments, we conclude that UT-A2s might have been periodically adjusted during or after cladogenesis and undergone periods of increased standing genetic variation, including mutations of small-to-modest effect that would allow adaptation to changing osmotic environments.

In summary, we found that a strong purifying selection plays a central role in the evolution of the urea transporter to maintain its important urea conduction function. We suggest that positive-destabilizing selection, relaxation of purifying selection and episodic diversification and play important roles in the process of cetaceans reinvading the sea and subsequently transitioning to different osmotic environments; this means that UT-A2 experienced standing genetic variation or mutations and has periodically adjusted during

or after cladogenesis. We suggest that the freshwater baiji and YFP have basically retained the molecular adaptations of their marine counterparts, although two common amino acid sites (sites 264 and 12) along with shifts of the properties P_{α} and E_t that differ from their homologous marine toothed whales were detected. We found that the UT-A2 protein has folding similarity with dvUT and UT-B, whereas some variants occurred in the functional So and Si regions of the selectivity filter. We also contribute to the knowledge of physiochemical and molecular adaptive evolution of UT-A2 by identifying several amino acid property shifts (pK', P_{α} , PH_i, E_t, et al.) that may generate mildly advantageous adaptations, which occurred at the entrance and surface of the UT-A2 protein. The conservation of amino acid residues within the selectivity filter of the urea conduction pore is likely necessary for urea conduction, whereas the radical amino acid replacements around the entrance and exit of the conduction pore could potentially affect the activity of the UT-A2 protein. Collectively, this study identified regions in UT proteins that are not as highly conserved as the selectivity filter region, but may play important roles in the activity of the UT protein. Further studies are necessary, however, to identify the function of N-terminus and to determine the effects of the radical amino acid changes.

Methods

Tissue collection. The kidney samples used for this study were collected from two freshly dead YFP (within 3 hours of death). One accidentally injured female YFP, with a body length of 128 cm, was found in the Shishou stretch of the Yangtze River; the other YFP, a male with a body length of 95 cm, was by-caught by a fisherman in the Honghu stretch. All possible rescue efforts had been strictly conducted following the Regulations of the People's Republic of China for the Implementation of Wild Aquatic Animal Protection (promulgated in 1993). Necropsy and sampling were conducted systematically in accordance with all ethical guidelines and legal requirements in China. The protocol of this study was approved by the Institutional Review Board of the Institute of Hydrobiology, Chinese Academy of Sciences.

Amplification and sequencing. Total RNA was extracted using TRIzol (Invitrogen), and then run on a 1.0% agarose gel to confirm its integrity and purity. First-strand cDNA was synthesized from the total RNA using a First Strand cDNA Synthesis Kit (Fermentas). The cDNA served as a template for PCR amplification, using the specific primers in Table S1 (the red bars in the schematic diagram Fig. 1A), which were designed on the basis of the sequences of UT-A2 cDNAs of several cetaceans and terrestrial mammals that were previously reported^{45,46}. The PCR amplification reaction conditions were as follows: initial denaturation at 94°C for 3 min, followed by 29 cycles of 94°C for 30 sec, 53°C for 30 sec, and 72°C for 75 sec, and ending with an additional 7 min at 72°C. Amplified PCR products were purified after running on an agarose gel and then cloned into a topTA2 vector (Toyobo). After extracting the plasmid DNA, nucleotide sequencing was conducted by Shanghai Majorbio Bio-Pharm Technology Company. Sequence data were assembled using the DNA Star Seq Man program and manually checked for accuracy.

Database searches for UT-A2 genes. We collected 10 publicly available UT-A2 coding sequences from the DDBJ (http://www.ddbj.nig.ac.jp/), NCBI (http://www.ncbi.nlm.nih.gov/), and Ensembl (http://www.ensembl.org/index.html) databases (Date of access: 01/09/2013) for analyses, including five marine cetaceans (*Globicephala macrothynchus*, GenBank Accession No. AY061881; Balaenoptera acutorostrata, AB266067; Balaenoptera borealis, AB266069; Balaenoptera brydei, AB266070; Tursiops truncatus, ENSTTRT0000007415), four terrestrial mammals (*Bos taurus*, ENSBTAT0000002712; *Homo sapiens*, NM_007baij163; *Rattus norvegicus*, U09957; *Mus musculus*, AF367359) and one river dolphin (*Lipotes vexillifer*, AUPI01117101.1)⁶³.

Phylogenetic Reconstruction. Nucleotide coding sequences and amino acid sequences of UT-A2 were aligned using CLUSTALX (Thompson, et al. 1997) and the BioEdit program (Hall, 1999) aided by manual checking. An 1194 bp alignment of 10 UT-A2 coding sequences was used for phylogenetic analyses. Alignments were unambiguous because these UT-A2 sequences contained no obvious insertions or deletions. Modeltest 3.7 (Posada and Crandall, 1998) and MrMTgui (Nuin, 2007) were introduced for the selection of the best-fitting evolution model and the Bayesian Information Criterion (BIC) was preferred as a model decision criterion. Then, phylogenetic tree reconstructions were performed using MEGA4 for MP and NJ analyses (Tamura, et al. 2007), and MrBayes-3.1.2 for Bayesian estimation (Ronquist and Huelsenbeck 2003). Both MP and NJ trees were evaluated using bootstrap testing with 1000 replicates.

Molecular evolutionary analysis. To investigate the possible role of positive selection on the evolution of UT-A2 gene in the cetacean clade, codeml (PAML) and DataMonkey were introduced. The codon substitution models were implemented in the PAML program with estimated transition/transversion rates and F3 imes 4codon frequencies algorithm (Yang, 2007). A likelihood ratio test (LRT) was used to compare the fit of null models and alternative models, and the significance of the LRT statistic was determined by using a χ^2 distribution. We applied various models, including branch-specific models, site-specific models and branch-site models to test the potential selection pressure acting on the UT-A2 gene. In these tests, the UT-A2 gene tree (Fig. 2) was introduced as the guide tree. In branch model testing, the LRT of the free-ratio model (an independent ω value for each branch) against the one-ratio model (a constant ω value across all branches) was performed to ensure variations in ω values among the branches. The two-ratio models were finally applied to test an *a*priori hypothesis and confirm selection pressure on cetacean-specific, baiji and YFP branches by setting a foreground ratio ω_1 while all other branches have the background ratio ω_0 , respectively. Also, the results were contrasted against a model of neutrality (fixed ω to 1). In the site model testing, M3 (discrete, fits k = 2 or $k = 3 \omega$ classes estimated from the data) against M0 (one average ω across all sites and all branches) was used to test for rate heterogeneity among amino acid sites; M7 (beta, which assumed a beta distribution of ω between 0 and 1) versus M8 (beta & ω , which added the extra $\omega > 1$) and M1a (neutral, assumes two sites classes $0 < \omega < 1$ and $\omega =$ 1) versus M2a (selection, adds an additional site class of $\omega > 1$) were used to identify the possible sites under selection. In the branch-site model testing, the cetaceanspecific branch, baiji and YFP were defined as the foreground while all the other branches were defined as the background, and then test 1 (modified model A versus site model M1a), test 2 (modified model A versus branch-site model A with $\omega_2 = 1$ fixed) and model B (ω_0 and ω_1 are estimated as free parameters) versus M3 (with two discrete classes of sites) were conducted to detect sites under positive selection along cetacean lineages. In the positive-selection site and branch-site model tests, the conservative Bayes Empirical Bayes (BEB) analyses were also implemented to test the possible selection sites and their posterior probabilities¹⁰¹. To compensate for local optima effects in the analyses, the program ran two times, starting with ω values both above and below one. The web interface DataMonkey (Pond and Frost, Datamonkey, http://www.DataMonkey.org, 2005. Date of access: 26/09/2013), which uses the HyPhy package as its processing engine, was implemented as a complementary approach to identify the codons and lineages under selection. First, the best-fitting nucleotide model (HKY85, AIC = 7058.196, p = 0.003) was automatically selected. SLAC (single-likelihood ancestor counting) analyses¹⁰² was used to estimate the number of synonymous and non-synonymous substitutions of each codon along the UT-A2 sequences. Exploratory analysis was also conducted by MEME¹⁰³ to assume that ω is not necessarily constant over time. To decrease the effect that purifying selection in some lineages might have to mask the signal of positive selection in others, and to measure the probability of observing a large d_N/d_S in random neutral data, high nominal α -levels were advised¹⁰². Here, the default cut-off value of P < 0.1 was chosen.

Amino acid properties and evolutionary conservation analyses. To detect significant physicochemical amino acid changes among the residues in the UT-A2 gene, selection tests were performed on the alignments and phylogenetic tree using TreeSAAP 3.2 (Woolley, et al. 2003), which is based on the MM01 mathematical model along with implemented modifications^{54,64,70}. In the program testing, universal genetic codes, the HKY85 evolution model and 31 quantitative biochemical properties were used to estimate adaptive selection (referred to as positive destabilizing selection, when the frequency of changes in magnitude categories 6, 7 and 8 were highly significant, z-scores > 3.09 and P < 0.001). Changes in magnitude (categories 6 to 8) that were also significant (p < 0.05) were first identified with an overall analysis of our data. Evaluation of several sliding window sizes showed that a window of 20 amino acids (incremented every single site) provided the best signal-tonoise ratio, and its statistical significance was determined using a Bonferroni correction ($\alpha = 0.001$; 99.9% confidence). The results of sliding window analyses were used to identify positive destabilizing selection regions for each property at a significance level of p < 0.001 (Z-Scores > 3.09)^{54,64,66}. To improve detection accuracy, the amino acid properties that tend to generate false positives were eliminated, and only those that had an accuracy of detection of selection higher than 85% were considered54.

It is well known that the three-dimensional (3D) structure of a protein determines its function. To understand the potential functional adaptation of the UT-A2 protein, we further investigated the evolutionary conserved and variable positions, as well as their physicochemical shifts, based on the known 3D crystal structures of UT. According to Levin and colleagues^{51,52}, the channel is lined by the residues from the highly conserved urea signature sequences¹⁰⁴ at the interface of the two halves (Pa,T1a T5a and Pb, T1b \sim T5b) (Fig. 4C). The channel contains a constricted selectivity filter that can accommodate urea molecules in single file and be divided into three regions: S_o, S_i and S_m^{51,52}. Given the high conservation of the "urea channel trimer" structure across the UT family^{51,52}, we superimposed the evolutionary conservation scores onto the 3D structures, Bos taurus UT-B (PDB ID: 4EZC)52 and Desulfovibrio vulgaris dvUT (PDB ID: 3K3F)51, to examine evolutionarily conserved and variable regions among these homologous sequences of this protein using the ConSurf webserver (Glaser, F., et al, The ConSurf Server, http://consurf.tau.ac.il/, 2003. Date of access 07/11/2013). In the analysis, the empirical Bayesian calculation method and LG substitution matrix were chosen to infer the scores. The scores were automatically divided into nine successive equally sized categories and assigned to different color codes according to the relative degree of conservation: grade 1 contains the most variable positions and grade 9 contains the most conserved positions. Regions under heavy selection detecting by TreeSAAP were correlated back to the 3D structure of the

Bos taurus UT-B complex using the program Pymol (Schrodinger, 2010). According to the evolutionary rate of each position of the protein and known mechanisms of urea conduction of dvUT and UT-B, the physicochemical evolutionary adaptation and potential mechanisms of UT-A2 were discussed.

- 1. Uhen, M. D. The origin(s) of whales. Annu Rev Earth Planet Sci 38, 189–219 (2010).
- Thewissen, J. G. M., Cooper, L. N., Clementz, M. T., Bajpai, S. & Tiwari, B. N. Whales originated from aquatic artiodactyls in the Eocene epoch of India. *Nature* 450, 1190–1194 (2007).
- Steeman, M. E. *et al.* Radiation of extant cetaceans driven by restructuring of the oceans. *Syst Biol* 58, 573–585 (2009).
- Berta, A., Sumich, J. L. & Kovacs, K. M. Marine mammals: evolutionary biology, 2edn., (Associated Press, San Diego, CA, 2006).
- Gingerich, P. D., von Koenigswald, W., Sanders, W. J., Smith, B. H. & Zalmout, I. S. New protocetid whale from the middle Eocene of Pakistan: birth on land, precocial development, and sexual dimorphism. *PLoS one* 4, e4366 (1–20) (2009).
- 6. Uhen, M. D. New protocetid whales from Alabama and Mississippi, and a new cetacean clade, Pelagiceti. J Vertebr Paleontol 28, 589–593 (2008).
- Thewissen, J. G. M. et al. Evolution of cetacean osmoregulation. Nature 381, 379–380 (1996).
- Uhen, M. D. Evolution of marine mammals: back to the sea after 300 million years. Anat Rec 290, 514–522 (2007).
- Luo, Z. & Gingerich, P. D. Terrestrial Mesonychia to aquatic Cetacea: transformation of the basicranium and evolution of hearing in whales. *Univ Mich Pap Paleontol* 31, 52–70 (1999).
- Ortiz, R. M. Osmoregulation in marine mammals. J Exp Biol 204, 1831–1844 (2001).
- Ridgway, S. H. Homeostasis in the aquatic environment. in *Mammals of the sea: biology and medicine* (ed. Ridgway, S. H.) 590–747 (Thomas Publisher, Limited, Charles C., 1972).
- Ridgway, S. & Venn-Watson, S. Effects of fresh and seawater ingestion on osmoregulation in Atlantic bottlenose dolphins (*Tursiops truncatus*). J Comp Physiol B 180, 563–576 (2010).
- Ortiz, R. M., Long, B., Casper, D., Ortiz, C. L. & Williams, T. M. Biochemical and hormonal changes during acute fasting and re-feeding in bottlenose dolphins (*Tursiops truncatus*). Mar Mamm Sci 26, 409–419 (2010).
- Birukawa, N. *et al.* Plasma and urine levels of electrolytes, urea and steroid hormones involved in osmoregulation of cetaceans. *Zoolog Sci* 22, 1245–1257 (2005).
- 15. Harrison, R. J. & Thurley, K. W. Fine structural features of delphinid epidermis. *J Anat* **111**, 498–500 (1972).
- Spearman, R. I. The keratinization of epidermal scales, feathers and hairs. *Biol Rev Camb Philos Soc* 41, 59–96 (1966).
- Sokolov, W. Some similarities and dissimilarities in the structure of the skin among the members of the suborders Odontoceti and Mystacoceti (Cetacea). *Nature* 185, 745–747 (1960).
- Sokolov, W. Adaptations of the mammalian skin to the aquatic mode of life. Nature 195, 464–466 (1962).
- Coulombe, H. N., Ridgway, S. H. & Evans, W. E. Respiratory water exchange in two species of porpoise. *Science* 149, 86–88 (1965).
- Perrin, W. F., Wursig, B. & Thewissen, J. G. M. Encyclopedia of marine mammals, (Academic Press, San Diego, CA, 2008).
- Wang, J. Z., Yu, X. Y., Guo, A. H., Hao, Y. J. & Wang, D. Research advances in cetacean osmoregulation. *Acta Theriologica Sinica* 32, 156–167 (2012).
- Hui, C. A. Seawater consumption and water flux in the common dolphin Delphinus delphis. Physiol Zool 54, 430–440 (1981).
- 23. Kjeld, M. Salt and water balance of modern baleen whales: rate of urine production and food intake. *Can J Zool* **81**, 606–616 (2003).
- Kjeld, M. Concentrations of electrolytes, hormones, and other constituents in fresh postmortem blood and urine of fin whales (*Balaenoptera physalus*). Can J Zool 79, 438–446 (2001).
- 25. Guo, A., Hao, Y., Wang, J., Zhao, Q. & Wang, D. Concentrations of osmotically related constituents in plasma and urine of finless porpoise (*Neophocaena asiaeorientalis*): implications for osmoregulatory strategies for marine mammals living in freshwater. *Zool Stud* 53, 10 (2014).
- Sands, J. M. Critical role of urea in the urine-concentrating mechanism. J Am Soc Nephrol 18, 670–671 (2007).
- Sands, J. M. & Layton, H. E. Advances in understanding the urine-concentrating mechanism. *Annu Rev Physiol* 76, 387–409 (2014).
- You, G. et al. Cloning and characterization of the vasopressin-regulated urea transporter. Nature 365, 844–847 (1993).
- Knepper, M. A. & Mindell, J. A. Structural biology: molecular coin slots for urea. Nature 462, 733–734 (2009).
- Grunewald, R. W., Reisse, C. H. & Muller, G. A. Characteristics of urea transport of cells derived from rabbit thick ascending limb of Henle's loop. *Kidney Int* 54, 152–159 (1998).
- Smith, C. P. & Rousselet, G. Facilitative urea transporters. J Membr Biol 183, 1–14 (2001).
- 32. Yancey, P. H., Clark, M. E., Hand, S. C., Bowlus, R. D. & Somero, G. N. Living with water stress: evolution of osmolyte systems. *Science* **217**, 1214–1222 (1982).

۲ ۲

- 33. Sands, J. M. Mammalian urea transporters. Annu Rev Physiol 65, 543–566 (2003).
- Sands, J. M. & Layton, H. E. The physiology of urinary concentration: an update. Semin Nephrol 29, 178–195 (2009).
- 35. Sands, J. M. & Layton, H. E. The urine concentrating mechanism and urea transporters. in *Seldin and Giebisch's the kidney: physiology and pathophysiology* (eds Alpern, R. J., Caplan, M. J. & Moe, O. W.) 1463–1510 (Academic Press, San Diego, 2013).
- 36. Smith, C. P. Mammalian urea transporters. Exp Physiol 94, 180-185 (2009).
- Knepper, M. A. & Roch-Ramel, F. Pathways of urea transport in the mammalian kidney. *Kidney Int* 31, 629–633 (1987).
- Knepper, M. A. & Star, R. A. The vasopressin-regulated urea transporter in renal inner medullary collecting duct. Am J Physiol 259, F393–F401 (1990).
- 39. Uchida, S. *et al.* Impaired urea accumulation in the inner medulla of mice lacking the urea transporter UT-A2. *Mol Cell Biol* **25**, 7357–7363 (2005).
- 40. Fenton, R. A., Chou, C. L., Stewart, G. S., Smith, C. P. & Knepper, M. A. Urinary concentrating defect in mice with selective deletion of phloretin-sensitive urea transporters in the renal collecting duct. *Proc Natl Acad Sci* 101, 7469–7474 (2004).
- 41. Fenton, R. A. & Knepper, M. A. Urea and renal function in the 21st century: insights from knockout mice. *J Am Soc Nephrol* **18**, 679–688 (2007).
- 42. Rafnar, T. *et al.* European genome-wide association study identifies SLC14A1 as a new urinary bladder cancer susceptibility gene. *Hum Mol Genet* **20**, 4268–4281 (2011).
- 43. Garcia-Closas, M. *et al.* A genome-wide association study of bladder cancer identifies a new susceptibility locus within SLC14A1, a urea transporter gene on chromosome 18q12.3. *Hum Mol Genet* **20**, 4282–4289 (2011).
- 44. Ranade, K. *et al.* Genetic variation in the human urea transporter-2 is associated with variation in blood pressure. *Hum Mol Genet* **10**, 2157–2164 (2001).
- 45. Janech, M. G. et al. Molecular and functional characterization of a urea transporter from the kidney of a short-finned pilot whale. Am J Physiol Regul Integr Comp Physiol 282, R1490–R1500 (2002).
- Birukawa, N. et al. Molecular cloning of urea transporters from the kidneys of baleen and toothed whales. Comp Biochem Physiol B Biochem Mol Biol 149, 227–235 (2008).
- 47. Promeneur, D. *et al.* Evidence for distinct vascular and tubular urea transporters in the rat kidney. J Am Soc Nephrol 7, 852–860 (1996).
- Karakashian, A. *et al.* Cloning and characterization of two new isoforms of the rat kidney urea transporter: UT-A3 and UT-A4. *J Am Soc Nephrol* 10, 230–237 (1999).
- Ashkar, Z. M., Martial, S., Isozaki, T., Price, S. R. & Sands, J. M. Urea transport in initial IMCD of rats fed a low-protein diet: functional properties and mRNA abundance. *Am J Physiol* 268, F1218–F1223 (1995).
- Shayakul, C., Steel, A. & Hediger, M. A. Molecular cloning and characterization of the vasopressin-regulated urea transporter of rat kidney collecting ducts. *J Clin Invest* 98, 2580–2587 (1996).
- Levin, E. J., Quick, M. & Zhou, M. Crystal structure of a bacterial homologue of the kidney urea transporter. *Nature* 462, 757–761 (2009).
- Levin, E. J. *et al.* Structure and permeation mechanism of a mammalian urea transporter. *Proc Natl Acad Sci* 109, 11194–11199 (2012).
- Raftery, A. E. Bayes factors and BIC: comment on 'a critique of the bayesian information criterion for model selection'. *Sociol Met Res* 27, 411–427 (1999).
- McClellan, D. A. & Ellison, D. D. Assessing and improving the accuracy of detecting protein adaptation with the TreeSAAP analytical software. *Int J Bioinform Res Appl* 6, 120–133 (2010).
- 55. Maciver, B., Smith, C. P., Hill, W. G. & Zeidel, M. L. Functional characterization of mouse urea transporters UT-A2 and UT-A3 expressed in purified Xenopus laevis oocyte plasma membranes. *Am J Physiol Renal Physiol* **294**, F956–F964 (2008).
- 56. Klein, J. D., Blount, M. A. & Sands, J. M. Urea Transport in the Kidney. Compr Physiol 1, 699–729 (2011).
- 57. Strugatsky, D. *et al.* Structure of the proton-gated urea channel from the gastric pathogen Helicobacter pylori. *Nature* **493**, 255–258 (2013).
- Kryazhimskiy, S. & Plotkin, J. B. The population genetics of dN/dS. *PLoS Genet* 4, e1000304(1–10) (2008).
- Yang, Z. Computational molecular evolution, (Oxford University Press, New York, 2006).
- 60. Kimura, M. Evolutionary rate at the molecular level. *Nature* **217**, 624–626 (1968).
- 61. Sharp, P. M. In search of molecular darwinism. *Nature* **385**, 111–112 (1997).
- McGowen, M. R., Grossman, L. I. & Wildman, D. E. Dolphin genome provides evidence for adaptive evolution of nervous system genes and a molecular rate slowdown. *Proc Biol Sci* 279, 3643–3651 (2012).
- 63. Zhou, X. *et al.* Baiji genomes reveal low genetic variability and new insights into secondary aquatic adaptations. *Nat Commun* **4**, 2708 (2013).
- 64. McClellan, D. A. *et al.* Physicochemical evolution and molecular adaptation of the cetacean and artiodactyl cytochrome b proteins. *Mol Biol Evol* **22**, 437–455 (2005).
- Streisfeld, M. A. & Rausher, M. D. Relaxed constraint and evolutionary rate variation between basic helix-loop-helix floral anthocyanin regulators in *Ipomoea. Mol Biol Evol* 24, 2816–2826 (2007).
- Porter, M. L., Cronin, T. W., McClellan, D. A. & Crandall, K. A. Molecular characterization of crustacean visual pigments and the evolution of pancrustacean opsins. *Mol Biol Evol* 24, 253–268 (2007).
- Hughes, A. L. Looking for Darwin in all the wrong places: the misguided quest for positive selection at the nucleotide sequence level. *Heredity* 99, 364–373 (2007).

- McClellan, D. Directional Darwinian Selection in proteins. BMC Bioinformatics 14, 1–8 (2013).
- Zhang, J., Nielsen, R. & Yang, Z. Evaluation of an improved branch-site likelihood method for detecting positive selection at the molecular level. *Mol Biol Evol* 22, 2472–2479 (2005).
- McClellan, D. A. & McCracken, K. G. Estimating the influence of selection on the variable amino acid sites of the cytochrome b protein functional domains. *Mol Biol Evol* 18, 917–925 (2001).
- 71. Conant, G. C. Neutral evolution on mammalian protein surfaces. *Trends Genet* **25**, 377–381 (2009).
- Woolley, S., Johnson, J., Smith, M. J., Crandall, K. A. & McClellan, D. A. TreeSAAP: selection on amino acid properties using phylogenetic trees. *Bioinformatics* 19, 671–672 (2003).
- Gromiha, M. M. & Ponnuswamy, P. K. Relationship between amino acid properties and protein compressibility. J Theor Biol 165, 87–100 (1993).
- 74. Oobatake, M. & Ooi, T. An analysis of non-bonded energy of proteins. *J Theor Biol* 67, 567–584 (1977).
- McNulty, R., Ulmschneider, J. P., Luecke, H. & Ulmschneider, M. B. Mechanisms of molecular transport through the urea channel of Helicobacter pylori. *Nat Commun* 4, 2900 (2013).
- 76. Weeks, D. L. & Sachs, G. Sites of pH regulation of the urea channel of Helicobacter pylori. *Mol Microbiol* **40**, 1249–1259 (2001).
- 77. Sayers, E. W. Amino Acid Explorer. http://www.ncbi.nlm.nih.gov/Class/ Structure/aa/aa_explorer.cgi, 2010. Date of access: 17/11/2013.
- Narhi, L. O. *et al.* Asn to Lys mutations at three sites which are N-glycosylated in the mammalian protein decrease the aggregation of Escherichia coli-derived erythropoietin. *Protein Eng* 14, 135–140 (2001).
- Chou, P. Y. & Fasman, G. D. Prediction of the secondary structure of proteins from their amino acid sequence. *Adv Enzymol Relat Areas Mol Biol* 47, 45–148 (1978).
- Rezus, Y. L. A. & Bakker, H. J. Effect of urea on the structural dynamics of water. Proc Natl Acad Sci 103, 18417–18420 (2006).
- Isogai, Y., Ota, M., Ishii, A., Ishida, M. & Nishikawa, K. Identification of amino acids involved in protein structural uniqueness: implication for de novo protein design. *Protein Eng* 15, 555–560 (2002).
- Poyurovsky, M. et al. COMT Val158Met polymorphism in schizophrenia with obsessive-compulsive disorder: a case-control study. Neurosci Lett 389, 21–24 (2005).
- Rojas, C. V. et al. A Met-to-Val mutation in the skeletal muscle Na⁺ channel [alpha]-subunit in hyperkalaemic periodic paralysis. Nature 354, 387–389 (1991).
- Nery, M., Arroyo, J. & Opazo, J. Accelerated evolutionary rate of the myoglobin gene in long-diving whales. J Mol Evol 76, 380–387 (2013).
- Burkin, D. J., Kim, J. E., Gu, M. & Kaufman, S. J. Laminin and alpha7beta1 integrin regulate agrin-induced clustering of acetylcholine receptors. *J Cell Sci* 113, 2877–2886 (2000).
- Baer, K. K. & McClellan, D. A. Molecular coevolution of the vertebrate cytochrome c(1) and Rieske iron sulphur protein in the cytochrome bc(1) complex. *Int J Bioinform Res Appl* 3, 456–470 (2007).
- 87. Shen, Y. Y. *et al.* Adaptive evolution of energy metabolism genes and the origin of flight in bats. *Proc Natl Acad Sci* **107**, 8666–8671 (2010).
- Ni, J. Y. & Zhou, K. Y. Rencular structural indices and urinary concentrating capacity of neophocaena phocaenoides. Acta Zool Sin 34, 243–250 (1988).
- 89. Ohta, T. Slightly deleterious mutant substitutions in evolution. *Nature* **246**, 96–98 (1973).
- Hughes, A. L. Relaxation of purifying selection on live attenuated vaccine strains of the family Paramyxoviridae. *Vaccine* 27, 1685–1690 (2009).
- Cassens, I. *et al.* Independent adaptation to riverine habitats allowed survival of ancient cetacean lineages. *Proc Natl Acad Sci* 97, 11343–11347 (2000).
- Nikaido, M. *et al.* Retroposon analysis of major cetacean lineages: the monophyly of toothed whales and the paraphyly of river dolphins. *Proc Natl Acad Sci* 98, 7384–7389 (2001).
- Hamilton, H., Caballero, S., Collins, A. G. & Brownell, R. L. Evolution of river dolphins. *Proc Biol Sci* 268, 549–556 (2001).
- 94. Fay, J. C. & Wu, C. I. Sequence divergence, functional constraint, and selection in protein evolution. *Annu Rev Genomics Hum Genet* **4**, 213–235 (2003).
- 95. Fay, J. C. & Wu, C.-I. The neutral theory in the genomic era. *Curr Opin Genet Dev* 11, 642–646 (2001).
- Zheng, J., Xia, J., He, S. & Wang, D. Population genetic structure of the Yangtze finless porpoise (*Neophocaena phocaenoides asiaeorientalis*): Implications for management and conservation. *Biochem Genet* 43, 307–320 (2005).
- Li, S. et al. Genome-wide SNP and population divergence of finless porpoises. Genome Biol Evol 5, 758–768 (2013).
- Yang, G., Guo, L. I., Bruford, M. W., Wei, F. & Zhou, K. Mitochondrial phylogeography and population history of finless porpoises in Sino-Japanese waters. *Biol J Linn Soc* 95, 193–204 (2008).
- Jefferson, T. A. & Wang, J. Y. Revision of the taxonomy of finless porpoises (genus Neophocaena): the existence of two species. J Mar Anim Ecol 4, 3–16 (2011).
- 100. Wang, J. Y., Frasier, T. R., Yang, S. C. & White, B. N. Detecting recent speciation events: the case of the finless porpoise (genus *Neophocaena*). *Heredity* 101, 145–155 (2008).
- Yang, Z., Wong, W. S. & Nielsen, R. Bayes empirical Bayes inference of amino acid sites under positive selection. *Mol Biol Evol* 22, 1107–1118 (2005).

- 102. Pond, S. L. K. & Frost, S. D. W. Not so different after all: a comparison of methods for detecting amino acid sites under selection. *Mol Biol Evol* 22, 1208–1222 (2005).
- Murrell, B. et al. Detecting individual sites subject to episodic diversifying selection. PLoS Genet 8, e1002764(1–10) (2012).
- 104. Rousselet, G., Ripoche, P. & Bailly, P. Tandem sequence repeats in urea transporters: identification of an urea transporter signature sequence. *Am J Physiol* 270, F554–F555 (1996).
- Zimmerman, J. M., Eliezer, N. & Simha, R. The characterization of amino acid sequences in proteins by statistical methods. *J Theor Biol* 21, 170–201 (1968).
 Bhaskaran, R. & Ponnuswamy, P. K. Dynamics of amino acid residues in globular
- proteins. Int J Pept Protein Res 24, 180-191 (1984). 107. Wang, P. L. Chinese cetaceans (Chemical Industry Press, Beijing, 2012).

Acknowledgments

We thank Dr. Zhao Huabin for his comments on the manuscript. A special thanks goes to Dr. Liu Zhen and Dr. Zhou Ming for their assistance in data analyses. We thank Nature Publishing Group Language Editing (NPG Language Editing) for language editing. The work was supported by the National Natural Science Foundation of China (grant numbers 30730018, 31000168 and 41306180), and the Knowledge Innovation Program of Chinese Academy of Sciences (grant number KSCX2-EW-Z-4).

Author contributions

Conceived and designed the experiments: D.W. and J.Z.W. Performed the experiments: J.Z.W., X.Y.Y. and B.H. Analyzed the data: J.Z.W. and X.Y.Y. Contributed reagents/ materials/analysis tools: J.S.Z., W.H.X. and Y.J.H. Wrote the paper: J.Z.W., W.H.L. and D.W.

Additional information

Supplementary information accompanies this paper at http://www.nature.com/ scientificreports

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Wang, J. *et al*. Physicochemical Evolution and Molecular Adaptation of the Cetacean Osmoregulation-related Gene UT-A2 and Implications for Functional Studies. *Sci. Rep.* **5**, 8795; DOI:10.1038/srep08795 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, with http://creativecommons.org/licenses/by/4.0/