



OPEN

SUBJECT AREAS:

PHOTOSYNTHESIS
ENVIRONMENTAL
MICROBIOLOGYReceived
11 January 2015Accepted
27 February 2015Published
30 March 2015

Correspondence and requests for materials should be addressed to Q.H. (qfhe@ualr.edu) or Q.W. (wangqiang@ihb.ac.cn)

* These authors contributed equally to this work.

A novel high light-inducible carotenoid-binding protein complex in the thylakoid membranes of *Synechocystis* PCC 6803

Soumana Daddy^{3*}, Jiao Zhan^{1,2*}, Saowarath Jantaro⁵, Chenliu He¹, Qingfang He^{3,4} & Qiang Wang¹

¹Key Laboratory of Algal Biology, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan 430072, China, ²University of Chinese Academy of Sciences, Beijing 100039, China, ³Department of Biology, University of Arkansas at Little Rock, Little Rock, AR 72204, ⁴Biotechnology Research Center, Shandong Academy of Agricultural Sciences, Jinan, Shandong 250100, China, ⁵Department of Biochemistry, Faculty of Science, Chulalongkorn University, Payathai Road, Patumwan, Bangkok 10330, Thailand.

Synechocystis sp. PCC 6803 is a model cyanobacterium extensively used to study photosynthesis. Here we reveal a novel high light-inducible carotenoid-binding protein complex (HLCC) in the thylakoid membranes of *Synechocystis* PCC 6803 cells exposed to high intensity light. Zeaxanthin and myxoxanthophyll accounted for 29.8% and 54.8%, respectively, of the carotenoids bound to the complex. Using Blue-Native PAGE followed by 2D SDS-PAGE and mass spectrometry, we showed that the HLCC consisted of Slr1128, IsiA, PsaD, and HliA/B. We confirmed these findings by SEAD fluorescence cross-linking and anti-PsaD immuno-coprecipitation analyses. The expression of genes encoding the protein components of the HLCC was enhanced by high light illumination and artificial oxidative stress. Deletion of these proteins resulted in impaired state transition and increased sensitivity to oxidative and/or high light stress, as indicated by increased membrane peroxidation. Therefore, the HLCC protects thylakoid membranes from extensive photooxidative damage, likely via a mechanism involving state transition.

Exposure to high-intensity light (HL) adversely affects the photosynthetic performance, cell growth, and viability of photosynthetic organisms. The damage is largely attributed to oxygen-dependent destruction of the photosynthetic apparatus and other cellular components^{1,2}.

Oxygenic photosynthetic organisms synthesize stress-associated proteins during exposure to HL. These proteins are often important for the acclimation of cells to HL. A family of HL-inducible genes, called *hli* or *scp* genes^{3,4}, encoding high light-inducible polypeptides (HLIP) with similarity to the light-harvesting chlorophyll *a/b*-binding proteins (LHCP) of plants, was shown to be critical for the survival of cyanobacteria under HL conditions⁵. All four HLIPs in *Synechocystis* PCC 6803, i.e., HliA, B, C, and D, are associated with photosystem I (PSI), and stabilize PSI trimers specifically under HL conditions. HliA and HliB also interact with Slr1128⁶. Another stress-associated protein, the iron stress-inducible chlorophyll-binding protein IsiA, stabilizes PSI trimers under HL conditions. Interestingly, IsiA, which is expressed at low levels under normal growth conditions and is induced by both HL and oxidative stress, is critical for the formation of PSI trimers in low-intensity light⁷. This is in contrast to the expression of PsaL and PsaI, both of which are essential for the formation of PSI trimers⁸, but are not strongly modified by stress conditions as compared to IsiA or HLIPs.

Even though the PSI complexes of green plants and cyanobacteria have similar structures and carry out similar functions, i.e., mediating the electron transfer between luminal plastocyanin (or cytochrome *c*₆) and stromal ferredoxin (Fd; or flavodoxin), there are several differences between them. One of the differences is that higher plants contain only monomeric PSI, whereas cyanobacteria harbor both monomeric and trimeric PSI^{9–11}. It is unclear why PSI trimers exist in cyanobacteria; however, studies in *Spirulina platensis* provided some insight into this phenomenon. In this cyanobacterium, the absorption spectra of the chlorophyll (Chl) components of the PSI trimers and monomers differ¹²; only the PSI trimers contain the extremely red-shifted Chl (the so-called red chlorophyll), which absorbs at 735 nm and gives rise to a 760-nm fluorescence emission peak (F760) at 77 K. The PSI monomer shows emission peaks only at 725–730 nm under the same conditions. In addition, the 760-nm emission band is only visible under reducing conditions, directly reflecting the redox state of P700^{12,13}. Red Chl is



thought to funnel light energy to P700^{14–16}, thereby increasing the cross-section of light absorption, or to dissipate excess energy into heat, thereby protecting PSI against photodestruction¹⁷.

The orange carotenoid protein (OCP), a 35-kDa two-domain soluble protein, was shown to trigger fluorescence quenching of phycobilisome (PBS) under blue-green light illumination¹⁸, and was proposed to be a photoactive protein that senses light intensity and triggers photoprotection¹⁹. The protein, which was first discovered by Holt and Krogman²⁰, contains one non-covalently bound carotenoid molecule^{20–24}. While the OCP is conserved among cyanobacteria except *Prochlorococci*²¹, the carotenoid differs in different species, being zeaxanthin in *Anacystis nidulans* and *Lyngbya wholei*²⁵, 3'-hydroxyechinenone in *Arthrospira maxima*, and a glycoside derivative of 3'-hydroxyechinenone in *Synechocystis* PCC 6803^{21,23,24,26}. Water-soluble OCP may function as a dimer in cells^{22,26}.

Here, we report the discovery of a novel high light-inducible carotenoid-binding protein complex (HLCC). This complex is normally concealed by trimeric PSI on a sucrose gradient, but was readily detected in a *Synechocystis* strain lacking PsaL (i.e., Δ PsaL) and hence lacking PSI trimers. The HLCC contains Slr1128, IsiA, PsaD, and HliA/B as its intrinsic protein components, binds zeaxanthin and myxothanxophyll, and is critical for the survival of cyanobacterial cells in HL conditions.

Results

Induction of a novel carotenoid-binding membrane protein complex by HL treatment. Thylakoid membranes were isolated from *Synechocystis* PCC 6803 cells grown in LL or HL, solubilized in mild detergent, and fractionated by sucrose gradient ultra-centrifugation (Fig. 1). Three pigmented fractions were detected in the samples of wild-type cells grown in LL. The uppermost orange fraction, F1, contained mainly free pigments; the middle green fraction, F2, contained PSI and PSII monomers; and the lower dark green fraction, F3, contained PSI trimers^{6,27}. In agreement with previous reports²⁸, the Δ PsaL strain grown in LL contained only F1 and F2, and lacked PSI trimers. Interestingly, when Δ PsaL cells were grown in HL for 24 h, an orange protein complex appeared at the gradient position of F3. The orange color of this membrane protein fraction indicates the presence of carotenoids.

We further purified the fraction by Blue-Native PAGE. As shown in Fig. 2A, a single orange band was detected, indicating that there is one carotenoid-containing protein complex in the sucrose gradient fraction. We named this orange protein complex high light-inducible carotenoid-binding complex (HLCC). The orange Blue-Native band




Figure 1 | Sucrose gradient fractions of thylakoid protein complexes from wild-type and Δ PsaL *Synechocystis* PCC 6803 strains. Thylakoid membranes were isolated from wild-type and PsaL deletion strains grown in LL or HL for 3 h and 24 h. Thylakoid protein complexes were separated by step sucrose gradient ultracentrifugation. Fractions are as indicated.




Figure 2 | Separation of the HLCC by Blue-Native PAGE followed by second dimension Tricine-SDS-PAGE. The HLCC fraction was carefully collected from the sucrose gradient and separated by BN PAGE (A), and the resulting single band was excised and denatured in 1.5 X SDS sample buffer and further separated on a 12–20% Tricine-SDS gel with 6 M urea (B). The proteins were visualized by silver staining and identified by mass spectrometry.

was then excised, denatured, and separated on a Tricine-SDS-PAGE gel (Fig. 2B), and four protein bands were revealed in the complex by silver staining. The bands were excised and identified by mass spectrometry. The three larger bands (from largest to smallest) contained the hypothetical protein Slr1128, the iron stress-induced chlorophyll-binding protein IsiA, and PsaD, a subunit of PSI, respectively. The fourth band at the lower end of the gel was identified as HliA and HliB.

To determine the pigment composition of the complex, the pigments were extracted and separated by HPLC according to²⁹. As shown in Fig. 3, myxoxanthophyll and zeaxanthin accounted for most of the pigment present in the HLCC, and there was a negligible amount of Chl a.

Slr1128, IsiA, PsaD, and HliA/B are intrinsic components of the HLCC. To characterize the HLCC further, we investigated whether IsiA, PsaD, and HliA/B are intrinsic components of the complex using anti-PsaD immuno-coprecipitation. Briefly, the sucrose gradient fraction containing the HLCC was collected and incubated first with anti-PsaD antibody and subsequently with protein G Sepharose beads. The beads were then loaded on an empty column and the protein complex was eluted and separated by SDS-PAGE. In addition to the heavy and light chains of the antibody, the anti-PsaD antibody identified Slr1128, IsiA, PsaD, and HliA/B as the major components of the HLCC (Fig. 4a).

We then confirmed these results by crosslinking analysis using SEAD (sulfosuccinimidyl-2-(7-azido-4-methylcoumarin-3-acetamido)-ethyl-1,3'-dithiopropionate), a crosslinker with a fluorescent spacer arm that fluoresces brightly when the disulfide bond between the crosslinked proteins is cleaved. Briefly, the non-covalent protein com-




Figure 3 | HPLC separation of the pigments in the sucrose gradient fractions. Pigments were identified by comparing their retention times and absorption spectra as described previously (Takaichi et al., 2001). Peaks are 1, myxoxanthophyll; 2, zeaxanthin; 3, hydroxyechinenone; 4, chlorophyll a; 5, echinenone; and 6, β -carotene.

plex was collected from F3 of the sucrose gradient, and the crosslinker was covalently incorporated into the complex by chemical bonding to free amino groups, mainly from lysine residues. The proteins of the complex were subsequently crosslinked by a photochemical reaction of the crosslinker's azido group. Cleavage of the disulfide group in the crosslinker by reduction released the fluorescently labeled proteins and removed the fluorescent group from proteins that were not cross-linked. The fluorescently labeled components of the protein complex were then separated by SDS-PAGE and detected by fluorescence imaging. As shown in Fig. 4B, when crosslinked with SEAD and separated by SDS-PAGE, the HLCC complex shows four fluorescent bands under UV light, which correspond to Slr1128, IsiA, PsaD, and

HliA/B, further demonstrating that these proteins are intrinsic components of the HLCC.

The HLCC is induced by high-intensity light, iron starvation, and oxidative stress. We previously showed that one of the major components of the HLCC discovered in the current study, IsiA, could be induced by iron depletion, HL stress, and rose bengal (RB) treatment⁷. To study the transcript induction of the other major component of the HLCC, i.e., Slr1128, we cultured both the wild-type and Δ PsaL strains in BG11, with or without iron, to the mid-logarithmic growth phase ($OD_{730} \sim 0.8$), and then diluted the cultures to $OD_{730} \sim 0.1$ with fresh medium and subjected them to




Figure 4 | Immunoprecipitation and SEAD crosslinkage of the HLCC. The HLCC collected from the sucrose gradient was (A) incubated with anti-PsaD and then Protein G resin, and the eluted proteins were separated by SDS PAGE, and (B) allowed to react in the dark with SEAD, quenched with lysine, photoactivated with UV light, and separated by SDS PAGE.

HL. Samples were taken at various time points for semi-quantitative RT-PCR analysis to examine the expression level of *slr1128* under various growth conditions. As shown in Fig. 5, *slr1128* is expressed in both the wild-type (Fig. 5A) and Δ PsaL (Fig. 5B) strains under standard conditions, and is further induced by HL and iron depletion (Fig. 5C). Expression levels plateaued after a 6-hour induction (Fig. 5D).

To further identify the conditions that induce transcription of the genes encoding components of the HLCC, we created artificial oxidative stress environments using the simple peroxide, hydrogen peroxide (H_2O_2); the lipid peroxide, decadienal (DD); the nitrite oxidant, peroxyxynitrite (PN); the redox-active compound, methyl viologen (MV); and the singlet oxygen generator, rose Bengal (RB). Fig. 6 shows that after a 12-hour treatment, *psaD* was strongly induced by all chemicals tested. Both *hliA* and *hliB*, which are normally not transcribed, were induced by H_2O_2 , MV, PN, and RB. Furthermore, *slr1128* was induced by H_2O_2 , MV, and RB. By contrast, as presented in our previous study, *isiA* was only induced by MV and RB⁷.

Deletion of *slr1128* resulted in increased sensitivity to oxidative stress due to membrane peroxidation. As shown above, all components of the HLCC were induced by oxidative stress (Fig. 6). We previously showed that PsaL, Slr1128, and IsiA are all important for *Synechocystis* survival upon exposure to HL^{6,7}. To evaluate the function of Slr1128 in oxidative stress conditions, we treated the deletion strain Δ Slr1128 and the wild type with RB and MV in the presence and absence of the chloroplast protein synthesis inhibitor chloramphenicol (Cm) for 12 h. We then measured the D1 protein level by immunoblot analysis as an indicator of oxidative stress. The Δ Slr1128 mutant strain was more sensitive than the wild type to both RB and MV treatment (Fig. 7), as indicated by the \sim 50% reduction in D1 protein after the treatment, and this sensitivity was enhanced in the presence of Cm (\sim 70% reduction in D1). These results suggest a role for Slr1128 in the response to oxidative stress.

To determine whether the enhanced light sensitivity of the Δ Slr1128 strain is due to HL-induced oxidative stress, we monitored the level of malondialdehyde (MDA), a product of lipid peroxidation, during HL treatment. The MDA level of all strains increased at the start of HL stress, and plateaued after 12 h of exposure to HL (Fig. 7B). After a 24-h HL treatment, MDA levels in the Δ Slr1128/

Δ PsaL double mutant increased by 160%, and the levels in all of the single mutants increased by about 110%, while those of the wild type increased by only 70%, indicating that the rate of membrane peroxidation was greater in the mutant than in the wild type under HL conditions. Furthermore, mutation of both *slr1128* and *psaL* had a greater effect than mutation of either gene alone, suggesting that SLR1128 and PsaL act synergistically. Thus, SLR1128 and PsaL play important roles in HL/oxidative stress.

Deletion of *slr1128* resulted in impaired state transition. State transition is a physiological adaptation in cyanobacteria that balances the distribution of light energy absorbed by phycobilisomes between PSI and PSII^{30,31}. Prompted by the previous observation that deletion of *isiA* altered the state transition capacity⁷, we examined whether state transition was also altered in the Δ Slr1128 strain. As shown in Fig. 8, the wild type has a fully functional state transition, as indicated by the fast and full relaxation of maximum fluorescence after high actinic light illumination (Fig. 8B and D). Therefore, the photoinhibitory effect of HL ($400 \mu\text{mol photon m}^{-2} \text{s}^{-1}$) on the wild type is negligible. By contrast, the Δ Slr1128 mutant exhibited a very low state transition level and resulted in substantial photoinhibitory quenching, as indicated by its poor recovery of maximum fluorescence (Fig. 8A and C).

Discussion

In the current study, we report the discovery of the novel high light-inducible carotenoid-binding protein complex (HLCC), which is normally concealed by trimeric PSI on a sucrose gradient, in the Δ PsaL strain of *Synechocystis* PCC 6803 subjected to HL stress. The HLCC contains Slr1128, IsiA, PsaD, and HliA/B as its intrinsic components, and binds to zeaxanthin and myxothanxophyll.

Slr1128 is a hypothetical protein highly conserved in cyanobacteria; the amino acid sequence identity between homologs of this protein in various species of cyanobacteria is generally above 80%. It also appears to be present in plants, with homologs in plants sharing 59–66% amino acid sequence identity with cyanobacterial Slr1128. Slr1128 has a transmembrane domain at its N-terminus and is likely an integral membrane protein, as it was also identified as a thylakoid membrane protein by other researchers³². Furthermore, Slr1128 exhibits high levels of similarity (around 35% amino acid sequence identity) to human and animal stomatins, which are thought to regulate an associated ion channel, and to a bacterial protease (HflC), but the similarity between Slr1128 and stomatins or HflC is not as high as that between the counterparts of these proteins in other cyanobacteria or plants. We previously reported that Slr1128 was associated with HliA/B, and suggested that it functions in photoprotection⁶. Here, we found that Slr1128 was also present in the HLCC, and appears to be closely associated with the HL-inducible proteins, HliA and HliB. This latter finding corresponds well with our previous work⁶.

The peripheral PsaD subunit is located at the stromal side of photosystem I³³, and is highly conserved in all photosynthetic organisms (including bacteria with Fe-S-type reaction centers). With PsaL, PsaD plays a critical role in the assembly and stability of PSI^{34–36} and mediates Fd electrostatic guidance and docking on PSI^{33,37–41}. Superoxide is formed by Fd at PSI through the Mehler reaction⁴².

In this study, we showed that PsaD, Slr1128, IsiA, and HliA/B formed a novel protein complex (the HLCC) in the cyanobacterium *Synechocystis* PCC 6803. Our earlier results showed that HliA and HliB stabilize PSI trimers, interact with Slr1128, and protect cells under HL conditions⁶. As our recent findings also showed that IsiA, which was identified here as one of the major components of the HLCC, forms a supercomplex with PSI and PSII, IsiA is important both for maintaining trimeric PSI integrity and for acclimation to HL and may protect cyanobacteria from HL and oxidative stresses through state transition⁷. Interestingly, we now found that the other major component of the HLCC, Slr1128, was also important for the




Figure 5 | Semi-quantitative PCR analysis of the induction of *slr1128* transcript during HL treatment in the presence and absence of iron. (A) Wild-type and (B) Δ PsaL *Synechocystis* strains were cultured in iron-deplete (Fe-) or iron-replete (Fe+) BG11 media to the mid-logarithmic growth phase ($OD_{730} \sim 0.8$ after 48 hours), diluted with fresh medium to $OD_{730} \sim 0.1$, and exposed to HL at 30°C. Samples for semi-quantitative PCR were taken at various time points as indicated; (C), Gel Bands quantification at time 0 as standardized with 16S; (D), Gel Bands quantification of (A) and (B), time 0 were set to 100 for easier comparison.

response to oxidative stresses and state transition (Fig. 7 and 8). These results suggest that Slr1128 interacts with IsiA to form the HLCC and protects cyanobacteria from oxidative stresses by mediating state transitions.

Thus, we propose that, by interacting with the peripheral PSI protein PsaD, which provides a docking site for Fd and interacts with PsaB, PsaC, PsaE, and PsaL, the HLCC complex stabilizes trimeric PSI and protects the photosystems, especially PSI, from HL and




Figure 6 | Semi-quantitative PCR analysis of genes encoding components of the HLCC upon 12 hours of exposure to chemically-induced oxidative stress. (A) Transcript levels upon treatment with the indicated artificial oxidants, and (B) quantification of the transcript levels shown in (A).




Figure 7 | Sensitivity of the HLCC mutants to oxidative stress. (A) D1 protein level in RB- and MV-treated cells in the presence (+Cm) or absence of chloramphenicol. Wild-type (Wt) and Δ Slr1128 (Δ Slr) cells in the mid-logarithmic growth phase (OD_{730} , 0.6–0.8) were diluted to an OD_{730} of 0.1 before exposure to HL for 12 h. Blots of the thylakoid membrane proteins were probed with polyclonal anti-D1 antibodies. (B) The level of lipid peroxidation in the HLCC mutant strains during HL treatment. Curves were generated by averaging the data obtained from three representative experiments.




Figure 8 | State transition of the wild-type and Δ Slr1128 strains. Representative traces of state transition analysis of the Δ Slr1128 (A) and wild-type (B) strains. Fast fluorescence induction kinetics of the Δ Slr1128 (C) and wild-type (D) strains at state 1 (S1), state 2 (S2), and state transition level (S').

oxidative stresses, probably through (1) direct scavenging of the superoxide and/or other reactive oxygen species (ROS) produced at Fd; and (2) mediating state transitions that are bridged by IsiA and Slr1128. The bound carotenoids, especially zeaxanthin, may function as antioxidants and directly dissipate excessive excitation energy.

Methods

Growth conditions and HL treatment. *Synechocystis* cells were cultivated in BG-11 medium with 10 mM TES, pH 8.2, at 30°C. The culture was bubbled with air under low-intensity light (LL; 40 $\mu\text{mol of photon m}^{-2} \text{s}^{-1}$) or high-intensity light (HL; 400 $\mu\text{mol of photon m}^{-2} \text{s}^{-1}$) conditions. For the experiments conducted under HL, cells in the mid-logarithmic growth phase ($\text{OD}_{730} \sim 0.8$) were diluted with fresh medium to $\text{OD}_{730} \sim 0.1$ and exposed to HL at 30°C.

Thylakoid membrane preparation and fractionation of membrane protein complexes. Thylakoid membranes were prepared as previously described⁴³ with some modifications. Briefly, cell pellets derived from cells grown to the mid-logarithmic phase were resuspended in ice-cold SMN thylakoid buffer (50 mM 3-(N-morpholino)-propanesulfonic acid (pH 7.0), 0.4 M sucrose, 10 mM NaCl, 5 mM MgCl_2 , and 1 mM freshly made phenylmethylsulfonyl fluoride). An equal volume of glass beads pre-wetted with thylakoid buffer was added to the cell suspension, and the cells were broken in a Bead-Beater with an ice-jacketed sample chamber in six breakage cycles at full speed (30 s of burst, followed by 5 min of chilling). The homogenate was centrifuged at 1,800 $\times g$ for 10 min to remove unbroken cells, cellular debris, and glass beads. The membranes in supernatant were then pelleted by centrifugation at 50,000 $\times g$ at 4°C for 60 min. After washing with 2 mM dodecyl maltoside to remove any remaining phycobilisomes, the membranes were washed twice and resuspended in thylakoid buffer to a chlorophyll a concentration of 1 mg/ml. The chlorophyll a concentration was estimated from the dimethylfluoride extract by the previously reported formula⁴⁴:

$$[\text{Chlorophyll a}] (\mu\text{g/ml}) = 12.1 \times \text{OD}_{664} - 0.17 \times \text{OD}_{625} \quad (\text{Eq. 1})$$

To fractionate membrane protein complexes, 150 μl of 10% dodecyl maltoside was added to the thylakoid membrane to achieve a detergent to chlorophyll ratio of 15:1. The membrane was solubilized at 4°C for 30 min before it was loaded onto a 10–30% (w/w) step sucrose gradient and centrifuged at 160,000 $\times g$ for 16 h at 4°C. Pigmented fractions were collected and stored at -80°C until use.

Blue-Native, Tricine-PAGE, and immuno-blotting. Blue-Native PAGE was performed as described⁴⁵. For electrophoresis in the second dimension, the single band of the blue native gel was excised and denatured in 1.5 X SDS sample buffer (50 mM Tris-HCl (pH 6.8), 3% SDS, 150 mM DTT, and 0.01% bromophenol blue) for 30 min at room temperature⁴⁶. The gel slice was then laid onto a 12–20% Tricine-SDS gel with 6 M urea as described. The proteins were visualized by silver staining⁴⁷. Polypeptides resolved by SDS-PAGE were electro-transferred onto a PVDF membrane. After the blocking step, the membranes were incubated with polyclonal primary antibodies (Agriser) followed by horseradish peroxidase-conjugated secondary antibody (Sigma-Aldrich). The reactive bands were detected by enhanced chemiluminescence detection reagents (GE healthcare). The bands were quantified using ImageJ Ver1.36 (National Institutes of Health, USA).

Database searching. All MS/MS samples were analyzed using Mascot (Matrix Science, London, UK; version 2.2.03). Mascot was set up to search the CBI nr_061307 database (selected for Bacteria, unknown version, 2419804 entries) assuming the

digestion enzyme trypsin. Mascot was searched with a fragment ion mass tolerance of 0.50 Da and a parent ion tolerance of 2.0 Da. The iodoacetamide derivative of cysteine was specified in Mascot as a fixed modification. S-carbamoylmethylcysteine cyclization (N-terminus) of the N-terminus, oxidation of methionine and acetylation of the N-terminus were specified in Mascot as variable modifications.

Criteria for protein identification. Scaffold (version Scaffold-01_07_00, Proteome Software Inc., Portland, OR) was used to validate MS/MS-based peptide and protein identifications. Peptide identifications were accepted if they could be established at greater than 80.0% probability as specified by the Peptide Prophet algorithm⁴⁸. Protein identifications were accepted if they could be established at greater than 95.0% probability and contained at least three identified peptides. Protein probabilities were assigned by the Protein Prophet algorithm⁴⁹. Proteins that contained similar peptides and could not be differentiated based on MS/MS analysis alone were grouped to satisfy the principles of parsimony.

SEAD crosslinking. The complex collected from the sucrose gradient was allowed to react in the dark with a 10-fold molar excess of freshly prepared SEAD. The reaction was quenched by adding a 30-fold molar excess of lysine, and the derivatized complex was separated from free reagent by gel filtration through spin chromatography columns (Bio-Rad, Hercules, CA) filled with porous polyacrylamide (Bio-Gel P-2 from Bio-Rad). Photoactivation was carried out for 15 min by exposing the sample to a black-ray, long-wave, 100-W ultraviolet lamp (Ultra Violet Products Inc., San Gabriel, CA). Then 4X Tris-glycine SDS reducing sample buffer was added at a ratio of 1:3 to the reaction mix, and the mix was incubated for 30 min at 22°C and then heated for 5 min at 75°C before electrophoresis.

Anti-PsaD Immuno-Coprecipitation. The complex collected from the sucrose gradient was allowed to react with anti-PsaD (Agriser) at 0.5–1 mg protein sample/1 μg antibody on ice for 90 min with occasional tube inversion.

The mixture was then added to the SMN-equilibrated Protein G Resin (Sigma-Aldrich), and allowed to react on ice for 1 h with occasional inversion. The Protein G Resin was loaded on a column and washed with 5X bed volume of SMN with 0.1% n-dodecyl-beta-D-maltoside (DM, Sigma-Aldrich). The protein complex was then eluted with elution buffer (Citric acid 0.1 M, pH 2.0) and immediately neutralized with 1 M Tris-HCl (pH 8.5 to pH 7.4), and dialyzed against 20 mM Tris-HCl, pH 7.4. The whole procedure was performed at 4°C and in dim light or in darkness, whenever possible.

Analysis of Carotenoid Composition by HPLC. Pigments extracted with methanol were subjected to HPLC analysis essentially as described²⁹. Pigments were identified by comparing their retention times and absorption spectra²⁹.

Analysis of state transition. Wild-type and mutant cells in the mid-logarithmic growth phase (OD_{730} , 0.6–0.8) were collected and resuspended to an OD_{730} of 0.6 with fresh BG-11 medium. State transition was analyzed by Chl fluorescence using a Dual-PAM-100 P700 & Chl Fluorescence Measuring System (Heinz Walz, Germany) as described previously⁷. A far-red light was first applied to the dark-adapted cyanobacterial cultures to fully oxidize PSI, which was followed by a series of saturation flashes to determine the S1 level. An actinic light of 100 $\mu\text{mol of photon m}^{-2} \text{s}^{-1}$ was subsequently turned on to induce photosynthesis. The S2 level was then assessed by a saturation pulse before the far-red light was applied again to drive the S2-S1 transition.

- Mittler, R. Oxidative stress, antioxidants and stress tolerance. *Trends in Plant Science* 7, 405–410 (2002).



2. Niyogi, K. K. Photoprotection revisited: Genetic and Molecular Approaches. *Annual Review of Plant Physiology and Plant Molecular Biology* **50**, 333–359, doi:10.1146/annurev.arplant.50.1.333 (1999).
3. Dolganov, N. A., Bhaya, D. & Grossman, A. R. Cyanobacterial protein with similarity to the chlorophyll a/b binding proteins of higher plants: evolution and regulation. *Proceedings of the National Academy of Sciences of the United States of America* **92**, 636–640 (1995).
4. Funk, C. & Vermaas, W. A Cyanobacterial Gene Family Coding for Single-Helix Proteins Resembling Part of the Light-Harvesting Proteins from Higher Plants. *Biochemistry* **38**, 9397–9404 (1999).
5. He, Q., Dolganov, N., Bjorkman, O. & Grossman, A. R. The High Light-inducible Polypeptides in *Synechocystis* PCC6803. Expression and function in high light. *J. Biol. Chem.* **276**, 306–314, doi:10.1074/jbc.M008686200 (2001).
6. Wang, Q. *et al.* The High Light-Inducible Polypeptides Stabilize Trimeric Photosystem I Complex under High Light Conditions in *Synechocystis* PCC 6803. *Plant Physiol.* **147**, 1239–1250, doi:10.1104/pp.108.121087 (2008).
7. Wang, Q., Hall, C. L., Al-Adami, M. Z. & He, Q. IsiA Is Required for the Formation of Photosystem I Supercomplexes and for Efficient State Transition in *Synechocystis* PCC 6803. *PLoS ONE* **5**, e10432 (2010).
8. Chitnis, V. P. *et al.* Targeted Inactivation of the Gene *PsaL* Encoding a Subunit of Photosystem-I of the Cyanobacterium *Synechocystis* Sp Pcc 6803. *J Biol Chem* **268**, 11678–11684 (1993).
9. Boekema, E. J. *et al.* Evidence for a Trimeric Organization of the Photosystem-I Complex from the Thermophilic Cyanobacterium *Synechococcus* Sp. *Febs Letters* **217**, 283–286 (1987).
10. Kruij, J., Bald, D., Boekema, E. & Rogner, M. Evidence for the Existence of Trimeric and Monomeric Photosystem-I Complexes in Thylakoid Membranes from Cyanobacteria. *Photosynthesis Research* **40**, 279–286 (1994).
11. Tsiotis, G., Haase, W., Engel, A. & Michel, H. Isolation and Structural Characterization of Trimeric Cyanobacterial Photosystem-I Complex with the Help of Recombinant Antibody Fragments. *European Journal of Biochemistry* **231**, 823–830 (1995).
12. Karapetyan, N. V., Dorra, D., Schweitzer, G., Bezsmertnaya, I. N. & Holzwarth, A. R. Fluorescence spectroscopy of the longwave chlorophylls in trimeric and monomeric photosystem I core complexes from the cyanobacterium *Spirulina platensis*. *Biochemistry* **36**, 13830–13837 (1997).
13. Shubin, V. V., Tsuprun, V. L., Bezsmertnaya, I. N. & Karapetyan, N. V. Trimeric Forms of the Photosystem-I Reaction-Center Complex Pre-Exist in the Membranes of the Cyanobacterium *Spirulina-Platensis*. *Febs Letters* **334**, 79–82 (1993).
14. Palsson, L. O., Dekker, J. P., Schlodder, E., Monshouwer, R. & vanGrondelle, R. Polarized site-selective fluorescence spectroscopy of the long-wavelength emitting chlorophylls in isolated Photosystem I particles of *Synechococcus elongatus*. *Photosynthesis Research* **48**, 239–246 (1996).
15. Shubin, V. V., Bezsmertnaya, I. N. & Karapetyan, N. V. Efficient Energy-Transfer from the Long-Wavelength Antenna Chlorophylls to P700 in Photosystem-I Complexes from *Spirulina-Platensis*. *Journal of Photochemistry and Photobiology B-Biology* **30**, 153–160 (1995).
16. Vangrondelle, R., Dekker, J. P., Gillbro, T. & Sundstrom, V. Energy-Transfer and Trapping in Photosynthesis. *Biochimica Et Biophysica Acta-Bioenergetics* **1187**, 1–65 (1994).
17. Trissl, H. W. Long-Wavelength Absorbing Antenna Pigments and Heterogeneous Absorption-Bands Concentrate Excitons and Increase Absorption Cross-Section. *Photosynthesis Research* **35**, 247–263 (1993).
18. Wilson, A. *et al.* A soluble carotenoid protein involved in phycobilisome-related energy dissipation in cyanobacteria. *Plant Cell* **18**, 992–1007 (2006).
19. Wilson, A. *et al.* A photoactive carotenoid protein acting as light intensity sensor. *Proceedings of the National Academy of Sciences* **105**, 12075–12080, doi:10.1073/pnas.0804636105 (2008).
20. Holt, K. T. & Krogmann, D. W. A carotenoid-protein from cyanobacteria. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **637**, 408–414 (1981).
21. Kerfeld, C. Structure and Function of the Water-Soluble Carotenoid-Binding Proteins of Cyanobacteria. *Photosynthesis Research* **81**, 215–225 (2004).
22. Polivka, T., Kerfeld, C. A., Pascher, T. & Sundstrom, V. Spectroscopic Properties of the Carotenoid 3'-Hydroxyechinenone in the Orange Carotenoid Protein from the Cyanobacterium *Arthrospira maxima*. *Biochemistry* **44**, 3994–4003 (2005).
23. Wu, Y. P. & Krogmann, D. W. The orange carotenoid protein of *Synechocystis* PCC 6803. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1322**, 1–7 (1997).
24. Kerfeld, C. A. Water-soluble carotenoid proteins of cyanobacteria. *Archives of Biochemistry and Biophysics* **430**, 2–9 (2004).
25. Divers-Pierluissi, M. & Krogmann, D. W. A zeaxanthin protein from *Anacystis nidulans*. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **933**, 372–377 (1988).
26. Kerfeld, C. A. *et al.* The Crystal Structure of a Cyanobacterial Water-Soluble Carotenoid Binding Protein. *Structure* **11**, 55–65 (2003).
27. Sun, J., Ke, A., Jin, P., Chitnis, V. P. & Chitnis, P. R. Isolation and functional study of photosystem I subunits in the cyanobacterium *Synechocystis* sp. PCC 6803. *Methods in enzymology* **297**, 124–139 (1998).
28. Chitnis, V. P. & Chitnis, P. R. *PsaL* Subunit Is Required for the Formation of Photosystem-I Trimers in the Cyanobacterium *Synechocystis* Sp Pcc-6803. *Febs Letters* **336**, 330–334 (1993).
29. Takaichi, S., Maoka, T. & Masamoto, K. Myxoxanthophyll in *Synechocystis* sp. PCC 6803 is Myxol 2'-Dimethyl-Fucoside, (3R,2'S)-Myxol 2'-(2,4-di-O-Methyl- α -l-Fucoside), not Rhamnoside. *Plant and Cell Physiology* **42**, 756–762, doi:10.1093/pcp/pce098 (2001).
30. Campbell, D., Hurry, V., Clarke, A. K., Gustafsson, P. & Oquist, G. Chlorophyll Fluorescence Analysis of Cyanobacterial Photosynthesis and Acclimation. *Microbiol. Mol. Biol. Rev.* **62**, 667–683 (1998).
31. Campbell, D. & Oquist, G. Predicting Light Acclimation in Cyanobacteria from Nonphotochemical Quenching of Photosystem II Fluorescence, Which Reflects State Transitions in These Organisms. *Plant Physiol.* **111**, 1293–1298, doi:10.1104/pp.111.4.1293 (1996).
32. Herranen, M. *et al.* Towards Functional Proteomics of Membrane Protein Complexes in *Synechocystis* sp. PCC 6803. *Plant Physiol.* **134**, 470–481, doi:10.1104/pp.103.032326 (2004).
33. Zilber, A. L. & Malkin, R. Organization and Topology of Photosystem I Subunits. *Plant Physiol.* **99**, 901–911, doi:10.1104/pp.99.3.901 (1992).
34. Chitnis, P. R., Reilly, P. A. & Nelson, N. Insertional inactivation of the gene encoding subunit II of photosystem I from the cyanobacterium *Synechocystis* sp. PCC 6803. *J. Biol. Chem.* **264**, 18381–18385 (1989).
35. Chitnis, V. P. & Chitnis, P. R. *PsaL* subunit is required for the formation of photosystem I trimers in the cyanobacterium *Synechocystis* sp. PCC 6803. *FEBS letters* **336**, 330–334 (1993).
36. Chitnis, V. P., Jung, Y.-S., Albee, L., Golbeck, J. H. & Chitnis, P. R. Mutational Analysis of Photosystem I Polypeptides. *J. Biol. Chem.* **271**, 11772–11780, doi:10.1074/jbc.271.20.11772 (1996).
37. Bottin, H. & Lagoutte, B. Ferredoxin and flavodoxin from the cyanobacterium *Synechocystis* sp PCC 6803. *Biochimica et Biophysica Acta (BBA) - Protein Structure and Molecular Enzymology* **1101**, 48–56 (1992).
38. Giuliana, Z. & Giuliana, M. Interaction between photosystem I and ferredoxin. *European Journal of Biochemistry* **169**, 143–146 (1987).
39. Lagoutte, B., Hanley, J. & Bottin, H. Multiple Functions for the C Terminus of the *PsaD* Subunit in the Cyanobacterial Photosystem I Complex. *Plant Physiol.* **126**, 307–316, doi:10.1104/pp.126.1.307 (2001).
40. Wynn, R. M., Omaha, J. & Malkin, R. Structural and functional properties of the cyanobacterial photosystem I complex. *Biochemistry* **28**, 5554–5560, doi:10.1021/bi00439a032 (2002).
41. Zilber, A. L. & Malkin, R. Ferredoxin Cross-Links to a 22 kD Subunit of Photosystem I. *Plant Physiol.* **88**, 810–814, doi:10.1104/pp.88.3.810 (1988).
42. Pfannschmidt, T. Chloroplast redox signals: how photosynthesis controls its own genes. *Trends in Plant Science* **8**, 33–41 (2003).
43. Shen, G. & Vermaas, W. F. Chlorophyll in a *Synechocystis* sp. PCC 6803 mutant without photosystem I and photosystem II core complexes. Evidence for peripheral antenna chlorophylls in cyanobacteria. *The Journal of biological chemistry* **269**, 13904–13910 (1994).
44. Moran, R. Formulas for Determination of Chlorophyllous Pigments Extracted with N,N-Dimethylformamide. *Plant Physiol* **69**, 1376–1381 (1982).
45. Schagger, H. & Vonjagow, G. Blue Native Electrophoresis for Isolation of Membrane-Protein Complexes in Enzymatically Active Form. *Analytical Biochemistry* **199**, 223–231 (1991).
46. Schagger, H. Tricine-SDS-PAGE. *Nature Protocols* **1**, 16–22 (2006).
47. Rabilloud, T., Carpentier, G. & Tarroux, P. Improvement and Simplification of Low-Background Silver Staining of Proteins by Using Sodium Dithionite. *Electrophoresis* **9**, 288–291 (1988).
48. Keller, A., Nesvizhskii, A. I., Kolker, E. & Aebersold, R. Empirical Statistical Model To Estimate the Accuracy of Peptide Identifications Made by MS/MS and Database Search. *Analytical Chemistry* **74**, 5383–5392, doi:10.1021/ac025747h (2002).
49. Nesvizhskii, A. I., Keller, A., Kolker, E. & Aebersold, R. A Statistical Model for Identifying Proteins by Tandem Mass Spectrometry. *Analytical Chemistry* **75**, 4646–4658 (2003).

Acknowledgments

This work was supported by the National Basic Research Program of China (973 Program, 2012CB224803), US National Science Foundation (MCB No. 1120153), Shandong Province 'Taishan Scholar' Foundation (No. tshw20091014) and the National Natural Science Foundation of China (Grant No. 31270094 and 31270102).

Author contributions

S.D. and J.Z. were responsible for study conception and design, data collection and analysis, manuscript writing and final approval of the manuscript; S.J. and C.H. for data collection and analysis, and final approval of the manuscript; Q.H. and Q.W. for conception and design, critical revision and manuscript writing, and final approval of the manuscript. All authors read and approved the final manuscript.



Additional information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Daddy, S. *et al.* A novel high light-inducible carotenoid-binding protein complex in the thylakoid membranes of *Synechocystis* PCC 6803. *Sci. Rep.* 5, 9480; DOI:10.1038/srep09480 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>