OPEN

SUBJECT AREAS:

ORGANOMETALLIC
CHEMISTRY

INORGANIC CHEMISTRY

ORGANIC CHEMISTRY

THEORETICAL CHEMISTRY

Received

30 November 2014

Accepted

23 February 2015

Published

9 April 2015

Correspondence and
requests for materials
should be addressed to
H.Z. (zh@xmu.edu.cn);
Z.L. (chzlin@ust.hk) or
H.X. (hpxia@xmu.
edu.cn)

Synthesis of Aromatic Aza-metallapentalenes from Metallabenzene via Sequential Ring Contraction/Annulation

Tongdao Wang¹, Feifei Han¹, Haiping Huang¹, Jinhua Li¹, Hong Zhang^{1,2}, Jun Zhu¹, Zhenyang Lin²
& Haiping Xia¹

¹State Key Laboratory of Physical Chemistry of Solid Surfaces and Collaborative Innovation Center of Chemistry for Energy Materials (iChEM), and Department of Chemistry, Xiamen University, Xiamen 361005, China, ²Department of Chemistry, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong.

The concept of aromaticity has long played an important role in chemistry and continues to fascinate both experimentalists and theoreticians. Among the archetypal aromatic compounds, heteroaromatics are particularly attractive. Recently, substitution of a transition-metal fragment for a carbon atom in the anti-aromatic hydrocarbon pentalene has led to the new heteroaromatic osmapentalenes. However, construction of the aza-homolog of osmapentalenes cannot be accomplished by a similar synthetic manipulation. Here, we report the synthesis of aza-osmapentalenes by sequential ring contraction/annulation reactions of osmabenzenes via osmapentafulvenes. Nuclear magnetic resonance spectra, X-ray crystallographic analysis, and DFT calculations all suggest that these aza-osmapentalenes exhibit aromatic character. Thus, the stepwise transformation of metallabenzenes to metallapentafulvenes and then aza-metallapentalenes provides an efficient and facile synthetic route to these bicyclic heteroaromatics.

As one of the most fundamental chemistry concepts, aromaticity has been continually investigated since Faraday's discovery of benzene in 1825¹ and Kekulé's famous alternating single- and double-bond cyclic structure proposed in 1865^{2,3}. The Hückel aromaticity rule⁴ is applied to cyclic compounds containing $4n + 2\pi$ electrons, but Möbius topologies favour $4n$ delocalized π electron counts^{5–8}. In general, aromatic compounds are substantially more stable thermodynamically and anti-aromatic compounds less stable than appropriate non-aromatic reference systems. Synergistic interplay of theoretical and experimental investigations has led to the synthesis and characterisation of numerous aromatic molecules with interesting structural features and properties^{9,10}. The incorporation of heteroatoms or related groups into aromatic hydrocarbons can result in various heteroaromatic compounds. These introduced heteroatoms are not limited to main-group atoms or related groups; they also include transition-metal fragments. As a result of the increasing structural and compositional variety of aromatic species, the definitions and criteria for characterising aromaticity have also been developing rapidly¹¹.

Metallaaromatics^{12–14} are organometallic species derived from the replacement of a (hydro)carbon unit in conventional aromatic hydrocarbons with a transition-metal fragment. As typical metallaaromatic compounds, metallabenzene complexes have been extensively studied by both experimentalists and theoreticians, dating back to the first computationally proposed metallabenzene complexes by Thorn and Hoffman¹⁵ in 1979. The increase in the variety of the metallaaromatic family is further evidenced by the recent discovery of metallapyridines^{16–18}, which are azaheterocyclic analogues of metallabenzenes. Very recently, efforts to expand the aromatic family has also led to the discovery of the first metallapentalene (osmapentalene)^{19,20}, which has been shown to exhibit aromaticity. Theoretical computations reveal that all these osmapentalenes with planar metallacycle benefit from Craig-Möbius aromaticity. The involvement of transition-metal d orbitals in the π conjugation switches the Hückel anti-aromaticity of pentalene into the Möbius aromaticity of metallapentalene. The introduction of nitrogen atoms into the pentalene system lead to the heterocyclic analogs of pentalene which are broadly defined as azapentalenes. Similar to the parent pentalenes, azapentalenes are anti-aromatic by virtue of a $8-\pi$ -electron system²¹. Some azapentalene molecules that contain nitrogen atoms in the bridgehead positions can be consid-

Figure 1 | The proposed structure of aza-metallapentalene.

ered as analogs of a pentalene dianion, and these molecules are actually aromatic²². The isolation of the metallapentalene prompted us to investigate whether an aza-metallapentalene exists and, if so, how the synthesis could be achieved. As shown in Fig. 1, the structure of an aza-metallapentalene resembles that of pentalene, which is generally considered to be an anti-aromatic molecule, prompting us to consider whether an aza-metallapentalene, which is structurally and electronically analogous to metallapentalene, is also aromatic. In this paper, we present the synthesis and characterisation of aza-metallapentalenes, which were obtained from metallabenzene via sequential ring contraction/annulation reactions.

Results

Synthesis of the first aza-metallapentalene. The recently reported osmapentalene^{19,20} was synthesised (derived) from the protonation of osmapentalyne (Fig. 2a). Experimentally, osmapentalyne²³ was observed to be thermally stable and easy to prepare via the reaction of osmium complexes with terminal alkynes (Fig. 2a). Inspired by this observation, we initially attempted to react osmium complexes with a number of nitriles such as benzonitrile or acetonitrile in the hope of obtaining the corresponding aza-osmapentalynes. However, the expected reactions did not occur, only osmabenzenes or other osmacycles were detected, which have been previously reported^{24,25}. Therefore, the preparation of an aza-metallapentalene via the synthesis of an aza-osmapentalyne followed by protonation is unfeasible.

Previously, we synthesised osmacyclopentadienes, each bearing an exocyclic C = N bond, by reacting an osmabenzene complex with amines²⁶ (Fig. 2b). We envisioned that an aza-osmapentalyne might be directly obtained by introducing a new carbon atom, with the intention of inducing a further annulation reaction yielding an osmacyclopentadiene monocyclic system, as illustrated by the retrosynthetic analysis shown in Fig. 2b.

We first modified an osmabenzene complex by ligand substitution to generate the more stable osmabenzene 1-I. The structure of osma-

benzene 1-I was verified by X-ray diffraction analysis (Fig. 3b; the relevant experimental details are presented in Supplementary Figures and Methods). Subsequent treatment of 1-I with aniline in the presence of sodium hexafluorophosphate afforded the expected osmacyclopentadiene 2-PF₆ in 95% yield (Fig. 3a). The complex 2-PF₆ was characterised by multinuclear nuclear magnetic resonance (NMR) spectroscopy, single-crystal X-ray diffraction analysis, and high-resolution mass spectroscopy. The structure of 2-PF₆ is shown in Fig. 3c. The X-ray diffraction analysis demonstrated that 2-PF₆ contains a coplanar metallacyclopentadiene ring with an exocyclic imino group. The structural parameters indicated that the metallacycle of 2-PF₆ could be represented by two resonance structures, 2-PF₆ and 2b-PF₆ (Fig. 3a), with 2-PF₆ as the more dominant structure. The structural data (averaged bond lengths) also indicated that both the metallacycles in 1-I and 2-PF₆ have delocalised structures. In addition, the six atoms Os1 and C1-C5 in each complex are approximately coplanar, which is reflected by their small mean deviation (0.0129 Å for 1-I; 0.0331 Å for 2-PF₆) from the least-squares plane.

As shown in Fig. 3a, the resonance structure 2b-PF₆ (metallapentafulvene) contributes to the overall structure of the complex 2-PF₆. Since pentafulvene molecules²⁷ are normally considered aromatic, we assume that the metallapentafulvene 2-PF₆ is also aromatic, as is indeed supported by our density functional theory (DFT) results (*vide infra*).

According to the retrosynthetic analysis shown in Fig. 2b, the complex 2-PF₆ could be used as the precursor to afford the desired aza-metallapentalene by introducing an additional carbon atom. To test this idea, reactions of 2-PF₆ with the propynols RCH(OH)C≡CH in the presence of Ag₂O were carried out in the hope of obtaining the desired aza-osmapentalyne complexes. Triethylamine was used in the reactions to ensure that an acetylide ligand could be easily delivered to the osmium metal centre (Warning: the reaction may proceed through the formation of silver acetylide, which decomposes violently on contact with moisture and water producing highly flammable and explosive acetylene gas, and causing fire and explosion

Figure 2 | Previous work and initial attempt. (a) Protonation of osmapentalyne produces aromatic osmapentalene. (b) Reactions of osmabenzene with amines, leading to the formation of osmacyclopentadiene with an exocyclic C = N bond.

Figure 3 | Synthesis of osmacyclopentadiene 2-PF₆. (a) The reaction of osmabenzene **1-I** with aniline produces osmacyclopentadiene **2-PF₆**. (b, c) X-ray structures of **1-I** (b) and **2-PF₆** (c) (50% probability level). Phenyl moieties in PPh₃, the counter anion and the solvent molecules have been omitted for clarity. Selected bond lengths (Å) for **1-I**: Os1-C1 2.065(7), Os1-C5 1.956(7), C1-C2 1.359(10), C2-C3 1.432(11), C3-C4 1.345(10), C4-C5 1.407(11), C4-I1 2.131(7), Os1-C01 1.925(8), C01-O01 1.145(9), Os1-I2 2.8161(8). Selected bond lengths (Å) for **2-PF₆**: Os1-C1 2.004(6), Os1-C4 2.183(7), Os1-I1 2.8347(5), C1-C2 1.372(9), C2-C3 1.430(9), C3-C4 1.386(9), C4-C5 1.418(9), C5-N1 1.309(8), Os1-C01 1.880(9), C01-O01 1.109(8).

hazard). As shown in Fig. 4a, treatment of **2-PF₆** with phenylpropynol in the presence of Ag₂O and triethylamine under reflux for approximately 3 h led to the formation of complex **3a-PF₆**, which was isolated in 81% yield. The complex **3a-PF₆** was characterised by NMR spectroscopy and HRMS, and its structure was determined by X-ray crystal structure analysis.

Complex **3a-PF₆** exhibits an essentially planar metal-bridged bicyclic structure (Fig. 4b). The mean deviation from the least-squares plane through Os1, N1 and C1-C6 is 0.0383 Å. The Os1-C6 bond length (2.180(9) Å) is appreciably longer than both the Os1-C1 bond length (2.040(8) Å) and the Os1-C4 bond length (2.100(7) Å). The three Os-C bond lengths of the metallabicyclic all

lie in the high end of the reported range for typical Os-C(vinyl)²⁸ bond lengths (1.859–2.359 Å) (on the basis of a search of the Cambridge Structural Database, CSD version 5.35, conducted in November, 2013). In addition, the structural parameters indicate a considerable bond distance alternation within the two fused five-membered rings. Therefore, the complex **3a-PF₆** is not the desired aza-osmapentalene, although the osmabicyclic framework is consistent with the suppositional aza-osmapentalene formula.

The hydroxyl group in **3a-PF₆** was easily removed to afford the desired complex **4a-(PF₆)₂** by treatment of **3a-PF₆** with NaPF₆. The complex **4a-(PF₆)₂** was isolated as a red solid in 98% yield and was characterised by multinuclear NMR spectroscopy and HRMS.

Figure 4 | Synthesis of aza-osmapentalene 4-(PF₆)₂. (a) The reaction of osmacyclopentadiene **2-PF₆** with substituted propynols followed by dehydroxylation produces aza-osmapentalene **4-(PF₆)₂**. (b, c) X-ray structures of **3a-PF₆** (b) and **4a-(PF₆)₂** (c) (50% probability level). Phenyl moieties in PPh₃, the counter anion and the solvent molecules have been omitted for clarity. Selected bond lengths (Å) for **3a-PF₆**: Os1-C1 2.040(8), Os1-C4 2.100(7), Os1-C6 2.180(9), C1-C2 1.381(11), C2-C3 1.450(11), C3-C4 1.373(12), C4-C5 1.377(13), C5-N1 1.313(11), C6-N1 1.483(11), C6-C7 1.338(12), C7-C8 1.489(12), C8-O1 1.430(12). Selected bond lengths (Å) for **4a-(PF₆)₂**: Os1-C1 2.041(4), Os1-C4 2.069(4), Os1-C6 2.082(4), C1-C2 1.407(5), C2-C3 1.410(5), C3-C4 1.394(5), C4-C5 1.381(5), C5-N1 1.374(5), C6-N1 1.396(5), C6-C7 1.465(5), C7-C8 1.337(5).

Single-crystal X-ray diffraction established the solid-state structure of **4a**-(PF₆)₂ unambiguously, demonstrating its aza-osmapentalene arrangement. As shown in Fig. 4c, the two fused five-membered rings of **4a**-(PF₆)₂ are nearly coplanar. The mean deviation from the least-squares plane through Os1, N1 and C1-C6 is only 0.0154 Å. All the bond distances within the two fused rings fall within the range observed for other typical metallaaromatics^{12–14}. The three Os–C distances are nearly equal (Os1–C1 2.041(4), Os1–C4 2.069(4), Os1–C6 2.082(4) Å), and the ring C–C/C–N distances are consistent with extensive delocalisation within the metallacycle (C1–C2 1.407(5), C2–C3 1.410(5), C3–C4 1.394(5), C4–C5 1.381(5), C5–N1 1.374(5), C6–N1 1.396(5) Å). The aromaticity in the metallacyclic rings is also clearly evident in the NMR spectra. In particular, the ¹H NMR spectrum shows three strongly downfield ¹H signals corresponding to the ring protons at δ = 14.3 (C1H), 10.0 (C3H), and 9.9 (C5H) ppm. With the aid of the ¹H–¹³C HSQC and ¹³C–DEPT 135 spectra, we observed the signals for the six carbon atoms of the osmabicyclic at δ = 248.5 (C1), 151.0 (C2), 171.9 (C3), 187.6 (C4), 163.0 (C5) and 234.1 (C6) ppm in the ¹³C{¹H} NMR spectrum.

Thus, using the strategy outlined in Fig. 2b, we successfully synthesised the first aza-metallapentalene complex in three steps starting from the six-membered osmabenzene complex **1**-I. Further investigations demonstrated that this synthetic strategy could be extended to prepare other aza-osmapentalene complexes with other substituents on the metallacycle. The vinyl-containing aza-osmapentalene **4b**-(PF₆)₂ was obtained from the reaction of **2**-PF₆ with propynol (Fig. 4a) and was isolated as a brown solid. Similarly, the butenylnyl-containing aza-osmapentalene **4c**-(PF₆)₂ was obtained from the reaction of **2**-PF₆ with penta-1,4-diyn-3-ol and was isolated as a dark-brown solid (Fig. 4a).

As previously mentioned, X-ray crystallography revealed a planar structure with no alternation in the carbon–carbon bond lengths of the osmabicyclic rings in the aza-osmapentalene **4a**-(PF₆)₂. Protons on the periphery of aromatic compounds are well known to have relatively large downfield NMR spectroscopic chemical shifts due to a diamagnetic ring current. Fig. 5a shows a comparison of the chemical shifts of three specific protons among four different metallacyclic

complexes. Compared with the structurally similar complex **3a**, complex **4a** gave considerable downfield chemical shifts because of its aromatic character. The ¹H NMR chemical shifts as well as the X-ray diffraction data indicate that the metallacycle in the cation of **4**-(PF₆)₂ has a delocalised structure, with contributions from the six resonance structures **4** to **4E**, as shown in Fig. 5b. Consistent with their aromaticity, all of the new aza-osmapentalenes exhibit remarkably high thermal stability. Solid samples of **4**-(PF₆)₂ were heated in air at 160°C for at least 5 h without noticeable decomposition. **4**-(PF₆)₂ was stable even at 80°C in 1,2-dichloro-ethane solution for 5 h. These new aza-osmapentalenes each contain a phosphonium substituent, which is also believed to partially contribute to the high stability²⁹.

DFT calculations of osmabenzene, osmacyclopentadiene and aza-osmapentalene. To gain more insight into the electronic structure and aromaticity of the aza-osmapentalenes **4**, we performed density functional theory (DFT, B3LYP/6-311++G(d,p)) calculations. The nucleus-independent chemical shift (NICS)^{30–32} values were computed for the metallacycle rings of complexes **1**-I, **2**-PF₆, and **4**-(PF₆)₂. In general, negative NICS values indicate aromaticity, whereas positive values suggest anti-aromaticity. As shown in Fig. 6a, the calculated NICS(1) values are –2.6 and –1.6 ppm for the rings in osmabenzene **1** and osmapentafulvene **2**, respectively. (When the environments at points 1 Å above and below the ring centres are not equivalent, the averaged values were used for NICS(1) values.) These values are comparable to those reported for other metallaaromatics^{19,20,23,31,33–36}. The NICS(1) values for the two five-membered rings of aza-osmapentalene **4a** are also negative (–6.7 and –6.9 ppm), in sharp contrast with those of azapentalene (+22.2 and +15.8 ppm). We also evaluated the aromatic stabilisation energy (ASE) by employing the isomerisation method introduced by Schleyer and Pühlhofer³⁷. For model complexes **1'**, **2'** and **4a'**, ASE values of 11.3, 10.1 and 26.2 kcal/mol were calculated, respectively; these values contrast sharply with the negative ASE value (–4.3 kcal mol^{–1}) calculated for azapentalene. The calculated ASE value for the aza-osmapentalene **4a'** is close to the

Figure 5 | Aromaticity of aza-osmapentalenes: downfield ¹H chemical shifts and resonance structures. (a) The proton chemical shifts (ppm vs. tetramethylsilane) of the ring protons of osmabenzene **1**, osmapentafulvene **2**, osmabicyclic **3a** and aza-osmapentalene **4a**. (b) Six possible resonance structures for the aza-osmapentalenes **4**.

previously reported values for osmapentalenes^{19,20}. The negative NICS values and the significant ASE values calculated for these model complexes indicate that aromaticity is closely associated with the metallacycle rings in complexes **1**, **2** and **4a**.

Additional evidence for the aromatic nature of aza-osmapentalene **4** was provided by examination of the molecular orbitals calculated for the further simplified model complex **4a''**. π Electrons are normally responsible for aromatic behaviour. Therefore, our attention is paid to those occupied orbitals. As shown in Fig. 7a, the five occupied π molecular orbitals (MOs) calculated for **4a''** (HOMO-1, HOMO-2, HOMO-3, HOMO-7 and HOMO-18) reflect the π -delocalisation along the perimeter of the bicyclic system. HOMO-1, HOMO-3 and HOMO-18 are derived from the orbital interactions between the p_π atomic orbitals of the C_6NH_6 unit (perpendicular to the bicycle plane) and the Os $5d_{xz}$ orbital, whereas HOMO-2 and HOMO-7 are formed by the orbital interactions between the p_π atomic orbitals of the C_6NH_6 unit and the Os $5d_{yz}$ orbital. The approximately uniform contribution of p_π orbitals from ring atoms in each of these occupied π molecular orbitals is typical for and consistent with a delocalised π molecular system. The four occupied π MOs calculated for the anti-aromatic azapentalene are also presented in Fig. 7b for comparison.

Conclusion

We synthesised the first examples of aza-metallapentalenes using metallabenzene as the starting material. The synthetic route to these aza-metallapentalenes is particularly interesting and remarkable. The bicyclic eight-membered ring complexes are derived from

sequential ring contractions, followed by annulation reactions of the six-membered metallabenzene via the five-membered metallapentafulvenes. Both experimental and theoretical studies suggest that the metallabenzene, metallapentafulvenes and aza-metallapentalenes all exhibit aromatic character. The present results not only expand the aromaticity concept but also open a promising avenue for the construction of new bicyclic metallaromatics containing main-group heteroatoms.

Methods

General considerations. Unless otherwise stated, all manipulations were carried out at room temperature under a nitrogen atmosphere using standard Schlenk techniques unless otherwise stated. Solvents were distilled from sodium/benzophenone (diethyl ether) or calcium hydride (dichloromethane) under N_2 prior to use. The starting material $[OsCl_2(PPh_3)_3]^{38}$, $HC \equiv CCH(OH)C \equiv CH^{39}$ and the osmabenzene $[Os\{CHC(PPh_3)CHCICH\}I_2(PPh_3)_2]$ was synthesised according to the previously published procedure²⁴. Other reagents were used as received from commercial sources without further purification. Further experimental details and the synthetic procedures for **1-I**, **3b-PF₆**, **3c-PF₆**, **4b-(PF₆)₂** and **4c-(PF₆)₂** are described in the Supplementary Methods. The procedures for **2-PF₆**, **3a-PF₆** and **4a-(PF₆)₂** are described below.

Synthesis of osmapentafulvene complex 2-PF₆. Aniline (60 μ L, 0.66 mmol) was added to a mixture of **1-I** (300 mg, 0.21 mmol) and sodium hexafluorophosphate (71 mg, 0.42 mmol) in CH_2Cl_2 (25 mL). The mixture was heated at reflux for approximately 10 h to afford a purple suspension. The solid suspension was removed by filtration, and the volume of the filtrate was reduced to approximately 2 mL under vacuum. The addition of ether (20 mL) to the solution then produced a purple solid that was collected by filtration, washed with diethyl ether (3×2 mL), and dried under vacuum. Yield: 282 mg, 95%. 1H NMR (500 MHz, CD_2Cl_2): δ = 11.9 (d, J (PH) = 15.6, 1 H, C'H), 10.8 (d, J (HH) = 14.6, 1 H, NH), 7.9 (d, J (HH) = 14.6, 1 H, C'H), 7.7

Figure 6 | Evaluation of aromaticity for osmabenzene **1, osmapentafulvene **2**, osmacycle **3**, and aza-osmapentalene **4a** by DFT calculations. (a) The NICS values calculated for the rings in osmabenzene **1**, osmapentafulvene **2**, osmacycle **3**, and aza-osmapentalene **4a**. (b) The ASE values calculated for the model complexes osmabenzene **1'**, osmapentafulvene **2'**, osmacycle **3'**, and aza-osmapentalene **4a'**. The energies computed at the B3LYP level using the LanL2DZ basis set for osmium and the 6-311++G(d,p) basis sets for carbon and hydrogen include zero-point energy corrections.**

Figure 7 | Molecular orbitals calculated for the aza-osmapentalene model 4a'' and azapentalene. The eigenvalues of the MO's are given in parentheses.

(br, 1 H, C³H), 6.6–7.8 ppm (m, 50H, Ph); ³¹P{¹H} NMR (202 MHz, CD₂Cl₂): δ = 13.0 (s, CPh₃), −3.1 (s, OsPPh₃), −144.4 ppm (septet, PF₆); ¹³C{¹H} NMR (126 MHz, CD₂Cl₂, plus ¹H-¹³C HSQC and ¹³C-dept 135): δ = 225.2 (br, C¹), 189.7 (br, C⁴), 175.5 (d, J(PC) = 22.0, C⁵), 169.3 (s, C⁵), 158.8 (br, Os(CO)), 137.2–118.1 (m, Ph), 122.0 (d, J(P,C) = 82.2 Hz, C²); HRMS (ESI): [M-PF₆]⁺ calcd for [C₆₆H₅₄NP₃IOOs]⁺, 1288.2072; found, 1288.2091; analysis (calcd., found for C₆₆H₅₄F₆INOOsP₄): C (55.35, 55.24), H (3.80, 3.72), N (0.98, 1.29).

Synthesis of osmabicyclic complex 3a-PF₆. A mixture of (Ph)CH(OH)C ≡ CH (40 mg, 0.30 mmol) Ag₂O (60 mg, 0.26 mmol), NEt₃ (1 mL) and 2-PF₆ (365 mg, 0.25 mmol) in dichloromethane (30 mL) was heated at reflux for 3 h to afford a blue suspension (Warning: the reaction may proceed through the formation of silver acetylide, which decomposes violently on contact with moisture and water producing highly flammable and explosive acetylene gas, and causing fire and explosion hazard). The solvent was removed under vacuum, and the residue was extracted with CH₂Cl₂ (3 × 5 mL). The volume of the filtrate was filtered through a Celite pad to remove the silver salt and was subsequently reduced to approximately 2 mL under vacuum. Diethyl ether (20 mL) was added slowly with stirring to afford a blue solid, which was collected by filtration, washed with hexane (3 × 3 mL), and dried under vacuum. Yield: 296 mg, 81%. ¹H NMR (500 MHz, CD₂Cl₂): δ = 11.4 (d, J(P,H) = 20.6 Hz, 1 H, C¹H), 7.5 (br, 1 H, C²H), 7.1 (s, 1 H, C⁵H), 6.9–7.9 (m, 55 H, Ph), 5.9 (d, J(H,H) = 8.8 Hz, 1 H, C⁷H), 5.3 (d, J(H,H) = 8.8 Hz, 1 H, C⁸H), 0.9 ppm (br, 1 H, OH); ³¹P{¹H} NMR (202 MHz, CD₂Cl₂): δ = 12.2 (s, CPh₃), 3.5 (d, J(P,P) = 218.6 Hz, OsPPh₃), 1.0 (J(P,P) = 218.6 Hz, OsPPh₃), −144.4 ppm (septet, PF₆); ¹³C{¹H} NMR (126 MHz, CD₂Cl₂, plus ¹³C-DEPT 135, ¹H-¹³C HSQC and ¹H-¹³C HMBC): δ = 233.0 (br, C¹), 196.0 (br, C⁴), 178.6 (br, C⁹), 169.0 (s, C⁵), 163.7 (br, C⁶), 161.5 (d, J(P,C) = 25.1 Hz, C³), 131.8 (s, C⁷), 123.7 (d, J(P,C) = 90.9 Hz, C²), 120.6–144.6 (m, Ph), 77.9 (s, C⁸); HRMS (ESI): [M-PF₆]⁺ calcd for [C₇₅H₆₁NP₃O₂Os]⁺, 1292.3524; found, 1292.3542; analysis (calcd., found for C₇₅H₆₁F₆NO₂OsP₄): C (62.71, 62.37), H (4.28, 4.19), N (0.98, 1.38).

Synthesis of aza-osmapentalene complex 4a-(PF₆)₂. A mixture of 3a-PF₆ (200 mg, 0.14 mmol) and sodium hexafluorophosphate (28 mg, 0.17 mmol) in CH₂Cl₂ (10 mL) was stirred at room temperature for 30 min to afford a red suspension. The solid suspension was removed by filtration, and the volume of the filtrate was reduced

to approximately 2 mL under vacuum. The addition of diethyl ether (20 mL) to the solution then produced a red solid that was collected by filtration, washed with diethyl ether (3 × 2 mL), and dried under vacuum. Yield: 213 mg, 98%. ¹H NMR (300 MHz, CD₂Cl₂): δ = 14.3 (d, J(P,H) = 16.5 Hz, 1 H, C¹H), 10.0 (br, 1 H, C³H), 9.9 (s, 1 H, C²H), 6.4–7.9 (m, 55 H, Ph), 7.7 (d, J(H,H) = 15.2, 1 H, C⁸H), 6.3 (d, J(H,H) = 15.2, 1 H, C⁷H); ³¹P{¹H} NMR (122 MHz, CD₂Cl₂): δ = 14.7 (s, CPh₃), 1.6 ppm (s, OsPPh₃), −144.4 ppm (septet, PF₆); ¹³C{¹H} NMR (76 MHz, CD₂Cl₂, and ¹³C-DEPT 135, ¹H-¹³C HSQC and ¹H-¹³C HMBC): δ = 248.5 (br, C¹), 234.1 (br, C⁶), 191.3 (br, C⁹), 187.6 (br, C⁴), 171.9 (d, J(P,C) = 21.4 Hz, C³), 163.0 (s, C⁵), 158.9 (s, C⁸), 151.0 (d, J(P,C) = 67.3 Hz, C²), 135.4 (s, C⁷), 117.9–140.5 (m, Ph); HRMS (ESI): [(M-2PF₆)/2]⁺ calcd for [(C₇₅H₆₀NP₃OOs)/2]⁺, 637.6746; found, 637.6763; analysis (calcd., found for C₇₅H₆₀F₁₂NOOsP₃): C (57.58, 57.86), H (3.87, 3.91), N (0.90, 1.30).

X-ray Crystallographic Analysis. Single crystals suitable for X-ray diffraction were grown from a dichloromethane solution layered with hexane. Diffraction data were collected on an Oxford Gemini S Ultra charge-coupled device (CCD) area detector (2-PF₆, 3a-PF₆, 4a-(PF₆)₂ and 4b-(PF₆)₂) or on a Bruker Apex CCD area detector (1-I) using graphite-monochromated Mo Kα radiation (λ = 0.71073 Å) or Cu Kα radiation (λ = 1.54178 Å). Semi-empirical or multi-scan absorption corrections (SADABS) were applied⁴⁰. All structures were solved by the Patterson function, completed by subsequent difference Fourier map calculations, and refined by full-matrix least-squares on F² with all the data using the SHELXTL program package⁴¹. All non-hydrogen atoms were refined anisotropically unless otherwise stated. Hydrogen atoms were placed at idealised positions and were refined using a riding model. See Supplementary Methods for detailed crystal data related to complexes 1-I, 2-PF₆, 3a-PF₆, 4a-(PF₆)₂ and 4b-(PF₆)₂.

Computational details. All structures were optimised at the B3LYP level of DFT^{42–44}. In addition, the frequency calculations were performed to confirm the characteristics of the calculated structures as minima. In the B3LYP calculations, the effective core potentials (ECPs) of Hay and Wadt with a double-ζ valence basis set (LanL2DZ) were used to describe the Os, P, and I atoms, whereas the standard 6-311++G(d,p) basis set was used for the C and H atoms⁴⁵ for all the ASE calculations. Polarisation functions were added for Os (ζ(f) = 0.886), I (ζ(d) = 0.266) and P (ζ(d) = 0.34)⁴⁶ in all of the calculations. NICS values were calculated at the B3-LYP-GIAO/6-311++G

(d,p) level. All the optimisations were performed with the Gaussian 03 software package⁴⁷. See Supplementary Materials for the Cartesian coordinates.

- Faraday, M. On new compounds of carbon and hydrogen and on certain other products obtained during the decomposition of oil by heat. *Philos. Trans. R. Soc. Lond.* **115**, 440–446 (1825).
- Kekulé, F. A. Sur la constitution des substances aromatiques. *Bull. Soc. Chim.* **3**, 98–111 (1865).
- Kekulé, F. A. Note sur quelques produits de substitution de la benzène. *Bull. Acad. R. Belg.* **19**, 551–563 (1865).
- Hückel, E. Quantum-theoretical contributions to the benzene problem. I. The electron configuration of benzene and related compounds. *Z. phys.* **70**, 204–286 (1931).
- Craig, D. P. A novel type of aromaticity. *Nature* **181**, 1052–1053 (1958).
- Heilbronner, E. Hückel molecular orbitals of Möbius-type conformations of annulenes. *Tetrahedron Lett.* **5**, 1923–1928 (1964).
- Rzepa, H. S. Möbius aromaticity and delocalization. *Chem. Rev.* **105**, 3697–3715 (2005).
- Mauksch, M., Gogonea, V., Jiao, H. & Schleyer, P. V. R. Monocyclic $(CH)_9^+$ — a Heilbronner Möbius aromatic system revealed. *Angew. Chem. Int. Ed.* **37**, 2395–2397 (1998).
- Garratt, P. J. *Aromaticity* (Wiley, 1986).
- Minkin, V. I., Glukhovtsev, M. N. & Simkin, B. Y. *Aromaticity and Antiaromaticity: Electronic and Structural Aspects* (Wiley, 1994).
- Schleyer, P. v. R. Thematic issue “Aromaticity” *Chem. Rev.* **101**, 115–1566 (2001).
- Cao, X.-Y., Zhao, Q., Lin, Z. & Xia, H. The chemistry of aromatic osmacycles. *Acc. Chem. Res.* **47**, 341–354 (2014).
- Chen, J. & Jia, G. Recent development in the chemistry of transition metal-containing metallabenzenes and metallabenzynes. *Coord. Chem. Rev.* **257**, 2491–2521 (2013).
- Frogley, B. J. & Wright, L. J. Fused-ring metallabenzenes. *Coord. Chem. Rev.* **270–271**, 151–166 (2014).
- Thorn, D. L. & Hoffmann, R. Delocalization in metallocycles. *Nouv. J. Chim.* **3**, 39–45 (1979).
- Weller, K. J., Filippov, I., Briggs, P. M. & Wigley, D. E. Pyridine degradation intermediates as models for hydrodenitrogenation catalysis: preparation and properties of a metalla-pyridine complex. *Organometallics* **17**, 322–329 (1998).
- Liu, B. *et al.* Osmapyridine and osmapyridinium from a formal [4 + 2] cycloaddition reaction. *Angew. Chem. Int. Ed.* **48**, 5430–5434 (2009).
- Wang, T. *et al.* Synthesis and characterization of a metalla-pyridyne complex. *Angew. Chem. Int. Ed.* **51**, 9838–9841 (2012).
- Zhu, C. *et al.* Planar Möbius aromatic pentalenes incorporating 16 and 18 valence electron osmiums. *Nat. Commun.* **5**, 3265–3271 (2014).
- Zhu, C. *et al.* A metal-bridged tricyclic aromatic system: synthesis of osmium polycyclic aromatic complexes. *Angew. Chem. Int. Ed.* **53**, 6232–6236 (2014).
- Hafner, K. & Schmidt, F. A Stable 2-Azapentalene. *Angew. Chem., Int. Ed. Engl.* **12**, 418–419 (1973).
- Elguero, J., Claramunt, R. M. & Summers, A. J. H. The chemistry of aromatic azapentalenes. *Adv. Heterocycl. Chem.* **22**, 183–320 (1978).
- Zhu, C. *et al.* Stabilization of anti-aromatic and strained five-membered rings with a transition metal. *Nat. Chem.* **5**, 698–703 (2013).
- Xia, H. *et al.* Osmabenzene from the Reactions of $HC \equiv CCH(OH)C \equiv CH$ with $OsX_2(PPh_3)_3$ ($X = Cl, Br$). *J. Am. Chem. Soc.* **126**, 6862–6863 (2004).
- Gong, L. *et al.* Osmabenzene from osmacycles containing an η^2 -coordinated olefin. *Chem. Eur. J.* **15**, 6258–6266 (2009).
- Wang, T. *et al.* cine-Substitution reactions of metallabenzene: an experimental and computational study. *Chem. Eur. J.* **19**, 10982–10991 (2013).
- Erker, G. Syntheses and reactions of fulvene-derived substituted aminoalkyl-Cp and phosphinoalkyl-Cp-Group 4 metal complexes. *Coord. Chem. Rev.* **250**, 1056–1070 (2006).
- Esteruelas, M. A., López, A. M. & Oliván, M. Osmium–carbon double bonds: formation and reactions. *Coord. Chem. Rev.* **251**, 795–840 (2007).
- Zhao, Q. *et al.* Stable isoosmabenzene from a formal [3 + 3] cycloaddition reaction of metal vinylidene with alkynols. *Angew. Chem. Int. Ed.* **50**, 1354–1358 (2011).
- Schleyer, P. V. R., Maerker, C., Dransfeld, A., Jiao, H. & Hommes, N. J. R. V. E. Nucleus-independent chemical shifts: a simple and efficient aromaticity probe. *J. Am. Chem. Soc.* **118**, 6317–6318 (1996).
- Chen, Z., Wannere, C. S., Corminboeuf, C., Puchta, R. & Schleyer, P. V. R. Nucleus-independent chemical shifts (NICS) as an aromaticity criterion. *Chem. Rev.* **105**, 3842–3888 (2005).
- Fallah-Bagher-Shadaei, H., Wannere, C. S., Corminboeuf, C., Puchta, R. & Schleyer, P. V. R. Which NICS aromaticity index for planar π rings is best? *Org. Lett.* **8**, 863–866 (2006).
- Mauksch, M. & Tsogoeva, S. B. Demonstration of “Möbius” aromaticity in planar metallocycles. *Chem. Eur. J.* **16**, 7843–7851 (2010).
- Iron, M. A., Lucassen, A. C. B., Cohen, H., van der Boom, M. E. & Martin, J. M. L. A computational foray into the formation and reactivity of metallabenzene. *J. Am. Chem. Soc.* **126**, 11699–11710 (2004).
- Periyasamy, G., Burton, N. A., Hillier, I. H. & Thomas, J. M. H. Electron delocalization in the metallabenzene: a computational analysis of ring currents. *J. Phys. Chem. A* **112**, 5960–5972 (2008).
- Han, F., Wang, T., Li, J., Zhang, H. & Xia, H. m-Metallaphenol: synthesis and reactivity studies. *Chem. Eur. J.* **20**, 4363–4372 (2014).
- Schleyer, P. V. R. & Pühlhofer, F. Recommendations for the evaluation of aromatic stabilization energies. *Org. Lett.* **4**, 2873–2876 (2002).
- Hoffmann, P. R. & Caulton, K. G. Solution structure and dynamics of five-coordinate d^6 complexes. *J. Am. Chem. Soc.* **97**, 4221–4228 (1975).
- Jones, E. R. H., Lee, H. H. & Whiting, M. C. Researches on acetylenic compounds. Part LXIV.* The preparation of conjugated octa- and deca-acetylenic compounds. *J. Chem. Soc.* 3483–3489 (1960).
- Sheldrick, G. M. SADABS, Program for Semi-Empirical Absorption Correction (University of Göttingen, Göttingen: Germany, 1997).
- Sheldrick, G. M. SHELXTL (Siemens Analytical X-ray Systems, Madison, Wisconsin, USA, 1995).
- Becke, A. D. Density-functional thermochemistry. III. The role of exact exchange. *J. Chem. Phys.* **98**, 5648–5652 (1993).
- Miehlich, B., Savin, A., Stoll, H. & Preuss, H. Results obtained with the correlation energy density functionals of Becke and Lee, Yang and Parr. *Chem. Phys. Lett.* **157**, 200–206 (1989).
- Lee, C., Yang, W. & Parr, R. G. Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Phys. Rev. B* **37**, 785–789 (1988).
- Hay, P. J. & Wadt, W. R. Ab initio effective core potentials for molecular calculations. Potentials for potassium to gold including the outermost core orbitals. *J. Chem. Phys.* **82**, 299–310 (1985).
- Huzinaga, S. *Gaussian Basis Sets for Molecular Calculations* (Elsevier, Amsterdam, 1984).
- Frisch, M. J. *et al.* Gaussian 03, Revision E.01 (Gaussian, Inc., Wallingford, CT, 2004).

Acknowledgments

This research was supported by the National Science Foundation of China (Nos. 21272193 and 21332002), and the Research Grants Council of Hong Kong (HKUST603313).

Author contributions

H.X. and H.Z. designed the research. T.W., H.H. and J.L. performed the experiments. T.W. and F.H. recorded all NMR data and solved all X-ray structures. H.X., H.Z. and T.W. analysed the experimental data. H.Z. performed the theoretical computations and analysed the theoretical data with assistance from Z.L. and J.Z. H.Z. wrote the paper with assistance from Z.L. All authors discussed the results and contributed to the preparation of the final manuscript.

Additional information

Accession codes: The X-ray crystal structure information is available at the Cambridge Crystallographic Data Centre (CCDC) under deposition numbers CCDC-1013726 (1-I), CCDC-1013727 (2-PF₆), CCDC-1013728 (3a-PF₆), CCDC-1013729 (4a-(PF₆)₂) and CCDC-1013730 (4b-(PF₆)₂). These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Wang, T. *et al.* Synthesis of Aromatic Aza-metallapentalenes from Metallabenzene via Sequential Ring Contraction/Annulation. *Sci. Rep.* **5**, 9584; DOI:10.1038/srep09584 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>