SCIENTIFIC REPORTS

Corrigendum: Estimation of the hydrogen concentration in rat tissue using an airtight tube following the administration of hydrogen via various routes

Chi Liu, Ryosuke Kurokawa, Masayuki Fujino, Shinichi Hirano, Bunpei Sato & Xiao-Kang Li

Scientific Reports 4:5485; doi: 10.1038/srep05485; published online 30 June 2014; updated on 20 July 2015

This Article contains an error in describing the unit of hydrogen concentration in Figures 2–5: ‘ppb/g’ should read ‘ppm/g’ in the Y-axes. The correct Figures 2–5 appear below as Figures 1, 2, 3, 4 respectively.

In the Results section:

“Furthermore, a C_{max} analysis showed that the oral administration of HSRW resulted in a high hydrogen concentration (more than 300 ppb/g) in the tissues of the spleen, small intestine and pancreas (Fig. 4A). Meanwhile, the intraperitoneal administration of HSRS resulted in a high hydrogen concentration in the spleen and pancreas (more than 300 ppb/g) (Fig. 4B). However, the hydrogen concentrations were not high (more than 35 ppb/g) in any of the tissues after intravenous treatment (Fig. 4C). Interestingly, the inhalation of hydrogen gas resulted in the highest hydrogen concentration in the muscle (more than 140 ppb/g; Fig. 4D).”

should read:

“Furthermore, a C_{max} analysis showed that the oral administration of HSRW resulted in a high hydrogen concentration (more than 300 ppm/g) in the tissues of the spleen, small intestine and pancreas (Fig. 4A). Meanwhile, the intraperitoneal administration of HSRS resulted in a high hydrogen concentration in the spleen and pancreas (more than 300 ppm/g) (Fig. 4B). However, the hydrogen concentrations were not high (more than 35 ppm/g) in any of the tissues after intravenous treatment (Fig. 4C). Interestingly, the inhalation of hydrogen gas resulted in the highest hydrogen concentration in the muscle (more than 140 ppm/g; Fig. 4D).”

In the Methods section under ‘Tissue dissociation and the determination of the hydrogen concentrations in the arterial blood and tissues’:

“The hydrogen concentration in each blood or tissue sample was defined as follows: Hydrogen concentration (ppb/g; V/V) = A/B × C; where A is the measurement value obtained using the Sensor Gas Chromatograph, B is the blood or tissue weight (g) and C is the volume of the gentleMACS tube (24 ml).”

should read:

“The hydrogen concentration in each blood or tissue sample was defined as follows: Hydrogen concentration (ppm/g; V/V) = A/B × C; where A is the measurement value obtained using the Sensor Gas Chromatograph, B is the blood or tissue weight (g) and C is the volume of the gentleMACS tube (24 ml).”

Figure 1.

Figure 2.

Figure 3.

Figure 4.