SCIENTIFIC REPORTS

OPEN

Flower evolution of alpine forbs in the open top chambers (OTCs) from the Qinghai-Tibet Plateau

Chan Zhang^{1,2,3,4}, Lin-Lin Wang^{1,2,3}, Yong-Ping Yang^{1,2,3} & Yuan-Wen Duan^{1,2,3}

Received: 10 October 2014

Accepted: 08 April 2015

Published: 22 May 2015

Effects of global changes on biodiversity have been paid more and more attention world widely, and the open top chambers (OTCs) are the most common tools to study the effects of climatic warming on plant diversity. However, it remains unclear how flowers evolve under environmental changes, which could help us to understand the changes of plant diversity in the OTCs. We compared the insect diversity and pollen:ovule (P/O) ratio of eight outcrossing species with different life histories inside and outside the OTCs on the Qinghai-Tibet Plateau, to examine the effects induced by OTCs on the evolution of floral traits. In the OTCs, P/O ratio decreased in annuals, but increased in perennials, indicating an overall trend toward selfing in annuals. We found that the insect diversity differed significantly inside and outside the OTCs, with decreases of dipteran insects and bees. We concluded that changes of P/O ratio in the studied plant species might result from pollination failure, which might be the results of mismatch between flowering time and pollinator activities. We also suggested annuals might be in a more extinction risk than perennials in OTCs, if strong inbreeding depression occurs in these annual outcrossing plants.

Global changes, especially warming, have been paid more and more attention world widely. The complex ecological consequences resulted from warming could change the biodiversity via altering number¹ and distribution² of plant species. For example, experimental warming induced by open top chambers (OTCs) caused a 26–36% decrease in species richness on the Qinghai-Tibet Plateau, although this decrease was generally dampened by experimental grazing³. Alpine and arctic ecosystems are known to be particularly sensitive to climate warming, and thus many experimental examinations on the effects of climatic warming were performed on alpine and arctic areas^{4,5}. Experimental warming suggested that flowering phenology of plant species could be advanced or delayed, depending on the flowering time of specific species before or after the peak of the summer heat⁶. These changes in flowering time might induce pollination failure of plant species in the warming plots due to the mismatch between plants and the corresponding pollinators, and the long time pollination failure might drive the evolution of mating systems^{7–9}. However, the effects of OTCs on plant mating system have rarely been examined.

Pollen:ovule (P/O) ratio is an important indicator of mating system and P/O ratios per hermaphroditic flowers of outcrossing species were generally higher than those of selfing plants^{10–12}. Therefore, changes in P/O ratio might reflect the evolutionary trend in mating system. A recent study suggested that variations in P/O ratio among taxa was mainly driven by pollinator efficiency¹³, which to a certain degree could indicate the variations of pollination visitation frequency among taxa. Additionally, intra-specific variations of P/O ratio were also observed within inflorescences, and these variations were attributed to

¹Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan, P. R. China. ²Plant Germplasm and Genomics Center, the Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan, P. R. China.

³Institute of Tibetan Plateau Research at Kunming, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan, P. R. China. ⁴Graduate University of the Chinese Academy of Sciences, Beijing 100049, Beijing, P. R. China. Correspondence and requests for materials should be addressed to Y.W.D. (email: duanyw@mail.kib.ac.cn) or Y.P.Y. (email: yangyp@mail.kib.ac.cn)

Figure 1. Changes of the number of pollen grains (A), ovule (B) and pollen:ovule ratio (C) in the plants species with different life histories from the Qinghai-Tibet Plateau in the open top chambers. Numbers indicated the comparison results (P values) of flowers between the OTCs and control using independent samples t test. Species number was shown in Table S1. Plants from No. 1 to 4 are annuals, and from No. 5 to 8 are perennials.

mating environment^{14–16} and/or resource allocation^{17,18}, indicating that experiments on changes of P/O ratio from flowers subjected to different treatments should be cautiously treated.

The Qinghai-Tibet Plateau is the largest and highest plateau in the world, and this region is predicted to experience “much greater than average” increases in surface temperature in the future. Studies have found that experimental warming causes large and rapid species loss³ and decreases rangeland quality¹⁹ in this region. In addition, the OTCs in the Qinghai-Tibet Plateau around this station have been established for more than ten years³. However, it remains unclear how flowers and the mating system of plant species evolve under environmental changes induced by the OTCs, and these results could help us to understand why some plant species disappeared under experimental warming in the OTCs. In the present study, we compared the insect diversity and P/O ratio in eight outcrossing species with different life histories inside and outside the OTCs, to examine the effects induced by OTCs on the evolution of floral traits. Specifically, we addressed the following questions: 1) what are the differences in the insect diversity between inside and outside the OTCs; 2) what are the differences in the P/O ratio between inside and outside the OTCs? Our results would be of great help to understand the evolutionary trend of mating systems of plant species in the OTCs, and provide a preliminary support on the priority of biodiversity conservation in the changing world.

Results

The number of pollen grains of the plant species was affected consistently by the OTCs, specified by the decrease in annual plants and increase in perennial plants, although the difference was not significant between inside and outside the OTCs across all the studied species (Fig. 1A). However, the ovule number did not show consistent changes across the eight plant species (Fig. 1B). Inside the OTCs, the ovule numbers of five plant species decreased, but the ovule number of *Gentiana leucomelaena* increased in the OTCs, and the ovule numbers of *Stellera chamaejasme* and *Allium cyaneum* were constant (Fig. 1B). For the P/O ratio, the effects of OTCs was conflicted for the annual and perennial plants, detailed by the decrease in annual plants and increase in perennial plants (Fig. 1C).

Figure 2. Insect diversity inside and outside the open top chambers.

Overall, our observations suggested that the OTCs changed the insect diversity significantly in comparison with the control outside the OTCs (Fig. 2, $\chi^2 = 14.3$, $p = 0.003$). Specifically, compared to the control plots, there were ca 4% increases of lepidopteran insects, 4.7% increases of hymenopteran insects, 14.2% increases of other insects and 22.8% decrease of dipteran insects in the OTCs. Furthermore, for the hymenopteran insects, ca 10.3% reduction of the bee visitations were found inside the OTCs in comparison with those outside the OTCs. Collectively, the insect diversity has been changed inside the OTCs, indicating the changed selective pressures on flowers from the pollinators.

Discussion

For the P/O ratio of our studied species in the OTCs, there was a general decrease in annuals, indicating an evolutionary trend towards selfing. While for the perennials, there was a general increase in P/O ratio in the OTCs, indicating the evolutionary trend towards outcrossing. Variations of P/O ratio were demonstrated recently to be governed by pollination efficiency¹³, and a low P/O ratio is necessary to guarantee seed set, which was generally found in plant species with mating systems of selfing^{20,21}. Accordingly, changes in P/O ratio could reflect the variations in selective pressures of pollinators and the evolutionary trend in mating system. In the OTCs, we found that insects did not disappear completely, despite of the facts that insect composition inside the OTCs has changed and the dominant pollinators of alpine ecosystems which consisted of bees and flies²², have decreased significantly. Thus it seemed that pollinator changes might contribute little to the changes of P/O ratio in the OTCs in our study site. In addition, intra-specific variations of P/O ratio from different positions within inflorescences were common in plant species^{14,15,17,18}, but both mating environment hypothesis and resource allocation cannot explain our results since our samples are from the same position on the inflorescences for each species in and outside of the OTCs. It has been demonstrated that experimental warming could advance or delay the flowering time of plant species⁶, which could induce the mismatch between plants and the corresponding pollinators and further reduce pollinator service of plant species in the OTCs. Collectively, changes of P/O ratio of the eight outcrossing plants in the OTCs could be induced by the pollination failure, although this argument needs future experimental supports.

Our results suggested the different responses for annual and perennial plants in the OTCs. There was a trend towards decreased ovule number, but no consistent response was found in the changes in both annuals and perennials (Fig. 1B). However, for the number of pollen grains, we found that there was a decreasing trend for the four annual plants and an increasing trend for the perennial plants (Fig. 1A), suggesting that the variations in P/O ratio of plants could be resultant from changes in the number of pollen grains, and indicating that resource allocation to male functions of plant species were more susceptible to environmental changes²³. Importantly, different evolutionary trends in P/O ratio of plants with different life histories could be explained in the following. For the annual plants, long-time pollination failure might drive the evolution of selfing, and then reduced P/O ratio was expected^{10,13}. However, for the perennials, because reproduction is costly in plants, current failures in seed production might induce the enhanced investments to future reproduction under the hypothesis on the trade-off between current and future reproduction²⁴. Furthermore, higher cost of reproduction in females than in males²⁵ could drive the future allocation to pollen production, resulting in the increased P/O ratio of

the perennials in the OTCs. Therefore, the life history traits could contribute to the different responses in P/O ratio of plants since the ten-year's treatment is far less than the history of angiosperm, although evolutionary trends were obvious in both annuals and perennials.

The temperature inside the OTCs has been elevated across the growing seasons²⁶, but the evolutionary trend towards selfing in annual plants could not be resultant from experimental warming. Furthermore, this evolutionary trend might increase the extinction risk of annual plants with mating system of outcrossing, if selfing occurs in these species and inbreeding depression is strong²⁷. Therefore, facing the global change, especially global warming, we tentatively suggested that we should give a conservation priority to those endangered annual plants with outcrossing, indicating the importance of studying breeding systems of endangered and endemic annual plants in the future conservation of biodiversity. Furthermore, we tentatively draw another conclusion that the widely used OTCs for experimental warming might not be an ideal method in studying the effects of warming on special level, especially when studying the relationships between plants and their mutualisms, e.g. pollinators, florivores, herbivores and seed predators.

Materials and Methods

Study site and materials. Our studies were carried around the Haibei Alpine Meadow Ecosystem Research Station, the Chinese Academy of Science. The station is located on the northeast corner of the Qinghai-Tibet Plateau, with lat. 37°29'–37°45' N, long. 101°12'–101°23' E, and alt. 3200 m. The average annual air temperature is −1.7°C with extremes of 27.6°C (maximum) and −37.1°C (minimum). Average annual precipitation ranges from 426 to 860 mm, 80% of which falls in the summer from late May to early September²⁸.

The open top chambers (OTCs) around this station have been established for more than ten years³. The OTCs, which were 1.5 m diameter and 40 cm height, were constructed of Sun-Lite HP (Solar Components Corporation, Manchester, NH, USA) 1.0 mm thick fiberglass and remained on the grassland year-round³. The OTCs consistently elevated growing season averaged mean daily air temperature by 1.0–2.0°C, maximum daily air temperature by 2.1–7.3°C and the diurnal air temperature range by 1.9–6.5°C, with mixed effects on minimum daily air temperature, and mean daily soil temperature and moisture²⁶.

The eight studied species and their characteristics are listed in Table S1, and all of the eight species are the common forbs inside and outside the open top chambers (OTCs). According to our observations, netting treatment and former studies^{29–32}, flowers of the eight plants cannot produce seeds without aids of pollinator, indicating (facultative) outcrossing mating systems in the eight plants. For example, all collected plant materials in Gentianaceae in the experiments are characteristics of protandry and/or herkogamy.

Methods. *Pollen and ovules.* For each of the eight species (Table S1), we selected twenty flower buds that will be open shortly inside and outside the OTCs, and fixed them in standard FAA solution (formalin:acetic acid:100% ethanol at a ratio of 5:5:90 by volume). To avoid the effects of architecture and resources from different positions on flower traits³³, only the terminal flowers were selected in this experiment. In the laboratory, we squashed all anthers from each flower and suspended all pollen grains in 5 mL of water with drops of detergent solution for full suspension³⁴. Total pollen in each flower was determined by counting the number of pollen grains in 20 drops of pollen solution (5 µL) with a light microscope. The number of ovules was determined under a stereoscope by dissecting all the carpels of the flower. We then calculated the pollen:ovule ratio for each flower of each species.

Insect observations. In the late May, 2011, we established ten plots adjacent to the OTCs, and each plot was similar to the OTC in size, and observed insects that entered the OTCs and the plots from early June to late August. In each month of the early-, middle- and late-stage, we monitored and recorded the insects in three sunny days without strong wind from 9:30 to 16:30 when the temperature was high, and we would cease the observations if rain occurred. Totally, the observation time amounted to 170 h. In light of our long time observations and experiments on pollination ecology of alpine plant around this station³⁵, the insects were classified as Lepidopteran, Hymenopteran, Dipteran and other insects, such as beetles.

Statistical analysis. The difference in the insect diversity inside and outside the OTCs were analyzed using a Chi-square test, and the number of pollen grains and ovules, and pollen:ovule ratio were ln-transformed before comparison, and independent sample *t* test was used to compare the differences inside and outside OTCs. The effects of OTCs were calculated as $\ln(X_T/X_C)$, where X_T and X_C are the averaged data of flowers in OTCs and control (outside OTCs), respectively³⁶.

References

1. Pimm, S. L. Climate disruption and biodiversity. *Curr. Biol.* **19**, R595–R601 (2009).
2. Høye, T. T., Post, E., Meltofte, H., Schmidt, N. M. & Forchhammer, M. C. Rapid advancement of spring in the high Arctic. *Curr. Biol.* **17**, R449–R451 (2007).
3. Klein, J. A., Harte, J. & Zhao, X.-Q. Experimental warming causes large and rapid species loss, dampened by simulated grazing, on the Tibetan Plateau. *Ecol. Lett.* **7**, 1170–1179 (2004).

4. Cornelissen, J. H. C. *et al.* Global negative vegetation feedback to climate warming responses of leaf litter decomposition rates in cold biomes. *Ecol. Lett.* **10**, 619–627 (2007).
5. Walker, M. D. *et al.* Plant community responses to experimental warming across the tundra biome. *Proc. Natl. Acad. Sci.* **103**, 1342–1346 (2006).
6. Sherry, R. A. *et al.* Divergence of reproductive phenology under climate warming. *Proc. Natl. Acad. Sci.* **104**, 198–202 (2007).
7. Kalisz, S., Vogler, D. W. & Hanley, K. M. Context-dependent autonomous self-fertilization yields reproductive assurance and mix mating. *Nature* **430**, 884–887 (2004).
8. Kalisz, S. & Vogler, D. W. Benefits of autonomous selfing under unpredictable pollinator environments. *Ecology* **84**, 2928–2942 (2003).
9. Kalisz, S. *et al.* The mechanism of delayed selfing in *Collinsia verna* (Scrophulariaceae). *Am. J. Bot.* **86**, 1239–1247 (1999).
10. Cruden, R. W. Pollen-ovule ratios: a conservative indicator of breeding systems in flowering plants. *Evolution* **31**, 32–46 (1977).
11. Scalone, R., Kolf, M. & Albach, D. C. Mating system variation in *Veronica* (Plantaginaceae): inferences from pollen/ovule ratios and other reproductive traits. *Nord. J. Bot.* **31**, 372–384 (2013).
12. Cruden, R. W. Pollen grains: why so many? *Plant Syst. Evol.* **222**, 143–165 (2000).
13. Gong, Y. B. & Huang, S. Q. Interspecific variation in pollen-ovule ratio is negatively correlated with pollen transfer efficiency in a natural community. *Plant Biol.* **16**, 843–847 (2014).
14. Brunet, J. Male reproductive success and variation in fruit and seed set in *Aquilegia caerulea* (Ranunculaceae). *Ecology* **77**, 2458–2471 (1996).
15. Huang, S. Q., Tang, L. L., Yu, Q. & Guo, Y. H. Temporal floral sex allocation in protogynous *Aquilegia yabeana* contrasts with protandrous species: Support for the mating environment hypothesis. *Evolution* **58**, 1131–1134 (2004).
16. Brunet, J. & Charlesworth, D. Floral sex allocation in sequentially blooming plants. *Evolution* **49**, 70–79 (1995).
17. Diggle, P. K. Architectural effects and the interpretation of patterns of fruit and seed development. *Annu. Rev. Ecol. Syst.* **26**, 531–552 (1995).
18. Diggle, P. K. Extreme preformation in alpine *Polygonum viviparum*: an architectural and developmental analysis. *Am. J. Bot.* **84**, 154–169 (1997).
19. Klein, J. A., Harte, J. & Zhao, X. Q. Experimental warming, not grazing, decreases rangeland quality on the Tibetan Plateau. *Ecol. Appl.* **17**, 541–557 (2007).
20. Duan, Y. W., Dafni, A., Hou, Q. Z., He, Y. P. & Liu, J. Q. Delayed Selfing in an Alpine Biennial *Gentianopsis paludosa* (Gentianaceae) in the Qinghai-Tibetan Plateau. *J. Integr. Plant Biol.* **52**, 593–599 (2010).
21. Duan, Y. W., He, Y. P., Zhang, T. F. & Liu, J. Q. Delayed selfing in an alpine species *Gentianopsis barbata*. *Chinese J. Plant Ecol.* **31**, 110–117 (2007).
22. Körner, C. *Alpine plant life*, 2nd edition. (Springer, 2003).
23. Schaeffer, R. N., Manson, J. S. & Irwin, R. E. Effects of abiotic factors and species interactions on estimates of male plant function: a meta-analysis. *Ecol. Lett.* **16**, 399–408 (2013).
24. Obeso, J. R. The costs of reproduction in plants. *New Phytol.* **155**, 321–348 (2002).
25. Delph, L. F. Sexual dimorphism in life history. In Geber M. A., Dawson T. E., & Delph L. F. (eds) *Gender and Sexual Dimorphism in Flowering Plants* (Springer-Berlin, 1999).
26. Klein, J. A., Harte, J. & Zhao, X. Q. Dynamic and complex microclimate responses to warming and grazing manipulations. *Global Change Biol.* **11**, 1440–1451 (2005).
27. O'Grady, J. J. *et al.* Realistic levels of inbreeding depression strongly affect extinction risk in wild populations. *Biol. Conserv.* **133**, 42–51 (2006).
28. Zhao, X. Q. & Zhou, X. M. Ecological basis of Alpine meadow ecosystem management in Tibet: Haibei Alpine Meadow Ecosystem Research Station. *Ambio* **28**, 642–647 (1999).
29. Duan, Y. W., Zhang, T. F. & Liu, J. Q. Interannual fluctuations in floral longevity, pollinator visitation and pollination limitation of an alpine plant (*Gentiana straminea* Maxim., Gentianaceae) at two altitudes in the Qinghai-Tibetan Plateau. *Plant Syst. Evol.* **267**, 255–265 (2007).
30. Duan, Y. W., He, Y. P. & Liu, J. Q. Reproductive ecology of the Qinghai-Tibet Plateau endemic *Gentiana straminea* (Gentianaceae), a hermaphrodite perennial characterized by herkogamy and dichogamy. *Acta Oecol.* **27**, 225–232 (2005).
31. He, Y. P. & Liu, J. Q. Pollination ecology of *Gentiana straminea* Maxim. (Gentianaceae), an alpine perennial in the Qinghai-Tibet Plateau. *Acta Ecol. Sin.* **24**, 215–220 (2004).
32. Zhang, Z. Q., Zhang, Y. H. & Sun, H. The reproductive biology of *Stellera chamaejasme* (Thymelaeaceae): A self-incompatible weed with specialized flowers. *Flora* **206**, 567–574, (2011).
33. Diggle, P. K. Ontogenetic contingency and floral morphology: The effects of architecture and resource limitation. *Int. J. Plant Sci.* **158**, S99 (1997).
34. Dafni, A., Pacini, E. & Nepi, M. Pollen and stigma biology. In Dafni A., Kevan P. G. & Husband B. C. (eds) *Practical Pollination Biology* (Enviroquest Ltd, 2005).
35. Liu, J. Q., Duan, Y. W., Hao, G., Ge, X. J. & Sun, H. Evolutionary history and underlying adaptation of alpine plants on the Qinghai-Tibet Plateau. *J. Syst. Evol.* **52**, 241–249 (2014).
36. Lin, D. L., Xia, J. Y. & Wan, S. Q. Climate warming and biomass accumulation of terrestrial plants: a meta-analysis. *New Phytol.* **188**, 187–198 (2010).

Acknowledgements

We are grateful to Prof. Xin-Quan Zhao for his supports to our filed work. This work was financially supported by 973 programs (2014CB954100) and NSFC (31270434).

Author Contributions

Y.W.D. and Y.P.Y. designed the research and wrote the manuscript; C.Z., L.L.W. and Y.W.D. performed experiments; Y.W.D. analysed data and prepared the figures and tables.

Additional Information

Supplementary Table S1: Traits of eight outcrossing plant species and their flowers. The ovule numbers of species 5 and 8 are constant.

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhang, C. *et al.* Flower evolution of alpine forbs in the open top chambers (OTCs) from the Qinghai-Tibet Plateau. *Sci. Rep.* **5**, 10254; doi: 10.1038/srep10254 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>