SCIENTIFIC REPORTS

OPEN

Novel anti-thrombotic agent for modulation of protein disulfide isomerase family member ERp57 for prophylactic therapy

Received: 30 November 2014

Accepted: 09 April 2015

Published: 03 June 2015

Guozhen Cui^{1,2,*}, Luchen Shan^{3,4,*}, Lin Guo³, Ivan Keung Chu⁵, Guohui Li⁵, Quan Quan³, Yun Zhao⁵, Cheong Meng Chong¹, Zaijun Zhang³, Pei Yu³, Maggie Pui Man Hoi¹, Yewei Sun³, Yuqiang Wang³ & Simon MingYuen Lee¹

Protein disulfide isomerase (PDI) family members including PDI and ERp57 emerge as novel targets for anti-thrombotic treatments, but chemical agents with selectivity remain to be explored. We previously reported a novel derivative of danshensu (DSS), known as ADTM, displayed strong cardioprotective effects against oxidative stress-induced cellular injury *in vitro* and acute myocardial infarct *in vivo*. Herein, using chemical proteomics approach, we identified ERp57 as a major target of ADTM. ADTM displayed potent inhibitory effects on the redox activity of ERp57, inhibited the adenosine diphosphate (ADP)-induced expressions of P-selectin and α IIb β 3 integrin, and disrupted the interaction between ERp57 and α IIb β 3. In addition, ADTM inhibited both arachidonic acid (AA)-induced and ADP-induced platelet aggregation *in vitro*. Furthermore, ADTM significantly inhibited rat platelet aggregation and thrombus formation *in vivo*. Taken together, ADTM represents a promising candidate for anti-thrombotic therapy targeting ERp57.

Cardiovascular diseases (CVDs) remain the leading causes of morbidity and mortality globally. The underlying causes of CVDs are complex and multi-factorial involving many risk factors such as abnormal blood lipid, high blood cholesterol levels, overweight and sedentary lifestyle^{1,2}. In particular, platelet dysfunction and aberrant thrombogenesis (thrombus formation) are final underlying pathophysiological processes in the development and progression in CVDs including atherosclerosis, thrombotic complications, stroke and myocardial infarction³. Current anti-platelet agents such as aspirin and clopidogrel are effective at decreasing platelet activation and subsequent risk of atherothrombotic disease, but these agents irreversibly inhibit platelet activation and lead to risks of unwanted hemorrhages^{4,5}. Furthermore, the multi-factorial nature of CVDs urges us to look for versatile pharmacological agents with fewer side effects.

We have reported previously that a novel cardioprotective compound known as ADTM was synthesized in our laboratory by conjugating two well-known active ingredients, DSS and tetramethylpyrazine (TMP), which are present in Chinese medicinal plants, danshen (*Salvia miltiorrhizae*) and chuanxiong (*Ligustium wallichii* Franch)⁶. ADTM displayed strong cardioprotective effects against oxidative stress-induced cellular injury and acute ischemic myocardial infarct in rat, partly through activation of

¹State Key Laboratory of Quality Research in Chinese Medicine and Institute of Chinese Medical Sciences, University of Macau, Macao, China. ²Department of Bioengineering, Zhuhai Campus of Zunyi Medical University, Guangdong, Zhuhai, China. ³Institute of New Drug Research, College of Pharmacy, Jinan University, Guangzhou, China. ⁴Key Laboratory of Cardiovascular Medicine Research, Ministry of Education, Harbin Medical University, Harbin, China. ⁵Department of Chemistry, The University of Hong Kong, Hong Kong, China. ⁶These authors contributed equally to this work. Correspondence and requests for materials should be addressed to M.P.M.H. (email: maghoi.um@gmail.com) or Y.W.S. (email: syw1984@163.com) or S.M.Y.L. (email: simonlee@umac.mo)

Figure 1. Chemical structures of DSS, TMP, ADTM, and BAA.

the anti-oxidative stress Nrf2/heme oxygenase-1 (HO-1) pathway⁷. Several registered pharmaceutical products containing danshen and/or chuanxiong such as “Danshen Dripping Pill” and “Guanxinjing Injection” have been widely used in China to treat CVDs. The “Danshen Dripping Pill” was approved by the US Food and Drug Administration (FDA) to undergo Phase III clinical trials in the treatment of CVDs⁸. Clinical studies indicated that these pharmaceutical products are effective in treating angina pectoris, a symptom of ischemic heart disease⁹. It is believed that active ingredients in danshen and chuanxiong induce beneficial therapeutic effects by the promotion of better blood circulation^{10,11}. Therefore, we postulated that ADTM might also display anti-platelet and anti-thrombotic properties.

Several members of the protein disulfide isomerases have been shown as important players in the regulation of platelet functions and thrombus formation including PDI, ERp72, ERp57 and ERp5^{12–14}. However, which member in the family represents the best anti-thrombotic drug target is largely unknown¹³. The prototypical member PDI is a multifunctional redox chaperone protein that plays critical roles in a variety of diseases including platelet aggregation and thrombus formation *in vivo* and *in vitro*^{15–17}. More recently, it is shown that blood platelets also secreted ERp57 (also known as GRP58 and PDIA3) in response to vascular injury to trigger platelet activation and aggregation and emerges as a novel target for anti-platelet aggregation and anti-thrombosis^{18–20}. Human ERp57 has 32% amino acid sequence homology with PDI, and exhibits 40% identity with the a-b-b'-a' domains of ERp72²¹. However, selective chemical agents for ERp57 are not yet available.

In this study, we further identified the drug target of ADTM as ERp57 by using a chemical proteomics approach analyzing liquid chromatography linked to tandem mass spectrometry (LC-MS/MS) profile of pull-down proteins bound to a biotinylated-ADTM analogue (BAA). The anti-platelet aggregation and anti-thrombosis activities of ADTM were determined both *in vitro* and *in vivo*. The results provided promising evidences for the potential of developing novel anti-CVD and anti-thrombotic agents with selectivity at the protein disulfide isomerases by using ADTM.

Results

Protein disulfide isomerases are identified as targets of ADTM by chemical proteomics. In order to isolate the protein targets that bound to ADTM, a biotin-conjugated ADTM analogue (BAA, Fig. 1) was designed and synthesized in our laboratory as described in Supplementary Methods (Supplemental Fig. 3). BAA (300 μ M) was incubated with rat blood platelet lysates and the BAA-protein complexes were pulled down with NeutrAvidin-agarose followed by protein profiling using LC-MS/MS. It was observed that BAA could bind to various proteins involved in platelet function, including glycoprotein 5 (a potential marker of thrombotic platelet activation^{22,23}), superoxide dismutase and xanthine oxidase (regulators of vascular redox signaling²⁴). In particular, four platelet aggregation-associated proteins were identified with >95% protein identification probability (Supplementary Table S1), including ERp72, ERp57, ERp5 and PDI, which are glycoprotein-specific members of the PDI family related to platelet function and redox homeostasis^{25,26}. We confirmed the identity of the pulled-down proteins as ERp72, ERp57 and ERp5 with their respective antibodies by using immunoblot analysis (Fig. 2b). Furthermore, the specificity of the binding between BAA and these targets (ERp72, ERp57 and ERp5) could be demonstrated by the competition with untagged ADTM in excess (Fig. 2a). The molecular docking analysis demonstrated that the free binding energy to the active cavity of ERp57 (PDB: 3F8U) is -6.5 kcal/mol. ADTM may form hydrogen bonds with side chains of ARG47, ILE453 and PHE450, respectively, in the catalytic center (CYS406/CYS409) of ERp57 (Supplemental Fig. 2). ERp72, ERp57, ERp5 and PDI are identified as important new targets in redox homeostasis of blood platelet aggregation¹², therefore we further evaluated the effects of ADTM on the reducing activities of the PDI family members. As shown in Fig. 2c, ADTM exhibited potent inhibition on the redox activity of ERp57 in a concentration-dependent manner ($IC_{50} = 100$ to 300 μ M). ADTM inhibited the activities of ERp72, ERp5 and PDI to a much lesser extent (18%, 38% and 30% of inhibition at 300 μ M, respectively), displaying differential inhibitory activity at ERp57 (order of potency as follows: ERp57 > ERp5 > PDI > ERp72).

Figure 2. Target validation and the inhibition of PDI members by ADTM. (a) Schematics of the target validation. (b) Proteins isolated from rat platelets were pretreated with DMSO or excess amount of untagged ADTM for 2 h, followed by treatment with BAA for another 2 h, and compound-protein-complexes were enriched with NeutrAvidin-agarose resin. The precipitates were detected by immunoblot for indicated proteins of PDI family. (c) ADTM selectively inhibits ERp57. The recombinant PDI, ERp57 and ERp72 proteins were incubated with different concentrations of ADTM for 100 min, and its catalytic activity was determined in insulin reduction assay.

Figure 3. Inhibitory effects of ADTM on ADP- and AA-induced platelet aggregation *in vitro*. Platelets were pretreated with or without the indicated concentrations of the tested compounds, ADTM (1–100 μM), DSS (100 μM), TMP (100 μM), aspirin (1–100 μM) and clopidogrel (1–100 μM) for 5 min at 37 °C and the percentage (%) of platelet aggregation inhibition normalized to DMSO treated samples (vehicle control) was determined as described in Materials and Methods. The platelets were stimulated with ADP (a), and AA (b). Data are expressed as mean ± SD, * $P < 0.01$ compared with the 100 μM DSS group, # $P < 0.01$ compared with the 100 μM TMP group, & $P < 0.01$ compared with the 100 μM DSS + TMP group, $n \geq 3$ /group. D = DSS, T = TMP, ASA = aspirin, clop = clopidogrel.

ERp57 inhibition by ADTM was irreversible after dilution of the ERp57-ADTM complex (Supplemental Figure 1), indicating that ADTM may covalently bind ERp57.

ADTM inhibits platelet aggregation *in vitro*. It was observed that ADTM exhibited concentration-dependent inhibition on ADP-induced platelet aggregation (Fig. 3a) as well as AA-induced platelet aggregation (Fig. 3b) *in vitro* and its effects were comparable to first-line clinical anti-platelet drugs, aspirin and clopidogrel. The parent compounds DSS and TMP, alone or in combination, could only inhibit AA-induced platelet aggregation but not ADP-induced platelet aggregation. These results demonstrated that ADTM exhibited broad-spectrum anti-platelet activities.

ADTM inhibits the expression of P-selectin. To further investigate the underlying mechanism of ADTM anti-platelet activity, we examined the effect of ADTM on the expression of P-selectin (also

Figure 4. ADTM inhibited the expression of P-selectin. Whole blood was treated with DMSO (0.1%) as a control or the indicated concentrations of ADTM (1–100 μM) for 5 min at 37°C. After that the platelets were activated using ADP (10 μM). Flow cytometry histogram (a) and the quantification of the data (b) show that ADTM inhibits P-selectin binding to ADP-activated platelets. $^{\#}P < 0.01$ compared with the control group, $^*P < 0.01$ compared with the ADP group. Data are representative of three independent experiments.

Figure 5. ADTM inhibits activation of α IIb β 3 integrin and disrupts the ERp57/ α IIb β 3 complex on platelets. Representative flow cytometry histogram (a) and the quantification of the data (b) show that ADTM inhibits PAC-1 binding to ADP-activated platelets. (c) ADTM disrupts the ERp57/ α IIb β 3 complex. Whole blood was treated with DMSO (0.1%) as a control or 100 μM ADTM for 5 min at 37°C. After that the platelets were activated using ADP (10 μM). Platelet lysates were prepared for immunoprecipitation (IP) and probed with anti-ERp57 antibody for immunoblot analysis (IB). The precipitates were also blotted with anti- α IIb β 3 antibody to ensure that equal amounts of α IIb β 3 were pulled down under each condition. $^{\#}P < 0.01$ compared with the control group, $^*P < 0.01$ compared with the ADP group. Data are representative of three independent experiments.

known as CD62P), which is an adhesion molecule expressed on the surface of platelet and is a critical marker of platelet activation, by using immunoreactivity and flow cytometry analysis. We observed that ADTM concentration-dependently inhibited ADP-induced expression of P-selectin and attenuated the expression of P-selectin by more than 65% at 100 μM (Fig. 4a and 4b).

ADTM inhibits activation of α IIb β 3 and disrupted ERp57/ α IIb β 3 interaction. Recently, ERp57 has been reported to directly interact with α IIb β 3 integrin and regulate its function in platelet aggregation²⁰. In line with this, we observed that ADTM could modulate α IIb β 3 and concentration-dependently inhibited ADP-induced α IIb β 3 activation, by using flow cytometry analysis with PAC-1, a monoclonal antibody that binds only the activated form of α IIb β 3 (Fig. 5a,b). We further examined the action of ADTM on the ERp57/ α IIb β 3 interaction by immunoprecipitation with anti- α IIb β 3 antibody, followed by detection of ERp57 immunoreactivity using immunoblotting. It was found that ADTM disrupted the interaction of ERp57 with α IIb β 3, both in the presence and absence of ADP (Fig. 5c).

ADTM increases vasodilator-stimulated phosphoprotein (VASP) phosphorylation and HO-1 protein level. In platelets, the phosphorylation of VASP is involved in the down-regulation of platelet

Figure 6. Effects of ADTM on VASP phosphorylation and HO-1 protein level. Rat platelets (1×10^8 platelets/ml) were incubated with DMSO and various concentrations of ADTM for 10 min followed by treatment with ADP for 5 min. Proteins were extracted from the rat platelets and then analyzed by immunoblot using antibodies against VASP and HO-1 as described in Materials and Methods. ADTM increased VASP phosphorylation (a) and HO-1 protein expression (b). * $P < 0.05$ compared with the ADP-stimulated group.

Figure 7. ADTM inhibits platelet aggregation-induced by ADP in a concentration-dependent manner *in vivo*. In order to compare the activities of ADTM, DSS, TMP, DSS + TMP and clopidogrel at a similar blood concentration, 20 mg/kg of ADTM (MW: 444), 10 mg/kg of DSS (MW: 220) and 6 mg/kg of TMP (MW: 136) alone and in combination, and 18 mg/kg of clopidogrel (MW: 322) were administered by IV injection once daily. This resulted in approximately the same blood concentration of ADTM, DSS, TMP, DSS + TMP and clopidogrel in rats *in vivo*. Five days after drug treatment, blood was withdrawn and ADP-induced platelet aggregation was measured. Data are expressed as mean $\pm$ SD, $n \geq 6$ /group. D = DSS, T = TMP, clop = clopidogrel, MW = molecular weight.

aggregation accompanied by inhibition of platelet integrin α IIB β 3 activation^{27,28}. As shown in Fig. 6a, pretreatment with ADTM (10 and 100 μ M) stimulated approximately three-fold increase in the phosphorylation of VASP. In addition, we also observed that ADTM increased the protein expression of HO-1, an anti-oxidative stress microsomal enzyme which when up-regulated could inhibit platelet-dependent thrombus formation (Fig. 6b)²⁹.

Inhibition of platelet aggregation and thrombus formation by ADTM *in vivo*. Various concentrations of ADTM were administered in rats *in vivo* (10–20 mg/kg, 5 days, intravenous, IV) and the platelet aggregation activity and the thrombus formation were evaluated. As shown in Fig. 7, ADP-induced platelet aggregation was significantly compromised (>40% reduction) in rats treated with ADTM (20 mg/kg). Similarly, ADTM also exhibited significant anti-thrombotic effect *in vivo* as shown in the ferric chloride (FeCl_3)-induced venous thrombosis (Fig. 8a). The inhibitory effects of ADTM in platelet aggregation and thrombus formation were comparable to clopidogrel. Furthermore, FeCl_3 induced significant decrease in plasma 6-Keto-PGF_{1 α} (an indicator of plasma PGI₂, an intrinsic inhibitor

Figure 8. Anti-thrombotic effect of ADTM in the FeCl₃-induced rat thrombosis model. The anti-thrombotic effect of ADTM in the FeCl₃-induced inferior vena cava thrombosis model (a). In order to compare the activities of ADTM, DSS, TMP and clopidogrel at similar blood concentrations, the compounds were administered by IV injection as described in Fig. 7. The effect of ADTM on plasma 6-Keto-PGF_{1α} concentration in the FeCl₃-induced thrombosis model (b). Data are expressed as mean ± SD, [#]*P* < 0.01 compared with the sham group, ^{*}*P* < 0.05 compared with the vehicle group, *n* = 5–7/group, D = DSS, T = TMP, clop = clopidogrel.

of platelet aggregation) and the treatment of ADTM (20 mg/kg, 5 days, IV) abolished the reduction of FeCl₃-induced 6-Keto-PGF_{1α} in a concentration-dependent manner (Fig. 8b), giving more evidence of the anti-thrombotic properties of ADTM.

Discussion

Anti-thrombotic therapy plays an essential role in the course of treatment of various cardiovascular disorders (CVDs) because platelet dysfunction and thrombogenesis are common underlying factors in the development and progression in CVDs, particularly atherosclerosis, myocardial infarction and ischemic stroke. Aspirin and clopidogrel are most routinely used anti-platelet agents for the management of CVDs, but these agents irreversibly inhibit platelet activation with potentially devastating unwanted effects such as hemorrhages. In present study we reported that ADTM, a novel cardioprotective compound synthesized in our own laboratory⁶, displayed specific binding for several members of the PDI family. By using a biotin-conjugated ADTM analogue (BAA) as bait and chemical proteomic analysis of the bound protein complexes, we identified eight potential interacting protein targets of ADTM (Supplementary Table S1). With further validation using immunoblot analysis the identity of major targets were confirmed as members of the PDI family, including ERp72, ERp57 and ERp5. In particular, ADTM showed highest potency in inhibiting the redox activity of ERp57 in enzymatic assay *in vitro*. Furthermore, untagged-ADTM could compete with BAA for the binding of ERp57, providing evidence that ADTM is a competitive ligand at ERp57. We further demonstrated that ADTM displayed concentration-dependent inhibition in ADP-induced platelet aggregation and AA-induced platelet aggregation *in vitro*. This result is in line with the observation that ERp57 has been recently implicated to play crucial roles in mediating platelet aggregation, homeostasis and thrombosis^{18–20}. It has also been reported that ERp57 present on platelet surface were activated by staphylococcal extracellular adhesive protein resulting in arteriosclerosis in *Staphylococcus aureus* infection, and the inhibition of ERp57 could suppressed platelet aggregation³⁰.

We further demonstrated that ADTM concentration-dependently inhibited the ADP-induced expression of P-selectin and activation of αIIbβ3 integrin *in vitro*. P-selectin is an adhesion molecule expressed on the surface of platelet and is an important marker for platelet activation. αIIbβ3 integrin is the most abundant adhesion receptor on the surface of circulating platelets and has been reported to

Figure 9. Proposed mechanism of ADTM anti-platelet aggregation. In part, ADTM binds with ERp57 and inhibits its redox activity, inhibits the activation of α IIb β 3 and the expression of P-selection and enhances the expression of HO-1 and phosphorylation of VASP. The crosstalk between ERp57 and α IIb β 3 is involved in the biological action of ADTM. In addition, ADTM reverses the decreased level of 6-Keto-PGF1 α , a stable metabolite of PGI2 in the FeCl $_3$ -induced thrombosis model *in vivo*.

mediate platelet aggregation through strong interactions with adhesive proteins such as fibrinogen, von Willebrand factor and fibronectin³¹. We further demonstrated that ERp57 bound to α IIb β 3 directly by using co-immunoprecipitation, and ADTM disrupted ERp57/ α IIb β 3 interaction, as shown by the dissociation of the complex (Fig. 5c). In line with this, ERp57 has been reported to directly interact with α IIb β 3 on platelet surface and regulate its function in platelet aggregation^{18–20}. Recent study has shown that disulfide bond exchanges in α IIb β 3 is required for activation and post-ligation signaling during clot retraction³², and the modulation of functional disulfide bonds is being explored for blood protein control³³. Our results suggested that ADTM produced its anti-thrombotic action via the modulation of disulfide bond by interacting with ERp57 (Fig. 9).

The inhibitory action of ADTM on platelet integrin α IIb β 3 activation was further evidenced by the increase of the phosphorylation of VASP after ADTM pretreatment. The phosphorylation VASP is involved in the down-regulation of platelet aggregation accompanied by inhibition of platelet integrin α IIb β 3 activation^{27,28}. We also observed that ADTM increased the expression of HO-1 in platelets. This is coherent with our previous finding that HO-1 activation has a key role in ADTM cardioprotective effect in H9c2 cardiomyoblast cells *in vitro* and in the myocardium *in vivo*⁷. HO-1 is a potential therapeutic target in treating thrombotic disease and HO-1 knockout mice demonstrated acute thrombus formation in response to hypoxia³⁴. Previous studies reported that dengue virus glycoprotein-1 (DV-NS1) antibody inhibited PDI activity and blocked platelet aggregation accompanied by an increase in HO-1^{35,36}. Furthermore, redox reactions have been shown to play crucial roles in platelet aggregation in a glutathione peroxidase-3 (GPx-3)-deficient mouse model. GPx-3 is an important anti-oxidant enzyme and redox sensor in plasma and GPx-3-deficiency impaired normal platelet inhibition and promoted thrombosis due to enhanced reactive oxygen species flux and decreased nitric oxide bioavailability³⁷. In our *in vivo* studies, we observed that ADP-induced platelet aggregation was significantly compromised (>40% reduction) in rats treated with ADTM (20 mg/kg). The anti-thrombotic effect of ADTM was further evidenced by the reduction of platelet activity and thrombus formation in the ferric chloride (FeCl $_3$)-induced venous thrombosis assay in the rat. The major metabolites of ADTM *in vivo* were 2-hydroxymethyl-3, 5, 6-trimethylpyrazin (TMP-OH) and DSS³⁸. We have checked the *in vivo* activity of DSS and TMP. Interestingly, the results showed that ADTM exhibited stronger antiplatelet and anti-thrombotic activities when compared to DSS and TMP, alone or in combination in ADP-induced platelet aggregation and FeCl $_3$ -induced thrombosis model, respectively (Figs. 7 and 8). The precise mechanisms of this action conferred by ADTM would be another interesting question that is worth investigating in the future. Recently there are accumulating reports that the inhibition of protein disulfide isomerases including PDI and ERp57 could block thrombus formation in various *in vivo* models and suggest that this protein family represents an important novel class of anti-thrombotic target^{39–43}. Preclinical studies demonstrate that deficiency in platelet ERp57 resulted in the increased tail bleeding times and delayed thrombus formation²⁰, while PDI is unable to compensate for the absence of platelet ERp57²⁰ and other study also showed that the blockade of ERp57 with specific antibody further inhibited platelet aggregation in PDI deficient platelets¹⁶. These independent studies further provided evidences that ERp57 and PDI have distinct roles in the mediation of platelet function.

Taken together, our present study reported the anti-thrombotic action of ADTM with both *in vitro* and *in vivo* data. ADTM exhibited comparable anti-thrombotic properties as aspirin and clopidogrel. The results suggested that the anti-thrombotic action of ADTM is mediated through disrupting the

interaction between ERp57 and α Ib β 3 possibly by blocking the action of ERp57 on disulfide bonding (Fig. 9). Our data provided a rationale for the further development of ADTM as anti-thrombotic agent targeting the underlying mechanism involving ERp57 and α Ib β 3. Furthermore, as a competitive ligand at ERp57, ADTM presents as a promising compound for the development of versatile anti-thrombotic agents. Our data also provide insights in novel strategies for the development of drugs targeted ERp57 for anti-thrombotic.

Methods

Materials. ADTM and BAA were synthesized at Jinan University, China. DSS and TMP were of analytical pure grade, and obtained from Xi'an Honson Biotechnology (China) and Shanghai Banghai Chemical Company (China), respectively. Human recombinant ERp57 was obtained from Abcam (Cambridge, UK), PDI and ERp72 were purchased from Enzo Life Sciences (Exeter, UK). Insulin, DTT, AA and ADP were obtained from Sigma Aldrich (St. Louis, MO, USA).

Preparation of platelet-rich plasma. All animal experiments were approved by the Animal Care and Experimentation Committee of Jinan University and were performed in accordance with the approved guidelines. Sprague-Dawley rats were anesthetized with 10% chloral hydrate, and blood was obtained by an arterial puncture. Whole blood was anticoagulated with citrate (3.8%; 1:9, v/v) and centrifuged at 200g for 8 min at room temperature to obtain platelet-rich plasma (PRP). The residue was centrifuged at 550g for 5 min to obtain platelet-poor plasma (PPP).

Protein preparation from platelets. Washed platelets were lysed using a dounce homogenizer in NP-40 lysis buffer (Beyotime, China) with 1 mM phenylmethylsulfonyl fluoride (PMSF). Platelet lysates were centrifuged at 12,500g for 20 min at 4°C, and the supernatant was collected and stored at −80°C until further analysis.

NeutrAvidin Agarose Resin pull-down with BAA. Platelet lysates (3 μg/μl) were exposed to NeutrAvidin Agarose Resin (Pierce Biotech., Rockford, IL, USA) for 2 h at 4°C and then centrifuged for 3 min at 2,500g. The resin was discarded to remove endogenous biotin. Aliquots of supernatants (300 μl) were first treated with DMSO or ADTM (600 μM) for 2 h, and then incubated with BAA (600 μM) for another 2 h. The supernatant was combined with NeutrAvidin Agarose Resin and allowed to shake overnight at 4°C as described previously⁴⁴. To eliminate the nonspecific combination with compound or agarose resin before protein analysis, 0.1% DMSO was added to one aliquot of lysate as a negative control.

Nano LC-MS/MS analysis. The compound-protein complex bound to agarose resin was digested with trypsin. The digested peptides were analyzed with 1D nano-RP LC-MS/MS. Acquired MS data were analyzed using the Paragon algorithm in ProteinPilot 4.0 software (Applied Biosystems, Framingham, MA, USA). The identified peptides from the Paragon algorithm were grouped into minimal non-redundant protein sets by the ProGroup algorithm of the software (details of this method can be found in Supplementary Methods).

Immunoblot analysis. Proteins eluted from the resin were subjected to SDS-PAGE and then transferred to a PVDF membrane for immunoblot (details in Supplementary Methods).

Insulin reduction assay. PDI activity was measured using an assay that measured the catalytic reduction of insulin, as described previously with minor modifications^{39,45}. Briefly, insulin (1 mg/ml) was freshly prepared in assay buffer (100 mM potassium phosphate, 1 mM EDTA (pH 7.4) and 10 μM dithiothreitol). 50 μl of insulin was incubated in the presence of 10 μl of enzyme and varying concentrations of compound, and then 10 μl of 0.1 mM dithiothreitol was added. The increase in turbidity was monitored at 620 nm using a microplate reader (Perkin-Elmer, Singapore) at 25°C for 60 min. PDI, ERp57, and ERp72 were all used at 0.04 mg/ml. PDI family enzyme inhibition in the presence of compound was determined by the following formula: enzyme inhibition (%) = $[1 - (\text{OD}_{[\text{compound} + \text{PDI} + \text{DTT}]} - \text{OD}_{[\text{DTT}]}) / (\text{OD}_{[\text{PDI} + \text{DTT}]} - \text{OD}_{[\text{DTT}]})] \times 100\%$.

Platelet aggregation *in vitro*. The inhibitory effect of ADTM was examined as previously described with modifications³⁹. Briefly, platelets were incubated with the indicated concentrations of compounds or 0.1% DMSO at 37°C for 5 min with shaking. Aggregation was induced by ADP (10 μM) or AA (200 μM) and measured using a platelet aggregometer (SC-2000, Saikexide Instrument Co. Ltd. China). Inhibition of aggregation was calculated using the following general formula: inhibition of aggregation = (rate of aggregation in vehicle group − rate of aggregation in treated group)/rate of platelet aggregation in vehicle group × 100%.

Determination of human platelet activation by flow cytometry. Flow cytometry and immunoprecipitation studies involving human platelets were approved by Jinan University with informed consent in accordance with the World Medical Association Declaration of Helsinki and carried out in accordance with the approved guidelines. Flow cytometry was performed as described in the Supplementary

Methods. PE conjugated anti-CD62P (P-selectin) antibody and FITC conjugated anti-PAC-1 (activated α Ib β 3) antibody were used to detect platelet activation markers after exposure to ADP in the presence of DMSO (0.1%) or different concentrations of ADTM. An isotype-matched IgG1 PE/FITC was used as a negative control.

Immunoprecipitation. To determine ERp57-bound α Ib β 3 contents, whole platelet lysates were incubated with anti- α Ib β 3 antibody overnight at 4°C under agitation. The antigen-antibody complex was pulled down after incubation for 4 h at 4°C with protein G-agarose (Cell Signaling Technology) as described previously⁴⁶. Immune complexes were boiled for 5 min with 2 × loading buffer followed by immunoblot analysis with antibodies for ERp57 and α Ib β 3.

Rat platelet aggregation-induced by ADP *in vivo*. To study the effects of ADTM on platelet aggregation *in vivo*, ADTM (5–20 mg/kg) in comparison with DSS (10 mg/kg), TMP (6 mg/kg), DSS (10 mg/kg) + TMP (6 mg/kg) and clopidogrel (18 mg/kg) were administered daily by IV injection for 5 days, respectively (detailed methodology in Supplementary Methods). Platelet aggregation was monitored for 5 min and the maximum aggregation inhibition (%) was calculated.

FeCl₃-induced inferior vena cava thrombosis in rat. Rats weighing 180 g to 220 g were anesthetized by intraperitoneal administration of 10% chloral hydrate. One hour post-treatment, the medioventral line in the rats was cut and the inferior vena cava was surgically exposed by blunt dissection. 1 × 1 cm filter paper, soaked in 50% FeCl₃ solution, was placed under and above the exposed inferior vena cava for 15 min. Subsequently, the thrombosed inferior vena cava was carefully dissected, and the weight (mg) of wet thrombus harvested from the inferior vena cava was measured as described previously⁴⁷.

Evaluation of plasma 6-Keto-PGF_{1 α} levels in rat. The concentration of 6-keto-PGF_{1 α} present in the supernatant was measured using a radioimmunoassay kit (New England Nuclear Dupont, Boston, MA) according to the manufacturer's instructions. Blood was collected into plastic tubes containing 10% EDTA-Na₂. Plasma was prepared by centrifuging blood at 550 g for 10 min. Plasma concentrations of 6-keto-PGF_{1 α} were determined by radioimmunoassay and expressed as pg/ml.

Statistical analysis. All assays were performed at least three times. Data are presented as mean ± SD. The groups were compared using one-way ANOVA followed by Tukey's multiple comparison tests using the statistics module of Graph Pad Prism 5.0. A value of $P < 0.05$ was considered statistically significant.

References

1. Roger V. L. *et al.* Heart disease and stroke statistics--2012 update: a report from the American Heart Association. *Circulation* **125**, e2–e220 (2012).
2. Martinelli L., Bucciarelli P., & Mannucci P. M. Thrombotic risk factors: basic pathophysiology. *Crit Care Med* **38**, S3–9 (2010).
3. Hansson G. K., & Klareskog L. Pulling down the plug on atherosclerosis: cooling down the inflammasome. *Nat Med* **17**, 790–791 (2011).
4. Investigators AWGoT *et al.* Clopidogrel plus aspirin versus oral anticoagulation for atrial fibrillation in the Atrial fibrillation Clopidogrel Trial with Irbesartan for prevention of Vascular Events (ACTIVE W): a randomised controlled trial. *Lancet* **367**, 1903–1912 (2006).
5. Investigators A *et al.* Effect of clopidogrel added to aspirin in patients with atrial fibrillation. *N Engl J Med* **360**, 2066–2078 (2009).
6. Cui Q. *et al.* Design, synthesis, and preliminary cardioprotective effect evaluation of danshensu derivatives. *Chem Biol Drug Des* **84**, 282–291 (2014).
7. Cui G. *et al.* A novel Danshensu derivative confers cardioprotection via PI3K/Akt and Nrf2 pathways. *Int J Cardiol* **168**, 1349–1359 (2013).
8. Tian P. Convergence: Where West meets East. *Nature* **480**, S84–86 (2011).
9. Jia Y., Huang F., Zhang S., & Leung S. W. Is danshen (Salvia miltiorrhiza) dripping pill more effective than isosorbide dinitrate in treating angina pectoris? A systematic review of randomized controlled trials. *Int J Cardiol* **157**, 330–340 (2012).
10. Cheng T. O. Cardiovascular effects of Danshen. *Int J Cardiol* **121**, 9–22 (2007).
11. Zhou L. M., Zuo Z., & Chow M. S. S. Danshen: An overview of its chemistry, pharmacology, pharmacokinetics, and clinical use. *J Clin Pharmacol* **45**, 1345–1359 (2005).
12. Essex D. W. Redox control of platelet function. *Antioxid Redox Signal* **11**, 1191–1225 (2009).
13. Furie B., Flaumenhaft R. Thiol isomerases in thrombus formation. *Circ Res* **114**, 1162–1173 (2014).
14. Cho J. Protein disulfide isomerase in thrombosis and vascular inflammation. *J Thromb Haemost* **11**, 2084–2091 (2013).
15. Cho J., Furie B. C., Coughlin S. R., & Furie B. A critical role for extracellular protein disulfide isomerase during thrombus formation in mice. *J Clin Invest* **118**, 1123–1131 (2008).
16. Kim K. *et al.* Platelet protein disulfide isomerase is required for thrombus formation but not for hemostasis in mice. *Blood* **122**, 1052–1061 (2013).
17. Reinhardt C. *et al.* Protein disulfide isomerase acts as an injury response signal that enhances fibrin generation via tissue factor activation. *J Clin Invest* **118**, 1110–1122 (2008).
18. Holbrook L. M. *et al.* The platelet-surface thiol isomerase enzyme ERp57 modulates platelet function. *J Thromb Haemost* **10**, 278–288 (2012).
19. Wu Y. *et al.* The disulfide isomerase ERp57 mediates platelet aggregation, hemostasis, and thrombosis. *Blood* **119**, 1737–1746 (2012).
20. Wang L. *et al.* Platelet-derived ERp57 mediates platelet incorporation into a growing thrombus by regulation of the α IIb β 3 integrin. *Blood* **122**, 3642–3650 (2013).
21. Ferrari D. M., Soling HD. The protein disulphide-isomerase family: unravelling a string of folds. *Biochem J* **339** (Pt 1), 1–10 (1999).

22. Ravanat C. *et al.* GPV is a marker of *in vivo* platelet activation--study in a rat thrombosis model. *Thromb Haemost* **83**, 327–333 (2000).
23. Moog S. *et al.* Platelet glycoprotein V binds to collagen and participates in platelet adhesion and aggregation. *Blood* **98**, 1038–1046 (2001).
24. Freedman J. E. Oxidative stress and platelets. *Arterioscler Thromb Vasc Biol* **28**, s11–16 (2008).
25. Corazzari M. *et al.* Targeting homeostatic mechanisms of endoplasmic reticulum stress to increase susceptibility of cancer cells to fenretinide-induced apoptosis: the role of stress proteins ERdj5 and ERp57. *Br J Cancer* **96**, 1062–1071 (2007).
26. Laurindo F. R., Pescatore L. A., Fernandes Dde C. Protein disulfide isomerase in redox cell signaling and homeostasis. *Free Radic Biol Med* **52**, 1954–1969 (2012).
27. Massberg S. *et al.* Enhanced *in vivo* platelet adhesion in vasodilator-stimulated phosphoprotein (VASP)-deficient mice. *Blood* **103**, 136–142 (2004).
28. Horstrup K. *et al.* Phosphorylation of Focal Adhesion Vasodilator-Stimulated Phosphoprotein at Ser157 in Intact Human Platelets Correlates with Fibrinogen Receptor Inhibition. *Eur J Biochem* **225**, 21–27 (1994).
29. Peng L. *et al.* Induction of heme oxygenase-1 expression inhibits platelet-dependent thrombosis. *Antioxid Redox Signal* **6**, 729–735 (2004).
30. Bertling A. *et al.* Staphylococcal extracellular adherence protein induces platelet activation by stimulation of thiol isomerases. *Arterioscler Thromb Vasc Biol* **32**, 1979–1990 (2012).
31. Hynes R. O. Integrins: bidirectional, allosteric signaling machines. *Cell* **110**, 673–687 (2002).
32. Mor-Cohen R., Rosenberg N., Averbukh Y., Seligsohn U., & Lahav J. Disulfide bond exchanges in integrins $\alpha\text{IIb}\beta\text{3}$ and $\alpha\text{v}\beta\text{3}$ are required for activation and post-ligation signaling during clot retraction. *Thromb Res* **133**, 826–836 (2014).
33. Butera D., Cook K. M., Chiu J., Wong J. W., & Hogg P. J. Control of blood proteins by functional disulfide bonds. *Blood* **123**, 2000–2007 (2014).
34. Wu M. L., Ho Y. C., & Yet S. F. A central role of heme oxygenase-1 in cardiovascular protection. *Antioxid Redox Signal* **15**, 1835–1846 (2011).
35. Cheng H. J. *et al.* Anti-dengue virus nonstructural protein 1 antibodies recognize protein disulfide isomerase on platelets and inhibit platelet aggregation. *Mol Immunol* **47**, 398–406 (2009).
36. Immenschuh S. *et al.* Antibodies against dengue virus nonstructural protein-1 induce heme oxygenase-1 via a redox-dependent pathway in human endothelial cells. *Free Radic Biol Med* **54**, 85–92 (2013).
37. Jin R. C. *et al.* Glutathione peroxidase-3 deficiency promotes platelet-dependent thrombosis *in vivo*. *Circulation* **123**, 1963–1973 (2011).
38. Li S. *et al.* Pharmacokinetic and Metabolic Studies of ADTM: A Novel Danshensu Derivative Confers Cardioprotection by HPLC-UV and LC-MS/MS. *J Chromatogr Sci*, doi: 10.1093 (2015).
39. Jasuja R. *et al.* Protein disulfide isomerase inhibitors constitute a new class of antithrombotic agents. *J Clin Invest* **122**, 2104–2113 (2012).
40. Kennedy D. R. *et al.* Development Of Second Generation Thiol Isomerase Inhibitors To Prevent Thrombus Formation. *Blood* **122**, 926–926 (2013).
41. Flaumenhaft R. Protein disulfide isomerase as an antithrombotic target. *Trends Cardiovasc Med* **23**, 264–268 (2013).
42. Loving G. S., Caravan P. Activation and retention: a magnetic resonance probe for the detection of acute thrombosis. *Angew Chem Int Ed Engl* **53**, 1140–1143 (2014).
43. Zhou J. *et al.* The disulfide isomerase ERp57 is required for fibrin deposition *in vivo*. *J Thromb Haemost* **12**, 1890–1897 (2014).
44. Liu C. X. *et al.* Adenanthin targets peroxiredoxin I and II to induce differentiation of leukemic cells. *Nat Chem Biol* **8**, 486–493 (2012).
45. Antoniou A. N. *et al.* The oxidoreductase ERp57 efficiently reduces partially folded in preference to fully folded MHC class I molecules. *EMBO J* **21**, 2655–2663 (2002).
46. Yang S. *et al.* Pellino3 targets RIP1 and regulates the pro-apoptotic effects of TNF- α . *Nat Commun* **4**, 2583 (2013).
47. Hennen J. K. *et al.* Effect of tiplaxtinin (PAI-039), an orally bioavailable PAI-1 antagonist, in a rat model of thrombosis. *J Thromb Haemost* **6**, 1558–1564 (2008).

Acknowledgements

This work was supported by grants from the Natural Science Fund of China (Grant No. U1032007, 81302776, 81328025, 81460552 and 8140339), the Science and Technology Development Fund Macao SAR, China (Grant No. 014/2011/A1), the “211” project of Jinan University, and Scientific and Technological Innovation Project of Department of Education of Guangdong Province (Grant No. 2012KJCX0018), and the Research Fund of University of Macau (Grant No. MYRG2015-00161-ICMS-QRCM).

Author Contributions

M.Y.L., P.M.H. and Y.Q.W. designed the project, supervised the research and revised the manuscript. G.Z.C. and L.S.S. performed research, analyzed data and wrote the manuscript. Y.W.S. and P.Y. synthesized the compounds. L.G., C.M.C. and Z.J.Z. help in animal study and platelet aggregation assay. I.K.C., G.H.L., Q.Q. and Y.Z. helped with the chemical proteomic assay.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Cui, G. *et al.* Novel anti-thrombotic agent for modulation of protein disulfide isomerase family member ERp57 for prophylactic therapy. *Sci. Rep.* **5**, 10353; doi: 10.1038/srep10353 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

SCIENTIFIC REPORTS

OPEN

Corrigendum: Novel anti-thrombotic agent for modulation of protein disulfide isomerase family member ERp57 for prophylactic therapy

Guozhen Cui, Luchen Shan, Lin Guo, Ivan Keung Chu, Guohui Li, Quan Quan, Yun Zhao, Cheong Meng Chong, Zaijun Zhang, Pei Yu, Maggie Pui Man Hoi, Yewei Sun, Yuqiang Wang & Simon MingYuen Lee

Scientific Reports 5:10353; doi: 10.1038/srep10353; published online 03 June 2015; updated on 02 September 2015

In this Article, Fig. 5 is a duplication of Fig. 7. The correct Fig. 5 appears below as Fig. 1.

Figure 1.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>