SCIENTIFIC REPORTS

OPEN

Fractal features of soil particle size distribution in newly formed wetlands in the Yellow River Delta

Received: 28 November 2014

Accepted: 17 April 2015

Published: 27 May 2015

Junbao Yu¹, Xiaofei Lv^{1,2}, Ma Bin¹, Huifeng Wu¹, Siyao Du³, Mo Zhou⁴, Yanming Yang⁵ & Guangxuan Han¹

The characteristic of particle size distribution (PSD) in the newly formed wetlands in coast has seldom been studied. We applied fractal-scaling theory in assessing soil particle size distribution (PSD) features of newly formed wetlands in the Yellow River Delta (YRD), China. The singular fractal dimensions (D) values ranged from 1.82 to 1.90, the capacity dimension (D_0) values ranged from 0.84 to 0.93, and the entropy dimension (D_1) values ranged from 0.66 to 0.84. Constrained corresponding analysis revealed that 43.5% of the variance in soil PSD can be explained by environmental factors, including 14.7% by seasonal variation, 8.6% by soil depth, and 8.0% by vegetation type. The fractal dimensions D and D_1 were sensitive with fine particles with size ranging less than 126 μm , and D_0 was sensitive with coarse particles with size ranging between 126 μm to 2000 μm . Fractal analysis makes full use of soil PSD information, and offers a useful approach to quantify and assess the soil physical attributes in the newly formed wetland.

Coastal wetlands play important roles in the coastal ecosystems, offering wildlife food and habitat, as well as functioning as important nutrient cycling capacity for maintaining water quality^{1–3}. Newly formed wetlands in estuary regions, where large quantities of silts have been deposited to form coastal wetlands, are generally found several hundreds of meters with vegetation changes gradually from barren land to halophytic vegetation and to hydrophyte vegetation^{4–6}. The attribute changes in newly formed coastal wetlands can influence the ecosystem service functions of the coastal habitats.

Soil particle size distribution (PSD) is one of the most fundamental physical attributes due to its great influence on soil hydraulic characteristics, soil moisture movement, contaminant transport, and soil erosion⁷. Accordingly, characterizing variations of soil PSD is an important way to understanding and quantifying soil structure and dynamics^{8–10}. The PSD characteristics in the newly formed wetlands are, however, still unclear.

Traditionally, textural triangle, the principal approach for soil PSD analysis, is restricted by the arbitrary definition of texture classes with incomplete information on the soil PSD¹¹. Moreover, there is a high fluctuation of PSD in each textural class. A better approach is combining laser diffraction method and fractal analysis, which can highly promote the research of soil texture^{10,12–15}. Fractal analysis offers the possibility of quantifying and integrating information on soil structure at different spatial scales^{16,17}. Soil fractal analysis is currently implemented in two ways, i.e. singular fractal analysis and multifractal analysis¹⁸. Singular fractal analysis can be used to quantitatively describe soil PSD characteristics, soil aggregate fragmentation, and other related soil properties¹⁶. Multifractal analysis is mainly employed

¹Key Laboratory of Coastal Environmental Processes and Ecological Remediation, Yantai Institute of Coastal, Zone Research (YIC), Chinese Academy of Sciences (CAS); Shandong Provincial Key Laboratory of Coastal Environmental Processes, YICCAS, Yantai 264003. ²University of Chinese Academy of Sciences, Beijing 100049, China. ³College of Environmental Science and Engineering, Ocean University of China, Qingdao 266100, P. R. China. ⁴Environment college, Northeast Normal University, Changchun 130024, P. R. China. ⁵Cadre School of Science and Technology of Shandong Province, Ji'nan 250101, P. R. China. Correspondence and requests for materials should be addressed to J.Y. (email: junbao.yu@gmail.com) or (email: jbyu@yic.ac.cn)

Figure 1. The location of study region and sampling sites (ARCGIS 9.3).

to capture the intrinsic variability of soil PSD to retain more detailed information^{13,19}. Consequently, there are a number of studies applying multifractal analysis to study soil particle-size distributions^{12,13,20}. Fractal theory is determined to the influence of different plant communities or land management on PSD^{21,22}.

In this study, fractal scale theory was employed for assessing spatial and temporal variation of the PSD characteristics in a successional series of newly formed wetlands in the Yellow River Delta (YRD). The Yellow River is regarded as the largest contributor of fluvial sediment load to the ocean in the world²³. The YRD is one of the most intensive land-ocean interaction regions among the large river deltas⁴, also has the youngest natural coastal wetland ecosystem in China. The objectives of this work were to 1) analyze the PSD and the fractal dimensions of soils in the newly formed wetlands of the YRD, and 2) explore the impacts of vegetation community on the spatial and temporal variation of soil PSD in the newly formed wetlands.

Methods

Study area. The investigation was conducted at the YRD located in northern part of Shandong Province, China. The YRD (118.6°E–119.3°E, 37.6°N–38.2°N) is characterized as a temperature, semi-humid continental monsoon climate. The average temperature is 11.7–12.6°C, and the average annual evaporation is 1900–2400 mm. The average annual precipitation is 530–630 mm, of which 70% is rainfall during June to September (in summer and fall). The soil is typical saline alluvial soil (Fluvisols, FAO). The net increase of delta shoreline length was ~61.64 km with annual increase of ~1.81 km, and net extension of newly formed wetland area was ~309.81 km² with rate of ~9.11 km² year⁻¹ in the YRD²⁴. The coastal wetlands in the YRD have clear horizontal distribution vegetation zones of ecosystems with the changes in soil salinity from seaside to inland (Fig. 1). A total of 400 plant species are recorded in the region, of which 55.1% vegetation is natural saline vegetation²⁵. The dominant plant species adapt the saline-alkaline are *Tamarix chinensis*, *Suaeda salsa* and *Phragmites australis* in the YRD.

Sampling and processing. We chose five sampling plots along the bank of the Yellow River in a successional series of newly formed wetlands (formed since 1976) based on vegetation community. The dominant plant species in sampling plots were no plant (P1), *Suaeda salsa* (P2), *Tamarix chinensis* (P3), *Phragmites australis* (P4) and *Typha orientalis* (P5), respectively (Fig. 1). Soil samples were collected from five random soil cores and 0–10, 10–20, 20–30 cm depth with in wet season (July, October 2012), and dry season (February, May 2013). The replicated samples were mixed homogeneously at each location to form a composite sample. We collected 60 samples totally in this study for the following analysis.

The visible roots and rocks were removed prior to further processing. Soil samples were air-dried and passed 2 mm sieve. After dispersed by sodium hexametaphosphate (NaHMP) and ultrasonic for 30 s, the samples were analyzed with Longbench Mastersizer2000 (Malvern Instruments, Malvern, England) to get particle size distribution ranging from 0.2 μm to 2000 μm. The interval of particle sizes (μm) $I_i = [0.2, 2000]$ were graded into 64 subintervals $I_i = [\phi_i, \phi_{i+1}]$, $i = 1, 2, \dots, 64$, and the lengths of subintervals followed a logarithmic scale such that $\log(\phi_{i+1}/\phi_i)$ is constant. Meanwhile, according to the United States Department of Agriculture classification of soil particle size, the soil size was partitioned into 3 grades, clay (0–2 μm), silt (2–50 μm) and sand (50–2000 μm). The soil organic matter (SOM) was determined by K₂CrO₇ routine colorimetric method²⁶.

The singular fractal analysis for soil PSD. The singular fractal dimension (D) of soil PSD was estimated from the following equation²⁷:

$$\frac{V(r < R_i)}{V_T} = \left(\frac{R_i}{R_{max}} \right)^{3-D} \quad (1)$$

Where $V(r < R_i)$ is the cumulative percentage of particles of i size r less than R_i , V_T is the total percentage ($V_T = 100$), R_i is the particle radius (mm) of the i size class, and R_{max} is the radius of the largest particle class ($R_{max} = 1$, in this study). The particle diameter is taken as the upper sieve sizes. Taking logarithms on both sides of Equation (1), the D value can be derived by the slopes of the logarithmic linear regression equation.

The multifractal analysis for soil PSD. The multifractal dimension (Rényi dimension) was calculated using Equation (2) and Equation (3) in this study. A number of cells with size of ε to cover the entire interval, and the cell number is N , the Rényi dimension is computed by the mass of soil particles in subinterval $\mu_i(\varepsilon)$, cell diameter ε and the parameter q ¹².

$$D_1 = \lim_{\varepsilon \rightarrow 0} \frac{\sum_{i=1}^{N(\varepsilon)} \mu_i(\varepsilon) \log \mu_i(\varepsilon)}{\log(\varepsilon)} \quad (q = 1) \quad (2)$$

$$D_q = \frac{1}{q-1} \lim_{\varepsilon \rightarrow 0} \frac{\log \sum_{i=1}^{N(\varepsilon)} \mu_i(\varepsilon)^q}{\log(\varepsilon)} \quad (q \neq 1) \quad (3)$$

D_q extracts the system parameters from different levels with the q value in the interval $[\infty, +\infty]$. The Rényi dimensions for $q=0$ and $q=1$ are known as capacity dimension D_0 and entropy dimension D_1 , respectively. In multifractal systems, D_0 is the capacity dimension is known as box-counting dimension. It provides average information of PSD system and reflects the range of a continuous distribution. In this study, all the PSDs are distributed continuously from $0.3 \mu\text{m}$ to $2000 \mu\text{m}$. So $D_0 = 1$ means the interval of particle sizes were all occupied by all scales. The PSDs with low D_0 hold a narrow range. The entropy dimension, D_1 , provides a measure of the heterogeneity of a PSD. The higher the value of D_1 is, the more heterogeneous the soil's PSD is, and the wider the range of PSD is. Considering that D_0 provides general information and D_1 measures the homogeneity of PSD system, D_1/D_0 is used to describe the heterogeneity in a distribution to obtain the relation between the two parameters. As D_1 takes values less than D_0 , the quotient D_1/D_0 is less than 1. The closer to 1 of D_1/D_0 refers the more evenly dispersed in the fractions over the set of sizes, and the more heterogeneous in the distribution.

Statistical analysis. Data in the figures and tables were mean values of each sample. All statistical analyses were implemented using various packages within the R statistical computing environment. One-way analysis of variance (ANOVA) procedures were used to detect the differences in measured parameters among plots. Constrained correspondence analysis (CCA) was used to assess the relationship between soil PSD profiles and environmental variables with the *vegan* package.

Network analyses can be used to show the composition of, and interactions between, multiple elements in communities. A matrix of correlation between all trait pairs was generated. Significant levels were set at $p < 0.05$. The total significant pairs were considered as a network in which a vertex corresponds to a trait and a link between two vertices corresponds to significant correlations. This network plot was then subjected based on the adjacency matrix with *igraph* package. Essentially, this algorithm divides the network into modules or groups of vertices that are more connected between themselves than to nodes from other modules, yielding a cartographic representation of a complex network.

Results

Soil particle size distribution. The soil texture changed from sand to silt loam with the soil organic matter concentration increasing (Fig. 2). The predominant soil particle was sand fraction, accounting for 62% of total soil volume (Fig. 3). The proportion of silt and clay particles was 31% and 6%, respectively.

The sand fraction in P4 (*Phragmites australis*) and P5 (*Typha orientalis*) were greater than those in P1 (no plant), P2 (*Suaeda aslsa*) and P3 (*Tamarix chinesis*) (Table 1). The proportions of sand in topsoil layers (0–10 cm) were smaller than those in underlying soil layers (10–30 cm). The seasonal variations of the clay and sand fractions were significant, but did not appeared in the silt fraction. The fine particles (clay and silt) proportion was the highest in May and the lowest in July.

Singular fractal dimension characteristics of soil PSD. The singular fractal dimension (D) values ranged from 1.82 to 1.90 (Table 1). Analysis of variance showed the significant differences in the singular fractal dimensions in different vegetation and soil layers. The D values of P2 and P3 were higher than those of other areas (P1, P4 and P5). The D values in the topsoil were higher than those in the underlying soil layers.

Figure 2. Texture of analyzed soil samples. The size of plots indicates the concentrations of soil organic matter.

Multifractal characterization of soil PSDs. The values of capacity dimension (D_0) and entropy dimension (D_1) varied from 0.84 to 0.93 and 0.66 to 0.84, respectively. The values of D_1/D_0 ranged from 0.76 to 0.94 (Fig. 4, Table 1). The multifractal dimensions were significantly different in different vegetation. The D_0 values were the highest in P3 and P4. The highest and the smallest values of D_1 appeared in P4 and P3, respectively. The D_1/D_0 values in P1, P2, and P4 were greater than those in P3 and P5 (Table 1). Soil layers significantly influenced D_0 and D_1 , but did not affect D_1/D_0 . The values of multifractal dimensions in 0–10 cm layer were greater than in underlying soils layers. The values of D_1 and D_1/D_0 were significantly different in different seasons (Table 1). The D_1 values and the D_1/D_0 values in July were lower than in other seasons. The D_1 and D_1/D_0 values in February and May were higher than those in October and February (Table 1).

Network analysis and CCA analysis. To test the association relationships between soil PSD and fractal dimensions, a combinatorial soil PSD- D_q network was constructed (Fig. 5). All significant correlations ($p < 0.05$) were visualized as edges. The figures in the dots mean the size (μm) of the soil particles. The resulting network contained three modules: one was large, strongly connected, and the other two were small weakly connected. The large and interconnected module contained soil size grades (clay, silt and sand) and soil particle with the size ranging from $0.40\mu\text{m}$ to $126\mu\text{m}$. The D and Rényi dimension spectra (D_q) for $q > 0$ were also included. One of the small loosely interconnected modules contained the capacity dimension D_0 and soil particle size ranging $126\text{--}2000\mu\text{m}$. The other small loosely connected module contained all of D_q for $q < -0.5$. The SOM connected the large module and the soil particle with size ranging between $224\mu\text{m}$ and $564\mu\text{m}$.

The CCA model revealed that 43.5% of the variance in PSD could be explained by environmental factors, including 14.7% for seasonal variation, 8.6% for soil depth and 8.0% for vegetation difference. The variation of PSD was significantly correlated with the seasonal variation ($R^2 = 0.147$ and $p = 0.009$) and the soil PSD were related with the depth ($R^2 = 0.086$, $p = 0.047$). The significant relation between PSD and vegetation was not observed in the study ($R^2 = 0.080$, $p = 0.06$).

Figure 3. Log-log plots of particle size distributions for soil samples.

Factor	Variation	D	D_0	D_1	D_1/D_0	clay	Silt	Sand
Vegetation	P1	1.84	0.87	0.76	0.87	6.55	36.58	56.87
	P2	1.88	0.86	0.76	0.88	6.78	36.37	56.85
	P3	1.87	0.88	0.74	0.84	6.73	36.39	56.88
	P4	1.85	0.88	0.77	0.88	4.80	23.54	71.66
	P5	1.85	0.87	0.74	0.85	4.67	23.81	71.51
	F values	2.95	2.95	2.51	3.08	3.82	4.32	4.34
	p values	0.03	0.03	0.05	0.02	0.01	0.01	0.01
Soil depth	0–10	1.90	0.89	0.78	0.88	6.87	38.97	54.15
	10–20	1.84	0.86	0.74	0.86	5.39	28.1	66.51
	20–30	1.83	0.86	0.74	0.86	5.46	26.94	67.59
	F values	22.36	22.36	12.45	1.715	3.85	6.48	6.19
	p values	0	0	0	0.19	0.03	0.01	0.01
Seasonal variation	Jul(2012)	1.84	0.87	0.73	0.84	4.77	24.92	70.3
	Oct(2012)	1.86	0.87	0.75	0.86	5.28	30.13	64.58
	Feb(2013)	1.85	0.87	0.76	0.88	6.35	36.39	59.67
	May(2013)	1.87	0.88	0.77	0.88	7.22	36.32	56.45
	F values	1.8	1.8	5.36	3.17	4.98	2.74	3.05
	p values	0.16	0.16	0.002	0.03	0.01	0.05	0.04

Table 1. Effect of vegetation, soil depth and seasonal variation on singular fractal dimensions (D), capacity dimension (D_0), entropy dimension (D_1), D_1/D_0 , soil texture in soil.

Figure 4. The Rényi dimensions spectra for soil samples.

Discussions

The range of D values was relatively low (1.82–1.90) in newly formed wetland of the YRD. In soils with high fertility and fine texture, the D values ranged from 2.60–2.80, however the D values ranged from 1.83–2.64 in soil with poor structure and coarse texture^{19,28,29}. Accordingly, the soils in newly formed wetlands of the YRD were relatively coarse and infertile. Although the abundance of plant is great in this area²⁵, the soil still has poor quality because the formation of the newly formed wetland is less than 40 years (since 1976). The sand fractions in soils near the riverbank were markedly greater than those in other plots. This result was similar to the results in mangrove forests and coast, where the sand fractions are higher in soils near the seawater³⁰. The water flow could transport fine particles in soils much easier, leading to coarse texture in soils near the river or seawater.

The fractal dimensions D and D_1 were highly correlated with particles less than $126\mu\text{m}$ and SOM contents. The fractal dimensions D_0 is correlated with coarse particle ranging from $126\mu\text{m}$ to $2000\mu\text{m}$. Liu *et al.*²⁹ found the significant correlation between singular fractal dimension and clay proportion. Wang *et al.*¹² confirmed that D_1 and D_1/D_0 were significantly positively correlated with fine particle content and SOM. Other studies have also obtained similar results that the high D values were observed in the soil with fine texture in different landscapes under contrasting climate conditions^{28,30–32}. However, the range of particle size correlated with D and D_1 values was not declared in former studies. Using networks analysis, we found that the size of particles correlated with D and D_1 values were less than $126\mu\text{m}$. In pedogenic processes, soil organic carbon and nutrients accumulation, soil structure development and some biological properties are accompanied by selective increasing in fine particle size fractions^{14,18}. The smaller particle sizes reflect the greater the spatial filling capacity of the soil corresponding higher fractal dimension values based on pore geometry²⁹. Accordingly, fractal dimensions (D_1 , D) that mainly

represent fine soil particles are high correlated with SOM content in soils. The fractal dimension D_0 can prove useful information about the coarse particles, which are neglected in fractal dimensions D and D_1 values.

The fractal dimensions in topsoil were higher than that in sub-layer soils because of well-developed structure and easily dispersed soil particle (fine particle) in topsoil. High fractal dimension on surface soil is associated with fine texture and high fertility^{18,28}. The soil organic matters and nitrogen concentrations are positively correlated with fractal dimensions²⁸. Accordingly, the higher soil organic matters and nitrogen concentrations in topsoil layers would result in greater fractal dimensions. The incidence of vegetation root activities could increase the soil fractal dimension³⁵, because the plant root activities could significantly increase topsoil aggregate stability³⁶.

The fractal dimension values in soils with vegetation were significantly higher than those in soils without plant covering. Our results agree with other studies that soils with plants have higher D values^{12,28,29}. Vegetation coverage can increase the ratio of fine particles, improve the soil structure, and decrease the

risk of soil wind erosion^{37,38}. The fractal dimensions in soils under various plant species are different. The root structures of various plant species might be a source of fractal dimensions difference³⁹.

Conclusions

The fractal dimensions were low in a newly formed wetland in the YRD. Different fractal dimensions of PSD were influenced by soil particles with different sizes. The fractal dimensions D and D_1 were sensitive with fine particles ($<126\mu\text{m}$) and the fractal dimension D_0 was sensitive with coarse particles ($126\text{--}2000\mu\text{m}$). The fractal dimensions were greater in wet season than in dry season. This result indicated that the influence of organic matters in soils was greater than that of soil water content. The fractal dimensions in topsoil layers were greater than subsoil layers because of the influence of root activities. Moreover, the fractal dimensions in soils with various plant species were obviously different and were greater than in soils without vegetation. Consequently, the soil particle size distribution is greatly affected by plant root activity in newly formed wetlands in the YRD. Vegetation management is the most effectively approach for increasing soil quality. Fractal analysis makes full use of soil PSD information, and offers a useful approach to quantify and assess the soil physical attributes in newly formed wetland.

References

- Sarcinelli, T. S., Schaefer, C. E. G. R., Fernandes, E. I., Mafia, R. G. & Neri, A. V. Soil modification by termites in a sandy-soil vegetation in the Brazilian Atlantic rain forest. *J. Trop. Ecol.* **29**, 439–448 (2013).
- Gomes, J. B. V., Fernandes, M. F., Barreto, A. C., de Araujo, J. C. & Curi, N. Soil Attributes under Agroecosystems and Forest Vegetation in the Coastal Tablelands of Northeastern Brazil. *Cienc. Agrotec.* **36**, 649–664 (2012).
- Miloshis, M. & Fairfield, C. A. Coastal wetland management: A rating system for potential engineering interventions. *Ecol. Engineer.* **75**, 195–198 (2015).
- Ye, Q. H. *et al.* Spatial-temporal characteristics in landscape evolution of the Yellow River Delta during 1855–2000 and a way out for the Yellow River estuary. *Chinese Sci. Bull.* **51**, 197–209 (2006).
- Peng, G., Xiang, N., Lv, S.-q. & Zhang, G.-c. Fractal characterization of soil particle-size distribution under different land-use patterns in the Yellow River Delta Wetland in China. *J. Soil. Sediment.* **14**, 1116–1122 (2014).
- Zhang, Y. N. *et al.* Soil microbiological variability under different successional stages of the Chongming Dongtan wetland and its effect on soil organic carbon storage. *Ecol. Engineer.* **52**, 308–315 (2013).
- Hillel, D. *Fundamentals of soil physics*. (Academic Press, Inc. (London) Ltd., 1980).
- Kravchenko, A. N., Boast, C. W. & Bullock, D. G. Multifractal analysis of soil spatial variability. *Agron. J.* **91**, 1033–1041 (1999).
- Gao, G.-L. *et al.* Fractal scaling of particle size distribution and relationships with topsoil properties affected by biological soil crusts. *Plos one* **9**, e88559 (2014).
- Ghanbarian, B. & Daigle, H. Fractal dimension of soil fragment mass-size distribution: A critical analysis. *Geoderma* **245**, 98–103 (2015).
- Montero, E. & Martin, M. I. Holder spectrum of dry grain volume-size distributions in soil. *Geoderma* **112**, 197–204 (2003).
- Wang, D., Fu, B. J., Zhao, W. W., Hu, H. F. & Wang, Y. F. Multifractal characteristics of soil particle size distribution under different land-use types on the Loess Plateau, China. *Catena* **72**, 29–36 (2008).
- Paz-Ferreiro, J., Vazquez, E. V. & Miranda, J. G. V. Assessing soil particle-size distribution on experimental plots with similar texture under different management systems using multifractal parameters. *Geoderma* **160**, 47–56 (2010).
- Martin, M. A., Garcia-Gutierrez, C. & Reyes, M. Modeling Multifractal Features of Soil Particle Size Distributions with Kolmogorov Fragmentation Algorithms. *Vadose. Zone J.* **8**, 202–208 (2009).
- Li, Y., Li, M. & Horton, R. Single and Joint Multifractal Analysis of Soil Particle Size Distributions. *Pedosphere* **21**, 75–83 (2011).
- Perfect, E. Fractal models for the fragmentation of rocks and soils: a review. *Engineer. Geol.* **48**, 185–198 (1997).
- Anderson, A. N., McBratney, A. B. & Crawford, J. W. Applications of fractals to soil studies. *Adv Agron* **63**, 1–76 (1997).
- Hu, H. C., Tian, F. Q. & Hu, H. P. Soil particle size distribution and its relationship with soil water and salt under mulched drip irrigation in Xinjiang of China. *Sci. China. Technol. Sci.* **54**, 1568–1574 (2011).
- Angelico, R. *et al.* Particle size, charge and colloidal stability of humic acids coprecipitated with ferrihydrite. *Chemosphere* **99**, 239–247 (2014).
- Miranda, J. G. V., Montero, E., Alves, M. C., Gonzalez, A. P. & Vazquez, E. V. Multifractal characterization of saprolite particle-size distributions after topsoil removal. *Geoderma* **134**, 373–385 (2006).
- Eghball, B., Schepers, J. S., Negahban, M. & Schlemmer, M. R. Spatial and temporal variability of soil nitrate and corn yield: Multifractal analysis. *Agron. J.* **95**, 339–346 (2003).
- Xie, X. J. & Wei, F. Q. Soil aggregates and fractal features under different land use types in a frequent debris flow area. *J. Mt. Sci.-Engl.* **10**, 437–444 (2013).
- Wang, Y. & Aubrey, D. G. The Characteristics of the China Coastline. *Cont. Shelf. Res.* **7**, 329–349 (1987).
- Yu, J. *et al.* Effects of water discharge and sediment load on evolution of modern Yellow River Delta, China, over the period from 1976 to 2009. *Biogeosciences* **8**, 2427–2435 (2011).
- Cui, B. S., Yang, Q. C., Yang, Z. F. & Zhang, K. J. Evaluating the ecological performance of wetland restoration in the Yellow River Delta, China. *Ecol. Engineer.* **35**, 1090–1103 (2009).
- Schulte, E. Recommended soil organic matter tests. Recommended soil testing procedures for the northeastern United States. *Northeast Regional Bull* **493**, 47–56 (1995).
- Tyler, S. W. & Wheatcraft, S. W. Fractal Scaling of Soil Particle-Size Distributions - Analysis and Limitations. *Soil Sci. Soc. Am. J.* **56**, 362–369 (1992).
- Xu, G. C., Li, Z. B. & Li, P. Fractal features of soil particle-size distribution and total soil nitrogen distribution in a typical watershed in the source area of the middle Dan River, China. *Catena* **101**, 17–23 (2013).
- Liu, X., Zhang, G. C., Heathman, G. C., Wang, Y. Q. & Huang, C. H. Fractal features of soil particle-size distribution as affected by plant communities in the forested region of Mountain Yimeng, China. *Geoderma* **154**, 123–130 (2009).
- Nguyen, H. Y. T., Cao, D. M. & Schmitt, K. Soil particle-size composition and coastal erosion and accretion study in Soc Trang mangrove forests. *J. Coast. Conserv.* **17**, 93–104 (2013).
- Eclesia, R. P., Jobbagy, E. G., Jackson, R. B., Biganzoli, F. & Pineiro, G. Shifts in soil organic carbon for plantation and pasture establishment in native forests and grasslands of South America. *Global Change Biol.* **18**, 3237–3251 (2012).
- Ghorbani, N., Raiesi, F. & Ghorbani, S. Bulk soil and particle size-associated C and N under grazed and ungrazed regimes in Mountainous arid and semi-arid rangelands. *Nutr. Cycl. Agroecosys.* **93**, 15–34 (2012).

33. Salako, F. K., Babalola, O., Hauser, S. & Kang, B. T. Soil macroaggregate stability under different fallow management systems and cropping intensities in southwestern Nigeria. *Geoderma* **91**, 103–123 (1999).
34. Zhang, G. H., Liu, G. B., Wang, G. L. & Wang, Y. X. Effects of Vegetation Cover and Rainfall Intensity on Sediment-Bound Nutrient Loss, Size Composition and Volume Fractal Dimension of Sediment Particles. *Pedosphere* **21**, 676–684 (2011).
35. Bai, E. *et al.* Spatial variation of soil delta C-13 and its relation to carbon input and soil texture in a subtropical lowland woodland. *Soil Biol. Biochem.* **44**, 102–112 (2012).
36. Pohl, M., Graf, F., Buttler, A. & Rixen, C. The relationship between plant species richness and soil aggregate stability can depend on disturbance. *Plant Soil* **355**, 87–102 (2012).
37. Zhang, G. H., Liu, G. B., Wang, G. L. & Wang, Y. X. Effects of vegetation cover and rainfall intensity on sediment-associated nitrogen and phosphorus losses and particle size composition on the Loess Plateau. *J. Soil Water Conserv.* **66**, 192–200 (2011).
38. Yan, Y. C. *et al.* Quantitative effects of wind erosion on the soil texture and soil nutrients under different vegetation coverage in a semiarid steppe of northern China. *Plant Soil* **369**, 585–598 (2013).
39. Reubens, B., Poesen, J., Danjon, F., Geudens, G. & Muys, B. The role of fine and coarse roots in shallow slope stability and soil erosion control with a focus on root system architecture: a review. *Trees-Struct Funct* **21**, 385–402 (2007).

Acknowledgments

We would like to thank the Project of National Science & Technology Pillar Program in “12th Five Year” period (2011BAC02B01), National Natural Science Foundation for Distinguished Young Scholar of Shandong Province (No.JQ201114), the National Science Foundation of China (41301333) and the CAS/SAFEA International Partnership Program for Creative Research Teams - “Representative environmental processes and resources effects in coastal zone”. We also would like to thank the Yellow River Delta Wetland Ecological Experimental Station, CAS, for providing field experimental and residential place for this study.

Author Contributions

X.L., B.M. and J.Y. designed and coordinated the study. X.L., S.D., M.Z., G.H. and Y.Y. performed the experiments. X.L. and B.M. were involved in analyzing the data. B.M. and H.W. contributed analysis tools. X.L. and B.M. wrote the paper and all the authors participated in revising. All authors have read and approved the final manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Yu, J. *et al.* Fractal features of soil particle size distribution in the newly formed wetlands of the Yellow River Delta estuary. *Sci. Rep.* **5**, 10540; doi: 10.1038/srep10540 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>