SCIENTIFIC REPORTS

OPEN

Circulating interleukin-6 and cancer: A meta-analysis using Mendelian randomization

Received: 10 December 2014

Accepted: 19 May 2015

Published: 22 June 2015

Geng Tian^{1,*}, Jia Mi^{1,*}, Xiaodan Wei^{1,*}, Dongmei Zhao², Lingyan Qiao³, Chunhua Yang¹, Xianglin Li¹, Shuping Zhang⁴, Xuri Li¹ & Bin Wang⁵

Interleukin-6 (IL-6) plays a contributory role in the progression and severity of many forms of cancer; it however remains unclear whether the relevance between circulating IL-6 and cancer is causal. We therefore meta-analyzed published articles in this regard using *IL-6* gene -174G/C variant as an instrument. Seventy-eight and six articles were eligible for the association of -174G/C variant with cancer and circulating IL-6, respectively. Overall analyses failed to identify any significance between -174G/C and cancer risk. In Asians, carriers of the -174CC genotype had an 1.95-fold increased cancer risk compared with the -174GG genotype carriers ($P = 0.009$). By cancer type, significance was only attained for liver cancer with the -174C allele conferring a reduced risk under allelic (odds ratio or OR = 0.74; $P = 0.001$), homozygous genotypic (OR = 0.59; $P = 0.029$) and dominant (OR = 0.67; $P = 0.004$) models. Carriers of the -174CC genotype (weighted mean difference or WMD = -4.23 pg/mL; $P < 0.001$) and -174C allele (WMD = -3.43 pg/mL; $P < 0.001$) had circulating IL-6 reduced significantly compared with the non-carriers. In further Mendelian randomization analysis, a reduction of 1 pg/mL in circulating IL-6 was significantly associated with an 12% reduced risk of liver cancer. Long-term genetically-reduced circulating IL-6 might be causally associated with a lower risk of liver cancer.

As a multifactorial cytokine, interleukin-6 (IL-6) is widely believed to play a role in the progression and severity of many forms of cancer. Several observational studies have suggested that circulating IL-6 can explain inter-individual variability in predisposition to cancer. Heikkilä and colleagues have written an excellent review, highlighting the involvement of elevated circulating IL-6 in human carcinogenesis¹. However, it remains unclear whether the relevance between circulating IL-6 and cancer is causal as the issue of confounding or reverse causation is often unavoidable in observational epidemiology.

Ideally, randomized controlled trial of the intervention that alters circulating IL-6 is considered as the gold standard for evaluating this causal relevance, but in some circumstances it is neither practical nor ethical to randomize human beings for such trials. As an alternative, a more promising method termed as 'Mendelian randomization' has been developed to exploit the impact of long-term exposure differences on disease risk by using a genetic instrument to account for the potential biases due to confounding and reverse causation². Like a randomized controlled trial, Mendelian randomization randomizes genotypes at conception according to Mendel's second law³. This method has been successfully applied to a variety of genetic exposures such as obesity and alcohol consumption in causal susceptibility to cancer⁴.

The most immediate determinant of circulating IL-6 is its encoding gene *IL-6*. The genomic sequence of *IL-6* gene is highly polymorphic and one of the most frequently evaluated variants is -174G/C

¹Medicine and Pharmacy Research Center, Binzhou Medical University, Laishan District, Yantai, Shandong, China. ²Institute of Anatomy, Binzhou Medical University, Laishan District, Yantai, Shandong, China. ³Clinic Institute, Binzhou Medical University, Laishan District, Yantai, Shandong, China. ⁴Institute of Pharmacology, Binzhou Medical University, Laishan District, Yantai, Shandong, China. ⁵Institute of Molecular Imaging, Binzhou Medical University, Laishan District, Yantai, Shandong, China. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to G.T. (email: tiangengshandong@yeah.net) or B.W. (email: wangbindoctor2015@163.com)

Figure 1. Flow diagram of search strategy and study selection.

(rs1800795) in the promoter region^{5,6}. *In vitro* studies have found the allele-specific impact of -174G/C variant on *IL-6* gene promoter activity, with the -174C allele corresponding to lower expression level⁷. Besides, carriers of the -174G allele had a higher level of circulating IL-6 than those with the -174CC genotype^{8,9}. Based on these observations, we thereby develop a completing hypothesis that the association between circulating IL-6 and cancer is causally related. To test this hypothesis, we employed Mendelian randomization technique to meta-analyze all available published articles in this regard by using *IL-6* gene -174G/C variant as an instrument.

Results

Eligible articles. The selection process of articles is schematized in Fig. 1. A total of 837 potentially relevant articles were identified after an initial literature search and 80 of them written in English language were finally analyzed^{5,6,8–84}. Article involving more than one independent study group was analyzed separately. Altogether, 78 articles with 87 study groups (45569 cancer patients and 57990 controls) were eligible for the association between *IL-6* gene -174G/C variant and cancer^{6,9–84}, and 6 articles with 9 study groups (1727 study subjects) were eligible for circulating IL-6 changes across -174G/C genotypes^{5,8,9,13,16,52}.

Study characteristics. Table 1 summarizes the baseline characteristics of 87 eligible study groups for the association between *IL-6* gene -174G/C variant and cancer. Supplementary Table S1 provides the quality assessment of 87 study groups and the genotype distributions of -174G/C variant. The quality

Author (year)	Ethnicity	Cancer type	Match	Source of controls	Study design	Sample size		Age (years)		Males		BMI (kg/m ²)		Smoking		Drinking	
						Cas-es	Cont's	Cas-es	Cont's	Cas-es	Cont's	Cases	Cont's	Cas-es	Cont's	Cas-es	Cont's
Slattery (2014)	Mixed	Breast	NA	Population	Prosp.	3567	4157	NA	NA	0.00	0.00	NA	NA	NA	NA	NA	NA
Mandal (2014)	Caucasian	Prostate	YES	Hospital	Retro-sp.	84	78	59.8	57.2	1.00	1.00	NA	NA	NA	NA	NA	NA
Mandal (2014)	African-American	Prostate	YES	Hospital	Retro-sp.	80	62	67.9	64.0	1.00	1.00	NA	NA	NA	NA	NA	NA
Cil (2014)	Mixed	Thyroid	YES	Hospital	Retro-sp.	190	216	47.2	46.0	0.23	0.26	25.55	25.70	0.34	0.32	NA	NA
Tindall (2012)	Caucasian	Prostate	YES	Population	Prosp.	818	734	NA	NA	1.00	1.00	NA	NA	NA	NA	NA	NA
Giannitrapani (2011)	Caucasian	Liver	YES	Population	Retro-sp.	95	98	NA	NA	0.53	NA	NA	NA	NA	NA	NA	NA
Giannitrapani (2011)	Caucasian	Liver	YES	Population	Retro-sp.	105	98	NA	NA	0.63	NA	NA	NA	NA	NA	NA	NA
Gaur (2011)	Mixed	Oral	YES	Hospital	Prosp.	140	120	51.4	51.4	0.85	0.85	NA	NA	NA	NA	NA	NA
Abuli (2011)	Caucasian	Colorectal	YES	Population	Prosp.	1405	1388	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cacev (2010)	Caucasian	Colorectal	NA	Population	Retro-sp.	160	160	64.5	63.1	0.53	0.54	NA	NA	NA	NA	NA	NA
Ognjanovic (2010)	Mixed	Colorectal	YES	Population	Retro-sp.	271	539	62.5	62.0	0.68	0.69	26.90	26.70	0.14	0.10	0.45	0.41
Hawken (2010)	Mixed	Colorectal	NA	Population	Retro-sp.	1133	1125	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Dossus (2010)	Mixed	Breast	YES	Population	Prosp.	6292	8135	63.1	63.1	0.00	0.00	NA	NA	NA	NA	NA	NA
Dossus (2010)	Mixed	Prostate	YES	Population	Prosp.	8008	8604	68.4	68.4	1.00	1.00	NA	NA	NA	NA	NA	NA
Tsilidis (2009)	Mixed	Colorectal	YES	Population	Prosp.	208	318	62.8	62.8	0.46	0.45	26.30	26.00	0.13	0.13	NA	NA
Ozgen (2009)	Mixed	Thyroid	NA	Hospital	Retro-sp.	42	340	43.1	43.8	0.19	0.19	NA	NA	NA	NA	NA	NA
Ognjanovic (2009)	Mixed	Liver	YES	Population	Retro-sp.	120	230	60.5	59.5	0.68	0.60	NA	NA	0.42	0.28	0.71	0.77
Gangwar (2009)	Asian	Cervical	YES	Hospital	Retro-sp.	160	200	45.0	46.0	0.00	0.00	NA	NA	0.34	0.10	0.06	0.01
Falleti (2009)	Caucasian	Liver	NO	Population	Retro-sp.	219	236	53.0	46.0	0.70	0.70	NA	NA	NA	NA	NA	NA
Cherel (2009)	Caucasian	Breast	NA	Hospital	Prosp.	293	112	NA	NA	0.00	0.00	NA	NA	NA	NA	NA	NA
Vasku (2009)	Caucasian	Colorectal	YES	Hospital	Retro-sp.	102	101	68.0	68.1	0.77	0.58	NA	NA	NA	NA	NA	NA
Talar-Wojnarowska (2009)	Caucasian	Pancreatic	YES	Hospital	Retro-sp.	97	50	NA	NA	0.57	0.57	NA	NA	NA	NA	NA	NA
Slattery (2009)	Mixed	Colorectal	NA	Population	Retro-sp.	1839	2014			0.55	0.54	NA	NA	NA	NA	NA	NA
Andrie (2009)	Caucasian	Lymphoma	YES	Hospital	Retro-sp.	81	81	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Aladzsyty (2009)	Caucasian	Myeloma	YES	Hospital	Retro-sp.	97	99	65.0	68.0	0.35	0.45	NA	NA	NA	NA	NA	NA
Birmann (2009)	Mixed	Myeloma	NA	Population	Prosp.	82	159	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Wilkening (2008)	Caucasian	Colorectal	YES	Population	Prosp.	308	585	56.8	56.8	0.44	0.44	NA	NA	NA	NA	NA	NA
Vairaktaris (2008)	Caucasian	Oral	YES	Population	Retro-sp.	162	168	58.5	54.7	0.80	0.75	NA	NA	NA	NA	NA	NA
Upadhyay (2008)	Asian	Esophageal	YES	Hospital	Retro-sp.	168	201	56.8	53.7	0.74	0.75	NA	NA	0.82	NA	0.37	NA
Continued																	

Author (year)	Ethnicity	Cancer type	Match	Source of controls	Study design	Sample size		Age (years)		Males		BMI (kg/m ²)		Smoking		Drinking	
						Cas-es	Cont's	Cas-es	Cont's	Cas-es	Cont's	Cases	Cont's	Cas-es	Cont's	Cas-es	Cont's
Slattery (2008) b	Caucasian	Breast	YES	Population	Retro-sp.	1176	1330	NA	NA	0.00	0.00	NA	NA	NA	NA	NA	NA
Slattery (2008) a	Mixed	Breast	YES	Population	Retro-sp.	576	727	NA	NA	0.00	0.00	NA	NA	NA	NA	NA	NA
Kesarwani (2008)	Asian	Prostate	YES	Population	Retro-sp.	200	200	62.5	59.5	1.00	1.00	NA	NA	0.32	0.30	0.60	0.69
Crusius (2008)	Caucasian	Gastric	YES	Population	Prosp.	439	1138	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Colakogul-lari (2008)	Mixed	Lung	YES	Population	Prosp.	44	58	60.0	NA	0.91	0.51	NA	NA	NA	NA	NA	NA
Bao (2008)	Asian	Prostate	YES	Hospital	Retro-sp.	136	120	62.8	62.3	1.00	1.00	NA	NA	NA	NA	NA	NA
Vogel (2008)	Caucasian	Lung	YES	Population	Retro-sp.	403	744	NA	NA	0.54	0.56	NA	NA	0.73	0.35	NA	NA
Kury (2008)	Caucasian	Colorec-tal	YES	Hospital	Retro-sp.	1023	1121	65.7	61.9	0.62	0.54	NA	NA	NA	NA	NA	NA
Ennas (2008)	Caucasian	Leukae-mia	NO	Population	Retro-sp.	40	113	61.8	56.5	0.73	0.40	NA	NA	NA	NA	NA	NA
Ahirwar (2008)	Asian	Bladder	YES	Population	Prosp.	136	200	61.6	58.3	0.88	0.88	NA	NA	NA	NA	NA	NA
Vishnoi (2007)	Asian	Gallblad-der	YES	Population	Retro-sp.	45	82	49.4	50.0	1.00	1.00	NA	NA	NA	NA	NA	NA
Vishnoi (2007)	Asian	Gallblad-der	YES	Population	Retro-sp.	79	118	49.4	50.0	0.00	0.00	NA	NA	NA	NA	NA	NA
Litovkin (2007)	Caucasian	Breast	YES	Population	Prosp.	73	143	55.0	33.0	0.00	0.49	NA	NA	NA	NA	NA	NA
Litovkin (2007)	Caucasian	Leiomy-oma	YES	Population	Prosp.	60	143	37.0	33.0	0.00	0.49	NA	NA	NA	NA	NA	NA
Gonullu (2007)	Mixed	Breast	YES	Population	Retro-sp.	38	24	47.0	39.0	0.00	0.00	NA	NA	NA	NA	NA	NA
Vogel (2007)	Caucasian	Breast	YES	Population	Prosp.	361	361	NA	NA	0.00	0.00	25.00	25.00	0.34	0.36	NA	NA
Vogel (2007)	Caucasian	Colorec-tal	YES	Population	Prosp.	355	753	59.0	56.0	0.56	0.56	26.00	26.00	0.37	0.35	NA	NA
Slattery (2007)	Mixed	Colorec-tal	YES	Population	Retro-sp.	1583	1979	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Slattery (2007)	Mixed	Colorec-tal	YES	Population	Retro-sp.	797	1011	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Nearman (2007)	Mixed	Leukae-mia	NA	Hospital	Retro-sp.	28	362	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Gatti (2007)	Mixed	Gastric	NA	Hospital	Retro-sp.	56	56	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Duch (2007)	Mixed	Myeloma	YES	Hospital	Retro-sp.	52	60	58.5	59.3	0.42	0.60	NA	NA	NA	NA	NA	NA
Deans (2007)	Caucasian	Gastric	NA	Population	Retro-sp.	203	224	71.0	39.2	0.66	0.53	NA	NA	NA	NA	NA	NA
Berkovic (2007)	Caucasian	Gastric	YES	Hospital	Retro-sp.	80	162	80.0	46.5	0.48	0.48	NA	NA	NA	NA	NA	NA
Vairaktaris (2006)	Caucasian	Oral	YES	Population	Retro-sp.	162	156	58.5	55.5	0.80	0.77	NA	NA	NA	NA	NA	NA
Theodor-opoulos (2006)	Caucasian	Colorec-tal	YES	Population	Prosp.	222	200	64.7	62.7	0.58	0.60	NA	NA	NA	NA	NA	NA
Nogueira (2006)	Mixed	Cervical	YES	Population	Retro-sp.	56	253	52.2	54.0	0.00	0.00	NA	NA	NA	NA	NA	NA
Michaud (2006)	Mixed	Prostate	YES	Population	Prosp.	503	652	67.1	66.6	1.00	1.00	NA	NA	0.08	0.11	NA	NA
Kamangar (2006)	Caucasian	Gastric	YES	Population	Prosp.	110	203	58.5	59.0	1.00	1.00	NA	NA	1.00	1.00	NA	NA
Continued																	

Author (year)	Ethnicity	Cancer type	Match	Source of controls	Study design	Sample size		Age (years)		Males		BMI (kg/m ²)		Smoking		Drinking	
						Cas-es	Cont's	Cas-es	Cont's	Cas-es	Cont's	Cases	Cont's	Cas-es	Cont's	Cas-es	Cont's
Gonzalez-Zuloeta (2006)	Caucasian	Breast	NA	Population	Prosp.	171	3651	67.8	70.8	0.00	0.00	26.70	27.10	NA	NA	NA	NA
Balasubramanian (2006)	Caucasian	Breast	NA	Hospital	Prosp.	197	490	63.0	57.0	0.00	0.00	NA	NA	NA	NA	NA	NA
Rothman (2006)	Caucasian	Lymphoma	NA	Population	Retro-sp.	2658	3068	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Lan (2006)	Mixed	Lymphoma	YES	Population	Retro-sp.	518	597	NA	NA	0.00	0.00	NA	NA	NA	NA	NA	NA
Gunter (2006)	Mixed	Colorectal	YES	Hospital	Retro-sp.	244	231	60.0	57.0	0.78	0.64	26.50	25.80	0.11	0.05	NA	NA
Gaustadnes (2006)	Caucasian	Colorectal	YES	Population	Retro-sp.	230	540	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cozen (2006)	Mixed	Myeloma	YES	Population	Retro-sp.	150	112	61.0	NA	0.61	0.58	NA	NA	NA	NA	NA	NA
Seifart (2005)	Caucasian	Lung	NA	Population	Retro-sp.	182	243	63.3	37.9	0.88	0.55	NA	NA	0.98	0.40	NA	NA
Migita (2005)	Asian	Liver	NO	Hospital	Prosp.	48	188	62.5	51.5	0.81	0.68	NA	NA	NA	NA	NA	NA
Leibovici (2005)	Caucasian	Bladder	YES	Hospital	Prosp.	465	450	NA	NA	NA	NA	NA	NA	0.74	0.53	NA	NA
Hefler (2005)	Caucasian	Breast	YES	Hospital	Retro-sp.	269	227	54.9	53.3	0.00	0.00	NA	NA	NA	NA	NA	NA
Basturk (2005)	Mixed	Renal cell	YES	Population	Retro-sp.	29	50	NA	NA	0.05	0.56	NA	NA	NA	NA	NA	NA
Snoussi (2005)	Caucasian	Breast	NA	Population	Prosp.	305	200	50.0	46.0	0.01	0.05	NA	NA	NA	NA	NA	NA
Skerrett (2005)	Mixed	Breast	NA	Population	Retro-sp.	88	102	49.2	NA	0.00	0.00	NA	NA	NA	NA	NA	NA
Mazur (2005)	Caucasian	Myeloma	NA	Hospital	Retro-sp.	54	50	62.0	NA	0.43	0.58	NA	NA	NA	NA	NA	NA
Festa (2005)	Caucasian	Basal cell	NA	Population	Prosp.	241	260	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Cordano (2005)	Caucasian	Lymphoma	YES	Population	Retro-sp.	408	349	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Campa (2005)	Caucasian	Lung	NA	Hospital	Retro-sp.	1995	1982	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Zhang (2004)	Caucasian	Basal cell	YES	Hospital	Retro-sp.	241	260	50.0	48.0	0.58	0.54	NA	NA	NA	NA	NA	NA
Smith (2004)	Caucasian	Breast	NO	Population	Retro-sp.	144	263	59.6	40.3	0.00	0.00	NA	NA	NA	NA	NA	NA
Campa (2004)	Caucasian	Lung	YES	Population	Prosp.	250	214	63.1	64.8	0.71	0.75	NA	NA	0.73	0.91	NA	NA
Bushley (2004)	Mixed	Ovarian	YES	Population	Retro-sp.	182	219	NA	NA	0.00	0.00	NA	NA	NA	NA	NA	NA
Landi (2003)	Mixed	Colorectal	YES	Hospital	Retro-sp.	377	326	NA	NA	0.60	0.53	NA	NA	0.15	0.18	0.67	0.56
El-Omar (2003)	Mixed	Esophageal	YES	Population	Retro-sp.	161	210	65.5	66.0	0.87	0.85	NA	NA	0.40	0.24	0.87	0.85
El-Omar (2003)	Mixed	Gastric	YES	Population	Retro-sp.	314	210	68.0	66.0	0.81	0.85	NA	NA	0.30	0.24	0.75	0.85
Hwang (2003)	Asian	Gastric	NA	Hospital	Retro-sp.	30	30	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Hwang (2003)	Caucasian	Gastric	NA	Hospital	Retro-sp.	30	30	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Howell (2003)	Caucasian	Melanoma	NA	Hospital	Prosp.	153	208	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Zheng (2000)	Caucasian	Myeloma	NA	Population	Retro-sp.	73	129	67.0	NA	0.45	NA	NA	NA	NA	NA	NA	NA

Table 1. The baseline characteristics of all study groups in this meta-analysis. Abbreviations: BMI, body mass index; cont's, controls; NA, not available; Prosp., prospective design; Retrosp., retrospective design.

Author (year)	Ethnicity	Status	Sample size	Circulating IL-6 (pg/mL) across -174G/C genotypes					
				Mean (GG)	SD (GG)	Mean (GC)	SD (GC)	Mean (CC)	SD (CC)
Malaponte (2013)	Caucasian	Cancer patients	130	22.10	4.30	11.90	2.60	5.30	1.50
Malaponte (2013)	Caucasian	Cancer patients	190	7.20	2.20	12.00	1.70	4.50	1.10
Malaponte (2013)	Caucasian	Healthy controls	215	2.80	0.70	2.70	0.50	2.60	0.70
Giannitrapani (2011)	Caucasian	Cancer patients	67	2.20	2.86	2.30	1.70	2.90	0.60
Giannitrapani (2011)	Caucasian	Cancer patients	80	4.80	4.25	3.60	3.00	0.87	1.97
Ognjanovic (2010)	Mixed	Cancer patients and controls	806	1.90	0.05	2.23	2.69	2.35	1.91
Talar-Wojnarowska (2009)	Caucasian	Cancer patients	97	65.00	10.00	27.00	10.00	33.00	10.00
Berkovic (2007)	Caucasian	Cancer patients	80	3.07	1.03	4.31	2.98	7.51	4.33
Belluco (2003)	Caucasian	Cancer patients	62	0.32	0.62	0.14	0.13	0.14	0.13

Table 2. Mean circulating IL-6 across IL-6 gene -174G/C genotypes in this meta-analysis. Abbreviations: SD, standard deviation.

score ranged from 4 to 11 and was averaged at 8.37. Cancer patients were older than controls (mean age: 59.26 versus 55.01 years, $P = 0.0007$). The percentage of smokers was slightly higher in cancer patients than healthy controls (42.24% versus 33.06%, $P = 0.038$). No significance was observed in mean body mass index and the percentages of male gender and drinkers between the two groups ($P > 0.05$).

Forty-six of 87 study groups were conducted in Caucasians, 31 in mixed populations, 9 in Asians and 1 in African-Americans. Sixteen study groups focused on colorectal cancer, 14 groups on breast cancer, 8 groups on gastric cancer, 7 groups on prostate cancer, 6 groups on myeloma, 5 groups respectively on lung and liver cancers, 4 groups on lymphoma, 3 groups on oral cancer, 2 groups respectively on esophageal, basal cell, bladder, cervical, leukatmia, gallbladder and thyroid cancers, 1 group respectively on melanoma, ovarian, renal cell, leiomyoma and pancreatic cancers. Thirty-three of 87 study groups had total sample size of at least 500. Age was reported to be matched in 60 study groups and unmatched in 4 study groups between cancer patients and controls. Fifty-seven study groups had controls enrolled from general populations and 30 study groups from hospitals. Fifty-nine and 28 study groups followed a retrospective and prospective design, respectively. Sixty-five of 87 study groups had genotype distributions of -174G/C variant in Hardy-Weinberg equilibrium at a significance level of 5%, 19 study groups in Hardy-Weinberg disequilibrium, and 3 study groups without mutation. The frequency of IL-6 gene -174C allele ranged from 0.0% to 52.94% in cancer patients and from 0.0% to 51.15% in healthy controls. By contrast, this frequency was significantly lower in Asians than in Caucasians (cancer patients: 0.0% to 28.0% versus 17.90% to 52.94%; healthy controls: 0.0% to 26.75% versus 13.33% to 51.15%).

Table 2 summarizes mean circulating IL-6 across -174G/C genotypes. All study groups involved Caucasians except for one with mixed descents. Seven of 9 study groups provided circulating IL-6 in cancer patients, and 1 study group respectively in healthy controls and in combined patients and controls.

Prediction of -174G/C variant for cancer risk. Overall and subgroup estimates of IL-6 gene -174G/C variant for cancer risk are provided in Table 3. Overall analyses failed to identify any significance for the -174C allele under allelic (OR = 1.02; 95% CI: 0.98 to 1.07; $P = 0.290$), homozygous genotypic (OR = 1.07; 95% CI: 0.99 to 1.16; $P = 0.103$) and dominant (OR = 1.02; 95% CI: 0.97 to 1.08; $P = 0.465$) models, with moderate heterogeneity ($I^2 = 66.6\%$, 54.0% and 65.1%, respectively). There was no indication of publication bias for three genetic models except for the homozygous genotypic model (Egger's test: $P = 0.075$) (Fig. 2). After restricting study groups with Hardy-Weinberg equilibrium, there was no material change in effect estimates.

Potential sources of heterogeneity were exploited by subgroup analyses according to ethnicity, cancer type, matching condition, source of controls, study design and sample size, respectively (Table 3). To avoid chance results, only subgroups with three or more study groups were considered in this meta-analysis. By ethnicity, no significance was attained in Caucasians under three genetic models, and contrastingly Asian carriers of -174CC genotype were observed to have 1.95-fold increased cancer risk compared with those with the -174GG genotype (95% CI: 1.95; 1.19 to 3.20; $P = 0.009$), even after the Bonferroni correction to control for the multiple testing ($P < 0.05/3$, here 3 refers to the total number of subgroups by ethnicity). There was no heterogeneity ($I^2 = 17.9\%$) for this significant association.

By cancer type, effect estimates were significant only for liver cancer with the -174C allele conferring a reduced risk under allelic (OR = 0.74; 95% CI: 0.61 to 0.89; $P = 0.001$), homozygous genotypic (OR = 0.59; 95% CI: 0.36 to 0.95; $P = 0.029$) and dominant (OR = 0.67; 95% CI: 0.52 to 0.88; $P = 0.004$) models, and this significance was less likely to be interpreted by heterogeneity ($I^2 = 0.0\%$, 0.0% and

Groups	Studies	Allelic model		Genotypic model		Dominant model	
		OR; 95% CI; P	I ²	OR; 95% CI; P	I ²	OR; 95% CI; P	I ²
Overall	87	1.02; 0.98–1.07; 0.290	66.6%	1.07; 0.99–1.16; 0.103	54.0%	1.02; 0.97–1.08; 0.465	65.1%
Ethnicity							
Caucasian	46	1.05; 0.98–1.12; 0.138	72.2%	1.09; 0.97–1.23; 0.136	59.5%	1.07; 0.98–1.17; 0.155	71.4%
Asian	9	1.03; 0.75–1.42; 0.870	68.8%	1.95; 1.19–3.20; 0.009	17.9%	0.90; 0.66–1.22; 0.483	52.7%
Mixed	31	0.98; 0.93–1.03; 0.374	50.6%	0.97; 0.88–1.06; 0.450	27.5%	0.97; 0.91–1.04; 0.430	53.0%
Sample size							
<500	54	1.03; 0.92–1.15; 0.615	73.7%	1.17; 0.94–1.45; 0.163	61.8%	1.03; 0.89–1.20; 0.654	72.7%
≥500	33	1.01; 0.98–1.04; 0.730	43.5%	1.02; 0.96–1.08; 0.615	29.2%	1.01; 0.96–1.05; 0.817	41.4%
Cancer type							
Myeloma	6	1.06; 0.89–1.28; 0.496	0.0%	1.13; 0.72–2.50; 0.592	0.0%	1.09; 0.84–1.40; 0.521	0.0%
Gastric	8	1.01; 0.79–1.28; 0.960	68.5%	1.12; 0.81–1.54; 0.498	28.2%	1.04; 0.72–1.51; 0.819	74.1%
Colorectal	16	1.00; 0.93–1.07; 0.941	63.3%	0.99; 0.88–1.13; 0.914	51.7%	1.01; 0.92–1.11; 0.850	58.1%
Lung	5	1.03; 0.95–1.11; 0.530	3.6%	1.06; 0.91–1.22; 0.463	0.0%	1.03; 0.86–1.25; 0.743	41.8%
Breast	14	0.99; 0.93–1.05; 0.716	44.0%	1.02; 0.89–1.16; 0.824	11.3%	0.99; 0.89–1.09; 0.773	58.2%
Lymphoma	4	1.00; 0.95–1.07; 0.888	0.0%	1.01; 0.89–1.15; 0.855	0.0%	1.00; 0.92–1.10; 0.940	0.0%
Liver	5	0.74; 0.61–0.89; 0.001	0.0%	0.59; 0.36–0.95; 0.029	0.0%	0.67; 0.52–0.88; 0.004	21.5%
Prostate	7	0.95; 0.80–1.14; 0.597	79.4%	0.94; 0.66–1.34; 0.724	76.3%	0.96; 0.81–1.13; 0.609	58.8%
Oral	3	1.49; 0.58–3.81; 0.409	95.0%	2.38; 0.34–16.93; 0.385	91.6%	1.98; 0.54–7.26; 0.303	95.4%
Matched							
NA	23	1.00; 0.94–1.06; 0.961	43.1%	0.99; 0.89–1.10; 0.821	18.1%	1.00; 0.92–1.08; 0.926	41.1%
YES	60	1.03; 0.98–1.09; 0.235	71.5%	1.11; 1.00–1.23; 0.044	59.9%	1.03; 0.96–1.11; 0.389	70.8%
NO	4	0.99; 0.68–1.45; 0.976	70.4%	1.00; 0.40–2.50; 0.999	75.0%	0.97; 0.69–1.36; 0.846	35.1%
Control source							
Population	57	1.01; 0.97–1.06; 0.592	61.5%	1.02; 0.94–1.11; 0.622	47.9%	1.02; 0.96–1.08; 0.569	63.4%
Hospital	30	1.02; 0.91–1.14; 0.701	73.3%	1.17; 0.96–1.43; 0.018	58.9%	1.00; 0.86–1.16; 0.973	68.7%
Study design							
Retrospective	59	1.02; 0.96–1.08; 0.546	69.0%	1.09; 0.97–1.23; 0.163	54.1%	1.01; 0.93–1.10; 0.751	68.8%
Prospective	28	1.03; 0.97–1.09; 0.316	61.6%	1.06; 0.95–1.19; 0.328	55.7%	1.04; 0.96–1.11; 0.350	54.8%

Table 3. Overall and stratified risk estimates of IL-6 gene -174G/C variant for cancer risk under three genetic models. Abbreviations: OR, odds ratio; 95% CI, 95% confidence interval; NA, not available.

21.5%, respectively). Even after the Bonferroni correction, significance was still preserved for the allelic and dominant models ($P < 0.05/9$, here 9 refers to the total number of subgroups by cancer type). Effect estimates of -174G/C variant for cancer by source of controls, study design and sample size did not deviate significantly from the unity under three genetic models (all $P > 0.05$), and there was no material improvement in heterogeneity within these subgroups.

Changes of circulating IL-6 across -174G/C genotypes. Carriers of the -174CC genotype (WMD = -4.23 pg/mL; 95% CI: -6.20 to -2.25 ; $P < 0.001$) and -174C allele (-174CC and -174GC genotypes) (WMD = -3.43 pg/mL; 95% CI: -4.94 to -1.93 ; $P < 0.001$) had significantly lowered circulating IL-6 when compared with the -174GG genotype carriers, yet with strong evidence of heterogeneity ($I^2 = 99.2\%$ and 99.1% , respectively) (Fig. 3).

Predicted causality of circulating IL-6 for cancer. Under the principles of Mendelian randomization, a reduction of 1 pg/mL in circulating IL-6 was significantly associated with an 12% reduced risk of liver cancer (95% CI: 0.64 to 0.99). However in Asians, this association was totally reversed with 1 pg/mL reduced circulating IL-6 corresponding to an 17% increased cancer risk (95% CI: 1.03 to 1.68). Considering that the unity was not included by above 95% CIs, it is safe to the reject the null hypothesis of none causal relevance between circulating IL-6 and certain cancer subtypes.

Sensitivity analysis. Sensitivity analysis confirmed the overall differences in risk estimates for the prediction of *IL-6* gene -174G/C variant for cancer risk and circulating IL-6 changes between -174G/C genotypes in both direction and magnitude by sequentially omitting each study once at a time and computing differential estimates for the remaining studies.

Discussion

In this meta-analysis of 80 qualified articles, we employed Mendelian randomization to test the completing hypothesis that the association between circulating IL-6 and cancer is causal. The most noteworthy finding of this study was that long-term genetically-reduced circulating IL-6 might be causally associated with a lower risk of liver cancer. To the authors' knowledge, this is the first comprehensive meta-analysis assessing the impact of long-term differences in circulating IL-6 on cancer risk.

Currently, targeted anti-IL-6 antibody therapy has been successfully applied in several clinical trials and found to be well tolerated in cancer patients⁸⁵. Evidence from epidemiological studies is accruing in favor of a contributory role of elevated circulating IL-6 in patients with advanced tumor stages of various cancers, such as non-small cell lung cancer, colorectal cancer and renal cell carcinoma^{86–88}. Currently, whether the progression and severity of cancer is due to elevated circulating IL-6 still remains an open question. Genetic association studies are deemed as more similar to randomized clinical trials than other types of observational epidemiological studies due to Mendelian randomization (Mendel's second law)⁸⁹. We therefore utilized Mendelian randomization to assess whether the relevance between circulating IL-6 and cancer is causal by selecting the most frequently evaluated variant -174G/C in *IL-6* gene as a genetic instrument to minimize residual confounding and reverse causation.

In this meta-analysis, risk estimates of *IL-6* gene -174G/C variant with cancer were heterozygous between Caucasians and Asians. Considering the multifactorial nature of cancer, divergent genetic backgrounds or linkage disequilibrium patterns might be the most likely explanation for such divergence⁹⁰. This is well exemplified in the present study with regard to the frequency of -174C allele, which was exceedingly lower in Asians than in Caucasians (Table 1). Even in some Asian populations, the mutation rate of this allele was zero^{40,67,81}. Generally, it is not uncommon for the same variant playing a different role in cancer susceptibility across different populations. This is the principal limitation of this Mendelian randomization meta-analysis, that is, if the other flanking variants within or near *IL-6* gene related to cancer risk are in linkage disequilibrium with -174G/C variant we have examined, this will confound our findings. What's more, such confounding is difficult to exclude completely; however it is unlikely that it would explain our finding that *IL-6* gene -174CC genotype was associated with lowered circulating IL-6 without predicting a low cancer risk. There is also evidence that a variant may be in close linkage with another nearby causal locus in one ethnic population but not in another⁹¹. In view of this limitation, it is necessary to establish an ethnicity-specific database of candidate genes and variants in susceptibility to cancer⁹². Another limitation of this study is that excluding the pleiotropy of *IL-6* gene -174G/C variant seems impractical for us since data on other inflammatory factors across -174G/C genotypes are rarely provided from most eligible articles, necessitating further confirmation using additional genetic variants and/or exposure outcomes.

It is also worth noting that *IL-6* gene -174G/C variant exhibited heterozygous association with different forms of cancer in this meta-analysis. For example, the -174C allele was observed to confer a significantly protective effect against liver cancer, yet a risk effect for oral cancer with no attainable significance. The identification of -174G/C variant affecting the significant risk of liver cancer allowed us to employ Mendelian randomization to account for potential biases due to residual confounding and reverse causation. Consistent with the findings of other studies^{8,9}, carriers of the -174C allele or -174CC genotype had lower circulating IL-6 than the non-carriers, supporting the plausibility of causal relevance between circulating IL-6 and liver cancer. Given the insufficient statistical power of this meta-analysis in some subgroups, far larger sample sizes than studied here will be required to produce enough power to evaluate the causality between circulating IL-6 and various forms of cancer.

Several limitations should be acknowledged in this meta-analysis. Firstly, only published articles were retrieved and the 'grey' literature (articles written in languages other than English) was not covered, leading to the possible existence of publication bias. However, the influence of publication bias on the gene-disease association is expected to result in an overestimation, rather than an underestimation. Secondly, as the majority of involved studies in this meta-analysis recruited cancer patients aged over 50 years for whom environmental factors are likely to contribute more prominently than a genetic component to the development of cancer, more large studies in a younger cancer population will of great interest. Thirdly, given the possible impact of drug regimens on circulating IL-6, the relationship between circulating IL-6 changes and *IL-6* gene -174G/C genotypes might be biased, calling for further validation of this relationship in healthy controls. Fourthly, nearly all involved studies had circulating IL-6 measured only once, which cannot reflect its long-term level in the development of cancer. Fifthly, this meta-analysis was based on summarized data, rather than individual participant data, precluding further gene-to-environment interaction. Sixthly, only one variant in *IL-6* gene was selected, and investigation on other candidate genes or polymorphisms involved in IL-6 regulation was highly encouraged, leaving a challengeable task to test whether this variant integrated with other risk determinants will

Allelic model

Homozygous genotypic model

Dominant model

Figure 2. The Begg's funnel plots of *IL-6* gene -174G/C variant for cancer under three genetic models.

Homozygous genotypic model

Dominant model

Figure 3. The funnel plots of circulating IL-6 changes across *IL-6* gene -174G/C genotypes under homozygous genotypic and dominant models.

enhance cancer risk prediction. The jury therefore must refrain from drawing a firm conclusion until large, well-designed studies to confirm our findings.

To sum up, there is possible evidence for causal association between long-term genetically-reduced circulating IL-6 and reduced risk of liver cancer. For practical reasons, we hope that this study will

advance our understanding of the role of circulating IL-6 leading to the progression and severity of liver cancer. Further studies to elucidate the specific role of IL-6 in cancer pathogenesis are required.

Methods

The implementation of this meta-analysis complied with the guidelines outlined in the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) statement (Please see the supplementary PRISMA Checklist)⁹³.

Search strategy. PubMed and Embase (excerpta medica database) were searched for potentially relevant articles from the earliest possible year to August 4, 2014. The key terms included ‘interleukin-6’, ‘IL6’, ‘IL-6’, ‘IL 6’, ‘cancer’, ‘carcinoma’, ‘neoplasia’, ‘tumor’, ‘adenoma’, ‘neoplasm’, ‘myeloma’, ‘melanoma’, ‘lymphoma’, ‘leukaemia’, ‘leiomyoma’, in combination with ‘level’, ‘concentration’, ‘polymorphism’, ‘variant’, ‘variation’, ‘mutation’, ‘SNP’. Citations from retrieved potential articles and reviews were also checked for eligibility.

The titles and abstracts of all retrieved relevant articles were independently reviewed by two investigators (Chunhua Yang and Xuri Li). In case of uncertainty for rejection, the full text and supplementary materials if available were downloaded to check whether information on the topic of interest was provided. If more than one article from a study group was published, data from the most recent or complete article were abstracted. The eligibility of each retrieved article was assessed in duplicate and independently by two authors (Chunhua Yang and Xuri Li). Any uncertainty over the eligibility was adjudicated by further joint inspection of the articles.

Inclusion/exclusion criteria. Inclusion criteria were to test the hypothesis that *IL-6* gene -174G/C variant was associated with cancer or circulating IL-6 and to provide detailed genotype or allele counts of this variant between cancer patients and healthy controls or the mean levels of circulating IL-6 across -174G/C genotypes. Articles were excluded if they assessed the progression, severity, phenotypic modification and response to treatment or survival of cancer, or if they lacked patients or controls, or if they were conference abstracts or proceedings, case reports or series, editorials, narrative reviews, and non-English articles.

Data extraction. Two authors (Chunhua Yang and Xuri Li) independently abstracted the following data from each qualified article according to a fixed protocol, including the first author’s last name, publication year, ethnicity, cancer type, matching condition, study design, sample size, the genotype or allele counts of the -174G/C variant between patients and controls, mean level of circulating IL-6 for each genotype carriers expressed as mean $\pm$ standard deviation, as well as some baseline characteristics of study populations where available, including age, gender, body mass index, the percentages of smokers and drinkers. The unit of circulating IL-6 was uniformly transferred into pg/mL in this meta-analysis.

Quality assessment. Criteria for quality assessment of the association between *IL-6* gene -174G/C variant and cancer risk were in agreement with the standards formulated by Thakkestian *et al.*⁹⁴ There were 7 criteria in total and summarized as a quality score, ranging from 0 (the worst) to 12 (the best). Quality assessment was independently conducted by two authors (Chunhua Yang and Xuri Li), and any disagreement was resolved by consensus.

Statistics. Risk estimates for the association of *IL-6* gene -174G/C variant with cancer were expressed as odds ratio (OR) and its corresponding 95% confidence interval (95% CI), and for the changes of circulating IL-6 between genotypes of this variant as weighted mean difference (WMD) and its 95% CI. Hardy–Weinberg equilibrium was tested by Chi-squared test. Differences at $P < 0.05$ were accepted as statistically significant. A random-effects model was employed to bring individual effect-size estimates together by using the DerSimonian and Laird method.

Heterogeneity between studies was quantified by the inconsistency index (I^2) statistic, which ranges from 0% to 100%. This statistic is defined as the percentage of the observed between-study variability that is due to heterogeneity rather than chance. A threshold of over 50% for I^2 statistic was treated as statistically significant heterogeneity.

Predefined subgroup analyses were explored to identify the potential sources of between-study heterogeneity according to ethnicity, cancer type, matching condition, source of controls (population-based controls and hospital-based controls), study design (prospective study and retrospective study) and sample size (< 500 subjects and ≥ 500 subjects). To assess the contribution of each individual studies to pooled effect estimates, sensitivity analyses were undertaken by sequentially omitting each study one at a time and computing differential estimates for remaining studies.

Publication bias was assessed by the Begg’s funnel plot and the Egger regression asymmetry test. The Egger test can identify the asymmetry of funnel plots by determining whether the intercept deviates significantly from zero in regressing the standardized effect estimates against their precision. A value of $P < 0.10$ was used to indicate statistical significance for Egger’s test.

Risk estimates in Mendelian randomization analysis were calculated as the ratio of the coefficient of the association between *IL-6* gene -174G/C variant and cancer to that of the association between -174G/C variant genotypes and circulating IL-6 as a reflection of the potential causal impact of circulating IL-6 on cancer.

All statistical analyses described above were completed with the StataCorp STATA version 12.0.

References

- Heikkilä, K., Ebrahim, S. & Lawlor, D. A. Systematic review of the association between circulating interleukin-6 (IL-6) and cancer. *Eur J Cancer* **44**, 937–945 (2008).
- Bochud, M. & Rousson, V. Usefulness of Mendelian randomization in observational epidemiology. *Int J Environ Res Public Health* **7**, 711–728 (2010).
- Timpson, N. J. *et al.* Does greater adiposity increase blood pressure and hypertension risk?: Mendelian randomization using the FTO/MC4R genotype. *Hypertension* **54**, 84–90 (2009).
- Brennan, P. *et al.* Obesity and cancer: Mendelian randomization approach utilizing the FTO genotype. *Int J Epidemiol* **38**, 971–975 (2009).
- Belluco, C. *et al.* -174 G>C polymorphism of interleukin 6 gene promoter affects interleukin 6 serum level in patients with colorectal cancer. *Clin Cancer Res* **9**, 2173–2176 (2003).
- Mandal, S., Abebe, F. & Chaudhary, J. -174G/C polymorphism in the interleukin-6 promoter is differently associated with prostate cancer incidence depending on race. *Genet Mol Res* **13**, 139–151 (2014).
- Moreira, P. R. *et al.* Interleukin-6 expression and gene polymorphism are associated with severity of periodontal disease in a sample of Brazilian individuals. *Clin Exp Immunol* **148**, 119–126 (2007).
- Malaponte, G. *et al.* IL-6-174 G>C and MMP-9-1562 C>T polymorphisms are associated with increased risk of deep vein thrombosis in cancer patients. *Cytokine* **62**, 64–69 (2013).
- Talar-Wojnarowska, R. *et al.* Clinical significance of interleukin-6 (IL-6) gene polymorphism and IL-6 serum level in pancreatic adenocarcinoma and chronic pancreatitis. *Dig Dis Sci* **54**, 683–689 (2009).
- Slattery, M. L. *et al.* Genetic variants in interleukin genes are associated with breast cancer risk and survival in a genetically admixed population: the Breast Cancer Health Disparities Study. *Carcinogenesis* **35**, 1750–1759 (2014).
- Cil, E. *et al.* Interleukin-10-1082 gene polymorphism is associated with papillary thyroid cancer. *Mol Biol Rep* **41**, 3091–3097 (2014).
- Tindall, E. A. *et al.* Interleukin-6 promoter variants, prostate cancer risk, and survival. *Prostate* **72**, 1701–1707 (2012).
- Giannitrapani, L. *et al.* IL-6 -174G/C polymorphism and IL-6 serum levels in patients with liver cirrhosis and hepatocellular carcinoma. *OMICS* **15**, 183–186 (2011).
- Gaur, P., Mittal, M., Mohanti, B. & Das, S. Functional variants of IL4 and IL6 genes and risk of tobacco-related oral carcinoma in high-risk Asian Indians. *Oral Dis* **17**, 720–726 (2011).
- Abuli, A. *et al.* Case-control study for colorectal cancer genetic susceptibility in EPICOLON: previously identified variants and mucins. *BMC Cancer* **11**, 339 (2011).
- Ognjanovic, S. *et al.* Serum CRP and IL-6, genetic variants and risk of colorectal adenoma in a multiethnic population. *Cancer Causes Control* **21**, 1131–1138 (2010).
- Hawken, S. J. *et al.* The utility and predictive value of combinations of low penetrance genes for screening and risk prediction of colorectal cancer. *Hum Genet* **128**, 89–101 (2010).
- Dossus, L. *et al.* PTGS2 and IL6 genetic variation and risk of breast and prostate cancer: results from the Breast and Prostate Cancer Cohort Consortium (BPC3). *Carcinogenesis* **31**, 455–461 (2010).
- Cacev, T., Jokic, M., Loncar, B., Krizanac, S. & Kapitanovic, S. Interleukin-6-174 G/C polymorphism is not associated with IL-6 expression and susceptibility to sporadic colon cancer. *DNA Cell Biol* **29**, 177–182 (2010).
- Vasku, A., Vokurka, J. & Bienertova-Vasku, J. Obesity-related genes variability in Czech patients with sporadic colorectal cancer: preliminary results. *Int J Colorectal Dis* **24**, 289–294 (2009).
- Tsilidis, K. K. *et al.* Association of common polymorphisms in IL10, and in other genes related to inflammatory response and obesity with colorectal cancer. *Cancer Causes Control* **20**, 1739–1751 (2009).
- Slattery, M. L. *et al.* Colon tumor mutations and epigenetic changes associated with genetic polymorphism: insight into disease pathways. *Mutat Res* **660**, 12–21 (2009).
- Ozgen, A. G. *et al.* The (-174) G/C polymorphism in the interleukin-6 gene is associated with risk of papillary thyroid carcinoma in Turkish patients. *J Endocrinol Invest* **32**, 491–494 (2009).
- Ognjanovic, S., Yuan, J. M., Chaptman, A. K., Fan, Y. & Yu, M. C. Genetic polymorphisms in the cytokine genes and risk of hepatocellular carcinoma in low-risk non-Asians of USA. *Carcinogenesis* **30**, 758–762 (2009).
- Gangwar, R., Mittal, B. & Mittal, R. D. Association of interleukin-6 -174G>C promoter polymorphism with risk of cervical cancer. *Int J Biol Markers* **24**, 11–16 (2009).
- Falletti, E. *et al.* Interleukin-6 polymorphisms and gender: relationship with the occurrence of hepatocellular carcinoma in patients with end-stage liver disease. *Oncology* **77**, 304–313 (2009).
- Cherel, M. *et al.* Molecular screening of interleukin-6 gene promoter and influence of -174G/C polymorphism on breast cancer. *Cytokine* **47**, 214–223 (2009).
- Birmann, B. M. *et al.* Insulin-like growth factor-1- and interleukin-6-related gene variation and risk of multiple myeloma. *Cancer Epidemiol Biomarkers Prev* **18**, 282–288 (2009).
- Andrie, E. *et al.* Genetic variants in immunoregulatory genes and risk for childhood lymphomas. *Eur J Haematol* **83**, 334–342 (2009).
- Aladzcity, I. *et al.* Comparative analysis of IL6 promoter and receptor polymorphisms in myelodysplasia and multiple myeloma. *Leuk Res* **33**, 1570–1573 (2009).
- Wilkening, S. *et al.* Interleukin promoter polymorphisms and prognosis in colorectal cancer. *Carcinogenesis* **29**, 1202–1206 (2008).
- Vogel, U. *et al.* Polymorphisms in genes involved in the inflammatory response and interaction with NSAID use or smoking in relation to lung cancer risk in a prospective study. *Mutat Res* **639**, 89–100 (2008).
- Vairaktaris, E. *et al.* Gene expression polymorphisms of interleukins-1 beta, -4, -6, -8, -10, and tumor necrosis factors-alpha, -beta: regression analysis of their effect upon oral squamous cell carcinoma. *J Cancer Res Clin Oncol* **134**, 821–832 (2008).
- Upadhyay, R., Jain, M., Kumar, S., Ghoshal, U. C. & Mittal, B. Association of interleukin-6 (-174G>C) promoter polymorphism with risk of squamous cell esophageal cancer and tumor location: an exploratory study. *Clin Immunol* **128**, 199–204 (2008).
- Slattery, M. L. *et al.* Modifying effects of IL-6 polymorphisms on body size-associated breast cancer risk. *Obesity (Silver Spring)* **16**, 339–347 (2008).

36. Kesarwani, P., Ahirwar, D. K., Mandhani, A. & Mittal, R. D. Association between -174 G/C promoter polymorphism of the interleukin-6 gene and progression of prostate cancer in North Indian population. *DNA Cell Biol* **27**, 505–510 (2008).
37. Ennas, M. G. *et al.* Interleukin-1B (IL1B) and interleukin-6 (IL6) gene polymorphisms are associated with risk of chronic lymphocytic leukaemia. *Hematol Oncol* **26**, 98–103 (2008).
38. Crusius, J. B. *et al.* Cytokine gene polymorphisms and the risk of adenocarcinoma of the stomach in the European prospective investigation into cancer and nutrition (EPIC-EURGAST). *Ann Oncol* **19**, 1894–1902 (2008).
39. Colakogullari, M. *et al.* The involvement of IL-10, IL-6, IFN-gamma, TNF-alpha and TGF-beta gene polymorphisms among Turkish lung cancer patients. *Cell Biochem Funct* **26**, 283–290 (2008).
40. Bao, S., Yang, W., Zhou, S. & Ye, Z. Relationship between single nucleotide polymorphisms in -174G/C and -634C/G promoter region of interleukin-6 and prostate cancer. *J Huazhong Univ Sci Technol Med Sci* **28**, 693–696 (2008).
41. Ahirwar, D., Kesarwani, P., Manchanda, P. K., Mandhani, A. & Mittal, R. D. Anti- and proinflammatory cytokine gene polymorphism and genetic predisposition: association with smoking, tumor stage and grade, and bacillus Calmette-Guerin immunotherapy in bladder cancer. *Cancer Genet Cytogenet* **184**, 1–8 (2008).
42. Vogel, U. *et al.* Peroxisome proliferator-activated [corrected] receptor-gamma2 [corrected] Pro12Ala, interaction with alcohol intake and NSAID use, in relation to risk of breast cancer in a prospective study of Danes. *Carcinogenesis* **28**, 427–434 (2007).
43. Vogel, U. *et al.* Prospective study of interaction between alcohol, NSAID use and polymorphisms in genes involved in the inflammatory response in relation to risk of colorectal cancer. *Mutat Res* **624**, 88–100 (2007).
44. Vishnoi, M. *et al.* Do TNFA -308 G/A and IL6 -174 G/C gene polymorphisms modulate risk of gallbladder cancer in the north Indian population? *Asian Pac J Cancer Prev* **8**, 567–572 (2007).
45. Litovkin, K. V., Domenyuk, V. P., Bubnov, V. V. & Zaporozhan, V. N. Interleukin-6 -174G/C polymorphism in breast cancer and uterine leiomyoma patients: a population-based case control study. *Exp Oncol* **29**, 295–298 (2007).
46. Gonullu, G. *et al.* Association of breast cancer and cytokine gene polymorphism in Turkish women. *Saudi Med J* **28**, 1728–1733 (2007).
47. Slattery, M. L., Wolff, R. K., Herrick, J. S., Caan, B. J. & Potter, J. D. IL6 genotypes and colon and rectal cancer. *Cancer Causes Control* **18**, 1095–1105 (2007).
48. Nearman, Z. P. *et al.* Immunogenetic factors determining the evolution of T-cell large granular lymphocyte leukaemia and associated cytopenias. *Br J Haematol* **136**, 237–248 (2007).
49. Gatti, L. L. *et al.* Interleukin-6 polymorphisms, Helicobacter pylori infection in adult Brazilian patients with chronic gastritis and gastric adenocarcinoma. *Arch Med Res* **38**, 551–555 (2007).
50. Duch, C. R. *et al.* Analysis of polymorphism at site -174 G/C of interleukin-6 promoter region in multiple myeloma. *Braz J Med Biol Res* **40**, 265–267 (2007).
51. Deans, C. *et al.* Host cytokine genotype is related to adverse prognosis and systemic inflammation in gastro-oesophageal cancer. *Ann Surg Oncol* **14**, 329–339 (2007).
52. Berkovic, M. C. *et al.* IL-6-174 C/G polymorphism in the gastroenteropancreatic neuroendocrine tumors (GEP-NETs). *Exp Mol Pathol* **83**, 474–479 (2007).
53. Vairaktaris, E. *et al.* Strong association of interleukin-6 -174 G>C promoter polymorphism with increased risk of oral cancer. *Int J Biol Markers* **21**, 246–250 (2006).
54. Theodoropoulos, G. *et al.* Relation between common polymorphisms in genes related to inflammatory response and colorectal cancer. *World J Gastroenterol* **12**, 5037–5043 (2006).
55. Nogueira de Souza, N. C. *et al.* Interleukin-6 polymorphisms and the risk of cervical cancer. *Int J Gynecol Cancer* **16**, 1278–1282 (2006).
56. Michaud, D. S. *et al.* Genetic polymorphisms of interleukin-1B (IL-1B), IL-6, IL-8, and IL-10 and risk of prostate cancer. *Cancer Res* **66**, 4525–4530 (2006).
57. Kamangar, F. *et al.* Polymorphisms in inflammation-related genes and risk of gastric cancer (Finland). *Cancer Causes Control* **17**, 117–125 (2006).
58. Gonzalez-Zuloeta Ladd, A. M. *et al.* Interleukin 6 G-174 C polymorphism and breast cancer risk. *Eur J Epidemiol* **21**, 373–376 (2006).
59. Balasubramanian, S. P. *et al.* Interleukin gene polymorphisms and breast cancer: a case control study and systematic literature review. *BMC Cancer* **6**, 188 (2006).
60. Rothman, N. *et al.* Genetic variation in TNF and IL10 and risk of non-Hodgkin lymphoma: a report from the InterLymph Consortium. *Lancet Oncol* **7**, 27–38 (2006).
61. Lan, Q. *et al.* Cytokine polymorphisms in the Th1/Th2 pathway and susceptibility to non-Hodgkin lymphoma. *Blood* **107**, 4101–4108 (2006).
62. Gunter, M. J. *et al.* Inflammation-related gene polymorphisms and colorectal adenoma. *Cancer Epidemiol Biomarkers Prev* **15**, 1126–1131 (2006).
63. Gaustadnes, M., Orntoft, T. F., Jensen, J. L. & Topping, N. Validation of the use of DNA pools and primer extension in association studies of sporadic colorectal cancer for selection of candidate SNPs. *Hum Mutat* **27**, 187–194 (2006).
64. Cozen, W. *et al.* Interleukin-6-related genotypes, body mass index, and risk of multiple myeloma and plasmacytoma. *Cancer Epidemiol Biomarkers Prev* **15**, 2285–2291 (2006).
65. Snoussi, K., Strosberg, A. D., Bouaouina, N., Ben Ahmed, S. & Chouchane, L. Genetic variation in pro-inflammatory cytokines (interleukin-1beta, interleukin-1alpha and interleukin-6) associated with the aggressive forms, survival, and relapse prediction of breast carcinoma. *Eur Cytokine Netw* **16**, 253–260 (2005).
66. Seifart, C. *et al.* TNF-alpha, TNF-beta, IL-6, and IL-10 polymorphisms in patients with lung cancer. *Dis Markers* **21**, 157–165 (2005).
67. Migita, K. *et al.* Cytokine gene polymorphisms in Japanese patients with hepatitis B virus infection—association between TGF-beta1 polymorphisms and hepatocellular carcinoma. *J Hepatol* **42**, 505–510 (2005).
68. Leibovici, D. *et al.* Polymorphisms in inflammation genes and bladder cancer: from initiation to recurrence, progression, and survival. *J Clin Oncol* **23**, 5746–5756 (2005).
69. Hefler, L. A. *et al.* Interleukin-1 and interleukin-6 gene polymorphisms and the risk of breast cancer in caucasian women. *Clin Cancer Res* **11**, 5718–5721 (2005).
70. Basturk, B. *et al.* Cytokine gene polymorphisms as potential risk and protective factors in renal cell carcinoma. *Cytokine* **30**, 41–45 (2005).
71. Skerrett, D. L., Moore, E. M., Bernstein, D. S. & Vahdat, L. Cytokine genotype polymorphisms in breast carcinoma: associations of TGF-beta1 with relapse. *Cancer Invest* **23**, 208–214 (2005).
72. Mazur, G. *et al.* IL-6 and IL-10 promoter gene polymorphisms do not associate with the susceptibility for multiple myeloma. *Immunol Lett* **96**, 241–246 (2005).
73. Festa, F. *et al.* Basal cell carcinoma and variants in genes coding for immune response, DNA repair, folate and iron metabolism. *Mutat Res* **574**, 105–111 (2005).

74. Cordano, P. *et al.* Effect of IL-6 promoter polymorphism on incidence and outcome in Hodgkin's lymphoma. *Br J Haematol* **128**, 493–495 (2005).
75. Campa, D. *et al.* Lack of association between polymorphisms in inflammatory genes and lung cancer risk. *Cancer Epidemiol Biomarkers Prev* **14**, 538–539 (2005).
76. Zhang, Z. *et al.* Use of pyrosequencing to detect clinically relevant polymorphisms of genes in basal cell carcinoma. *Clin Chim Acta* **342**, 137–143 (2004).
77. Smith, K. C., Bateman, A. C., Fussell, H. M. & Howell, W. M. Cytokine gene polymorphisms and breast cancer susceptibility and prognosis. *Eur J Immunogenet* **31**, 167–173 (2004).
78. Campa, D. *et al.* Association of a common polymorphism in the cyclooxygenase 2 gene with risk of non-small cell lung cancer. *Carcinogenesis* **25**, 229–235 (2004).
79. Bushley, A. W. *et al.* Polymorphisms of interleukin (IL)-1alpha, IL-1beta, IL-6, IL-10, and IL-18 and the risk of ovarian cancer. *Gynecol Oncol* **95**, 672–679 (2004).
80. Landi, S. *et al.* Association of common polymorphisms in inflammatory genes interleukin (IL)6, IL8, tumor necrosis factor alpha, NFKB1, and peroxisome proliferator-activated receptor gamma with colorectal cancer. *Cancer Res* **63**, 3560–3566 (2003).
81. Hwang, I. R. *et al.* Interleukin-6 genetic polymorphisms are not related to Helicobacter pylori-associated gastroduodenal diseases. *Helicobacter* **8**, 142–148 (2003).
82. Howell, W. M., Turner, S. J., Theaker, J. M. & Bateman, A. C. Cytokine gene single nucleotide polymorphisms and susceptibility to and prognosis in cutaneous malignant melanoma. *Eur J Immunogenet* **30**, 409–414 (2003).
83. El-Omar, E. M. *et al.* Increased risk of noncardia gastric cancer associated with proinflammatory cytokine gene polymorphisms. *Gastroenterology* **124**, 1193–1201 (2003).
84. Zheng, C. *et al.* Interleukin 6, tumour necrosis factor alpha, interleukin 1beta and interleukin 1 receptor antagonist promoter or coding gene polymorphisms in multiple myeloma. *Br J Haematol* **109**, 39–45 (2000).
85. Puchalski, T., Prabhakar, U., Jiao, Q., Berns, B. & Davis, H. M. Pharmacokinetic and pharmacodynamic modeling of an anti-interleukin-6 chimeric monoclonal antibody (siltuximab) in patients with metastatic renal cell carcinoma. *Clin Cancer Res* **16**, 1652–1661 (2010).
86. Guo, Y., Xu, F., Lu, T., Duan, Z. & Zhang, Z. Interleukin-6 signaling pathway in targeted therapy for cancer. *Cancer Treat Rev* **38**, 904–910 (2012).
87. Jacobson, C. M., Rosenfeld, B., Pessin, H. & Breitbart, W. Depression and IL-6 blood plasma concentrations in advanced cancer patients. *Psychosomatics* **49**, 64–66 (2008).
88. Mitsunaga, S. *et al.* Serum levels of IL-6 and IL-1beta can predict the efficacy of gemcitabine in patients with advanced pancreatic cancer. *Br J Cancer* **108**, 2063–2069 (2013).
89. Salanti, G., Sanderson, S. & Higgins, J. P. Obstacles and opportunities in meta-analysis of genetic association studies. *Genet Med* **7**, 13–20 (2005).
90. Zhou, X. C., Dowdy, S. C., Podratz, K. C. & Jiang, S. W. Epigenetic considerations for endometrial cancer prevention, diagnosis and treatment. *Gynecol Oncol* **107**, 143–153 (2007).
91. Yu, K. *et al.* Methionine synthase A2756G polymorphism and cancer risk: a meta-analysis. *Eur J Hum Genet* **18**, 370–378 (2010).
92. Pan, H. *et al.* Contributory role of five common polymorphisms of RAGE and APE1 genes in lung cancer among Han Chinese. *PLoS One* **8**, e69018, doi:10.1371/journal.pone.0069018 (2013).
93. Moher, D., Liberati, A., Tetzlaff, J. & Altman, D. G. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* **339**, b2535 (2009).
94. Thakkestian, A. *et al.* Systematic review and meta-analysis of the association between beta2-adrenoceptor polymorphisms and asthma: a HuGE review. *Am J Epidemiol* **162**, 201–211 (2005).

Acknowledgements

Grant support: Taishan Scholars Construction Engineering; National Natural Science Foundation of China (81400771 and 81171303), Shandong provincial natural science foundation (ZR2014HL028 and ZR2010HM091) and Binzhou Medical University Scientific Research Funds (BY2013KYQD17, BY2013KYQD18).

Author Contributions

G.T. and B.W. conceived and designed the experiments; G.T., J.M. and X.D.W. performed the experiments; X.D.W. and C.H.Y. analyzed the data; X.R.L., X.L.L., D.M.Z., L.Y.Q. and S.P.Z. contributed materials/analysis tools; G.T., C.H.Y. and B.W. wrote and revised the manuscript. All authors reviewed and approved the manuscript prior to submission.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Tian, G. *et al.* Circulating interleukin-6 and cancer: A meta-analysis using Mendelian randomization. *Sci. Rep.* **5**, 11394; doi: 10.1038/srep11394 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>