

SCIENTIFIC REPORTS



OPEN

Deletion of integrin $\alpha 7$ subunit does not aggravate the phenotype of laminin $\alpha 2$ chain-deficient mice

Kinga I. Gawlik & Madeleine Durbeej

Received: 17 March 2015

Accepted: 10 August 2015

Published: 10 September 2015

Laminin-211 is a major constituent of the skeletal muscle basement membrane, exerting its biological functions by binding to cell surface receptors integrin $\alpha 7\beta 1$ and dystroglycan (the latter is part of the dystrophin-glycoprotein complex). The importance of these molecules for normal muscle function is underscored by the fact that their respective deficiency leads to different forms of muscular dystrophy with different severity in humans and animal models. We recently demonstrated that laminin $\alpha 2$ chain and members of the dystrophin-glycoprotein complex have overlapping but non-redundant roles despite being part of the same adhesion complex. To analyse whether laminin-211 and integrin $\alpha 7$ subunit have non-redundant functions we generated mice deficient in laminin $\alpha 2$ chain and integrin $\alpha 7$ subunit (*dy^{3K}/itga7*). We show that lack of both molecules did not exacerbate the severe phenotype of laminin $\alpha 2$ -chain deficient animals. They displayed the same weight, survival and dystrophic pattern of muscle biopsy, with similar degree of inflammation and fibrosis. These data suggest that laminin-211 and integrin $\alpha 7\beta 1$ have intersecting roles in skeletal muscle.

The mechanical stability and function of skeletal muscle are largely dependent on the association of the extracellular matrix with the muscle cell membrane and the intracellular cytoskeleton. The significance of this cell-matrix continuity is highlighted in humans as mutations in the genes encoding extracellular matrix, cell surface receptors or cytoskeleton-interacting proteins cause various types of muscular dystrophy¹. A central element of the extracellular matrix-muscle fiber linkage is laminin-211, a heterotrimeric cell adhesion molecule (composed of $\alpha 2$, $\beta 1$ and $\gamma 1$ subunit) that strengthens sarcolemmal stability, protects muscles from damage, controls myofiber survival and regeneration and regulates muscle function^{2–6}. The deficiency of laminin $\alpha 2$ chain leads to a severe form of congenital muscular dystrophy (type 1A, referred to as MDC1A)^{7–9}. Moreover, complete deficiency of laminin $\alpha 2$ chain in mice (*dy^{3K}/dy^{3K}* animals used in this study) results in the most severe phenotype among mouse models lacking proteins of cell adhesion complexes^{2,10–12}.

Integrin $\alpha 7\beta 1$ is one of the major cell surface receptors that binds laminin-211 in the musculoskeletal system^{13,14}. Integrin $\alpha 7\beta 1$ is localized both at the sarcolemma and enriched at the neuromuscular and myotendinous junctions¹⁵. In addition to providing anchorage and structural framework integrin $\alpha 7\beta 1$ mediates signalling transduction cues^{16–18}. This dual role has been suggested to be an engine for numerous processes: muscle development (myoblast migration, fusion, differentiation *in vitro*), survival, growth, regeneration and force generation^{5,15,18–22}. Yet, skeletal muscle develops normally in integrin $\alpha 7$ -null mice. Additionally, these mice are presented with a mild form of muscular dystrophy that mostly affects myotendinous junction^{23,24}. Similarly, the loss of integrin $\alpha 7$ subunit in humans causes an extremely rare form of congenital myopathy^{25,26}.

Another protein complex that interacts with laminin-211 and comprises the molecular link between extracellular matrix and cytoskeleton is the dystrophin-glycoprotein complex (DGC). Numerous research investigations have addressed the role of DGC in preservation of the structural and functional integrity of skeletal muscle^{5,9,27,28}.

Department of Experimental Medical Science, Muscle Biology Unit, Lund University, Sweden. Correspondence and requests for materials should be addressed to K.I.G. (email: kinga.gawlik@med.lu.se)

In spite of the relatively mild phenotype of integrin $\alpha 7$ -deficient muscle, several studies in mice and humans indicated that integrin $\alpha 7$ subunit could play a significant role in modifying the disease progression in muscular dystrophies involving the absence of DGC components (dystrophin- and γ -sarcoglycan-deficient muscular dystrophies)^{29–34}. These studies suggested that integrin $\alpha 7\beta 1$ and the DGC could have redundant functions. Yet, the separate roles for these two complexes are also evident^{5,35}. Even the interplay between laminin $\alpha 2$ chain and integrin $\alpha 7\beta 1$ seems to be complex and its role in skeletal muscle has not been fully deciphered. Reduced expression of integrin $\alpha 7$ subunit^{20,29,36} and its aberrant deposition at the sarcolemma³⁷ accompany the laminin $\alpha 2$ chain loss. Consequently, over-expression of integrin $\alpha 7$ subunit in laminin $\alpha 2$ chain-deficient mice improves muscle pathology³⁸. Nonetheless, the phenotypes of integrin $\alpha 7$ and laminin $\alpha 2$ chain-deficient mice and patients are markedly different^{9,11,23}.

We have recently demonstrated that the absence of laminin $\alpha 2$ chain in dystrophin or β -sarcoglycan deficiency drastically aggravates the phenotype of *mdx* and *Sgcb* mice, respectively¹². Our results pointed out that laminin $\alpha 2$ chain, dystrophin, and β -sarcoglycan have complementary, but non-redundant roles in spite of being part of the same adhesion complex. In the current study we generated laminin $\alpha 2$ chain-integrin $\alpha 7$ chain double knockout mice (*dy^{3K}/itga7*) in order to establish the relationship between these molecules independently of the DGC and to determine if the main role of integrin $\alpha 7\beta 1$ is to mediate laminin-211 interactions in muscle.

Results

Expression profile of integrins in laminin $\alpha 2$ chain- and integrin $\alpha 7$ -deficient muscles.

Dy^{3K}/dy^{3K} mice exhibit very severe muscular dystrophy^{10,12}. In order to determine if the deletion of integrin $\alpha 7$ subunit exacerbates the phenotype of laminin $\alpha 2$ chain-deficient dystrophic mice, we generated mice lacking both molecules (*dy^{3K}/itga7*) (by series of heterozygous breeding, see materials and methods). The genotype was confirmed by PCR (data not shown) and immunofluorescence (Fig. 1). In accordance with previous studies, laminin $\alpha 2$ chain was normally expressed in the absence of integrin $\alpha 7$ subunit³⁶ (Fig. 1). Integrin $\alpha 7$ chain, on the other hand, was reduced in the *dy^{3K}/dy^{3K}* laminin $\alpha 2$ chain-deficient muscle³⁷ (Fig. 1). Yet, compared with *itga7* knockout mice and double knockout muscles, integrin $\alpha 7$ subunit was not completely absent from the sarcolemma of laminin $\alpha 2$ chain-deficient mice and it was also present in vessels (Fig. 1). Furthermore, the expression of integrin $\alpha 7$ subunit in younger (2-week-old) *dy^{3K}/dy^{3K}* muscles was maintained at the sarcolemma (Fig. 1). Taken together, these results clearly distinguish *dy^{3K}/dy^{3K}* muscles from *dy^{3K}/itga7* double knockout muscles in terms of integrin $\alpha 7$ chain expression. Additionally, integrin $\beta 1$ subunit, which forms a dimer with integrin $\alpha 7$ chain in healthy muscle, was expressed in all three knockout models³⁷ (Fig. 1). Hence, we investigated the expression of integrin α subunits that could potentially dimerize with integrin $\beta 1$ chain at the sarcolemma. Integrin $\alpha 6\beta 1$ is a laminin-211 receptor and integrin $\alpha 6$ subunit has been demonstrated to be upregulated in mice with partial deficiency of laminin $\alpha 2$ chain (*dy/dy* mouse model)³⁹. Integrin $\alpha 5$ chain is another candidate to form a dimer with integrin $\beta 1$ subunit in muscle, as it has been shown to contribute to maintenance of normal muscle morphology⁴⁰. We found moderate upregulation of integrin $\alpha 6$ subunit in *dy^{3K}/dy^{3K}* and *dy^{3K}/itga7* muscles compared to wild-type and *itga7* muscles (Fig. 1). Integrin $\alpha 6$ was mostly present in smaller regenerating fibers (Fig. 1, arrows), but it was also occasionally found in a few bigger muscle fibers, though its expression was rather weak and not continuous throughout the sarcolemma (Fig. 1, arrowheads). Integrin $\alpha 5$ chain was expressed only in large vessels and in streaks of interstitial connective tissue in wild-type and integrin $\alpha 7$ -null muscle (Fig. 1). It was not deposited at the sarcolemma of *dy^{3K}/dy^{3K}* and *dy^{3K}/itga7* muscles, but instead it was massively upregulated at the interstitial space within muscle fibers (Fig. 1), suggesting a role for integrin $\alpha 5$ in inflammation and fibrosis.

Integrin $\alpha 7\beta 1$ binds also laminin $\alpha 4$ and $\alpha 5$ chains¹⁴. Those laminin chains are upregulated upon laminin $\alpha 2$ chain loss in skeletal muscle⁴¹. Accordingly, we detected their upregulation in the extrasynaptic basement membranes in *dy^{3K}/dy^{3K}* and *dy^{3K}/itga7* muscles (Supplementary Fig. S1). Additionally, we found laminin $\alpha 4$ and laminin $\alpha 5$ chains to be weakly expressed at the myotendinous junction (MTJ) in wild-type mice. Their expression was preserved (laminin $\alpha 5$) or increased (laminin $\alpha 4$) at this site in all three analysed mutants, which could partially explain maintained expression of integrin $\alpha 7\beta 1$ at the MTJ in *dy^{3K}/dy^{3K}* mice and integrin $\beta 1$ presence in double knockout mice (Supplementary Fig. S1). Integrin $\alpha 5$ and $\alpha 6$ have also been found to be moderately expressed at the MTJ in all three knockout mice (Supplementary Fig. S1).

Taken together, the expression pattern of integrin $\alpha 5\beta 1$, $\alpha 6\beta 1$, laminin $\alpha 4$ and $\alpha 5$ chains is the same between *dy^{3K}/dy^{3K}* and *dy^{3K}/itga7* muscle, indicating no additional compensatory mechanisms in double knockout mice.

Overall phenotype of *dy^{3K}/itga7* double knockout mice is not aggravated compared with *dy^{3K}/dy^{3K}* single knockouts.

Approximately 60% of *itga7* null-mice are embryonic lethal due to cerebrovascular defects⁴². Thus, the number of *dy^{3K}/itga7* mice used in this study was limited. Integrin $\alpha 7$ -deficient animals that survive embryonic development have near normal life span, weight and are fertile. Double knockout mice that were born did not die earlier than *dy^{3K}/dy^{3K}* mice and both mutants lived up to 7–8 weeks after birth (Fig. 2b) (*dy^{3K}/dy^{3K}* mice usually die at around 3–4 weeks of age, but we observed increased survival on the B6;129-*Itga7^{tm1Burk/J}* background). Likewise, the overall health of




Figure 1. Immunostaining with antibodies against laminin $\alpha 2$ chain (green, LM $\alpha 2$) and integrin $\alpha 7$ chain (red, Itg $\alpha 7$) confirms the complete absence of both proteins in *dy*^{3K}/*itga7* double knockout muscles. Expression of integrin $\alpha 7$ subunit is reduced at the sarcolemma of 5-week-old *dy*^{3K}/*dy*^{3K} laminin $\alpha 2$ chain-deficient muscles. However, integrin $\alpha 7$ chain is present at the sarcolemma of the majority of 2-week-old *dy*^{3K}/*dy*^{3K} muscle fibers. White arrowheads show maintained expression of integrin $\alpha 7$ in vessels. Laminin $\alpha 2$ chain is not reduced in integrin $\alpha 7$ knockout muscles. Expression of integrin $\beta 1D$ subunit (Itg $\beta 1D$, green) is maintained in muscles from all three mutants. Integrin $\alpha 6$ chain (Itg $\alpha 6$, green) is expressed in big and small vessels and peripheral nerves (white star), but not at the sarcolemma of normal muscle and *itga7*-null muscle. In addition to normal expression in vessels and nerves, integrin $\alpha 6$ subunit is present in muscle cell membranes in small regenerating *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* muscle fibers (white arrows) and weakly expressed at the sarcolemma in some bigger muscle fibers (white arrowheads). Sections were co-stained with the antibody against dystrophin (dystr, red) and DAPI (blue). Integrin $\alpha 5$ subunit (Itg $\alpha 5$, red) was massively upregulated in fibrotic lesions and/or sites of inflammation in laminin $\alpha 2$ chain-null mice and laminin $\alpha 2$ chain/integrin $\alpha 7$ double knockout animals (white asterisk), but was not found to be deposited at the sarcolemma. In normal muscle and in integrin $\alpha 7$ -deficient muscle integrin $\alpha 5$ chain is expressed only in bigger vessels (pink arrowheads) and in interstitial connective tissue (pink arrows). Muscle sections were co-stained with dystrophin antibody (dystr, green). Scale bars, 40 μ m.




Figure 2. General health of laminin $\alpha 2$ chain/integrin $\alpha 7$ -deficient mice is not worsened compared to severely affected laminin $\alpha 2$ chain single knockout animals. (a) Photos of 4-week-old dy^{3K}/dy^{3K} and $dy^{3K}/itga7$ mice (left picture) with a normal littermate (right picture). Both mutants are severely emaciated and display dystrophic phenotype, with severe loss of muscle tissue. (b) Lifespan of $dy^{3K}/itga7$ is not shortened compared to dy^{3K}/dy^{3K} mice (survival curves are not significantly different, $p = 0.2331$, log-rank Cox-Mantel test). Note that much fewer double knockout mice were taken under consideration due to partial embryonic lethality of $itga7$ mice. (c) The whole body weight is not significantly different between dy^{3K}/dy^{3K} and $dy^{3K}/itga7$ mice ($p = 0.3175$, Mann-Whitney test). The numbers of animals used are indicated in the graph.

$dy^{3K}/itga7$ mice was not worsened compared to dy^{3K}/dy^{3K} animals. They displayed comparable growth retardation, muscle wasting, tremor, scoliosis and reduced eagerness to move (Fig. 2a). Accordingly, the weight of $dy^{3K}/itga7$ mice was not significantly different from single knockout dy^{3K}/dy^{3K} animals (Fig. 2c).

Dystrophic features of laminin $\alpha 2$ chain-deficient muscle are not dramatically changed upon integrin $\alpha 7$ deletion. Laminin $\alpha 2$ chain-deficient muscle displays multiple defects. Cell death and degeneration/regeneration cycles are concomitant with massive infiltration of inflammatory cells and subsequent robust production of extracellular matrix components² (Fig. 3a). These pathogenic processes result in severe muscle wasting and loss of muscle function. Integrin $\alpha 7$ -null mice, on the other hand, develop mild myopathy with defects concentrated around the myotendinous junction – with its disrupted structure and function as well as with myofiber necrosis, muscle regeneration and inflammation around the myotendinous junctions^{23,24} (Fig. 3a).

Histological analysis of limb muscle sections from 5–7-week-old double knockout $dy^{3K}/itga7$ mice did not reveal additional changes in muscle morphology compared to the severe defects seen in dy^{3K}/dy^{3K} muscles (Fig. 3a). Muscle fascicles were equally severely disrupted in both mutants, with large areas of focal apoptosis/necrosis and mononuclear cells infiltrates (Fig. 3a). Regenerating fibers with centrally located nuclei and loose atrophic fibers were also present. Nevertheless, quantification of centrally nucleated muscle cells showed decreased number of regenerating fibers in $dy^{3K}/itga7$ muscles (triceps and quadriceps) compared to dy^{3K}/dy^{3K} muscles (Fig. 3b). We next estimated the regenerative capacity of $dy^{3K}/itga7$ muscle by immunostaining using the antibody against the embryonic myosin heavy chain. Newly regenerating fibers expressing embryonic form of myosin heavy chain were present in both dy^{3K}/dy^{3K} and $dy^{3K}/itga7$ muscles (Fig. 3c), indicating that muscle regeneration is not exhausted in double knockout mice. Yet, the regenerative capacity could still be altered/slowed down in the absence of integrin $\alpha 7$ subunit, as indicated by decreased numbers of regenerating cells.

A general examination of the $dy^{3K}/itga7$ diaphragm did not reveal a dramatic aggravation of the phenotype either (Fig. 4a). Both dy^{3K}/dy^{3K} and $dy^{3K}/itga7$ diaphragm showed dystrophic features (Fig. 4a, hematoxylin&eosin staining) and rich collagen deposition (Fig. 4a, Sirius red/fast green staining). As it has been shown before, heart muscle from dy^{3K}/dy^{3K} mice was not affected³⁵ (Fig. 4b), despite laminin $\alpha 2$ chain being abundantly deposited in heart basement membranes. Integrin $\alpha 7$ subunit is also strongly




Figure 3. Muscular dystrophy hallmarks are not exacerbated in limb muscles from *dy*^{3K}/*itga7* mice. (a) Hematoxylin and eosin staining of quadriceps and triceps muscles from 5–7-week-old mice shows mild myopathy (white arrows) at the myotendinous junctions (blue asterisk) in *itga7*^{-/-} mice and severe muscular dystrophy with robust inflammation (orange arrows), muscle damage (single degenerating fibers or areas with a group of damaged fibers/fiber debris are indicated with green arrow or green star, respectively), muscle regeneration (fibers with centrally located nucleus) and connective tissue build-up (black arrows) in *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* animals. (b) Quantification of centrally nucleated fibers shows significant decrease in number of regenerating fibers in *dy*^{3K}/*itga7* quadriceps (vastus intermedius) and triceps compared with *dy*^{3K}/*dy*^{3K} corresponding muscles ($p = 0.0317$; $p = 0.0159$, respectively; Mann-Whitney test). The numbers of animals used are indicated in the graph. (c) Embryonic myosin heavy chain staining (eMyoH, green) reveals ongoing regeneration in both *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* muscles. Sections were costained with DAPI (blue) and laminin $\gamma 1$ (LM $\gamma 1$, red) to visualize nuclei and delineate muscle fibers, respectively. Wild-type muscle treated with cardiotoxin (CTX) is shown as a positive control for embryonic myosin heavy chain staining. Scale bars, 30 μm .

expressed in heart⁴³. However, in addition to laminin $\alpha 2$ chain loss, the absence of integrin $\alpha 7$ subunit in *dy*^{3K}/*itga7* heart did not result in appearance of cardiomyopathy features and visibly increased collagen content (Fig. 4b).

Although the dystrophic phenotype of muscles from double knockout mice is not drastically worsened compared to *dy*^{3K}/*dy*^{3K} muscles (Fig. 3a), slightly impaired/slower regeneration (Fig. 3b) could result in more rapid loss of muscle fibers and in consequence, give rise to increased fibrotic build-up. Thus,




Figure 4. No differences in the phenotype of diaphragm and heart between *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* mice. (a) Hematoxylin and eosin staining (H&E) and Sirius red/fast green staining (SR-FG) demonstrate dystrophic features in diaphragm from *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* mice. Loss of muscle fibers, inflammation and connective tissue build up are evident (H&E) in these mutants. Presence of fibrosis was confirmed by Sirius red staining (collagen deposition, dark pink). No dystrophic changes were observed in *itga7*^{-/-} diaphragm. (b) Hematoxylin and eosin staining (H&E) and Sirius red/fast green staining (SR-FG) do not show cardiomyopathy features in hearts from single and double knockout mice. Scale bars, 50 μm.

we analysed the expression of collagen III and fibronectin, two major components of fibrotic lesions, in triceps and quadriceps muscle (Fig. 5). Integrin $\alpha 7$ -deficient muscles did not exhibit fibrotic changes, whereas *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* muscles showed increased collagen III and fibronectin production (Fig. 5a). The quantitative comparison of areas affected with collagen III and fibronectin deposition in *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* muscles (triceps and quadriceps) did not demonstrate a significant difference in fibronectin and collagen content between the genotypes (Fig. 5b). Yet, a trend for marginally enhanced fibrotic tissue build-up in double knockout triceps was observed (Fig. 5b, $p = 0.0635$), which could indeed be matched with moderate regeneration impairment in *dy*^{3K}/*itga7* muscles. Nevertheless, it does not change the fact that muscles from both laminin $\alpha 2$ chain-null mice and double knockout mice are severely dystrophic and do not differ substantially between each other. This was also confirmed by analyses of apoptosis and inflammation (Fig. 6). Apoptosis is a hallmark of laminin $\alpha 2$ chain-deficient muscle and integrin $\alpha 7\beta 1$ signalling is involved in maintenance of muscle survival^{6,10,16,20}. We analysed apoptosis in both single and double knockout muscles (Fig. 6a). No increase in apoptosis was detected




Figure 5. Fibrotic lesions are equally abundant in muscles from *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* mice.

(a) Immunofluorescence with antibodies against collagen III (coll III) and fibronectin (FN) demonstrate extensive production of fibrotic proteins in *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* muscles. (b) Fibronectin and collagen III deposition was not statistically different between these two groups in triceps ($p = 0.0635$ and $p = 0.0635$, respectively; Mann-Whitney) and quadriceps (vastus intermedius) ($p = 0.1905$ and $p = 0.1905$, respectively; Mann-Whitney). Yet, the p values for triceps muscle could indicate a trend for slightly larger areas of collagen III and fibronectin in *dy*^{3K}/*itga7* triceps in comparison with *dy*^{3K}/*dy*^{3K} specimens. The numbers of animals used are indicated in the graph. Scale bar, 300 μ m.

in double knockout muscles compared to *dy*^{3K}/*dy*^{3K} muscles as indicated by caspase-3 immunostaining and quantification of caspase-3 positive fibers ($p = 0.6828$) (Fig. 6a,c). Integrin α 7-deficient mice did not display apoptotic muscle cells (Fig. 6a).

Since inflammation is a feature of laminin α 2 chain-deficient muscular dystrophy^{12,44–46}, we assessed the inflammatory response in muscles from all three mouse models used in this study (Fig. 6b). CD11b immunostaining depicting macrophages revealed equally strong inflammatory response in both *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* muscles ($p = 0.7143$) (Fig. 6b,d) and showed almost no inflammation in *itga7*-null muscles (Fig. 6b).

Discussion

Although numerous studies with muscular dystrophy patients and with genetically modified dystrophic animals have been performed, the molecular puzzle of relationships between different adhesion complexes in skeletal muscle has not been entirely deciphered. Laminin α 2 chain is a bridging element between two adhesion complexes – integrin α 7 β 1 and the DGC. The DGC remains intact in laminin α 2 chain-deficient muscle and we have recently demonstrated non-redundant functions of laminin α 2 chain and the DGC components dystrophin and β -sarcoglycan, as loss of these molecules significantly exacerbated the phenotype of *dy*^{3K}/*dy*^{3K} mice¹². Integrin α 7 chain expression, on the other hand, is secondarily




Figure 6. Muscles from *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* are equally affected with apoptosis and inflammation. (a) Caspase-3 immunostaining (green) reveals spread apoptotic fibers or group of dying cells in both *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* muscles. Collagen IV antibody (red staining) and DAPI (blue) were used to delineate muscle fibers and show nuclei. (b) CD11b staining (red) showed infiltration of macrophages into *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* dystrophic triceps muscle. DAPI (blue) depicts cell nuclei. (c) Percentage of apoptotic fibers was not increased in double knockout triceps muscle compared to *dy*^{3K}/*dy*^{3K} triceps ($p = 0.6828$, Mann-Whitney test). (d) Amount of macrophages was not significantly different between *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* triceps ($p = 0.7143$, Mann-Whitney test). The numbers of animals used are indicated in the graph. Scale bar, 50 μm .

reduced in MDC1A^{20,36,37} and therefore we reasoned that mice deficient in both laminin $\alpha 2$ chain and integrin $\alpha 7$ chain should have a similar phenotype as laminin $\alpha 2$ chain-null mice¹². In the current study we test this hypothesis by deleting the integrin $\alpha 7$ subunit in laminin $\alpha 2$ chain-deficient *dy*^{3K}/*dy*^{3K} mice.

In general, the phenotype of double knockout mice did not exceed the severity of laminin $\alpha 2$ chain single knockout mice, which represent one of the most severe muscular dystrophy mouse models¹². The fact that mice deficient in both laminin $\alpha 2$ chain and β -sarcoglycan or dystrophin (*dy*^{3K}/*Sgcb* and *dy*^{3K}/*mdx*, respectively) exhibited a dramatically deteriorated dystrophic phenotype¹² indicates that integrin $\alpha 7\beta 1$ -laminin-211 axis could have an inferior role to the DGC-laminin-211 association in skeletal muscle. On the other hand, mice lacking integrin $\alpha 7$ subunit and the DGC components showed a very severe phenotype^{32,33,34} (albeit less severe than *dy*^{3K}/*mdx* and *dy*^{3K}/*Sgcb* mice¹²), implying that integrin $\alpha 7\beta 1$ also contributes to the functional integrity of skeletal muscle. Consequently, we have demonstrated that truncated laminin that cannot bind dystroglycan but binds integrin $\alpha 7\beta 1$ is sufficient to adequately maintain muscle function³⁵. It could be that laminin $\alpha 2$ chain binding to integrin $\alpha 7\beta 1$ has a more important role in signalling than in providing structural support to muscle cells⁵. However, the signalling cascades involving integrin $\alpha 7\beta 1$ that are fundamental for muscle function are yet to be identified and this remains an important task.

We also sought to elucidate if compensatory upregulation of other α integrin subunits could be an alternative reason behind the not substantially aggravated phenotype of double knockout mice. Overlapping functions of different $\beta 1$ integrins have been indicated in numerous studies¹⁵. Accordingly, integrin $\beta 1$ subunit is not absent from cell membranes in laminin $\alpha 2$ chain-null and integrin $\alpha 7$ /laminin $\alpha 2$ chain-deficient muscle (Fig. 1 and³⁷). However, the undistinguishable expression pattern of integrin $\alpha 6$ and integrin $\alpha 5$ subunits in *dy*^{3K}/*dy*^{3K} and *dy*^{3K}/*itga7* muscles excludes the possibility that integrin $\alpha 6\beta 1$ and/or integrin $\alpha 5\beta 1$ prevent further deterioration of dystrophic phenotype in double knockout animals, even if integrin $\alpha 5$ subunit has been shown to protect muscle from damage⁴⁰. What is more,

neither of those integrin complexes seems to be involved in inhibiting the dystrophic phenotype of single knockout dy^{3K}/dy^{3K} mice. Contrary, the upregulation of integrin $\alpha 5$ chain, a major fibronectin receptor expressed in fibroblasts, has been shown to impact various fibrotic conditions^{47–51} and inflammation^{52,53}. It is not excluded that upregulation of integrin $\alpha 5$ in laminin $\alpha 2$ chain-deficient muscles promotes fibrosis, as integrin $\alpha 5$ subunit does not bind laminin-211, but interacts with fibronectin.

Although the phenotype of double knockout $dy^{3K}/itga7$ muscles is to large extent the same as the phenotype of dy^{3K}/dy^{3K} muscles, a more detailed analysis revealed a slight exacerbation of muscle defects, such as somewhat impaired regeneration and tendency for marginally increased fibrosis. These results match the data published by Doe *et al.*, where the overexpression of integrin $\alpha 7$ subunit has been shown to moderately improve the phenotype of dy^W/dy^W mice³⁸. On the other hand, reduced inflammation has also been suggested to be controlled by integrin $\alpha 7$ overexpression in dy^W/dy^W muscle³⁸. Yet, the deletion of integrin $\alpha 7$ subunit in dy^{3K}/dy^{3K} mice did not exacerbate the inflammatory response, pointing toward no significant role for integrin $\alpha 7\beta 1$ in inflammation regulation, at least in dy^{3K}/dy^{3K} muscle. Collectively, results obtained in Burkin's laboratory and our data suggest that the upregulation of integrin $\alpha 7$ subunit in laminin $\alpha 2$ chain-deficiency could only affect the muscle phenotype to some extent, most likely influencing muscle regeneration and perhaps muscle survival²¹. Indeed, there is a line of evidence for the role of integrin $\alpha 7\beta 1$ in muscle regeneration and satellite cell activation/myogenic function^{22,31,33,54,55}. Besides MDC1A, the integrin $\alpha 7\beta 1$ upregulation approach has also been tested for Duchenne muscular dystrophy. Both transgenic overexpression and AVV-mediated delivery of integrin $\alpha 7\beta 1$ have shown rather good effects in dystrophin/utrophin-deficient mice (*mdx/utr*) and dystrophin-null mice (*mdx*), respectively^{30,31,56,57}. Thus, targeting integrin $\alpha 7\beta 1$ expression may show more promise for treatment of Duchenne muscular dystrophy.

In summary, our previous results indicate that DGC and laminin $\alpha 2$ chain have complementary but non-redundant functions in skeletal muscle while the current study shows that integrin $\alpha 7\beta 1$ does not play any other roles in skeletal muscle than mediating the laminin $\alpha 2$ chain interaction. Moreover, integrin $\alpha 7\beta 1$ probably does not bind any additional vital ligands. Laminin $\alpha 2$ chain, on the other hand, may interact with other receptors than dystroglycan and integrin $\alpha 7\beta 1$. Potential receptors may include integrin $\alpha 9\beta 1$, which is expressed in skeletal muscle⁵⁸ and has been shown to bind laminin-111⁵⁹. Also, proteins that were originally identified in Schwann cells as laminin-binding molecules (galactosyl-sulfatides⁶⁰ and adhesion G protein-coupled receptors⁶¹) may serve as laminin $\alpha 2$ chain binding receptors in skeletal muscle. In order to further understand the molecular pathology of MDC1A and other muscular dystrophies it will be crucial to validate whether laminin-211 interacts with additional receptors in striated muscle.

Methods

Mouse models, double knockout mice generation. Laminin $\alpha 2$ chain-null dy^{3K}/dy^{3K} mice were previously described¹⁰. *Itga7/+* mice (B6;129-*Itga7*^{tm1Burk/J})⁴² were obtained from Jackson laboratory. $Dy^{3K}/+$ males or females were bred with *itga7/+* females or males, respectively. The resulting $dy^{3K}/+; itga7/+$ mice were mated to generate wild-type, *itga7* knockout, dy^{3K}/dy^{3K} and $dy^{3K}/itga7$ double knockout mice. Control animals were: wild-type or $dy^{3K}/+$. Mice were maintained in the animal facilities of Biomedical Center (Lund) according to the animal care guidelines. All experimental procedures involving animals were approved by the Malmö/Lund (Sweden) Ethical Committee for Animal Research (the ethical permit number: M15-12 and M152-14) in accordance with the guidelines approved by the Swedish Board of Agriculture.

Histology and morphometric analysis. Quadriceps femoris, triceps brachii, diaphragm and heart muscles were isolated from 5–7-week-old mice (single knockouts, double knockouts and control animals), embedded in OCT and frozen rapidly in liquid nitrogen. Cryosections (7 μ m) were stained with hematoxylin and eosin¹² or picosirius red/fast green⁴⁴. Stained cross-sections were scanned using Aperio's Scanscope CS2 (with Scanscope console v.8.2.0.1263) and images were created using Aperio software.

Centrally nucleated fibers representing regenerating muscle cells and peripherally nucleated normal muscle cells were counted in triceps brachii and vastus intermedius (quadriceps femoris) from dy^{3K}/dy^{3K} and $dy^{3K}/itga7$ mice using ImageJ software version 143u (NIH, Bethesda, MD). A whole area of each muscle cross section (from both legs) was considered. Percentage of centrally nucleated fibers in each muscle was calculated and averaged between two collateral muscles from the same animal. Mann-Whitney test was used for statistical analysis ($p < 0.05$).

Immunofluorescence. Cryosections were subjected to immunofluorescence labeling¹² with antibodies against: laminin $\alpha 2$ chain (rat monoclonal 4H8-2, 1:100, Alexis Biochemicals), integrin $\alpha 7\beta$ subunit (rabbit polyclonal U31, 1:300, kindly provided by Dr. U. Mayer), integrin $\beta 1D$ subunit (mouse monoclonal 2B1, 1:80, Millipore), integrin $\alpha 5$ subunit (rat monoclonal 5H10-27 phycoerythrin-conjugated, 1:100, Abcam), integrin $\alpha 6$ subunit (rat monoclonal GoH3, 1:200, Abcam), CD11b (rat monoclonal M1/70, 1:250, BD Pharmingen), collagen III (goat polyclonal, 1:100, Southern Biotech), fibronectin (rabbit polyclonal, 1:1000, Abcam), embryonic myosin heavy chain (mouse monoclonal F1.652, 1:10; Developmental Studies Hybridoma Bank), caspase-3 (mouse monoclonal 46, 1:100, BD Transduction Laboratories), collagen IV (rabbit polyclonal, 1:100, Millipore), dystrophin (rabbit polyclonal, 1:100, Abcam), laminin $\gamma 1$,

laminin $\alpha 4$ and laminin $\alpha 5$ chain (all rabbit polyclonal, 1:100, kindly provided by Dr. T. Sasaki). Primary antibodies were detected with proper secondary antibodies (Molecular Probes). Stained cross-sections were analysed using a Zeiss Axioplan fluorescence microscope. Images were taken using with an ORCA 1394 ER digital camera and Openlab 4 software, at the same exposure times for all genotypes. The area corresponding to CD11b, collagen III and fibronectin labeling was quantified in relation to the entire area of vastus intermedius (quadriceps femoris) and/or triceps brachii cross-sections from dy^{3K}/dy^{3K} and $dy^{3K}/itga7$ mice (for each animal the average percentage of stained area was calculated for muscles from collateral legs). ImageJ software version 143u (NIH, Bethesda, MD) was used. Mann-Whitney test was used for statistical analysis ($p < 0.05$). For collagen III and fibronectin staining, myotendinous junctions were excluded from quantification.

Caspase-3 positive fibers and normal muscle fibers were counted in triceps from dy^{3K}/dy^{3K} and $dy^{3K}/itga7$ mice using ImageJ software version 143u (NIH, Bethesda, MD). Percentage of caspase-3 positive fibers was calculated. Mann-Whitney test was used for statistical analysis ($p < 0.05$).

Statistical analysis. All statistical analyses were performed with GraphPad Prism software version 6 (La Jolla, CA). Averaged data were reported as means \pm SEM. Mann-Whitney test was used. Statistical significance was accepted for $p < 0.05$.

References

- Carmignac, V. & Durbeej, M. Cell-matrix interactions in muscle disease. *J Pathol* **226**, 200–218 (2012).
- Gawlik, K. I. & Durbeej, M. Skeletal muscle laminin and MDC1A: pathogenesis and treatment strategies. *Skelet Muscle* **1**, 9 (2011).
- Holmberg, J. & Durbeej, M. Laminin-211 in skeletal muscle function. *Cell Adh Migr* **7**, 111–121 (2013).
- Kuang, W., Xu, H., Vilquin, J. T. & Engvall, E. Activation of the lama2 gene in muscle regeneration: abortive regeneration in laminin alpha2-deficiency. *Lab Invest* **79**, 1601–1613 (1999).
- Han, R. *et al.* Basal lamina strengthens cell membrane integrity via the laminin G domain-binding motif of α -dystroglycan. *Proc Natl Acad Sci USA* **106**, 12573–12579 (2009).
- Vachon, P. H., Loechel, F., Xu, H., Wewer, U. M. & Engvall, E. Merosin and laminin in myogenesis; specific requirements for merosin in myotubal stability and survival. *J Cell Biol* **134**, 1483–1497 (1996).
- Helbling-Leclerc, A. *et al.* Mutations in the laminin $\alpha 2$ chain gene (LAMA2) cause merosin-deficient muscular dystrophy. *Nat Genet* **11**, 216–218 (1995).
- Allamand, V. & Guicheney, P. Merosin-deficient muscular dystrophy, autosomal recessive (MDC1A, MIM#156225, LAMA2 gene coding for $\alpha 2$ chain of laminin). *Eur J Hum Genet* **10**, 91–94 (2002).
- Voit, T. & Tomé, F. S. The congenital muscular dystrophies in *Myology*. (ed Engel, A., Franzini-Armstrong, C.) 1203–1238 (McGraw-Hill Inc, 2004).
- Miyagoe, Y. *et al.* Laminin $\alpha 2$ chain-null mutant mice by targeted disruption of the Lama2 gene: a new model of merosin (laminin 2)-deficient congenital muscular dystrophy. *FEBS Lett* **415**, 33–39 (1997).
- Guo, L. T. *et al.* Laminin alpha2 deficiency and muscular dystrophy; genotype-phenotype correlation in mutant mice. *Neuromuscul Disord* **13**, 207–215 (2003).
- Gawlik, K. I., Holmberg, J. & Durbeej, M. Loss of Dystrophin and β -Sarcoglycan, Respectively, Significantly Exacerbates the Phenotype of Laminin $\alpha 2$ Chain-Deficient Animals. *Am J Pathol* **184**, 740–752 (2014).
- Song, W. K., Wang, W., Foster, R. E., Bielser, D. A. & Kaufman, S. J. H36-alpha 7 is a novel integrin alpha chain that is developmentally regulated during skeletal myogenesis. *J Cell Biol* **117**, 643–657 (1992).
- von der Mark, H. *et al.* Alternative splice variants of $\alpha 7\beta 1$ integrin selectively recognize different laminin isoforms. *J Biol Chem* **277**, 6012–6016 (2002).
- Mayer, U. Integrins: redundant or important players in skeletal muscle? *J Biol Chem* **278**, 14587–14590 (2003).
- Laprise, P., Poirier, E. M., Vézina, A., Rivard, N. & Vachon, P. H. Merosin-integrin promotion of skeletal myofiber cell survival: Differentiation state-distinct involvement of p60Fyn tyrosine kinase and p38alpha stress-activated MAP kinase. *J Cell Physiol* **191**, 69–81 (2002).
- Boppert, M. D., Burkin, D. J. & Kaufman, S. J. Alpha7beta1-integrin regulates mechanotransduction and prevents skeletal muscle injury. *Am J Physiol Cell Physiol* **290**, C1660–1665 (2006).
- Boppert, M. D., Burkin, D. J. & Kaufman, S. J. Activation of AKT signaling promotes cell growth and survival in $\alpha 7\beta 1$ integrin-mediated alleviation of muscular dystrophy. *Biochim Biophys Acta* **1812**, 439–446 (2011).
- Burkin, D. J. & Kaufman, S. J. The alpha7beta1 integrin in muscle development and disease. *Cell Tissue Res* **296**, 183–190 (1999).
- Vachon, P. H. *et al.* Integrins ($\alpha 7\beta 1$) in muscle function and survival. Disrupted expression in merosin-deficient congenital muscular dystrophy. *J Clin Invest* **11**, 1870–1881 (1997).
- Liu, J., Burkin, D. J. & Kaufman, S. J. Increasing alpha 7 beta 1-integrin promotes muscle cell proliferation, adhesion, and resistance to apoptosis without changing gene expression. *Am J Physiol Cell Physiol* **294**, C627–640 (2008).
- Rooney, J. E., Gurpur, P. B., Yablonka-Reuveni, Z. & Burkin, D. J. Laminin-111 restores regenerative capacity in a mouse model for $\alpha 7$ integrin congenital myopathy. *Am J Pathol* **174**, 256–264 (2009).
- Mayer, U. *et al.* Absence of integrin $\alpha 7$ causes a novel form of muscular dystrophy. *Nat Genet* **17**, 18–23 (1997).
- Nawrotzki, R., Willem, M., Miosge, N., Brinkmeier, H. & Mayer, U. Defective integrin switch and matrix composition at alpha 7-deficient myotendinous junctions precede the onset of muscular dystrophy in mice. *Hum Mol Genet* **12**, 483–495 (2003).
- Hayashi, Y. K., Chou, F. L., Engvall, E., Ogawa, M. & Matsuda, C. Mutations in the integrin alpha7 gene cause congenital myopathy. *Nat Genet* **19**, 94–97 (1998).
- Pegoraro, E., Cepollaro, F., Prandini, P., Marin, A. & Fanin, M. Integrin alpha 7 beta 1 in muscular dystrophy/myopathy of unknown etiology. *Am J Pathol* **160**, 2135–2143 (2002).
- Cohn, R. D. & Campbell, K. P. Molecular basis of muscular dystrophies. *Muscle Nerve* **23**, 1456–1471 (2000).
- Durbeej, M. & Campbell, K. P. Muscular dystrophies involving the dystrophin-glycoprotein complex: an overview of current mouse models. *Curr Opin Genet Dev* **12**, 349–361 (2002).
- Hodges, B. L. *et al.* Altered expression of the $\alpha 7\beta 1$ integrin in human and murine muscular dystrophies. *J Cell Sci* **110**, 2873–2881 (1997).
- Burkin, D. J., Wallace, G. Q., Nicol, K. J., Kaufman, D. J. & Kaufman, S. J. Enhanced expression of the $\alpha 7\beta 1$ integrin reduces muscular dystrophy and restores viability in dystrophic mice. *J Cell Biol* **152**, 1207–1218 (2001).

31. Burkin, D. J. *et al.* Transgenic expression of $\alpha 7\beta 1$ integrin maintains muscle integrity, increases regenerative capacity, promotes hypertrophy, and reduces cardiomyopathy in dystrophic mice. *Am J Pathol* **166**, 253–263 (2005).
32. Allikian, M., Hack, A. A., Mewborn, S., Mayer, U. & McNally, E. M. Genetic compensation for sarcoglycan loss by integrin $\alpha 7\beta 1$ in muscle. *J Cell Sci* **117**, 3821–3830 (2004).
33. Guo, C. *et al.* Absence of $\alpha 7$ integrin in dystrophin-deficient mice causes a myopathy similar to Duchenne muscular dystrophy. *Hum Mol Genet* **15**, 989–998 (2006).
34. Rooney, J. E. *et al.* Severe muscular dystrophy in mice that lack dystrophin and $\alpha 7$ integrin. *J Cell Sci* **119**, 2185–2195 (2006).
35. Gawlik, K. I., Åkerlund, M., Carmignac, V., Elamaa, H. & Durbeej, M. Distinct roles for laminin globular domains in laminin $\alpha 1$ chain mediated rescue of murine laminin $\alpha 2$ chain deficiency. *PLoS ONE* **5**, e11549 (2010).
36. Cohn, R. D. *et al.* Secondary reduction of $\alpha 7\beta$ integrin in laminin $\alpha 2$ deficient congenital muscular dystrophy supports an additional transmembrane link in skeletal muscle. *J Neurol Sci* **163**, 140–152 (1999).
37. Gawlik, K. I. *et al.* Laminin $\alpha 1$ chain mediated reduction of laminin $\alpha 2$ chain deficient muscular dystrophy involves integrin $\alpha 7\beta 1$ and dystroglycan. *FEBS Lett* **580**, 1759–1765 (2006).
38. Doe, J. A. *et al.* Transgenic overexpression of the $\alpha 7$ integrin reduces muscle pathology and improves viability in the dy(W) mouse model of merosin-deficient congenital muscular dystrophy type 1A. *J Cell Sci* **124**, 2287–2297 (2011).
39. Sorokin, L. M. *et al.* Laminin $\alpha 4$ and integrin $\alpha 6$ are upregulated in regenerating dy/dy skeletal muscle: comparative expression of laminin and integrin isoforms in muscles regenerating after crush injury. *Exp Cell Res* **256**, 500–514 (2000).
40. Taverna, D. *et al.* Dystrophic muscle in mice chimeric for expression of $\alpha 5$ integrin. *J Cell Biol* **143**, 849–859 (1998).
41. Patton, B. L., Miner, J. H., Chiu, A. Y., & Sanes, J. R. Distribution and functions of laminins in the neuromuscular system of developing, adult, and mutant mice. *J Cell Biol.* **139**, 1507–1521 (1997).
42. Flintoff-Dye, N. L. *et al.* Role for the $\alpha 7\beta 1$ integrin in vascular development and integrity. *Dev Dyn* **234**, 11–21 (2005).
43. Velling, T. *et al.* Distinct $\alpha 7\beta 1$ and $\alpha 7\beta 2$ integrin expression patterns during mouse development: $\alpha 7\beta 1$ is restricted to skeletal muscle but $\alpha 7\beta 2$ is expressed in striated muscle, vasculature, and nervous system. *Dev Dyn* **207**, 355–371 (1996).
44. Holmberg, J., Alajbegovic, A., Gawlik, K. I., Elowsson, L. & Durbeej, M. Laminin $\alpha 2$ Chain-Deficiency is Associated with microRNA Deregulation in Skeletal Muscle and Plasma. *Front Aging Neurosci* **6**, 155 (2014).
45. Pegoraro, E. *et al.* Congenital muscular dystrophy with primary laminin $\alpha 2$ (merosin) deficiency presenting as inflammatory myopathy. *Ann Neurol* **40**, 782–791 (1996).
46. Wardrop, K. E. & Dominov, J. A. Proinflammatory signals and the loss of lymphatic vessel hyaluronan receptor-1 (LYVE-1) in the early pathogenesis of laminin $\alpha 2$ -deficient skeletal muscle. *J Histochem Cytochem* **59**, 167–179 (2011).
47. Intengan, H. D. & Schiffrin, E. L. Vascular remodeling in hypertension: roles of apoptosis, inflammation, and fibrosis. *Hypertension* **38**, 581–587 (2001).
48. Weston, B. S., Wahab, N. A. & Mason, R. M. CTGF mediates TGF- β -induced fibronectin matrix deposition by upregulating active $\alpha 5\beta 1$ integrin in human mesangial cells. *J Am Soc Nephrol* **14**, 601–610 (2003).
49. Zhou, X. *et al.* Expression of fibronectin receptor, integrin $\alpha 5\beta 1$ of hepatic stellate cells in rat liver fibrosis. *Chin Med J (Engl)* **113**, 272–276 (2000).
50. Raghov, R. The role of extracellular matrix in postinflammatory wound healing and fibrosis. *FASEB J* **8**, 823–831 (1994).
51. Taniguchi, M. *et al.* Expression profiling of muscles from Fukuyama-type congenital muscular dystrophy and laminin- $\alpha 2$ deficient congenital muscular dystrophy; is congenital muscular dystrophy a primary fibrotic disease? *Biochem Biophys Res Commun* **342**, 489–502 (2006).
52. Klein, S. *et al.* $\alpha 5\beta 1$ integrin activates an NF- κ B-dependent program of gene expression important for angiogenesis and inflammation. *Mol Cell Biol* **22**, 5912–5922 (2002).
53. Accorsi, A., Mehuron, T., Kumar, A., Rhee, Y. & Girgenrath, M. Integrin dysregulation as a possible driver of matrix remodeling in Laminin-deficient congenital muscular dystrophy (MDC1A). *Journal of Neuromuscular Diseases* **2**, 51–61 (2015).
54. Kuang, S., Kuroda, K., Le Grand, F. & Rudnicki, M. A. Asymmetric self-renewal and commitment of satellite stem cells in muscle. *Cell* **129**, 999–1010 (2007).
55. Ozeki, N., Lim, M., Yao, C. C., Tolar, M. & Kramer, R. H. $\alpha 7$ integrin expressing human fetal myogenic progenitors have stem cell-like properties and are capable of osteogenic differentiation. *Exp Cell Res* **312**, 4162–4180 (2006).
56. Liu, J., Milner, D. J., Boppart, M. D., Ross, R. S. & Kaufman S. J. $\beta 1$ chain increases $\alpha 7\beta 1$ integrin and laminin and protects against sarcolemmal damage in mdx mice. *Hum Mol Genet* **21**, 1592–1603 (2012).
57. Heller, K. N. *et al.* AAV-mediated overexpression of human $\alpha 7$ integrin leads to histological and functional improvement in dystrophic mice. *Mol Ther* **21**, 520–525 (2013).
58. Palmer, E. L., Rüegg, C., Ferrando, R., Pytela, R. & Sheppard D. Sequence and tissue distribution of the integrin $\alpha 9$ subunit, a novel partner of $\beta 1$ that is widely distributed in epithelia and muscle. *J Cell Biol* **123**, 1289–1297 (2002).
59. Forsberg, E., Ek, B., Engström, A. & Johansson S. Purification and characterization of integrin $\alpha 9\beta 1$. *Exp Cell Res* **213**, 183–190 (1994).
60. Li, S. *et al.* Laminin-sulfatide binding initiates basement membrane assembly and enables receptor signaling in Schwann cells and fibroblasts. *J Cell Biol* **169**, 179–189 (2005).
61. Petersen, S. C. *et al.* The adhesion GPCR GPR126 has distinct, domain-dependent functions in Schwann cell development mediated by interaction with laminin-211. *Neuron* **85**, 755–769 (2015).

Acknowledgements

This work was supported by: The Swedish Research Council, Crafoord's Foundation, The Royal Physiographic Society, Lars Hierta's Memorial Foundation, Alfred Österlund's Foundation, Greta & Johan Kock's Foundation, Fanny Ekdahl's Foundation and Linnéa & Josef Carlsson's Foundation.

Author Contributions

K.I.G. and M.D. designed the study; K.I.G. performed the experiments; K.I.G. and M.D. analysed the results; K.I.G. and M.D. wrote the manuscript; all authors reviewed the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Gawlik, K. I. and Durbejj, M. Deletion of integrin $\alpha 7$ subunit does not aggravate the phenotype of laminin $\alpha 2$ chain-deficient mice. *Sci. Rep.* **5**, 13916; doi: 10.1038/srep13916 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>