SCIENTIFIC REPORTS

OPEN

A moment from before 365 Ma frozen in time and space

Błażej Błażejowski¹, Piotr Gieszczy², Carlton E. Brett³ & Marcin Binkowski⁴

Received: 12 May 2015

Accepted: 27 July 2015

Published: 18 September 2015

This study presents a detailed analysis of an exceptionally well-preserved articulated specimen of the trilobite *Trimerocephalus* from the Late Devonian of the Holy Cross Mountains in Poland. X-ray microtomography reveals the oldest direct evidence for a moulting episode known from the fossil record. The process of moulting as well as associated features observed in the investigated specimen are interpreted by comparison with extinct and extant Xiphosurida arthropods, which survived global P/T extinction and are among the closest extant relatives of trilobites. A very special moment frozen in time and space millions years ago provides rare insights into the behavior and physiology of these long-extinct arthropods.

Trilobites, a group of marine arthropods which came to prominence in the ‘Cambrian explosion’, finally disappeared during the end Permian Period mass extinction, about 250 million years ago, after a prolonged decline in diversity and abundance. The trilobites were among the most successful of all the early animals, roaming the oceans for over 270 million years. The recent discovery of a three-dimensionally preserved articulated specimen of the blind trilobite *Trimerocephalus chopini* from the Late Devonian (Early Famennian; ~365 Ma) at the Kowala Quarry, near Kielce, Poland (Fig. 1a,b) adds significantly to our understanding of the subphylum Trilobitomorpha. In order to put these observations in a broader context, we extended our study to another group of arthropods – the xiphosurid horseshoe crabs, commonly regarded as the closest extant relatives of trilobites¹. For a comparison, we chose an exceptionally well-preserved horseshoe crab *Crenatolimulus* sp. nov. from the latest Jurassic (Late Tithonian) sediments at the Owadów-Brzezinki Quarry, near Tomaszów Mazowiecki in Central Poland (Fig. 1a,c). The aim of this paper is to present evidence about the mode of preservation, behavior and physiological effects of moulting and its relationship to arthropod survivability.

The blind phacopine genus *Trimerocephalus*² has an almost worldwide distribution in the Famennian³, except for North America. Numerous recent finds of *Trimerocephalus*, commonly forming a line (or ‘queue’) of a few to more than 20 individuals aligned one after another on the upper surfaces of shale beds, are reported from Kowala Quarry, in the Holy Cross Mountains, South-central Poland^{3–4}. From this collection, several specimens were chosen for detailed investigation with X-ray microcomputed tomography (XMT). This provides precise information about the shape and dimensions of internal structures preserved in the rock matrix, giving an opportunity to study valuable fossils without a risk of damage during preparation^{5–6}. Tomographic data after digital processing enable the construction of an isosurface-based and volume-based 3-D model (a ‘virtual fossil’), which can be manipulated, dissected or measured interactively. A sample visualization of the discussed articulated specimen of the *Trimerocephalus chopini* is available at http://www.paleo.pan.pl/people/Blazejowski/SOM_T_chopini_TIME_SPACE.mpg.

Under X-ray tomography the specimen of *Trimerocephalus chopini* is seen to consist of two adjacent exoskeletons, of which the upper one is an articulated thorax and pygidium with the cephalon associated but detached and rotated slightly upward. These skeletal remains are interpreted as a moult

¹Institute of Paleobiology, Polish Academy of Sciences, Twarda 51/55, 00-818 Warszawa, Poland. ²Association of Polish Climatologists, Krakowskie Przedmieście 30, 00-927 Warszawa, Poland. ³Department of Geology, University of Cincinnati, Cincinnati, Ohio, USA, 45221-0013. ⁴Department of Computer Science and Materials Science, University of Silesia, Katowice, Poland. Correspondence and requests for materials should be addressed to B.B. (email: bblazej@twarda.pan.pl)

Figure 1. Contour of Poland with the location of Kowala (50° 47' 46.21" N, 20° 33' 57.52" E) and Owadów-Brzezinki (51° 22' 34.53" N, 20° 89' 07.86" E) quarries. Drawings by B. Błażejowski.

ensemble (exuvium) preserved *in situ* shortly after separating from the moulting individual recognizable as the lower articulated exoskeleton (Fig. 2a,b). The latter appears to reflect the “soft-shelled” individual that died exactly after time of moulting. The increased number of segments visible in the “soft-shelled” trilobite after moulting illustrates the process of growth typical of all arthropods (Fig. 2b,c). A new segment is developed under the old, loosening exoskeleton, and until the moult is loosened, it remains compressed, limited by the total volume of the old exoskeleton (Fig. 2b,c,f). As in extant arthropods the individual inflates to reach new dimensions directly after moulting.

The preservation of this extraordinary composite moult and carcass of *Trimerocephalus* raises intriguing issues of taphonomy and provides further insights into trilobite moult behavior. The articulated preservation of both the carcass and the exuviated exoskeleton requires not only mortality of this individual immediately following moulting but also synchronous entombment of both skeletons within the sediment. This could have happened as a result of rapid, live burial but, in that case, the carcass exoskeleton and exuvium would probably have remained pancaked rather than separated by a layer of sediment. A more plausible scenario was suggested by newly documented specimens of the Silurian–Early Devonian phacopid trilobite *Paciphacops*⁷ from Argentina⁸. In this case, specimens from concretions show the cephalon to be slightly disarticulated and rotated upward from the thoracopygidium⁸. Rustan *et al.* surmised that these trilobites moulted while burrowed shallowly within the sediment. Similar specimens of *Phacops* from the Lower Devonian of Morocco show the cephalon detached and “floating” above the articulated thoracopygidium⁹. In all of these cases, the “soft shelled” trilobite was not apparently preserved, presumably because the animals crawled out of the sediment following successful moulting, leaving the moult parts suspended within the mud. In the illustrated *Trimerocephalus* specimen, the upper, moulted exoskeleton is quite similar in configuration to the other exoskeletons that were inferred to have been left within the sediment. In this specimen, however, the moulting was evidently not successful and the animal died within the sediment shortly after extricating itself from the old exoskeleton. We infer that *Trimerocephalus* also moulted within the sediment and, in this case, left its moult slightly before it also died within the sediment. It is notable that the inferred soft-shelled carcass is preserved below and somewhat forward of the exuvium as might be expected of an animal that had extricated itself from its old exoskeleton (Fig. 2). The cause of mortality is, of course, not known but might be due to stress of moulting or possibly, given the context, death by hypoxia or hydrogen sulfide toxicity within the sediment.

It is also pertinent to enquire how the newly moulted “soft-shelled” trilobite could theoretically be preserved at all. However, it is evident that some individuals did have enough mineralization following moulting to be preserved. Speyer and Brett¹⁰ illustrated a specimen of a very thin, wrinkled exoskeleton of *Eldredgeops rana* found adjacent to other individuals with typical thick, dark calcitic exoskeletons; a few specimens of this sort have been noted and interpreted as carcasses of soft shelled individuals. This is in line with the wrinkling observed here. It is also possible that this was a case of delayed moulting so that the new exoskeleton was more strongly developed than in most “soft shelled” trilobites; hence, its strongly defined preservation.

Figure 2. *Trimerocephalus chopini* from Early Famennian marly shales at Kowala Quarry, Holy Cross Mountains, central Poland. (a). oblique lateral view (SEM); (b–d). oblique lateral view (XMT), (e). dorsal view (SEM); (f). dorsal view (XMT).

Wrinkles and scars

Observed features of the “soft-shelled” trilobite specimen, such as noticeable compression and the presence of wrinkles in the area of the anterior cephalon (Fig. 2e) clearly indicate that the death of this individual occurred just after the process of moulting. Moulting is a common feature of all arthropods including extant and extinct horseshoe crabs, therefore we searched for analogies among an exceptionally well-preserved collection of Late Jurassic horseshoe crabs^{11–12}. These are among the rarest of macrofossils and it is noteworthy that the collection consists of only very well preserved groups of juvenile specimens found¹³. In the latter case, we observed very similar wrinkles in many (but not all) investigated horseshoe crab specimens (Fig. 3). The wrinkling along the anterior prosoma has been interpreted previously as an effect of sediment compaction¹⁴, and this can be the reason in many cases of partly or entirely flattened specimens, but the specimen presented here is very well three-dimensionally preserved (Fig. 3). Taking into consideration the above mentioned course of moulting, another explanation is suggested: when an individual dies during or immediately following moulting, “decompression” does not occur (or occurs only to a limited extent) and results in a form of ‘wrinkles’ that can be observed in the fossilised state. Such wrinkling may be especially well preserved when, as in the case of the *Trimerocephalus* specimen, the animal died within the sediment. Similar features were observed in laboratory environment, where a 5–6 year old juvenile of modern Atlantic horseshoe crab (*Limulus polyphemus*) died in a difficult moulting episode, which resulted in deformation in limbs and wrinkles on its dorsal exoskeleton (Y. Iwasaki, pers. obs.). Possibly the cause of death in all these cases is the same: the effort and energetic loss during the moulting could be so excessive under degraded environmental conditions, that many weak juveniles simply cannot survive the event. This neatly explains only juvenile specimens are commonly found in the fossil record.

The preservation of many specimens very close to each other may be the effect of behavioral phenomena observed in recent horseshoe crabs. Young individuals spend their first years rapidly increasing in size, prior to venturing out into deeper waters, in the so called “nursery grounds^{15–16}”. By analogy, this may indicate that the depositional area of fossil horseshoe crabs could have acted as a nursery ground

Figure 3. Three-dimensionally preserved representatives of Late Jurassic horseshoe crabs from Owadów-Brzezinki, Poland. *Crenatolimulus* sp. nov. (ZPAL X.1/O-B/14.1). (a) Negative and (b) positive (rock slab with imprint).

for numerous individuals, which moulted under somewhat adverse conditions associated with their rapid burial and did not survive the process, forming concentrations of fossils. This may be the one of the reasons why we often observe concentrations of juvenile fossil arthropods. The lack of preserved adult individuals (known only from trackways^{17–19}) in all collections of Jurassic horseshoe crabs is a strong premise supporting that assumption.

A somewhat similar scenario could apply in the case of the Late Devonian *Trimerorophalus*. As noted, this specimen is associated with ‘queues’ of conspecific trilobites, which may represent rows of trilobites that migrated into certain areas, possibly deep offshore setting to breed. An abundance of minute individuals (meraspids?) in association with the rows may represent recently metamorphosed juveniles growing in a “nursery ground”. It is possible, that in a restricted setting, low oxygen and/or high hydrogen sulfide may have been more detrimental to mature trilobites, many of which, including the specimen described herein, appear to have expired literally in their tracks.

References

- Shuster, C. N. Jr. & Anderson, L. I. *A history of skeletal structure: clues to relationships among species*. 154–188. In: C. N. Shuster, R. B. Barlow & H. J. Brockman (eds), *The American Horseshoe Crab*. Harvard University Press, Cambridge, Massachusetts (2003).
- McCoy, F. On the classification of some British fossil Crustacea with notices of new forms in the university collection at Cambridge. *Annals and Magazine of Natural History* **2**, 392–414 (1849).
- Kin, A. & Błażejowski, B. A new species of blind phacopid trilobite from the Late Devonian (early Famennian) of Poland. *Zootaxa* **3626**(3), 345–355 (2013).
- Radwański, A., Kin, A. & Radwańska, U. Queues of blind phacopid trilobites *Trimerorophalus*: A case of frozen behaviour of Early Famennian age from the Holy Cross Mountains, Central Poland. *Acta Geol Polonica*. **59**(4), 459–481 (2009).
- Błażejowski, B., Binkowski, M., Bitner, M. A. & Gieszczyński, P. X-ray microtomography (XMT) of fossil brachiopod shell interior for taxonomy. *Acta Palaeontologica* **56**(2), 427–428 (2011).
- Sutton, M.D., Rahman, I. & Garwood, R. J. *Techniques for Virtual Palaeontology*. Wiley-Blackwell, 208 pp (2014).
- Maksimova, Z. A. New Devonian trilobites of the Phacopoidea [Novye devonskie trilobity Phacopoidea]. *Paleontologicheskij Zhurnal* **14**(1), 88–95 (1972).
- Rustan, J. J., Balsiero, D., Waisfeld, B., Fogalia, R. D. & Vaccari, N. E. Infaunal moulting in Trilobita and escalatory responses against predation. *Geology* **39**(5), 495–498 (2011).
- Brett, C. E., Zambito, J. J., Hunda, B. & Schindler, E. Mid-Paleozoic trilobite Lagerstätten: Models of diagenetically enhanced obrution deposits. *Palaos* **27**, 326–345 (2012).
- Speyer, S. E. & Brett, C. E. Clustered trilobite assemblages in the Middle Devonian Hamilton Group. *Lethaia* **18**, 85–103 (1986).
- Kin, A., Gruszczynski, M., Martill, D., Marshall, J. & Błażejowski, B. Palaeoenvironment and taphonomy of a Late Jurassic (Late Tithonian) Lagerstätte from central Poland. *Lethaia* **46**, 71–81 (2013).
- Błażejowski, B. The oldest species of the genus *Limulus* from the Late Jurassic of Poland. In: Carmichael, R. H., Botton, M. L., Shin, P. K. S. & Cheung, S. G. (ed.), *Changing global perspectives on biology, conservation, and management of horseshoe crabs*. Springer, New York (2015).
- Kin, A. & Błażejowski, B. The Horseshoe Crab of the genus *Limulus*: living fossil or stabilomorph? *PLoS ONE* **9**(10), e108036. doi: 10.1371/journal.pone.0108036 (2014).
- Shuster, C. N., Barlow, R. B. & Brockmann, H. J. *The American Horseshoe Crab*. Harvard University Press, Cambridge, Massachusetts (2003).
- Carmichael, R. H., Rutecki, D. & Valiela, I. Abundance and population structure of the Atlantic horseshoe crab *Limulus polyphemus* in Pleasant Bay, Cape Cod. *Mar Ecol Prog Ser* **246**, 225–239 (2003).
- Chiu, H. M. C. & Morton, B. The Behaviour of Juvenile Horseshoe Crabs, *Tachypleus tridentatus* (Xiphosura), on a Nursery Beach at Shui Hau Wan, Hong Kong. *Hydrobiologia* **523**(1–3), 29–35 (2004).
- Schweigert, G. Die Spurenfauna des Nusplinger Plattenkalks (Oberjura, Schwäbische Alb). *Stutt Beitr Natur B* **262**, 1–47 (1998).
- Gaillard, C. A giant limulid trackway (*Kouphichnium lithographicum*) from the lithographic limestones of Cerin (Late Kimmeridgian, France): ethological and environmental implications. *Swiss J Geosci* **103**, 1–16 (2011).

19. Moreau, J.-D., Fara, E., Gand, G., Lafaurie, G. & Baret, L. Gigantism among Late Jurassic limulids: New ichnological evidence from the Causses Basin (Lozère, France) and comments on body-size evolution among horseshoe crabs. *Geobios* **47**(4), 237–253. doi: 10.1016/j.geobios.2014.06.005 (2014).

Acknowledgements

This work was completed while the first author was a recipient of a grant from the Polish National Science Centre (NCN): Grant number 2012/07/B/ST10/04175. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. No additional external funding were received for this study. We would like to sincerely thank Peter Walsh (University of Silesia) for many useful suggestions and linguistic correction. We express our thank to Yumiko Iwasaki and Mark Botton (both from Fordham University at Lincoln Center), Ruth H. Carmichael (University of South Alabama), Jennifer H. Mattei (Sacred Heart University), David Rudkin (Royal Ontario Museum), H. Jane Brockmann (University of Florida) and Gary Kreamer (Delaware Division of Fish and Wildlife) for many helpful suggestions related to living horseshoe crabs during the early phase of this investigation. And last but not least, we wish to acknowledge Michał Walczak (University of Silesia, Katowice) for making segmentation and visualisation of trylobite based on XMT images.

Author Contributions

B.B. had the idea for the project, conceived and designed research. M.B. performed X-ray microtomography imaging and imaginary data processing. B.B., P.G. and C.E.B. analysed and discussed the results and wrote the manuscript. B.B. prepared maps and figures.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Błażejowski, B. *et al.* A moment from before 365 Ma frozen in time and space. *Sci. Rep.* **5**, 14191; doi: 10.1038/srep14191 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>