

SCIENTIFIC REPORTS

OPEN

Quantitative assessment of gene expression network module-validation methods

Received: 20 March 2015

Accepted: 21 September 2015

Published: 16 October 2015

Bing Li^{1,2}, Yingying Zhang¹, Yanan Yu¹, Pengqian Wang¹, Yongcheng Wang³, Zhong Wang¹ & Yongyan Wang¹

Validation of pluripotent modules in diverse networks holds enormous potential for systems biology and network pharmacology. An arising challenge is how to assess the accuracy of discovering all potential modules from multi-omic networks and validating their architectural characteristics based on innovative computational methods beyond function enrichment and biological validation. To display the framework progress in this domain, we systematically divided the existing *Computational Validation Approaches based on Modular Architecture (CVAMA)* into topology-based approaches (TBA) and statistics-based approaches (SBA). We compared the available module validation methods based on 11 gene expression datasets, and partially consistent results in the form of homogeneous models were obtained with each individual approach, whereas discrepant contradictory results were found between TBA and SBA. The TBA of the Zsummary value had a higher Validation Success Ratio (VSR) (51%) and a higher Fluctuation Ratio (FR) (80.92%), whereas the SBA of the approximately unbiased (AU) p-value had a lower VSR (12.3%) and a lower FR (45.84%). The Gray area simulated study revealed a consistent result for these two models and indicated a lower Variation Ratio (VR) (8.10%) of TBA at 6 simulated levels. Despite facing many novel challenges and evidence limitations, *CVAMA* may offer novel insights into modular networks.

Modularity is a common characteristic of omics-based biological networks^{1–3}. Module-based analyses that investigate or deconstruct omics-based biological networks have become a hot topic in recent years^{4,5}. Various types of algorithms have been proposed to identify modules (also known as communities, clusters, and subnetworks), including network clustering^{6,7}, heuristic search^{8,9}, seed extension¹⁰, topology network^{11,12}, and matrix decomposition^{13,14}. However, in contrast to the large number of module detection methods⁴, there are few methods for module validation and evaluation. How to evaluate the accuracy and validity of modules has become a new challenge for researchers. Most previous studies used function enrichment methods to evaluate modules based on functional annotations, such as GO, MIPS, and KEGG^{15–24}. However, some modules may be enriched with too many functions, whereas others may be enriched without any functions, and the background annotation database itself is constantly being updated. Other studies used molecular biological experimental techniques to verify the co-expression, transcription regulation or other interaction relationships among members of a given module^{25–30}. However, this method is only suitable for small modules that consist of only a few nodes, and it is nearly impossible to perform this method for a larger module. Thus, in the era of Big Data and omics revolution, an arising challenge is to explore rational strategies to validate biological network modules.

¹Institute of Basic Research in Clinical Medicine, China Academy of Chinese Medical Sciences, 16 Nanxiaojie, Dongzhimennei, Beijing 100700, China. ²Institute of Information on Traditional Chinese Medicine, China Academy of Chinese Medical Sciences, 16 Nanxiaojie, Dongzhimennei, Beijing 100700, China. ³Brightech International, LLC, 285 Davidson Ave #504, Somerset, NJ 08873, USA. Correspondence and requests for materials should be addressed to Z.W. (email: zhonwtcm@sina.com) or Y.W. (email: wangyongyan@sina.cn)

Several published studies have employed *Computational Validation Approaches based on Modular Architecture* (CVAMA) to evaluate modules' authenticity, reproducibility, and significance or to identify phenotype-related functional modules^{31–35}. These approaches are not limited by module size and supporting databases. With an increasing number of omics technologies and module analysis methods, CVAMA may become the new focus. In this paper, we summarized the available CVAMA methods, which were divided into topology-based approaches (TBA) and statistics-based approaches (SBA). One representative method of each was selected to validate modules obtained from genomic datasets, and comparative analyses were performed to illuminate the feasibility and challenges in CVAMA.

Results

Topology-based approaches (TBA) for module validation. A module may have several topological features, such as modularity², connectivity³⁶, density^{36,37}, clustering coefficient³⁷, degree³⁸, and edge betweenness³⁹. Module detection methods may focus on one or a few topological criteria, and it is also essential to determine whether the identified modules have a modular structure. Therefore, we may use a single or composite topological index to evaluate whether a module is valid (Table 1). Any single topological index used to validate a module should be independent of the methods used to identify the module, such as the network perplexity index of *Entropy*^{40,41}. The entropy increases when the data are more uniformly distributed; therefore, a good quality module is expected to have a low entropy^{42,43}. Topological indexes, including intra-modular connectivity⁴⁴ and *NB* value²⁶, have been applied to evaluate whether the intra-modular structure is different from other parts of the whole network. Other indexes, such as compactness⁴⁵ and weak community⁴⁶, can be used to select good clusters from integrated clustering results. Because a single topological index is not likely to provide a global evaluation of the modular structure, an alternative choice is to combine multiple topological indexes into an integrated measure to assess a module's validity. Both internal and external indexes, such as density, connectivity, and tabulation-based module preservation statistics, can be integrated to validate the existence of a module^{35,47}. Based on a global view of the modular structure, it may be advantageous to aggregate multiple module evaluation statistics into summary preservation statistics. In our study, we validated five preserved modules whose *Zsummary* value (an integrated index) was greater than 2.

Statistics-based approaches (SBA) for module validation. In addition to topological criteria, a module should also be statistically significant, which means that the modular architecture distribution ought to be highly unlikely to be obtained by chance in a randomized network. Moreover, exploring the relationship between modules and various phenotypes or identifying consistent modules may also require significance testing^{31,33,48,49}. For this reason, SBA is an important process to assess a module's stability, phenotypic correlation or significance of consistency (Table 1). For responsive modules or module biomarker identification^{50–52}, binary or mixed integer linear programming models can be used to validate the causal or dependent relations between network modules and biological phenotypes^{34,53,54}. In phylogeny, resampling approaches are defined as a confidence measure for splits in a phylogenetic tree and are used to calculate consensus trees⁵⁵, which can also be used to assess the robustness of modules in network analysis^{25,56}. A permutation test with a *p* value calculated by empirically estimating the null distribution can be adopted to determine whether the module composition is higher than expected by chance or associated with the disease being investigated^{57,58}. Moreover, given two or more networks, comparative network analysis is often used to identify modules across networks or species, and these modules are defined as consensus or conserved modules^{31,59}. Moreover, a module's "reproducibility" can also be assessed, i.e., to what extent a module obtained from one network is compatible with modules in another network^{46,55,60–62}. In our study, by using hierarchical clustering, we identified 66 statistically significant modules based on approximately unbiased (AU) *p*-values.

Module identification based simply on topological criteria or statistical significance may not discover certain types of biologically meaningful modules⁶³. Because disparate results can be obtained from the same network with different algorithms, functional validation can be used to evaluate the performance of different module identification methods⁶⁴. Although it is not our focus in this study, we summarize and list functional module validation methods reported in the published literature (Supplementary Table 1). Typically, the most widely used functional validation method is functional homogeneity evaluation^{65,66}, with indexes such as functional enriched *p* value^{7,45,67} and *R* score^{7,68}. Furthermore, the index of quantitative score based on function enrichment analysis may be applied to assess a module's confidence level²¹ or disease relationship⁶⁹. Moreover, known protein complex matching can also provide functional evidence for modules, and the commonly used indexes include the overlapping score (OS)^{7,70} and *F*-measure^{67,71}. Other measurements, such as the positive predictive value (PPV), accuracy, and separation, can also be used⁷². For small modules, the experimental techniques of molecular biology, such as real-time quantitative PCR (qPCR), western blotting, and siRNA knock-down, may be applied to validate the co-expression, co-regulation or other interaction relations among the genes or proteins within a module^{25–28}.

Homogeneity of different models of TBA on the same dataset. Both *Zsummary* and *medianRank* are integrated topological indexes of module preservation. We applied these two models to evaluate modules identified from the same dataset (GSE24001), which was derived from 30 newly diagnosed infant acute lymphoblastic leukemia samples. Modules were identified by the Weighted Gene

No.	Type	Index	Equation	Criteria	Application	Test data	Ref.
Topological validation							
1	Integrated index	Z_{summary}	$Z_{\text{summary}} = \frac{Z_{\text{density}} + Z_{\text{connectivity}}}{2}$	≥ 10 , strongly preserved; 2~10, moderately preserved; ≤ 2 , no preservation	Composite preservation statistics to validate whether a module is significantly preserved in another network. Apply to correlation networks (e.g., co-expression networks)	yes	35,40,41
2		$Z_{\text{summaryADJ}}$	$Z_{\text{summary}} = \frac{Z_{\text{densityADJ}} + Z_{\text{connectivityADJ}}}{2}$	≥ 10 , strongly preserved; 2~10, moderately preserved; ≤ 2 , no preservation	Same as above. Apply to general networks (e.g., adjacency matrix networks)	yes	35
3		medianRank	$\text{medianRank} = \frac{\text{medianRank.density} + \text{medianRank.connectivity}}{2}$	The lower the better	Same as above.	yes	35
4	Single index	Entropy	$\text{Entropy}(M) = -\sum_{j \in \text{bins}} p_j \log_2 p_j$	The smaller the better	Access the quality of identified modules. A good quality module is expected to have a low entropy.	no	42,43
5		M_{pres}	$M_{\text{pres}} = \text{cor}(k^l, k^m)$	The closer to 1, the better	Describe the preservation of intra-modular connectivity across two networks. A p-value can be assigned to evaluate the reproducibility of modules.	yes	44,45
6		NB value	$\text{NB} = \frac{\sum e(i)}{\sum d(i)}$	$\text{NB} \geq 0.5$	A ratio of edges within a module and the total number of edges between modules is used to select modules with high intra-modular connectivity.	no	26
7		CS (S)	$\text{CS}(S) = \frac{\text{the number of closed walks whose step is three in } G[S]}{\text{the average shortest path length in } G[S]}$	$\text{CS}(S) > 0$, the higher the better	Describe the compactness and neighboring conditions of a cluster. Apply to select good clusters from integrated clustering results	no	46
8		LS (S)	$L(S) = \frac{1}{ S } \times \frac{ E(G[S]) }{\binom{ S }{2}} \sum_{i \in S} \frac{k_i^{\text{in}}(S) - k_i^{\text{out}}(S)}{d_i \times s_i}$	The higher the better	Judge the quality of a cluster S in a graph G and help to select good clusters from integrated clustering results.	no	47
9		Modularity	$Q = \sum_i (e_{ii} - a_i^2) = T_r(e) - \ e^2\ $	$0.3 \leq Q \leq 0.7$	Evaluate the level of modular structure and the best split of a network into modules.	no	2,39,102
Statistical validation							
1	Integer linear programming	$C \cdot (X_1, X_2, \dots, X_k)$	$\ S - \bar{S}_{\text{control}}\ _2^2 - \ S - \bar{S}_{\text{case}}\ _2^2 < 0$, for $S \in S_{\text{control}}$, $\ S - \bar{S}_{\text{case}}\ _2^2 - \ S - \bar{S}_{\text{control}}\ _2^2 < 0$ for $S \in S_{\text{case}}$,	$C \leq 0$, the smaller the better	A classifier and integer linear programming model to select modules based on the activity of the module in case and control samples.	yes	49,50
2	Bootstrap resampling	P-value	NULL	$P \leq 0.05$	P-value is derived from multiscale bootstrap resampling to assess the uncertainty of clustering analysis and search for significant modules.	no	25,33
3		Consensus score	$S_f^p(v) = (\sum_{i=1}^I \{v\} \cap V_i) - \rho$, $S_f^p(e) = (\sum_{i=1}^I \{e\} \cap E_i) - \rho$,	$\geq \rho$, the higher the better	A jackknife resampling procedure is used to assess the accuracy and robustness of functional modules resulting in an ensemble of optimal modules.	no	56
4	Permutation test	Combinatorial p-value	NULL	Combinatorial criteria: (1) $P_{(Zm)} < 0.05$; (2) $P_{GL} > P_{nSNPs}$, $P_{\text{topo}} < 0.05$; (3) $P_{\text{emp}} < 0.05$ Additional criteria: $P_{(Zm(\text{eval}))}$ and/or $P_{\text{emp}(\text{eval})} < 0.05$	Significance and permutation tests are used to calculate the P value of module scores. Appropriate for GWAS data; multiple GWAS datasets are needed when using additional criteria.	yes	52
5		coClustering (q)	$\text{coClustering}(q) = \sum_{q'=1}^{Q[\text{test}]} \text{propCoClustering}(q, q')$	$\geq 95\%$	A cross-tabulation-based statistic for determining whether modules in the reference dataset are preserved in a test dataset, a permutation test to determine the p value.	yes	45
Continued							

No.	Type	Index	Equation	Criteria	Application	Test data	Ref.
6	Modular compatibility	Compatibility Score (C_p)	$C_p = \frac{1}{2} \left[\sum_{i=1}^M \frac{C_{1i}}{T_{1i} + P_{1i} + C_{1i}} + \sum_{j=1}^N \frac{C_{2j}}{T_{2j} + P_{2j} + C_{2j}} \right]$	The closer to 1, the better	An indication of agreement or overlap between two sets of modules to measure the network modular compatibility between two networks.	yes	53
7		Matching p-value	NULL	$P < 0.05$	Modified hypergeometric test-derived p-values with Bonferroni correction to measure modules' conservation between any two species or networks.	yes	54
8		IGP	$IGP(u, X) = \frac{\# \{j \mid Class_X(j) = Class_X(j^N) = u\}}{\# \{j \mid Class_X(j) = u\}}$	The closer to 1, the better	Defined to validate an individual cluster's reproducibility and prediction accuracy.	yes	55

Table 1. Topology- and statistics-based methods for module validation. The topology-based methods (TBA) and statistics-based methods (SBA) for module validation. The columns reports the types, index names, equations, criteria (the cut-off value to evaluate modules), applicable conditions, test data (whether this method requires an additional test network to validate a module) and references.

Co-Expression Network Analysis (WGCNA) R package⁷³, setting 3 as the minimum module gene number. Each module was detected based on a hierarchical cluster tree and was labeled by colors (Fig. 1A). The validation results are shown in Fig. 1B,C. Five preserved modules ($Z_{summary} \geq 2$) all had a relatively low *medianRank* values, and the most strongly preserved module ($Z_{summary} = 13$) had the lowest *medianRank* value. Similarly, the two modules that had the highest *medianRank* value (*medianRank* = 10) were both unpreserved.

Example modules validated by two models of TBA. Among the 5 preserved modules, the turquoise module, which had 250 nodes and 30,833 edges, was most strongly preserved ($Z_{summary} = 13$), having the lowest *medianRank* value (*medianRank* = 2) (Fig. 2A). The preserved yellow module, which was composed of 64 nodes and 1,970 edges, had a low *medianRank* value (*medianRank* = 5) (Fig. 2B). Among the 4 unpreserved modules, the red module, consisting of 23 nodes and 253 edges (Fig. 2C), and the magenta module, consisting of 5 nodes and 9 edges (Fig. 2D), had the highest *medianRank* values (both *medianRank* = 10).

Variability of TBA ($Z_{summary}$) and SBA (AU P -value) results on the same dataset. Based on the same dataset (GSE24001), we compared the validation results of TBA and SBA, choosing $Z_{summary}$ and AU p -value as the representative methods for each approach. For $Z_{summary}$, 5 preserved modules were validated from 9 modules. The $Z_{summary}$ value was dependent on the module size, which was consistent with previous studies³⁵ (Fig. 1D). The AU p -value used to access the modular architecture distributional probability was computed to search for significant modules (clusters). As shown in Fig. 1E, 66 modules with 3 or more genes and an AU p -value larger than 0.95 are highlighted by rectangles, which are strongly supported by the gene expression data. Different numbers of detected modules and valid modules were obtained through the two methods.

Multiple comparisons of $Z_{summary}$ and *medianRank* on 10 datasets. For the same modules identified by WGCNA, we applied the integrated topological indexes $Z_{summary}$ and *medianRank* to 10 datasets to further compare the results of module preservation evaluation. The average size of modules obtained from the 10 datasets is shown in Fig. 3A. Because *medianRank* was a relative preservation index without a cutoff value, we compared the top 10 ranked modules validated by $Z_{summary}$ and *medianRank* (Fig. 3B). We failed to obtain a valid $Z_{summary}$ value in two datasets (GSE6448, GSE29230) when the minimum module size was set at 3. For the other 8 datasets, overlapping preserved modules validated by $Z_{summary}$ and *medianRank* were found in 7 datasets, and consistent ranked modules were observed in 6 datasets, demonstrating the consistency of the two indexes. However, no overlapping modules in the top 10 preserved modules were found in one dataset (GSE4882).

$Z_{summary}$ analysis was impeded by small module size. Because we failed to obtain a valid $Z_{summary}$ value in two datasets (GSE6448, GSE29230) with a minimum module size of 3, we changed this cutoff value from 4 to 10. Then, valid $Z_{summary}$ values were acquired in both datasets, and the percent of preserved modules was stable (Fig. 3C) due to the too small density or connectivity, leading to invalid $Z_{summary}$ values when the minimum module size was set at 3.




Figure 1. (A) Hierarchical cluster tree showing coexpression modules identified by WGCNA. Each leaf in the tree represents one gene. The major tree branches constitute 9 modules labeled by different colors. (B) The medianRank preservation statistics (y-axis) of the modules. Each point represents a module, labeled by color and names. Low numbers on the y-axis indicate high preservation. (C) The Zsummary preservation statistics (y-axis) of the modules. The modules are labeled as in panel (B). The dashed blue and green lines indicate the thresholds. A Zsummary value over 2 represents a moderately preserved module, and a value over 10 provides strong evidence of module preservation. (D) Scatter plots showing the correlation between the Zsummary (y-axis) and module size (x-axis). (E) The cluster dendrogram with approximately unbiased (AU) P-values. The AU p-values are displayed in red, and clusters with an AU p-value lower than 0.05 are highlighted by rectangles. The calculations and the drawn figure were performed using the pvclust R package.

Comparison of TBA (*Zsummary*) and SBA (AU *P-value*) results for 10 datasets. As mentioned above, *Zsummary* is a TBA index and AU *P-value* is an SBA index. Based on 10 datasets, we compared the performance of these two types of index. In this application, modules with 3 or more genes were considered as valid modules. The proportions of valid (preserved or significant) modules obtained by these two indexes from 10 datasets are shown in Fig. 3D,E. For different methods and datasets, both the module number and the proportion of valid modules varied greatly. Overall, the Validation Success Ratios (VSR) of *Zsummary* and AU *P-value* were 51% and 12.3%, respectively (Fig. 3F). This indicated that *Zsummary* obtained a higher ratio of valid modules (invalid values from two datasets were deemed as zero). A prior study adopted *Zsummary* to validate CASTxB6 female liver modules with 9 other expression datasets and revealed an average VSR of 86.44%⁷⁴. However, *Zsummary* also had a higher Fluctuation Ratio (FR, 80.92%) than AU *P-value* (45.84%), indicating that the stability of the AU *P-value* results was superior to that of *Zsummary* (Fig. 3F).

Correlation between the network parameters and the ratio of valid modules. To further determine which network parameters influence the ratio of valid modules of the *Zsummary* and AU *P-value* methods, we selected 6 main network parameters of the 10 datasets (Supplementary Table 2), i.e., modularity, density, clustering coefficient, characteristic path length, network heterogeneity, and network centralization. Linear regression analysis indicated that none of these 6 parameters was correlated with the valid module ratio of the *Zsummary* and AU *P-value* methods (Fig. 4). This implied that the valid module ratio of genomic networks may not be influenced by a single network parameter.




Figure 2. The preserved and un-preserved modules. Each node is a gene, and each edge is the co-expression relationship. (A) The turquoise module. (B) The yellow module. (C) The red module. (D) The magenta module.

Impact of gray area variation on 8 datasets. The gray area was the region of gray genes that was not assigned into any module and labeled in gray by WGCNA. Except for the two datasets (GSE6448 and GSE29230) without valid *Zsummary* values, 8 datasets were simulated by changing the gray area genes' expression levels to 0.1, 0.5, 0.9, 1.1, 1.5, and 2 times that of the original dataset. Based on each simulated dataset, WGCNA (an R package used to compute *Zsummary*) and pvclust (an R package used to compute *AU p-value*) were performed for module identification and validation. For WGCNA, 5 datasets (GSE2283, GSE12148, GSE6738, GSE12520, and GSE4882) had no changes at the 6 simulated levels compared with the original datasets (a change in *Zsummary* less than the cutoff value was considered as no change). The changes in the number of modules or gray genes (only for WGCNA) and Variation Ratio (VR) for the remaining 3 datasets are shown in Fig. 5A–C. For pvclust, changes were observed in all 8 datasets compared with the original datasets, and the changes in the module number and VR are shown in Fig. 5D–K. No correlation was found between the VR and the simulated levels in the changed datasets.

Comparison of TBA and SBA by Gray area simulation. For the 8 datasets, the average VRs of WGCNA and pvclust at the 6 simulated levels are shown in Fig. 6A. With regard to WGCNA, only the VRs in the changed datasets were calculated. The VRs of WGCNA and pvclust in all simulated datasets can be seen in Fig. 6B. WGCNA had a higher VR (8.43%) for module identification but a lower VR (8.10%) for module validation. By contrast, pvclust had a lower VR (1.29%) for module identification and a higher VR (14.06%) for module validation. Thus, the gray area changes had different impacts on the two models in terms of module identification and validation. Moreover, the VSR and FR of TBA (*Zsummary*) and SBA (*AU p-value*) were stable at each simulated level (Fig. 6C). *Zsummary* had a higher VSR and FR at each simulated level (2 datasets with invalid *Zsummary* values were not included). When data at all 6 simulated levels were aggregated, the VSR of *Zsummary* and *AU p-value* was 63.82% and 12.30%, and the FR was 55.84% and 51.42%, respectively.




Figure 3. The comparison of TBA (Zsummary and medianRank) and SBA (AU p-value) on 10 datasets. (A) A mean module size of 10 datasets. The y-axis is the mean module size (nodes), and the x-axis is each dataset number. (B) Comparison of the top 10 preserved modules validated by Zsummary and medianRank. The red spots represent the number of consistently ranked modules. The green spots represent the number of overlapping modules. The blue spots represent the number of non-overlapping modules. (C) The effect of changing the minimum module size setting from 4 to 10 on two datasets. The y-axis is the percentage of preserved modules, and the x-axis is the different cutoff value settings. (D) The percentage of preserved modules validated by Zsummary on 10 datasets. (E) The percentage of significant modules validated by AU P-value on 10 datasets. In (D,E) the blue bars indicate the number of all the modules detected, and the red bars indicate the number of preserved or significant modules. (F) The VSR and FR of Zsummary and AU P-value on 10 datasets. The red bars represent Zsummary, and the blue bars represent the AU P-value.

Discussion

Functional enrichment and biological experiments based on module validation methods may not satisfy the rising demands of various omics networks. As an alternative choice, CVAMA can potentially provide an analytical assessment of the structure and stability of modules captured by various partitioning methods and should be considered as a crucial tool in the interpretation of network modules. The feasibility of CVAMA was demonstrated in our applications of TBA and SBA-based module validation methods in genomic network modules. As a TBA, Zsummary had a high VSR (51%) but also a high FR (80.92%) and was impeded by small module size. As an SBA, AU p-value had a low FR (45.84%) but also a low VSR (12.3%). The Gray area simulated study showed that the VSR and FR of both TBA (Zsummary) and SBA (AU p-value) remained stable at each simulated level. Meanwhile, TBA (Zsummary) had a lower VR (8.10%) for module validation at the 6 simulated levels, indicating that the gray gene changes had little impact on the topology-based models. Although different validation results were obtained by the two types of method with different gene expression datasets and module detection methods, one may choose an appropriate validation index based on the topological structure or stability of the modules. For example, if we focus on whether a module is structurally different from the rest of the network, we may use Zsummary to assess its preservation. If we focus on whether a module is stable or robust, we may choose AU P-value to assess its confidence. Taken together, the existing methods are not ideal, and further improvement is justified.

Modular analysis in genomic networks is a complicated process that involves various factors. In general, there is no “golden standard” for assessing the validity and quality of modules, and different algorithms for module identification with different parameters may produce disparate module partition results^{4,7,76}. Thus, it is difficult to determine which module is “correct” and which module partition method outperforms others. As our application demonstrated, different types of module validation indexes for the same network may generate different outcomes. Generally speaking, each type of module validation method may have its own advantages and disadvantages, and some methods may require certain conditions (e.g.,




Figure 4. The relationship between the percentage of valid (preserved or significant) modules (y-axis) and the network parameters (x-axis). The network parameters were calculated by plugins in Cytoscape. (A,B) MCL versus valid module percentage. (C,D) Density versus valid module percentage. (E,F) Clustering coefficient versus valid module percentage. (G,H) Characteristic path length versus valid module percentage. (I,J) Network heterogeneity versus valid module percentage. (K,L) Network centralization versus valid module percentage. The red line added to each plot is the linear regression line with intercept 0 and slope 1.

data type, network pattern, or module identification method), limiting their applicability and flexibility. There is no universally acceptable approach that can perform well for all types of data under all scenarios, which results in challenges to make a clear-cut prescription for genomic module validation.

As for the possible discordance between topological criteria and biological meaning in module identification⁷⁷, methods combining both function and structure have been proposed to identify functional modules⁷⁸. Similarly, function and topology are also the two aspects of module validation. CVAMA may neglect the biological meaning and directly assess the correlation of modules with known functional annotation, which may deviate from the densely connected property. It is assumed that fusion of functional and topological evaluation may lead to a high quality selection of better modules. However, most of the existing methods in published literature focus on either function or topology, and researchers may only be interested in their own subject or module identification algorithm. Therefore, module validation methods that integrate both functional and topological indexes and are independent of the vagaries of module detection algorithms need to be further explored.

For the modules obtained by clustering algorithms, internal and external cluster validation methods for assessing the quality of clustering results have been discussed in previous studies^{47,79}. The internal validation attempts to measure how well a given partitioning corresponds to the natural cluster structure of the data, and such indexes include compactness, connectedness, separation, combinations, and stability⁴¹. External validation attempts to compare the recovered structure to a priori knowledge and to quantify the match between them, and such indexes include unary measures and binary measures⁷⁹. In addition to cluster quality assessment, how to estimate the optimal number of clusters has also been discussed^{40,80}. Generally, if a module is known to be consistent with the known knowledge, it would show stronger evidence of preservation than a module without a priori evidence, such as a known pathway or co-transcriptional regulation^{17,81–83}.

Gene interactions are dynamic in regulating the functioning of cells and organisms^{84,85}. A number of studies have focused on dynamic module identification, as well as the dynamic behavior of modules in networks^{11,86,87}. As such, module validation should not be constrained to a static situation. Modular dynamics may involve time-series molecular interaction⁸⁸, environment changes⁸⁹, phenotypic




Figure 5. The changes of modules or gray genes (only for WGCNA) on 8 datasets by Gray area simulation. The blue spots represent the number of modules identified by WGCNA or pvcust. The red spots represent the number of valid (preserved or significant) modules. The gray spots are the number of gray genes. The numerical value above the spots is the VR at 6 simulated levels compared with the original dataset.




Figure 6. Comparison of WGCNA (Zsummary) and pvcust (AU p-value) on Grey area simulated datasets. (A) The average VR of the modules and genes of 8 datasets. (B) The average VR of WGCNA and pvcust of all simulated datasets. The red bars represent WGCNA, and the blue bars represent pvcust. (C) The VSR and FR of Zsummary and AU P-value of 8 datasets at 6 simulated levels. The red bars represent Zsummary, and the blue bars represent AU P-value.

changes^{90,91}, and ontogenetic and phylogenetic time^{92,93}. Dynamic evaluation of modules in certain dynamic processes may generate more comprehensive results, which cannot be obtained in a static state. The more meaningful consequence is intertwined with greater challenges in dynamic CVAMA.

Therefore, to address the challenge of omics-based module validation, computing-based methods are an easy and feasible choice. Despite these challenges, CVAMA, in addition to functional enrichment and biological experiments, offers novel insights into module network research and may become a new paradigm in modular analysis.

Materials and Methods

Datasets and samples. Gene expression datasets were obtained from the GEO database (<http://www.ncbi.nlm.nih.gov/geo/>). Eleven spotted DNA/cDNA datasets from different organisms and experiment platforms were downloaded, with the sample size ranging from 28 to 592, and the gene number ranging from 448 to 3,520. Because a test dataset was needed for *Zsummary* and *medianRank*, we selected half of the samples in each dataset as reference data and the other half as test data. The raw gene-expression information is shown in Supplementary Table 2.

Network construction and network parameter calculation. For each genomic dataset, weighted gene co-expression network analysis (WGCNA)⁷³ was used to construct a network and to detect modules. The freely available WGCNA R package and R tutorials were described in⁷³. After network construction, we exported each network to Cytoscape software⁹⁴. Network analysis was conducted with the NetworkAnalyzer⁹⁵ plugin to calculate the network topological parameters. Network modularity was calculated by the CommFinder⁹⁶ plugin in Cytoscape. The network topological parameters and modularity are listed in Supplementary Table 2.

***Zsummary* and *medianRank* module preservation statistics.** *Zsummary* is an integrated statistic implemented in functional module preservation in the WGCNA R package³⁵. It is composed of 4 statistics related to density and 3 statistics related to connectivity that can quantitatively assess whether the density and connectivity patterns of modules defined in a reference dataset are preserved in a test dataset. A *Zsummary* value between 2 and 10 indicates moderate module preservation, whereas a *Zsummary* > 10 provides strong support for module preservation³⁵. Another integrated index, *medianRank*³⁵, is also composed of statistics related to density and connectivity. It is a rank-based measure to compare the relative preservation among multiple modules; a module with lower *medianRank* tends to exhibit stronger observed preservation.

Approximately unbiased (AU) p-value. The AU *p-value*, computed by multi-scale bootstrap resampling⁹⁷, was selected as a representative SBA index. The AU *p-value* is often used to assess the uncertainty of clustering analysis⁹⁸. In our application, the AU *p-value* was computed using the R package pvclust⁹⁹ with 1,000 times resampling, varying the bootstrap sample size from 0.5 to 1.4-fold the real sample size of the gene expression data. We set clusters with an AU *p-value* larger than 0.95 as significant modules⁹⁹.

Validation success ratio (VSR) and fluctuation ratio (FR). The VSR and FR were defined for comparing the two types of module validation approaches on multiple datasets. The VSR (Eq. 1) was defined as the average percentage of valid modules against all modules available on multiple datasets. The FR (Eq. 2) was defined as the stability or variation degree of the percentage of valid modules on multiple datasets; a lower FR indicated that the percentage of valid modules was more stable.

$$VSR = \frac{(R_1 + R_2 + \dots + R_n)}{N_{\text{datasets}}} \quad (1)$$

$$FR = \left(\frac{\sqrt{\frac{\sum_{i=1}^n (R_i - \bar{R})^2}{N-1}}}{\bar{R}} \right) \times 100\% \quad (2)$$

where R is the ratio of valid modules on one dataset, and N is the number of all available datasets.

Simulated comparisons by changing the gray area gene expression level. In WGCNA, genes that were not assigned into any module were labeled in gray. We called these gray gene regions the gray area. The gray area represented genes whose profiles were simulated to be independent (i.e., without any correlation structure). To illustrate the impact of gray area changes on module identification and validation by TBA and SBA, we changed the gray area genes' expression levels to 0.1, 0.5, 0.9, 1.1, 1.5, and 2 times that of the original datasets to obtain simulated datasets. Based on the simulated datasets, we compared the Variation Ratio (VR, Eq. 3) of TBA (WGCNA) and SBA (pvclust) for module identification and validation.

$$VR = \frac{(\Delta X_1 + \Delta X_2 \dots \dots + \Delta X_n)}{n} \times 100\% \quad (3)$$

where ΔX is the changed number of modules or genes relative to the original data, and n is the number of the simulated datasets.

References

- Lorenz, D. M., Jeng, A. & Deem, M. W. The emergence of modularity in biological systems. *Physics of life reviews* **8**, 129–160, doi: 10.1016/j.plrev.2011.02.003 (2011).
- Newman, M. E. Modularity and community structure in networks. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 8577–8582, doi: 10.1073/pnas.0601602103 (2006).
- Ziv, E., Middendorff, M. & Wiggins, C. Information-theoretic approach to network modularity. *Physical Review E* **71**, doi: 10.1103/PhysRevE.71.046117 (2005).
- Chen, Y., Wang, Z. & Wang, Y. Spatiotemporal positioning of multipotent modules in diverse biological networks. *Cellular and molecular life sciences: CMLS* **71**, 2605–2624, doi: 10.1007/s00018-013-1547-2 (2014).
- Sridharan, G. V., Yi, M., Hassoun, S. & Lee, K. Metabolic flux-based modularity using shortest retroactive distances. *BMC systems biology* **6**, 155, doi: 10.1186/1752-0509-6-155 (2012).
- Ihmels, J., Bergmann, S., Berman, J. & Barkai, N. Comparative gene expression analysis by differential clustering approach: application to the *Candida albicans* transcription program. *PLoS genetics* **1**, e39 (2005).
- Wang, J., Li, M., Deng, Y. & Pan, Y. Recent advances in clustering methods for protein interaction networks. *BMC genomics* **11** Suppl 3, S10, doi: 10.1186/1471-2164-11-S3-S10 (2010).
- Wu, Z., Zhao, X. & Chen, L. Identifying responsive functional modules from protein-protein interaction network. *Molecules and cells* **27**, 271–277, doi: 10.1007/s10059-009-0035-x (2009).
- Dittrich, M. T., Klau, G. W., Rosenwald, A., Dandekar, T. & Muller, T. Identifying functional modules in protein-protein interaction networks: an integrated exact approach. *Bioinformatics* **24**, i223–231, doi: 10.1093/bioinformatics/btn161 (2008).
- Ulitsky, I. & Shamir, R. Identifying functional modules using expression profiles and confidence-scored protein interactions. *Bioinformatics* **25**, 1158–1164, doi: 10.1093/bioinformatics/btp118 (2009).
- Wang, X., Dalkic, E., Wu, M. & Chan, C. Gene module level analysis: identification to networks and dynamics. *Current opinion in biotechnology* **19**, 482–491, doi: 10.1016/j.copbio.2008.07.011 (2008).
- Chin, C. H., Chen, S. H., Ho, C. W., Ko, M. T. & Lin, C. Y. A hub-attachment based method to detect functional modules from confidence-scored protein interactions and expression profiles. *BMC bioinformatics* **11** Suppl 1, S25, doi: 10.1186/1471-2105-11-S1-S25 (2010).
- Zhang, W., Edwards, A., Fan, W., Zhu, D. & Zhang, K. svdPPCS: an effective singular value decomposition-based method for conserved and divergent co-expression gene module identification. *BMC bioinformatics* **11**, 338, doi: 10.1186/1471-2105-11-338 (2010).
- Li, H., Sun, Y. & Zhan, M. The discovery of transcriptional modules by a two-stage matrix decomposition approach. *Bioinformatics* **23**, 473–479, doi: 10.1093/bioinformatics/btl640 (2007).
- Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nature genetics* **25**, 25–29, doi: 10.1038/75556 (2000).
- Lysenko, A. *et al.* Assessing the functional coherence of modules found in multiple-evidence networks from Arabidopsis. *BMC bioinformatics* **12**, 203, doi: 10.1186/1471-2105-12-203 (2011).
- Li, W. *et al.* Integrative analysis of many weighted co-expression networks using tensor computation. *PLoS computational biology* **7**, e1001106, doi: 10.1371/journal.pcbi.1001106 (2011).
- Colak, R. *et al.* Module discovery by exhaustive search for densely connected, co-expressed regions in biomolecular interaction networks. *PloS one* **5**, e13348, doi: 10.1371/journal.pone.0013348 (2010).
- Minguez, P. & Dopazo, J. Assessing the biological significance of gene expression signatures and co-expression modules by studying their network properties. *PloS one* **6**, e17474, doi: 10.1371/journal.pone.0017474 (2011).
- Hou, L. *et al.* Modular analysis of the probabilistic genetic interaction network. *Bioinformatics* **27**, 853–859, doi: 10.1093/bioinformatics/btr031 (2011).
- Marras, E., Travaglione, A., Chaurasia, G., Futschik, M. & Capobianco, E. Inferring modules from human protein interactome classes. *BMC systems biology* **4**, 102, doi: 10.1186/1752-0509-4-102 (2010).
- Reimand, J., Tooming, L., Peterson, H., Adler, P. & Vilo, J. GraphWeb: mining heterogeneous biological networks for gene modules with functional significance. *Nucleic acids research* **36**, W452–459, doi: 10.1093/nar/gkn230 (2008).
- Kim, W., Li, M., Wang, J. & Pan, Y. Biological network motif detection and evaluation. *BMC systems biology* **5** Suppl 3, S5, doi: 10.1186/1752-0509-5-S3-S5 (2011).
- Hodgkinson, L. & Karp, R. M. Optimization criteria and biological process enrichment in homologous multiprotein modules. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 10872–10877, doi: 10.1073/pnas.1308621110 (2013).
- Wang, X., Castro, M. A., Mulder, K. W. & Markowetz, F. Posterior association networks and functional modules inferred from rich phenotypes of gene perturbations. *PLoS computational biology* **8**, e1002566, doi: 10.1371/journal.pcbi.1002566 (2012).
- Ozaki, S. *et al.* Coexpression analysis of tomato genes and experimental verification of coordinated expression of genes found in a functionally enriched coexpression module. *DNA research: an international journal for rapid publication of reports on genes and genomes* **17**, 105–116, doi: 10.1093/dnares/dsq002 (2010).
- Romanoski, C. E. *et al.* Network for activation of human endothelial cells by oxidized phospholipids: a critical role of heme oxygenase 1. *Circulation research* **109**, e27–41, doi: 10.1161/CIRCRESAHA.111.241869 (2011).
- Pradhan, M. P., Prasad, N. K. & Palakal, M. J. A systems biology approach to the global analysis of transcription factors in colorectal cancer. *BMC cancer* **12**, 331, doi: 10.1186/1471-2407-12-331 (2012).
- Bundy, J. G. *et al.* Evaluation of predicted network modules in yeast metabolism using NMR-based metabolite profiling. *Genome research* **17**, 510–519, doi: 10.1101/gr.5662207 (2007).
- de Jong, S. *et al.* A gene co-expression network in whole blood of schizophrenia patients is independent of antipsychotic-use and enriched for brain-expressed genes. *PloS one* **7**, e39498, doi: 10.1371/journal.pone.0039498 (2012).
- Koyuturk, M., Szpankowski, W. & Grama, A. Assessing significance of connectivity and conservation in protein interaction networks. *Journal of computational biology: a journal of computational molecular cell biology* **14**, 747–764, doi: 10.1089/cmb.2007.R014 (2007).
- He, D., Liu, Z. P. & Chen, L. Identification of dysfunctional modules and disease genes in congenital heart disease by a network-based approach. *BMC genomics* **12**, 592, doi: 10.1186/1471-2164-12-592 (2011).
- Chen, L., Xuan, J., Riggins, R. B., Wang, Y. & Clarke, R. Identifying protein interaction subnetworks by a bagging Markov random field-based method. *Nucleic acids research* **41**, e42, doi: 10.1093/nar/gks951 (2013).
- Gaire, R. K. *et al.* Discovery and analysis of consistent active sub-networks in cancers. *BMC bioinformatics* **14** Suppl 2, S7, doi: 10.1186/1471-2105-14-S2-S7 (2013).
- Langfelder, P., Luo, R., Oldham, M. C. & Horvath, S. Is my network module preserved and reproducible? *PLoS computational biology* **7**, e1001057, doi: 10.1371/journal.pcbi.1001057 (2011).
- Dong, J. & Horvath, S. Understanding network concepts in modules. *BMC systems biology* **1**, 24, doi: 10.1186/1752-0509-1-24 (2007).

37. Georgii, E., Dietmann, S., Uno, T., Pagel, P. & Tsuda, K. Enumeration of condition-dependent dense modules in protein interaction networks. *Bioinformatics* **25**, 933–940, doi: 10.1093/bioinformatics/btp080 (2009).
38. Gerlee, P., Lizana, L. & Sneppen, K. Pathway identification by network pruning in the metabolic network of *Escherichia coli*. *Bioinformatics* **25**, 3282–3288, doi: 10.1093/bioinformatics/btp575 (2009).
39. Girvan, M. & Newman, M. E. Community structure in social and biological networks. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 7821–7826, doi: 10.1073/pnas.122653799 (2002).
40. Zhang, Y., Xuan, J., de los Reyes, B. G., Clarke, R. & Ressom, H. W. Reverse engineering module networks by PSO-RNN hybrid modeling. *BMC genomics* **10** Suppl 1, S15, doi: 10.1186/1471-2164-10-S1-S15 (2009).
41. Pihur, V., Datta, S. & Datta, S. Weighted rank aggregation of cluster validation measures: a Monte Carlo cross-entropy approach. *Bioinformatics* **23**, 1607–1615, doi: 10.1093/bioinformatics/btm158 (2007).
42. Kim, Y. A., Salari, R., Wuchty, S. & Przytycka, T. M. Module cover—a new approach to genotype-phenotype studies. *Pacific Symposium on Biocomputing. Pacific Symposium on Biocomputing*, 135–146, doi: 10.1142/9789814447973_0014 (2013).
43. Rau, C. D. *et al.* Maximal information component analysis: a novel non-linear network analysis method. *Frontiers in genetics* **4**, 28, doi: 10.3389/fgene.2013.00028 (2013).
44. Dewey, F. E. *et al.* Gene coexpression network topology of cardiac development, hypertrophy, and failure. *Circulation. Cardiovascular genetics* **4**, 26–35, doi: 10.1161/CIRCGENETICS.110.941757 (2011).
45. Chin, C. H. *et al.* Spotlight: assembly of protein complexes by integrating graph clustering methods. *Gene* **518**, 42–51, doi: 10.1016/j.gene.2012.11.087 (2013).
46. Lázár, A., Ábel, D. & Vicsek, T. Modularity measure of networks with overlapping communities. *EPL (Europhysics Letters)* **90**, 18001, doi: 10.1209/0295-5075/90/18001 (2010).
47. Yona, G., Dirks, W. & Rahman, S. Comparing algorithms for clustering of expression data: how to assess gene clusters. *Methods in molecular biology* **541**, 479–509, doi: 10.1007/978-1-59745-243-4_21 (2009).
48. Hsu, J. T., Peng, C. H., Hsieh, W. P., Lan, C. Y. & Tang, C. Y. A novel method to identify cooperative functional modules: study of module coordination in the *Saccharomyces cerevisiae* cell cycle. *BMC bioinformatics* **12**, 281, doi: 10.1186/1471-2105-12-281 (2011).
49. Street, N. R., Jansson, S. & Hvidsten, T. R. A systems biology model of the regulatory network in *Populus* leaves reveals interacting regulators and conserved regulation. *BMC plant biology* **11**, 13, doi: 10.1186/1471-2229-11-13 (2011).
50. Hendrix, W. *et al.* DENSE: efficient and prior knowledge-driven discovery of phenotype-associated protein functional modules. *BMC systems biology* **5**, 172, doi: 10.1186/1752-0509-5-172 (2011).
51. Schmidt, M. C. *et al.* Efficient alpha, beta-motif finder for identification of phenotype-related functional modules. *BMC bioinformatics* **12**, 440, doi: 10.1186/1471-2105-12-440 (2011).
52. Shi, Z., Derow, C. K. & Zhang, B. Co-expression module analysis reveals biological processes, genomic gain, and regulatory mechanisms associated with breast cancer progression. *BMC systems biology* **4**, 74, doi: 10.1186/1752-0509-4-74 (2010).
53. Wen, Z., Liu, Z. P., Liu, Z., Zhang, Y. & Chen, L. An integrated approach to identify causal network modules of complex diseases with application to colorectal cancer. *Journal of the American Medical Informatics Association: JAMIA* **20**, 659–667, doi: 10.1136/amiajnl-2012-001168 (2013).
54. Wen, Z. *et al.* Identifying responsive modules by mathematical programming: an application to budding yeast cell cycle. *PLoS one* **7**, e41854, doi: 10.1371/journal.pone.0041854 (2012).
55. Sanderson, M. J. & Wojciechowski, M. F. Improved bootstrap confidence limits in large-scale phylogenies, with an example from *Neo-Astragalus* (Leguminosae). *Systematic biology* **49**, 671–685 (2000).
56. Beisser, D. *et al.* Robustness and accuracy of functional modules in integrated network analysis. *Bioinformatics* **28**, 1887–1894, doi: 10.1093/bioinformatics/bts265 (2012).
57. He, D., Liu, Z. P., Honda, M., Kaneko, S. & Chen, L. Coexpression network analysis in chronic hepatitis B and C hepatic lesions reveals distinct patterns of disease progression to hepatocellular carcinoma. *Journal of molecular cell biology* **4**, 140–152, doi: 10.1093/jmcb/mjs011 (2012).
58. Jia, P. *et al.* Network-assisted investigation of combined causal signals from genome-wide association studies in schizophrenia. *PLoS computational biology* **8**, e1002587, doi: 10.1371/journal.pcbi.1002587 (2012).
59. Sharan, R. & Ideker, T. Modeling cellular machinery through biological network comparison. *Nature biotechnology* **24**, 427–433, doi: 10.1038/nbt1196 (2006).
60. Zhang, M. & Lu, L. J. Investigating the validity of current network analysis on static conglomerate networks by protein network stratification. *BMC bioinformatics* **11**, 466, doi: 10.1186/1471-2105-11-466 (2010).
61. Zinman, G. E., Zhong, S. & Bar-Joseph, Z. Biological interaction networks are conserved at the module level. *BMC systems biology* **5**, 134, doi: 10.1186/1752-0509-5-134 (2011).
62. Kapp, A. V. & Tibshirani, R. Are clusters found in one dataset present in another dataset? *Biostatistics* **8**, 9–31, doi: 10.1093/biostatistics/kxj029 (2007).
63. Padmanabhan, K., Wang, K. & Samatova, N. F. Functional annotation of hierarchical modularity. *PLoS one* **7**, e33744, doi: 10.1371/journal.pone.0033744 (2012).
64. Hsu, C. H., Wang, T. Y., Chu, H. T., Kao, C. Y. & Chen, K. C. A quantitative analysis of monochromaticity in genetic interaction networks. *BMC bioinformatics* **12** Suppl 13, S16, doi: 10.1186/1471-2105-12-S13-S16 (2011).
65. Ma, N. & Zhang, Z. G. Evaluation of clustering algorithms for gene expression data using gene ontology annotations. *Chinese medical journal* **125**, 3048–3052 (2012).
66. King, A. D., Przulj, N. & Jurisica, I. Protein complex prediction via cost-based clustering. *Bioinformatics* **20**, 3013–3020, doi: 10.1093/bioinformatics/bth351 (2004).
67. Ma, X. & Gao, L. Discovering protein complexes in protein interaction networks via exploring the weak ties effect. *BMC systems biology* **6** Suppl 1, S6, doi: 10.1186/1752-0509-6-S1-S6 (2012).
68. Pereira-Leal, J. B., Enright, A. J. & Ouzounis, C. A. Detection of functional modules from protein interaction networks. *Proteins* **54**, 49–57, doi: 10.1002/prot.10505 (2004).
69. Sun, P. G., Gao, L. & Han, S. Prediction of human disease-related gene clusters by clustering analysis. *International journal of biological sciences* **7**, 61–73 (2011).
70. Wang, J., Ren, J., Li, M. & Wu, F. X. Identification of hierarchical and overlapping functional modules in PPI networks. *IEEE transactions on nanobioscience* **11**, 386–393, doi: 10.1109/TNB.2012.2210907 (2012).
71. Li, M., Chen, J. E., Wang, J. X., Hu, B. & Chen, G. Modifying the DPPlus algorithm for identifying protein complexes based on new topological structures. *BMC bioinformatics* **9**, 398, doi: 10.1186/1471-2105-9-398 (2008).
72. Brohee, S. & van Helden, J. Evaluation of clustering algorithms for protein-protein interaction networks. *BMC bioinformatics* **7**, 488, doi: 10.1186/1471-2105-7-488 (2006).
73. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis. *BMC bioinformatics* **9**, 559, doi: 10.1186/1471-2105-9-559 (2008).
74. Langfelder, P. *et al.* A systems genetic analysis of high density lipoprotein metabolism and network preservation across mouse models. *Biochimica et biophysica acta* **1821**, 435–447, doi: 10.1016/j.bbailip.2011.07.014 (2012).

75. Frings, O., Alexeyenko, A. & Sonnhhammer, E. L. MGclus: network clustering employing shared neighbors. *Molecular bioSystems* **9**, 1670–1675, doi: 10.1039/c3mb25473a (2013).
76. Cui, G., Shrestha, R. & Han, K. ModuleSearch: finding functional modules in a protein-protein interaction network. *Computer methods in biomechanics and biomedical engineering* **15**, 691–699, doi: 10.1080/10255842.2011.555404 (2012).
77. Pinkert, S., Schultz, J. & Reichardt, J. Protein interaction networks—more than mere modules. *PLoS computational biology* **6**, e1000659, doi: 10.1371/journal.pcbi.1000659 (2010).
78. Wang, Y. & Qian, X. A novel subgradient-based optimization algorithm for blockmodel functional module identification. *BMC bioinformatics* **14** Suppl 2, S23, doi: 10.1186/1471-2105-14-S2-S23 (2013).
79. Handl, J., Knowles, J. & Kell, D. B. Computational cluster validation in post-genomic data analysis. *Bioinformatics* **21**, 3201–3212, doi: 10.1093/bioinformatics/bti517 (2005).
80. Mar, J. C., Wells, C. A. & Quackenbush, J. Defining an informativeness metric for clustering gene expression data. *Bioinformatics* **27**, 1094–1100, doi: 10.1093/bioinformatics/btr074 (2011).
81. Liu, X., Liu, Z. P., Zhao, X. M. & Chen, L. Identifying disease genes and module biomarkers by differential interactions. *Journal of the American Medical Informatics Association: JAMIA* **19**, 241–248, doi: 10.1136/amiajnl-2011-000658 (2012).
82. Qi, J., Michoel, T. & Butler, G. An integrative approach to infer regulation programs in a transcription regulatory module network. *Journal of biomedicine & biotechnology* **2012**, 245968, doi: 10.1155/2012/245968 (2012).
83. Prieto, C., Risueno, A., Fontanillo, C. & De las Rivas, J. Human gene coexpression landscape: confident network derived from tissue transcriptomic profiles. *PloS one* **3**, e3911, doi: 10.1371/journal.pone.0003911 (2008).
84. Jin, R. *et al.* Identifying dynamic network modules with temporal and spatial constraints. *Pacific Symposium on Biocomputing. Pacific Symposium on Biocomputing*. 203–214 (2009).
85. Przytycka, T. M., Singh, M. & Slonim, D. K. Toward the dynamic interactome: it's about time. *Briefings in bioinformatics* **11**, 15–29, doi: 10.1093/bib/bbp057 (2010).
86. Li, W. *et al.* Dynamical systems for discovering protein complexes and functional modules from biological networks. *IEEE/ACM transactions on computational biology and bioinformatics/IEEE, ACM* **4**, 233–250, doi: 10.1109/TCBB.2007.070210 (2007).
87. Li, Z., Srivastava, S., Findlan, R. & Chan, C. Using dynamic gene module map analysis to identify targets that modulate free fatty acid induced cytotoxicity. *Biotechnology progress* **24**, 29–37, doi: 10.1021/bp070120b (2008).
88. Goncalves, J. P., Aires, R. S., Francisco, A. P. & Madeira, S. C. Regulatory Snapshots: integrative mining of regulatory modules from expression time series and regulatory networks. *PloS one* **7**, e35977, doi: 10.1371/journal.pone.0035977 (2012).
89. Kim, D., Kim, M. S. & Cho, K. H. The core regulation module of stress-responsive regulatory networks in yeast. *Nucleic acids research* **40**, 8793–8802, doi: 10.1093/nar/gks649 (2012).
90. Dimitrakopoulou, K. *et al.* Revealing the dynamic modularity of composite biological networks in breast cancer treatment. Conference proceedings Annual International Conference of the IEEE Engineering in Medicine and Biology Society. *IEEE Engineering in Medicine and Biology Society. Conference* **2012**, 5432–5436, doi: 10.1109/EMBC.2012.6347223 (2012).
91. Hernandez-Hernandez, V., Niklas, K. J., Newman, S. A. & Benítez, M. Dynamical patterning modules in plant development and evolution. *The International journal of developmental biology* **56**, 661–674, doi: 10.1387/ijdb.120027mb (2012).
92. Taylor, I. W. *et al.* Dynamic modularity in protein interaction networks predicts breast cancer outcome. *Nature biotechnology* **27**, 199–204, doi: 10.1038/nbt.1522 (2009).
93. Winther, R. G. Varieties of modules: kinds, levels, origins, and behaviors. *The Journal of experimental zoology* **291**, 116–129, doi: 10.1002/jez.1064 (2001).
94. Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome research* **13**, 2498–2504, doi: 10.1101/gr.1239303 (2003).
95. Assenov, Y., Ramirez, F., Schelhorn, S. E., Lengauer, T. & Albrecht, M. Computing topological parameters of biological networks. *Bioinformatics* **24**, 282–284, doi: 10.1093/bioinformatics/btm554 (2008).
96. Ruan, J. & Zhang, W. Identifying network communities with a high resolution. *Physical review. E, Statistical, nonlinear, and soft matter physics* **77**, 016104 (2008).
97. Felsenstein J. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* **39**, 783–791, (1985).
98. Shimodaira, H. An approximately unbiased test of phylogenetic tree selection. *Systematic biology* **51**, 492–508, doi: 10.1080/10635150290069913 (2002).
99. Suzuki, R. & Shimodaira, H. Pvcust: an R package for assessing the uncertainty in hierarchical clustering. *Bioinformatics* **22**, 1540–1542, doi: 10.1093/bioinformatics/btl117 (2006).
100. Kommadath, A. *et al.* Gene co-expression network analysis identifies porcine genes associated with variation in Salmonella shedding. *BMC genomics* **15**, 452, doi: 10.1186/1471-2164-15-452 (2014).
101. Tong, M. *et al.* A modular organization of the human intestinal mucosal microbiota and its association with inflammatory bowel disease. *PloS one* **8**, e80702, doi: 10.1371/journal.pone.0080702 (2013).
102. Narayanan, T., Gersten, M., Subramaniam, S. & Grama, A. Modularity detection in protein-protein interaction networks. *BMC research notes* **4**, 569, doi: 10.1186/1756-0500-4-569 (2011).

Acknowledgements

The authors' work is funded by the National Natural Science Foundation of China (90209015) and the Foundation of 'Eleventh Five' National Key Technologies R&D Programmer (2006BAI08B04-06). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

Z.W. and Y.W. designed and coordinated the study. Y.Y., Y.Z., P.W. and B.L. performed data analysis. B.L. and Y.W. wrote and modified the manuscript. All authors reviewed, edited and approved the final manuscript.

Additional Information

Accession numbers: Raw gene-expression data together with information related to demographics, disease state, and technical covariates are available via the GEO database (GEO accession number GSE24001, GSE4882, GSE2283, GSE6448, GSE6738, GSE12148, GSE12520, GSE5373, GSE5316, GSE29230, GSE5834) (<http://www.ncbi.nlm.nih.gov/geo/>).

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Li, B. *et al.* Quantitative assessment of gene expression network module-validation methods. *Sci. Rep.* **5**, 15258; doi: 10.1038/srep15258 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>