SCIENTIFIC REPORTS

OPEN

Insight into 2 α -Chloro-2'-(2',6')-(Di) Halogenopicrodophyllotoxins Reacting with Carboxylic Acids Mediated by BF₃·Et₂O

Received: 12 August 2015
Accepted: 09 October 2015
Published: 17 November 2015

Lingling Fan^{1,*}, Xiaoyan Zhi^{1,*}, Zhiping Che¹ & Hui Xu^{1,2}

Stereospecific nucleophilic substitution at the C-4 α position of 2 α -chloro-2'-(2',6')-(di) halogenopicrodophyllotoxin derivatives with carboxylic acids mediated by BF₃·Et₂O was described. Interestingly, this stereoselective products were completely controlled by the reaction time. That is, if the reaction time was prolonged to 24.5–31 h, the resulting compounds were all transformed into the unusual C-ring aromatization products. Additionally, it demonstrated that BF₃·Et₂O and reaction temperature were the important factors for C-ring aromatization, and AlCl₃ could be substituted for BF₃·Et₂O as a Lewis acid for C-ring aromatization. Halogenation of E-ring of 2 β -chloropicrodophyllotoxins with NCS or NBS also led to the same C-ring aromatization compounds. Especially compounds 5c, 6g and 7b exhibited insecticidal activity equal to that of toosendanin.

Podophyllotoxin (**1**, Fig. 1), an important naturally occurring cyclolignan isolated from the roots and rhizomes of *Podophyllum* species, has been widely used as a lead compound for preparation of potential anticancer agents such as etoposide (VP-16, **2**, Fig. 1), teniposide (VM-26, **3**, Fig. 1), etoposide phosphate, GL331 and TOP53^{1,2}. On the other hand, compound **1** also exhibited its interesting insecticidal and antifungal activities^{3–5}. To find more potent podophyllotoxin derivatives, besides total synthesis of podophyllotoxin and its derivatives^{6–11}, structural modifications of them have also attracted much attention in recent years¹². To solve the instable trans-fused lactone of podophyllotoxin when exposed to mild base, a chlorine/bromine atom was introduced at its C-2 α /3 position to give 2 α /3-halogenopicrodophyllotoxins (**4**, Fig. 1)^{13–16}. More recently, we found that once a chlorine or bromine atom was firstly introduced at the C-2' position on the E-ring of podophyllotoxin, 2 α -chloro-2'-(2',6')-(di)halogenopicrodophyllotoxins (**5**, Fig. 1) were then stereoselectively produced, and especially some 4 α -acyloxy-2 α -chloro-2'-(2',6')-(di) halogenopicrodophyllotoxins (**6**, Fig. 1) displayed more potent insecticidal activity than toosendanin, a commercial insecticide derived from *Melia azedarach*¹⁷. Additionally, during the long period of plants evolution, plant secondary metabolites are produced due to the interaction between plants and environment (life and non-life). It is considered that pesticides originating from plant secondary metabolites may result in less or slower resistance development and lower pollution^{18,19}. Recently, discovery of new insecticidal components directly from plant secondary metabolites, or by using them as the lead-compounds for structural modifications has been one of the important procedures for research and development of new pesticides^{20–23}.

In continuation of our program aimed at the discovery and development of natural-product-based insecticidal agents, we envisaged to prepare a series of 4 β -acyloxy-2 α -chloro-2'-(2',6')-(di)halogenopicrodophyllotoxins (**6'**) by S_N1 reaction of **5** with carboxylic acids in the presence of BF₃·Et₂O. To our surprise, only compounds **6** were obtained, whereas compounds **6'** were not produced at all. Moreover,

¹Research Institute of Pesticidal Design & Synthesis, College of Sciences, Northwest A&F University, Yangling 712100, Shaanxi Province, P. R. China. ²State Key Laboratory of Crop Stress Biology for Arid Areas, Northwest A&F University, Yangling 712100, Shaanxi Province, P. R. China. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to H.X. (email: orgxuhui@nwsuaf.edu.cn)

Figure 1. Chemical structures of podophyllotoxin (1) and its derivatives (2–7).

the above $\text{BF}_3 \cdot \text{Et}_2\text{O}$ -mediated reaction was completely controlled by the reaction time. If the reaction time was prolonged, compounds **6** were all transformed into the unusual C-ring aromatization compounds **7**.

Methods

Materials and Instruments. All chemical reagents and solvents were of reagent grade or purified according to standard methods before use. Analytical thin-layer chromatography (TLC) and preparative thin-layer chromatography (PTLC) were performed with silica gel plates using silica gel 60 GF₂₅₄ (Qingdao Haiyang Chemical Co., Ltd., Qingdao, China). Silica gel column chromatography was performed with silica gel 200–300 mesh (Qingdao Haiyang Chemical Co., Ltd., Qingdao, China). Melting points were determined on a XT-4 digital melting-point apparatus (Beijing Tech Instrument Co., Ltd., Beijing, China) and were uncorrected. Infrared spectra (IR) were recorded on a Bruker TENSOR 27 spectrometer. Proton nuclear magnetic resonance spectra (¹H NMR) and carbon nuclear magnetic resonance spectra (¹³C NMR) were recorded in CDCl_3 or $\text{DMSO}-d_6$ on a Bruker Avance DMX 400 or 500 MHz instrument using tetramethylsilane (TMS) as the internal standard. High-resolution mass spectra (HR-MS) were carried out with IonSpec 4.7 Tesla FTMS instrument. Compounds **5a–c** were prepared in the same way as in our previous report¹⁴.

General procedure for synthesis of 6a–l. To a mixture of **5a–c** (0.15 mmol) and carboxylic acids (0.18 mmol) in dry CH_2Cl_2 (5 mL) at -15°C , a solution of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.18 mmol) in dry CH_2Cl_2 (5 mL) was added dropwise to keep the temperature below -15°C . After adding, the reaction temperature was raised from -15°C to -5 or 0°C , and the reaction process was checked by TLC analysis. When the reaction was complete, the mixture was diluted by CH_2Cl_2 (30 mL), washed by water (20 mL), HCl (0.1 mol/L, 20 mL), 5% aq. NaHCO_3 (20 mL) and brine (20 mL), dried over anhydrous Na_2SO_4 , concentrated *in vacuo*, and purified by PTLC to give **6a–l** (40–93% yields) and **7a–c** (6–38% yields) as the white solids. The example data of **6a–c** and **7a–c** are shown as follows, whereas data of **6d–l** can be found in the Supporting Information.

Data for 6a: White solid; m.p. $150\text{--}151^\circ\text{C}$; $[\alpha]_D^{20} = -78$ (c 3.5 mg/mL, CHCl_3); IR cm^{-1} (KBr): 3050, 2936, 1786, 1735, 1487, 1399, 1109, 1016, 867; ¹H NMR (500 MHz, CDCl_3) δ : 6.68 (s, 1H, H-5), 6.64 (s, 1H, H-8), 6.58 (s, 1H, H-6'), 5.93–5.95 (m, 3H, H-4, OCH_2O), 5.52 (s, 1H, H-1), 4.80–4.81 (m, 2H, H-11), 3.94 (s, 3H, 3'- OCH_3), 3.88 (s, 3H, 5'- OCH_3), 3.76 (s, 3H, 4'- OCH_3), 2.97–2.98 (m, 1H, H-3), 2.16 (s, 3H, COCH_3); ¹³C NMR (125 MHz, CDCl_3) δ : 171.9, 170.6, 152.0, 149.9, 149.2, 148.2, 142.6, 132.1, 130.8, 123.9, 122.0, 108.8, 108.6, 108.3, 101.7, 75.1, 73.2, 66.8, 61.1, 61.1, 56.1, 49.3, 44.7, 21.1; HRMS (ESI): Calcd for $\text{C}_{24}\text{H}_{22}\text{O}_9\text{Cl}_2\text{Na}$ ($[\text{M} + \text{Na}]^+$), 547.0533; found, 547.0529.

Figure 2. Compounds 5a–c reacting with carboxylic acids in the presence of BF₃·Et₂O.

Figure 3. Compounds 5a–c reacting with carboxylic acids in the presence of BF₃·Et₂O for a prolonged time.

Data for 6b: White solid; m.p. 179–180 °C; [α]_D²⁰ = −80 (c 2.9 mg/mL, CHCl₃); IR cm^{−1} (KBr): 3057, 2935, 1788, 1724, 1486, 1400, 1108, 1033, 868; ¹H NMR (400 MHz, CDCl₃) δ : 6.67 (s, 1H, H-5), 6.62 (s, 1H, H-8), 6.57 (s, 1H, H-6'), 5.93–5.94 (m, 3H, OCH₂O, H-4), 5.51 (s, 1H, H-1), 4.80–4.81 (m, 2H, H-11), 3.93 (s, 3H, 3'-OCH₃), 3.88 (s, 3H, 5'-OCH₃), 3.75 (s, 3H, 4'-OCH₃), 2.96–2.97 (m, 1H, H-3), 2.35–2.42 (m, 2H, CH₃CH₂), 1.20 (t, *J* = 7.6 Hz, 3H, CH₃CH₂); ¹³C NMR (100 MHz, CDCl₃) δ : 174.2, 171.9, 152.1, 150.0, 149.2, 148.2, 142.8, 132.2, 130.7, 124.2, 122.1, 108.8, 108.6, 108.5, 101.7, 74.9, 73.1, 66.8, 61.1, 61.1, 56.2, 49.4, 44.7, 27.5, 9.0; HRMS (ESI): Calcd for C₂₅H₂₄O₉Cl₂Na ([M + Na]⁺), 561.0689; found, 561.0691.

Data for 6c: White solid; m.p. 79–80 °C; [α]_D¹⁹ = −66 (c 3.6 mg/mL, CHCl₃); IR cm^{−1} (KBr): 3062, 2936, 1789, 1737, 1486, 1399, 1110, 1034, 864, 697; ¹H NMR (400 MHz, CDCl₃) δ : 7.27–7.39 (m, 5H), 6.67 (s, 1H, H-5), 6.57 (s, 1H, H-8), 6.51 (s, 1H, H-6'), 5.93 (d, *J* = 1.2 Hz, 2H, OCH₂O), 5.89 (d, *J* = 3.2 Hz, 1H, H-4), 5.50 (s, 1H, H-1), 4.77–4.78 (m, 2H, H-11), 3.93 (s, 3H, 3'-OCH₃), 3.89 (s, 3H, 5'-OCH₃), 3.75 (s, 3H, 4'-OCH₃), 3.67 (d, *J* = 6.8 Hz, 2H, PhCH₂), 2.99–3.00 (m, 1H, H-3); ¹³C NMR (100 MHz, CDCl₃) δ : 171.8, 171.6, 152.0, 150.0, 149.2, 148.2, 142.8, 132.6, 132.1, 130.6, 129.0, 128.9, 127.7, 123.9, 122.1, 108.7, 108.6, 108.4, 101.7, 75.5, 72.9, 66.7, 61.1, 61.1, 56.3, 49.2, 44.6, 41.1; HRMS (ESI): Calcd for C₃₀H₂₆O₉Cl₂Na ([M + Na]⁺), 623.0846; found, 623.0849.

Data for 7a: White solid; m.p. 168–169 °C; IR cm^{−1} (KBr): 3064, 2934, 2845, 1762, 1466, 1109, 1036, 1012, 883; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 8.01 (s, 1H), 7.56 (s, 1H), 6.77 (s, 1H), 6.71 (s, 1H), 6.20 (s, 2H, OCH₂O), 5.44–5.53 (m, 2H, H-11), 3.90 (s, 3H, OCH₃), 3.87 (s, 3H, OCH₃), 3.74 (s, 3H, OCH₃); ¹³C NMR (125 MHz, DMSO-*d*₆) δ : 168.8, 151.8, 149.7, 149.1, 148.6, 142.3, 140.0, 135.2, 134.2, 129.4, 128.7, 120.1, 118.8, 118.3, 109.9, 130.7, 102.1, 101.5, 68.2, 60.9, 60.6, 56.0; HRMS (ESI): Calcd for C₂₂H₁₇O₇ClNa ([M + Na]⁺), 451.0555; found, 451.0570.

Data for 7b: White solid; m.p. 218–219 °C; IR cm^{−1} (KBr): 3052, 2942, 2911, 1745, 1467, 1207, 1010, 895; ¹H NMR (500 MHz, DMSO-*d*₆) δ : 8.05 (s, 1H), 7.58 (s, 1H), 6.75 (s, 1H), 6.22 (s, 2H, OCH₂O), 5.53 (s, 2H, H-11), 4.02 (s, 3H, 4'-OCH₃), 3.89 (s, 6H, 3', 5'-OCH₃); ¹³C NMR (125 MHz, DMSO-*d*₆) δ : 168.8, 149.9, 149.1, 148.8, 147.5, 140.1, 134.5, 132.3, 128.5, 128.2, 122.7, 120.8, 118.8, 103.9, 102.3, 100.7, 68.5, 61.2, 61.0; HRMS (ESI): Calcd for C₂₂H₁₆O₇Cl₂Na ([M + Na]⁺), 485.0165; found, 485.0170.

Data for 7c: White solid; m.p. 237–238 °C; IR cm^{−1} (KBr): 3062, 2920, 1758, 1619, 1460, 1249, 1101, 895. ¹H NMR (400 MHz, CDCl₃) δ : 7.74 (s, 1H), 7.22 (s, 1H), 6.84 (s, 1H), 6.59 (s, 1H), 6.09 (s, 2H, OCH₂O), 5.35–5.45 (m, 2H, H-11), 4.01 (s, 3H, OCH₃), 3.98 (s, 3H, OCH₃), 3.79 (s, 3H, OCH₃); ¹³C

Entry	Compound	Acid	<i>T</i> [°C]	<i>t</i> [h]	Isolated yield [%]	
					6	7
1	5a	acetic acid	−15 ~ −5	0.5	6a (75)	7a (27)
2	5a	propionic acid	−15 ~ −5	0.5	6b (47)	7a (27)
3	5a	phenylacetic acid	−15 ~ −5	0.5	6c (66)	7a (21)
4	5a	1-naphthylacetic acid	−15 ~ −5	0.5	6d (53)	7a (22)
5	5b	acetic acid	−15 ~ −5	0.5	6e (64)	7b (14)
6	5b	propionic acid	−15 ~ −5	0.5	6f (40)	7b (23)
7	5b	phenylacetic acid	−15 ~ −5	0.5	6g (78)	7b (11)
8	5b	1-naphthylacetic acid	−15 ~ 0	1	6h (50)	7b (38)
9	5c	acetic acid	−15 ~ −5	0.5	6i (93)	7c (trace)
10	5c	propionic acid	−15 ~ −5	0.5	6j (66)	7c (6)
11	5c	phenylacetic acid	−15 ~ 0	1	6k (72)	7c (27)
12	5c	1-naphthylacetic acid	−15 ~ 0	1	6l (50)	7c (29)

Table 1. Investigation of 5a-c Reacting with Carboxylic Acids Mediated by $\text{BF}_3 \cdot \text{Et}_2\text{O}$.

Figure 4. X-ray crystal structure of compound 6b. Drawing by Hui Xu.

NMR (100 MHz, CDCl_3) δ : 169.2, 152.7, 151.1, 150.1, 148.9, 143.0, 139.7, 138.4, 134.6, 131.5, 129.7, 119.6, 119.3, 110.3, 109.7, 103.8, 103.0, 101.9, 68.3, 61.3, 61.2, 56.1; HRMS (ESI): Calcd for $\text{C}_{22}\text{H}_{18}\text{O}_7\text{Br}$ ($[\text{M} + \text{H}]^+$), 473.0230; found, 473.0235.

Biological assay¹⁷. The insecticidal activity of **5a-c**, **6a-l** and **7a-c** against the pre-third-instar larvae of *Mythimna separata* Walker was assessed by leaf-dipping method. For each compound, 30 larvae (10 larvae per group) were used. Acetone solutions of compounds **5a-c**, **6a-l**, **7a-c**, and toosendanin (used as a positive control) were prepared at the concentration of 1 mg/mL. Fresh wheat leaves were dipped into the corresponding solution for 3 s, then taken out, and dried in a room. Leaves treated with acetone alone were used as a blank control group. Several treated leaves were kept in each dish, where every 10 larvae were raised. If the treated leaves were consumed, additional treated leaves were added to the dish. After 48 h, untreated fresh leaves were added to all dishes until adult emergence. The experiment was carried out at 25 ± 2 °C and relative humidity (RH) 65–80%, and on 12 h/12 h (light/dark) photoperiod.

Entry	Compound	Acid	<i>t</i> [h]	Isolated yield [%]
1	5a	acetic acid	25.5	7a (75)
2	5a	propionic acid	25.5	7a (80)
3	5a	phenylacetic acid	30.5	7a (47)
4	5a	1-naphthylacetic acid	24.5	7a (93)
5	5b	acetic acid	30.5	7b (90)
6	5b	propionic acid	28.5	7b (40)
7	5b	phenylacetic acid	24.5	7b (41)
8	5b	1-naphthylacetic acid	26	7b (91)
9	5c	acetic acid	25.5	7c (99)
10	5c	propionic acid	30.5	7c (50)
11	5c	phenylacetic acid	25	7c (78)
12	5c	1-naphthylacetic acid	31	7c (64)

Table 2. Compounds 5a–c Reacting with Carboxylic Acids Mediated by BF₃·Et₂O for a Prolonged Time.

Figure 5. X-ray crystal structure of compound 6i. Drawing by Hui Xu.

The insecticidal activity of the tested compounds against the pre-third-instar larvae of *M. separata* was calculated by the following formula:

$$\text{corrected mortality rate (\%)} = (T - C) \times 100 / (100\% - C)$$

where *T* is the mortality rate in the treated group expressed as a percentage and *C* is the mortality rate in the untreated group expressed as a percentage.

Results and Discussion

As shown in Fig. 2 and Table 1, when compounds 5a–c were allowed to react with carboxylic acids in the presence of BF₃·Et₂O at –15 °C to –5 °C for 0.5 h (or at –15 °C to 0 °C for 1 h), the expected compounds 6' were not obtained, whereas compounds 6a–l were stereoselectively afforded in 40–93% yields. Meanwhile, the corresponding by-products 7a–c, containing the aromatized C-ring, were also afforded. Interestingly, the amount of 7a–c would be increased with the advance of time; on the contrary, the amount of 6a–l would be decreased accordingly. Subsequently, compounds 5a–c reacting

Figure 6. X-ray crystal structure of compound **6j**. Drawing by Hui Xu.

Figure 7. X-ray crystal structure of compound **7a**. Drawing by Hui Xu.

with carboxylic acids in the presence of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ for a prolonged time was investigated as described in Fig. 3 and Table 2. It generally took 24.5–31 h for **6a–l** all transformed into the corresponding compounds **7a–c** in 40–99% yields. The assignment of configuration of acyloxy at the C-4 position of **6c**, **6d**, **6k** and **6l** (containing the *cis*-lactone, and an halogen atom on the E-ring) was according to our previous research results: if $J_{3,4} \approx 2.0$ Hz, it indicates that H-3 and H-4 is *trans* relationship, that is, the substituent at the C-4 position of picropodophyllotoxin is α configuration¹⁵. The $J_{3,4}$ values of H-4 of **6c**, **6d**, **6k** and **6l** were 3.2 Hz, therefore, the substituents at the C-4 position of **6c**, **6d**, **6k** and **6l** were α configuration. Because the NMR spectra of **6e–h** (bearing the *cis*-lactone, and two chlorine atoms on the E-ring) were the same as those of **6e'–h'** (compounds **6e'–h'** were prepared from **5b** in the presence of DCC and DMAP with acetic acid, propionic acid, phenylacetic acid, and 1-naphthylacetic acid, respectively. See Supporting Information), so the configuration of acyloxy at the C-4 position of **6e–h** was α . The configuration of **6b**, **6i**, **6j**, and **7a–c** was confirmed by the X-ray crystallography (Figs 4–9). Six crystallographic data (excluding structure factors) for the structures of **6b**, **6i**, **6j**, and **7a–c** have been deposited with the Cambridge Crystallographic Data Centre as supplementary

Figure 8. X-ray crystal structure of compound **7b**. Drawing by Hui Xu.

Figure 9. X-ray crystal structure of compound **7c**. Drawing by Hui Xu.

publication number CCDC 918891, 922185, 922186, 918697, 918890, and 922184, respectively. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge CB21EZ, UK [fax: + 44 (0)1223 336033 or e-mail: deposit@ccdc.cam.ac.uk]. It clearly demonstrated that the propionyloxy of **6b**, acetyloxy of **6i**, and propionyloxy of **6j** at the C-4 position all were present in α configuration. The C-ring of **7a–c** was all aromatized. Meanwhile, based on the X-ray crystallography (Figs 4–6), if the acyloxy group at the C-4 position adopted β configuration, big steric effects would be observed between the lactone and the acyloxy group. Hence the acyloxy group at the C-4 position of **6a–l** were all present in α configuration.

Furthermore, compounds **5a**, **5b** or **5c** with $\text{BF}_3 \cdot \text{Et}_2\text{O}$ in CH_2Cl_2 was also examined. As described in Fig. 10, when the reaction time was prolonged to 16–24 h, compounds **5a–c** were smoothly transformed into the corresponding compounds **7a–c** in 51–98% yields by C-ring aromatization. However, when only compound **5a** in CH_2Cl_2 was stirred at room temperature for two weeks, except **5a**, no product was produced. Additionally, when compound **5c** with $\text{BF}_3 \cdot \text{Et}_2\text{O}$ in CH_2Cl_2 was stirred at -78°C for 5.5 h, -40°C for 20 h, or -15°C for 24 h, except **5c**, the target compound **7c** was not obtained. It demonstrated that $\text{BF}_3 \cdot \text{Et}_2\text{O}$ and reaction temperature were the important factors for C-ring aromatization of **5a–c**.

Figure 10. Investigation of 5a–c mediated by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ or not.

Figure 11. Investigation of 5c mediated by AlCl_3 .

Subsequently, as shown in Fig. 11, when compound **5c** with AlCl_3 in CH_2Cl_2 was stirred at -15°C to r.t. for 24 h, compound **7c** was obtained in 77% yield. It suggested that AlCl_3 could be substituted for $\text{BF}_3 \cdot \text{Et}_2\text{O}$ as a Lewis acid for C-ring aromatization. It might involve a $\text{BF}_3 \cdot \text{Et}_2\text{O}$ -promoted dehydrochlorination, followed by a $\text{BF}_3 \cdot \text{Et}_2\text{O}$ -promoted dehydration or dehydroacylation. Meanwhile, as shown in Fig. 12, when 2 β -chloropodophyllotoxin (**8a**) in the presence of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ was stirred at -15°C to r.t. for 48 h, C-ring aromatization product **9** was obtained in 52% yield (see Supporting Information). However, 4 α -acetyloxy-2 β -chloropodophyllotoxin (**8b**) in the presence of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ was stirred at -15°C to 9°C for 24 h, or at 40°C for 48 h, the proposed product **9** was not detected.

On the other hand, as shown in Fig. 13, we envisaged whether 2 β -chloro-2'-(2',6')-(di)halogenopodophyllotoxin derivatives (**10**, the diastereoisomers of **6**), could be prepared by direct halogenation of E-ring of 2 β -chloropodophyllotoxins. As described in Table 3, the reaction of 2 β -chloropodophyllotoxins

Figure 12. Investigation of **8a** and **8b** mediated by $\text{BF}_3 \cdot \text{Et}_2\text{O}$.

(**8a–c**) with NCS or NBS was further examined. When compound **8a** reacted with 1.1 equiv of NCS at 25°C for 24 h, no product was obtained (entry 1). When the reaction temperature was raised to 30°C , the target E-ring halogenation product was not produced; interestingly, whereas the C-ring aromatization compounds such as **7a** and **7b** were obtained in 35% and 15% yields, respectively (entry 2). And when at 40°C for 4 h, compounds **7a** and **7b** were obtained in 54% and 12% yields, respectively (entry 3). However, when the reaction temperature was raised to 50°C , the yields of **7a** and **7b** were not increased (entry 4). When compound **8a** reacted with 2.2 equiv of NCS at 40°C for 11 h, only **7b** was obtained in 52% yield (entry 5). Similarly, when compound **8a** reacted with 1.1 equiv of NBS at 40°C for 5 h, compound **7c** was obtained in 50% yield (entry 7); when compound **8c** reacted with 1.1 equiv of NCS at 40°C for 19 h, compounds **7a** and **7b** were obtained in 34% and 25% yields, respectively (entry 12). But when compound **8b** reacted with NCS or NBS, the reaction temperature should be raised. For example, when compound **8b** reacted with 1.1 equiv of NCS at 40°C for 48 h, compound **7a** was obtained only in 12% yield (entry 8); whereas the reaction temperature was raised to 60°C for 24 h, compound **7a** was obtained in 94% (entry 9). When compound **8b** reacted with 2.2 equiv of NCS for 17 h or 1.1 equiv of NBS for 24 h at 60°C , compounds **7b** and **7c** were obtained in 87% and 91% yields, respectively (entries 10 and 11). All in all, the reaction temperature was very important for 2 β -chloropodophyllotoxins reacting with NCS or NBS to give **7a–c**. According to our previous results¹⁴, reaction of **8a–c** with NCS or NBS might involve E-ring halogenation of 2 β -chloropodophyllotoxins, followed by the dehydrochlorination and dehydration or dehydroacylation.

Finally, compounds **5a–c**, **6a–l**, and **7a–c** were evaluated as insecticidal agents against the pre-third-instar larvae of oriental armyworm, *Mythimna separata* (Walker) at the concentration of 1 mg/mL. Among **5a–c** and **6a–l**, as described in our previous paper¹⁷, only compounds **5c** and **6g** exhibited insecticidal activity equal to that of toosendanin, and the final corrected mortality rates of **5c**, **6g** and toosendanin were 51.9%, 55.6% and 51.9%, respectively. The final corrected mortality rates of **1** and **7a–c** were 37%, 40.7%, 51.9%, and 37%, respectively.

Conclusion

In summary, we have developed a $\text{BF}_3 \cdot \text{Et}_2\text{O}$ -mediated stereoselective synthesis 4 α -acyloxy-2 α -chloro-2'-(2',6')-(di)halogenopodophyllotoxin derivatives, which was completely controlled by the reaction time. If the reaction time was prolonged to 24.5–31 h, the target compounds were all transformed into the C-ring aromatization compounds. Additionally, it demonstrated that $\text{BF}_3 \cdot \text{Et}_2\text{O}$ and reaction temperature were the important factors for C-ring aromatization, and AlCl_3 could be substituted for $\text{BF}_3 \cdot \text{Et}_2\text{O}$ as a Lewis acid for C-ring aromatization. However, in the presence of NCS or NBS, 2 β -chloropodophyllotoxins could also be transformed into the same C-ring aromatization compounds. Notably, compounds **5c**, **6g** and **7b** exhibited insecticidal activity equal to that of toosendanin.

Figure 13. 2β-Chloropodophyllotoxins (8a–c) reacting with NCS or NBS.

Entry	Compound	Amount of NCS or NBS	T [°C]	t [h]	Isolated yield [%]
1	8a	NCS (1.1 equiv)	25	24	7a (0)
2	8a	NCS (1.1 equiv)	30	7	7a (35) + 7b (15)
3	8a	NCS (1.1 equiv)	40	4	7a (54) + 7b (12)
4	8a	NCS (1.1 equiv)	50	4	7a (40) + 7b (7)
5	8a	NCS (2.2 equiv)	40	11	7b (52)
6	8a	NBS (1.1 equiv)	25	24	7c (0)
7	8a	NBS (1.1 equiv)	40	5	7c (50)
8	8b	NCS (1.1 equiv)	40	48	7a (12)
9	8b	NCS (1.1 equiv)	60	24	7a (94) + 7b (trace)
10	8b	NCS (2.2 equiv)	60	17	7b (87)
11	8b	NBS (1.1 equiv)	60	24	7c (91)
12	8c	NCS (1.1 equiv)	40	19	7a (34) + 7b (25)

Table 3. Investigation of 2β-Chloropodophyllotoxins (8a–c) Reacting with NCS or NBS.

References

1. You, Y. J. Podophyllotoxin derivatives: Current synthetic approaches for new anticancer agents. *Curr. Pharm. Des.* **11**, 1695–1717 (2005).
2. Gordaliza, M., Garcia, P. A., Miguel Del Corral, J. M., Castro, M. A. & Gomez-Zurita, M. A. Podophyllotoxin: distribution, sources, applications and new cytotoxic derivatives. *Toxicon* **44**, 441–459 (2004).
3. Kumar, K. A., Singh, S. K., Kumar, B. S. & Doble, M. Synthesis, anti-fungal activity evaluation and QSAR studies on podophyllotoxin derivatives. *Cent. Eur. J. Chem.* **5**, 880–897 (2007).
4. Miyazawa, M., Fukuyama, M., Yoshio, K., Kato, T. & Ishikawa, Y. Biologically active components against *Drosophila melanogaster* from *Podophyllum hexandrum*. *J. Agric. Food Chem.* **47**, 5108–5110 (1999).
5. Xu, H., Lv, M. & Tian, X. A review on hemisynthesis, biosynthesis, biological activities, mode of action, and structure-activity relationship of podophyllotoxins: 2003–2007. *Curr. Med. Chem.* **16**, 327–349 (2009).
6. Stadler, D. & Bach, T. Concise stereoselective synthesis of (–)-podophyllotoxin by an intermolecular iron (III)-catalyzed Friedel-Crafts alkylation. *Angew. Chem. Int. Ed.* **47**, 7557–7559 (2008).
7. Engelhardt, U., Sarkar, A. & Linker, T. Efficient enantioselective total synthesis of (–) epipodophyllotoxin. *Angew. Chem. Int. Ed.* **42**, 2487–2489 (2003).
8. Reynolds, A. J., Scott, A. J., Turner, C. I. & Sherburn, M. S. The intramolecular carboxyarylation approach to podophyllotoxin. *J. Am. Chem. Soc.* **125**, 12108–12109 (2003).
9. Jeedimalla, N., Johns, J. & Roche, S. P. Mechanistic investigation and implications of a sacrificial aniline for the tandem cascade synthesis of 4-aza-podophyllotoxin analogues. *Tetrahedron Lett.* **54**, 5845–5848 (2013).
10. Merchan Arenas, D. R., Rojas Ruiz, F. A. & Kouznetsov, V. V. Highly diastereoselective synthesis of new heterolignan-like 6,7-methylenedioxy-tetrahydroquinolines using the clove bud essential oil as raw material. *Tetrahedron Lett.* **52**, 1388–1391 (2011).
11. Ting, C. P. & Maimone, T. J. C-H bond arylation in the synthesis of aryltetralin lignans: A short total synthesis of podophyllotoxin. *Angew. Chem. Int. Ed.* **53**, 3115–3119 (2014).
12. Lv, M. & Xu, H. Recent advances in semisynthesis, biosynthesis, biological activities, mode of action, and structure-activity relationship of podophyllotoxins: an update (2008–2010). *Mini-Rev. Med. Chem.* **11**, 901–909 (2011).
13. Glinski, M. B., Freed, J. C. & Durst, T. Preparation of 2-substituted podophyllotoxin derivatives. *J. Org. Chem.* **52**, 2749–2753 (1987).
14. Che, Z. P., Yu, X., Zhi, X. Y., Fan, L. L. & Xu, H. Synthesis of novel 4α-(acyloxy)-2′(2′,6′)-(di)halogenopodophyllotoxin derivatives as insecticidal agents. *J. Agric. Food Chem.* **61**, 8148–8155 (2013).
15. Xu, H., Wang, Q. T. & Guo, Y. Stereoselective synthesis of 4α-alkyloxy-2-α/β-bromopodophyllotoxin derivatives as insecticidal agents. *Chem. Eur. J.* **17**, 8299–8303 (2011).

16. Xu, H., Xiao, X. & Wang, Q. T. Natural products-based insecticidal agents 7. Semisynthesis and insecticidal activity of novel 4 α -alkyloxy-2-chloropodophyllotoxin derivatives against *Mythimna separata* Walker *in vivo*. *Bioorg. Med. Chem. Lett.* **20**, 5009–5012 (2010).
17. Fan, L. L., Guo, Y., Zhi, X. Y., Y, X. & Xu, H. Stereoselective synthesis of 2 α -chloropicropodophyllotoxins and insecticidal activity of their esters against oriental armyworm, *Mythimna separata* Walker. *J. Agric. Food Chem.* **62**, 3726–3733 (2014).
18. Isman, M. B. Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annu. Rev. Entomol.* **51**, 45–66 (2006).
19. Seiber, J. N., Coats, J., Duke, S. O. & Gross, A. D. Biopesticides: state of the art and future opportunities. *J. Agric. Food Chem.* **62**, 11613–11619 (2014).
20. Caboni, P. *et al.* Nematicidal activity of mint aqueous extracts against the root-knot nematode *meloidogyne incognita*. *J. Agric. Food Chem.* **61**, 9784–9788 (2013).
21. Meepagala, K. M., Bernier, U. R., Burandt, C. & Duke, S. O. Mosquito repellents based on a natural chromene analogue with longer duration of action than *N,N*-diethyl-meta-toluamide (DEET). *J. Agric. Food Chem.* **61**, 9293–9297 (2013).
22. Oliver, M. P., Crouse, G. D., Demeter, D. A. & Sparks, T. C. Synthesis and insecticidal activity of spinosyns with C9-O-benzyl bioisosteres in place of the 2',3',4'-tri-O-methyl rhamnose. *J. Agric. Food Chem.* **63**, 5571–5577 (2015).
23. Gerwick, B. C. & Sparks, T. C. Natural products for pest control: an analysis of their role, value and future. *Pest Manag. Sci.* **70**, 1169–1185 (2014).

Acknowledgements

The present research was supported by National Natural Science Foundation of China (No. 31071737), and the Special Funds of Central Colleges Basic Scientific Research Operating Expenses, Northwest A&F University (No. YQ2013008, 2452015096).

Author Contributions

L.F. and X.Z. performed experiments, and analysed data; C.Z. analysed data; H.X. designed experiments, analysed data and wrote the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Fan, L. *et al.* Insight into 2 α -Chloro-2'-(2',6')-(Di) Halogenopicropodophyllotoxins Reacting with Carboxylic Acids Mediated by BF₃·Et₂O. *Sci. Rep.* **5**, 16285; doi: 10.1038/srep16285 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>