SCIENTIFIC REPORTS

OPEN

Elasticity of Ferropericlasite across the Spin Crossover in the Earth's Lower Mantle

Received: 29 July 2015

Accepted: 26 October 2015

Published: 01 December 2015

Jing Yang¹, Xinyue Tong¹, Jung-Fu Lin^{1,2}, Takuo Okuchi³ & Naotaka Tomioka³

Knowing the elasticity of ferropericlasite across the spin transition can help explain seismic and mineralogical models of the lower-mantle including the origin of seismic heterogeneities in the middle to lowermost parts of the lower mantle^{1–4}. However, the effects of spin transition on full elastic constants of ferropericlasite remain experimentally controversial due to technical challenges in directly measuring sound velocities under lower-mantle conditions^{1–5}. Here we have reliably measured both V_p and V_s of a single-crystal ferropericlasite ($(\text{Mg}_{0.92}\text{Fe}_{0.08})\text{O}$) using complementary Brillouin Light Scattering and Impulsive Stimulated Light Scattering coupled with a diamond anvil cell up to 96 GPa. The derived elastic constants show drastically softened C_{11} and C_{12} within the spin transition at 40–60 GPa while C_{44} is not affected. The spin transition is associated with a significant reduction of the aggregate V_p/V_s via the aggregate V_p softening because V_s softening does not visibly occur within the transition. Based on thermoelastic modelling along an expected geotherm, the spin crossover in ferropericlasite can contribute to 2% reduction in V_p/V_s in a pyrolite mineralogical model in mid lower-mantle. Our results imply that the middle to lowermost parts of the lower-mantle would exhibit enhanced seismic heterogeneities due to the occurrence of the mixed-spin and low-spin ferropericlasite.

Seismic wave studies of the lower mantle have established relatively reliable seismic models including compressional and shear wave velocities (V_p and V_s) in one-, two-, and three-dimensional tomographic models (e.g., PREM, AK135, S40RTS)^{6–8}. Thus far, it has been shown that most of the lower mantle, except the lowermost mantle such as the D'' layer, exhibit relatively smooth changes in seismic parameters that have been commonly interpreted as a result of the high pressure-temperature (P - T) effects on physical properties of candidate minerals in the region. The lower mantle is mostly believed to be seismically and chemically homogeneous and likely consisting of approximately 75% bridgmanite (silicate perovskite $(\text{Mg,Fe})(\text{Al,Fe,Si})\text{O}_3$; Pv), 20% ferropericlasite $(\text{Mg,Fe})\text{O}$; Fp), and 5% calcium perovskite (CaSiO_3) by volume in a pyrolite compositional model^{9,10}. However, a number of seismic studies have shown that possible thermal and/or chemical heterogeneities, especially in the middle to lower part of the lower mantle ranging from approximately 1500 km to 2800 km in depth, are needed to reconcile differences between our current understanding of seismic models and mineral physics results^{11–15}. Other than the bridgmanite to post-perovskite structural transition at the D'' zone region¹⁶, which may be responsible for the seismic discontinuities in the lowermost mantle, the electronic spin transitions of iron in lower-mantle minerals have been suggested to affect our understanding of mid to lowermost lower-mantle seismic heterogeneities.

The electronic spin transition of iron in lower-mantle bridgmanite and ferropericlasite has been recently reported to affect physical and chemical properties of the host minerals, including changes in elasticity, iron partitioning, and electrical and thermal conductivities^{1,4,17–22}, that may contribute to

¹Department of Geological Sciences, Jackson School of Geosciences, The University of Texas at Austin, Austin, TX 78712, USA. ²Center for High Pressure Science and Technology Advanced Research (HPSTAR). ³Institute for Study of the Earth's Interior, 827 Yamada, Misasa, Tottori, 682-0193 Japan. Correspondence and requests for materials should be addressed to J.F.L. (email: afu@jsg.utexas.edu)

seismic heterogeneities of the region. It has been shown that a broad spin crossover occurs in ferropericlase at conditions ranging from 1700 km to 2700 km in depth²³, while the Fe^{3+} in the octahedral site of bridgmanite undergoes a high-spin to low-spin transition at P - T conditions relevant to the top lower mantle³. Of particular interest to our understanding of deep-mantle seismology and geodynamics are the effects of the spin transition on the elasticity (e.g., sound velocities, equation of states (EoS), and seismic anisotropies) of lower-mantle minerals, because a thorough knowledge of their elastic properties is essential for interpreting seismic observations as well as for constraining the chemical composition and mineralogy of the region¹⁴.

In recent years, there have been a number of experimental and theoretical studies on the elasticity of single-crystal ferropericlase across the spin transition at high pressures using various techniques, including Impulsive Stimulated Light Scattering (ISS), Brillouin Light Scattering (BLS), Inelastic X-ray Scattering (IXS), and Density Function Theory (DFT)^{1,3,4,24,25}. DFT calculations have shown that ferropericlase exhibits significant softening in V_p as well as in the C_{11} and C_{12} elastic constants across the spin crossover at lower-mantle P - T conditions, although V_s and C_{44} are not affected by the transition²⁵. On the other hand, experimental results on the elasticity across the spin transition differ drastically^{1–4}. In particular, ISS measurements on $(\text{Mg}_{0.94}\text{Fe}_{0.06})\text{O}$ up to 60 GPa showed a remarkable reduction in both V_p and V_s across the spin transition¹, although the reliability of deriving the V_s from the interfacial wave has been questioned. BLS measurements on $(\text{Mg}_{0.9}\text{Fe}_{0.1})\text{O}$ up to 81 GPa showed no substantial reduction of the directly-measured V_s across the spin transition, while the combination of the V_s values from BLS and the EoS parameters from X-ray diffraction have shown a V_p softening up to 17% within the spin transition². High-pressure IXS experiments up to 80 GPa on $(\text{Mg}_{0.83}\text{Fe}_{0.17})\text{O}$, which used acoustic phonon dispersions at very high frequencies to extract the velocities, did not reveal any reduction of either V_p or V_s within the spin transition³. Furthermore, some of these previous studies have indicated that the spin transition can markedly enhance the elastic V_s splitting anisotropy of ferropericlase such that the low-spin state becomes much more elastically anisotropic as compared to its high-spin counterpart^{3,4}. The full elastic constants of single-crystal ferropericlase across the spin transition and in the low-spin state can help provide new insight into the effects of the spin transition on thermodynamic and seismic parameters of the sample, but these essential experimental data remain unclear due to the aforementioned technical difficulties. Since different experimental results point to entirely different scenarios for seismic and geochemical models of the lower mantle, seismically homogeneous vs. heterogeneous lower mantle, it remains unclear if the effect of the spin transition on the elasticity of the lower-mantle ferropericlase should be taken into account in our understanding of the seismic models of the lower mantle. Reliable experimental results can also help elucidate recent theoretical predictions of the elasticity and thermodynamics of ferropericlase at high pressures²⁵.

Here we have directly measured V_p , V_s , and the pressure-volume (P - V) relation of a single-crystal ferropericlase $(\text{Mg}_{0.92}\text{Fe}_{0.08})\text{O}$ in order to solve for its full elastic constants (C_{ij}) using combined BLS, ISS, and X-ray diffraction (XRD) measurements in a diamond anvil cell (DAC) up to 96 GPa (see Methods for details). The combined experimental results overcome previous technical difficulties and permit direct and reliable evaluation of the full elastic constants and thermoelastic parameters across the spin transition at lower-mantle pressures. Our results show that ferropericlase with 8 at% iron undergoes a spin transition at 40–60 GPa that is associated with changes in elastic constants: C_{11} softening by a maximum of 16%, and C_{12} by 70%, whereas C_{44} does not show any observable reduction across the transition. Within the spin transition, the aggregate K_s and V_p calculated from Voigt-Reuss-Hill average of the elastic constants reduces by 38% and 13%, respectively, and the aggregate V_s is not visibly affected by the spin transition. The V_p anisotropy changes to a maximum value of 11% (± 0.7) at 50 GPa, which is midway between the spin transition as compared to 9% (± 0.2) in the V_p anisotropy for the extrapolated high-spin state. The low-spin state also exhibits some elastic behavior distinct from that of the high-spin and mixed-spin states. To decipher the geophysical and geochemical consequences of the spin crossover in the deep mantle, we have modelled elastic and seismic parameters of ferropericlase along an expected lower-mantle geotherm²⁶. Our results show that the velocity abnormalities and elastic softening remain significantly strong across the spin crossover in the lower mantle and that the low-spin ferropericlase exhibits significantly enhanced V_p and V_s profiles from that of the extrapolated high-spin state. These results are applied to understand potential seismic and/or chemical heterogeneities induced by the spin transition in the deep lower mantle.

Results and Discussion

Experiments and Thermoelastic Modelling. P - V relations of single-crystal ferropericlase $(\text{Mg}_{0.92}\text{Fe}_{0.08})\text{O}$ in the (100) platelet were measured using synchrotron X-ray diffraction up to 91 GPa at room temperature in a DAC. These results are used to evaluate the EoS parameters, the width of the spin transition, and the fraction of the high-spin (HS) and low-spin (LS) states in ferropericlase^{20,23} (Fig. S1 and Fig. S2) (See Methods and SI for details of the experiments and modelling). Analysis of the measured P - V curve shows that the spin transition occurs over pressures ranging between 40 GPa and 60 GPa, and is associated with a density increase of 1.2% (0.1%). The derived isothermal bulk modulus at ambient conditions (K_{T0}) and its pressure derivative (K'_{T0}) are: $K_{T0} = 152.5$ (2.4) and $K'_{T0} = 4.1$ (0.2) for the HS state, and $K_{T0} = 161.6$ (7.1) with a fixed K'_{T0} of 4 for the LS state, consistent with previous studies²⁷ (Fig. S3).

Figure 1. Representative Brillouin light scattering (BLS), impulsive stimulated scattering (ISS), and power spectra of the single-crystal ferropericlasite ($\text{Mg}_{0.92}\text{Fe}_{0.08}\text{O}$) along [110] crystallographic axis at high pressures. The BLS spectra were used to derive the V_s , while the V_p was detected in the ISS spectra. The ISS spectra in the time domain were analyzed and Fourier-transformed to the power spectra in the velocity (frequency) domain to derive the acoustic waves of the sample at high pressures. Neon medium was also observed in the ISS spectra at pressures up to approximately 70 GPa.

The single-crystal platelet was also used for simultaneously measuring V_s in the BLS experiments and V_p in the ISS experiments along principle [100] and [110] crystallographic axes up to 96 GPa in the Mineral Physics Laboratory of The University of Texas at Austin (Figs 1 and 2); at relatively lower pressures, the V_p and V_s velocities of the platelet were also measured as a function of the azimuthal angles in order to confirm the orientation of the platelets and to further assure the reliability of our measurements as compared with previous studies (Fig. S4). Together with P - V results from XRD measurements, the measured V_p and V_s velocities of single-crystal ferropericlasite permit direct derivations of the full elastic constants (C_{11} , C_{12} , C_{44}) at high pressures via Christoffel's equations (Fig. 2). Using the Eulerian finite-strain theory²⁸ and a thermoelastic model for the cubic system²⁵, we have modelled the elastic constants within the spin transition using formulations reported previously²⁵ (See SI for details). Specifically, the elastic compliances S_{ij} of the crystal across the spin transition are given by:

$$S^{ij}V = n_{LS}S_{LS}^{ij}V_{LS} + (1 - n_{LS})S_{HS}^{ij}V_{HS} - \left(\frac{\partial G_{LS}}{\partial \sigma_j} - \frac{\partial G_{HS}}{\partial \sigma_j} \right) \frac{\partial n_{LS}}{\partial \sigma_i}, \quad (1)$$

where V is the volume, n_{LS} is the LS fraction, σ_i and σ_j are the i th and j th stress component, respectively, in the Voigt notation, and G is the Gibbs free energy. In this modelling, the low-spin fraction (n_{LS}) and the unit cell volume (V) derived from the equation of state, and the elastic constants (C_{ij}) are used to constrain the elastic compliances S_{ij} according to the relationship between C_{ij} and S_{ij} (See SI for details). To further obtain the pressure-dependent EoS parameters for the HS and LS states, respectively, the elastic constants and the aggregate bulk and shear moduli as a function of pressure are derived by fitting the results to the third-order Eulerian finite-strain equation of state (Figs 2 and 3).

Examination of the directly-measured V_p and V_s velocities show that the V_p softens by $\sim 10\%$ maximum in both the [100] and [110] directions within the spin transition, while the V_s along the [100] is slightly enhanced, but the V_s along the [110] is not noticeably affected (Fig. 2A) (see SI for comparison with previous results). Furthermore, the C_{11} and C_{12} elastic constants are significantly softened by a maximum of 16% and 70%, respectively, across the spin transition, but C_{44} is not affected by the spin transition (Fig. 2B). The maximum softening for these parameters occurs at approximately 50 GPa, which is midway within the spin transition where the fraction of the LS state is about 50% (Fig. S2). As reported in previous theoretical calculations²⁵, the C_{11} and C_{12} softening can be explained by the addition of an energy abnormality as a result of the HS and LS mixing (shown in the last term in Equation (1)). Since $n(\sigma_4)$ is an even function and $\partial n / \partial \sigma_4|_{\sigma_4=0} = 0$, the last term in Equation (1) vanishes such that the C_{44} softening is not expected to occur across the spin crossover²⁵. Our results thus confirm theoretical predictions of the elasticity across the spin transition in ferropericlasite at high pressures. The derived elastic constants are used to calculate V_{s1} , V_{s2} , and V_p velocities as a function of propagation directions ($\mathbf{n}$) by

Figure 2. Elasticity of single-crystal ferropericlase ($\text{Mg}_{0.92}\text{Fe}_{0.08}\text{O}$) as a function of pressure at 300 K. (A) Compressional and shear wave velocities along the [100] and [110] crystallographic axes as a function of pressure. Compressional wave velocities were measured using the ISS technique, while shear wave velocities were measured using the BLS technique. Open circles: experimental data; solid lines: modelled velocity profiles using thermoelastic equations (see SI for details). (B) Elastic constants (C_{ij}) as a function of pressure. Open circles: C_{ij} directly derived from measured compressional and shear wave velocities via Christoffel's equations; solid lines: modelled C_{ij} profiles. Vertical dashed lines are plotted to guide the eyes for the high-spin (HS), mixed-spin (MS; HS + LS), and low-spin (LS) regions, respectively (see Supplementary Fig. S2 for details).

Figure 3. Aggregate bulk and shear moduli K_s and G , aggregate velocities, elastic anisotropies and aggregate V_p/V_s ratio of ferropericlase ($\text{Mg}_{0.92}\text{Fe}_{0.08}\text{O}$) at high pressure and 300 K. (A) Adiabatic bulk and shear modulus from Voigt-Reuss-Hill average; (B) Aggregate compressional V_p and shear wave velocities V_s , where $V_p = \sqrt{K_s + 4/3G/\rho}$ and $V_s = \sqrt{G/\rho}$; (C) Compressional and shear wave anisotropy as a function of pressure; dashed lines are the extrapolated anisotropies for the HS state that are plotted for comparison; (D) aggregate V_p/V_s ratio. Grey shaded areas represent uncertainties calculated from standard error propagations using the experimentally derived elastic constants. Vertical dashed lines are plotted to guide the eyes for the HS, MS, and LS regions, respectively.

solving Christoffel's equation $\det|C_{ijkl}n_jn_l - \rho V^2\delta_{ik}| = 0$ (Fig. S7). At ambient conditions, the single-crystal ferropericline exhibits the slowest V_p in the [100] direction and the fastest V_p in the [111] direction, while the V_s is slowest in the [110] direction and fastest in the [100] direction. This anisotropic behavior is reversed at pressures of approximately 21 GPa, above which the V_p minimum and maximum are along the [111] and [100], respectively, while the V_s minimum and maximum are along the [100] and [110] directions. The velocities vary significantly with the propagation direction at pressures above approximately 40 GPa, indicating that single-crystal ferropericline exhibits strong V_p and V_s anisotropy. The velocity anisotropy factor (A) is defined as $A = (V_{\max} - V_{\min})/2(V_{\max} + V_{\min}) \times 100\%$, where $V_{\max}$ and $V_{\min}$ are the maximum and minimum velocities, respectively²⁹. For V_p , the anisotropy is defined as the difference between the maximum and minimum velocities in corresponding propagation directions, while the V_s splitting anisotropy is defined as the velocity difference between the maximum and minimum velocities for two orthogonally polarized V_s velocities along corresponding propagation directions. Our results show that the V_p anisotropy and the V_s splitting anisotropy are 11.5% and 23.5% at ambient conditions, decreasing to almost zero at approximately 21 GPa, and then increasing monotonically up to 40 GPa. The V_p anisotropy increases to ~11% at 50 GPa, which is midway between the spin transition, while the anisotropy of the extrapolated HS state is approximately ~9%. Furthermore, the V_s splitting anisotropy continuously increases with increasing pressure but slightly deviates from the extrapolated HS state counterpart starting with the spin transition (Fig. 3C and Fig. S8). Within the experimental uncertainties in this study, the effect of the spin transition on the V_s splitting anisotropy can be considered negligible (Fig. 3C). Contrary to previous studies⁴, our results show that the spin transition is actually associated with a slightly enhanced V_p anisotropy (Fig. S8). Furthermore, our results clearly show that the LS ferropericline exhibits some unique elastic behavior distinct from that of the HS state, including an enhanced pressure derivative of C_{11} , C_{12} , and C_{44} , implying an enhanced pressure dependence of aggregate K_s , V_p , and G (Figs 2 and 3A,B; Table S2). In particular, the spin transition is associated with a significant reduction of the aggregate V_p/V_s ratio via the aggregate V_p softening since V_s softening does not visibly occur within the spin transition; this ratio is reduced from 1.75 at approximately 40 GPa at the onset of the transition to 1.6 at approximately 50 GPa midway within the transition (Fig. 3B,D). That is, the LS state exhibits a maximum of approximately 12% reduction in the V_p/V_s ratio as compared to that of the extrapolated HS state within the spin transition. Such a reduction in the V_p/V_s ratio manifests an abnormal Poisson's ratio within the spin transition and into the LS state.

Elasticity of ferropericline along an expected geotherm: implication for lower-mantle seismic heterogeneities.

To understand the effects of the spin transition on the elasticity of ferropericline at relevant P - T conditions of the lower mantle^{22,30}, we have modelled the elastic constants of ferropericline along an expected lower-mantle geotherm up to approximately 125 GPa using thermoelastic models and a previously reported spin crossover diagram^{23,25}. The geotherm profile at the core-mantle boundary conditions has not been considered in our modelling here. The thermal EoS parameters of ferropericline with 25 at% iron in a previous experimental report are linearly scaled back for the compositional effects of FeO in MgO to construct the spin crossover diagram for our ferropericline with 8 at% iron at high P - T (See SI for details). Our modelled results show that the spin crossover of ferropericline with 8 at% iron occurs between 65 and 105 GPa along the geotherm. The temperature derivatives of the elastic constants for pure MgO³¹ are combined with our high-pressure elasticity results (Table S1 and Table S2) to account for the high P - T effects of FeO solid solution for the HS and LS states (Fig. S9). These modelled results show that the effects of the spin crossover on the elastic and seismic parameters along an expected adiabatic geotherm remain profound, even though the spin crossover is broadened by high temperatures (Fig. 4 and Fig. S9). In particular, C_{11} and C_{12} exhibit 15% and 60% maximum reduction, respectively, within the spin crossover at approximately 85 GPa that corresponds to 1900 km in depth. The V_p anisotropy increases to 18.6% at ~85 GPa within the spin transition (a 23% increase in the anisotropy as compared to the extrapolated HS state reference), while the pressure-dependent V_s anisotropy is lower than that of the HS counterpart in the LS state (Fig. S9 B). Compared to the HS state reference, the aggregate V_p decreases by 10% while the V_p/V_s ratio drops by 13% within the spin crossover (Fig. 4). On the other hand, the aggregate V_p and V_s profiles of the LS ferropericline are significantly higher than that of their HS state counterparts (Fig. 4A). Using the HS state as the reference, we have calculated the deviations of a number of seismic parameters across the spin crossover along an expected mantle geotherm. These results show that the spin crossover produces V_p and V_s velocities and anisotropies, V_p/V_s ratio, and Poisson's ratio that vary as a function of the low-spin fraction as compared to the extrapolated HS state. Specifically, the V_p , V_p/V_s ratio, and Poisson's ratio are significantly reduced within the spin crossover, whereas the LS state exhibits enhanced V_p and V_s velocities as well as reduced V_p/V_s and Poisson's ratio (Fig. S10).

Our modelled velocity profiles show that the V_p profile of ferropericline with 8 at% iron is significantly reduced by a maximum of 10% within the spin crossover at approximately 1900 km in depth along an expected mantle geotherm, while the V_s profile is slightly enhanced with increasing fraction of the LS state as compared to the extrapolated HS state (Fig. 4). That is, the V_p/V_s ratio is also significantly reduced with the spin crossover. On the other hand, the V_p and V_s profiles of the LS ferropericline are higher than those of the extrapolated HS state as well as the PREM model toward the lower parts of

Figure 4. Modelled seismic velocities and V_p/V_s ratio of ferropericlase ($(\text{Mg}_{0.92}\text{Fe}_{0.08})\text{O}$) along an expected lower-mantle geotherm. (A) Aggregate compressional and shear wave velocities. These results are calculated from the single-crystal elastic constants. (B) Calculated V_p/V_s ratio. Solid lines: modelled seismic parameters with uncertainties shown as grey areas; black dashed lines: modelled parameters for the high-spin state; green dotted lines: theoretical V_p and V_s values of ferropericlase ($(\text{Mg}_{0.875}\text{Fe}_{0.125})\text{O}$)²⁵; crosses: experimental results with 17% iron¹⁴. PREM seismic parameters are plotted as blue dotted dashed lines for comparison⁶; The magenta lines are the modelled velocity profiles assuming that the lower mantle is composed of 20% ferropericlase (fp8) and 80% bridgmanite (Pv)¹⁴.

the lower mantle. Considering that ferropericlase in the lower mantle may contain approximately 20% FeO and account for approximately 20 vol.% of the lower-mantle in a pyrolite model⁹, our calculated V_p and V_s profiles with 20 vol.% of ferropericlase and 80 vol.% of bridgmanite are fairly consistent with the PREM model at depths from uppermost to mid lower-mantle. On the other hand, the spin crossover in ferropericlase is expected to contribute a few percent reduction in V_p and V_p/V_s ratio in the middle parts of the lower mantle compared to the PREM seismic model, while the occurrence of the LS ferropericlase would result in enhanced V_p and V_s profiles compared to the seismic model (Fig. 4). Since the V_p and V_s velocities of ferropericlase behave quite distinctly across the spin transition, our results here also indicate V_p/V_s ratio can be used as a more sensitive seismic indicator for probing the spin transition-induced heterogeneities in the lower mantle (Figs 3 and 4).

Based on V_s profiles of polycrystalline ferropericlase and bridgmanite at high P - T conditions, it has been proposed that the lower mantle is predominantly made of bridgmanite by 93% in volume, called the perovskitic lower mantle, and that ferropericlase may only account for 7% of the lower-mantle mineralogy¹⁴. In this scenario, the contributions of the elastic and seismic anomalies of ferropericlase across the spin crossover would play a much smaller role on the overall seismic profiles of the lower mantle such that the associated effects may become seismically insignificant. In our modelling, we have considered the lower-mantle P - T conditions along an expected adiabatic geotherm, but the possibility of having a super-adiabatic lower mantle with a steeper thermal gradient than the adiabatic geotherm model is possible¹⁴. The relatively higher geotherm would widen the spin crossover leading to smaller velocity anomalies within the spin crossover and slower V_p and V_s profiles of the LS ferropericlase in the lower mantle. It also remains to be seen as to how the changes in the partitioning coefficient of iron between bridgmanite and ferropericlase ($K_D^{\text{Pv-Fp}} = (X_{\text{Fe}}^{\text{Pv}} / X_{\text{Mg}}^{\text{Pv}}) / (X_{\text{Fe}}^{\text{Fp}} / X_{\text{Mg}}^{\text{Fp}})$) across the spin transition can influence the abnormal elasticity in ferropericlase reported here. Previous studies have shown that the $K_D^{\text{Pv-Fp}}$ decreases from approximately 0.85 at ~750 km depth to 0.2 at ~1800 km depth in the pyrolitic composition, indicating that Fe^{2+} preferentially partitions into ferropericlase in the middle to lower part of the lower-mantle conditions^{32,33}. That is, the iron content of lower mantle minerals can be largely influenced by the change of the iron partitioning associated with the occurrence of the spin transition.

In this case, the spin crossover of ferropericlase can even occur over a wider range of *P-T* conditions with variable amounts of iron involved, thus spreading seismic and chemical anomalies in ferropericlase that may become too broad to be seismically detectable with our current seismic resolution. Thus far, seismic studies have yet revealed reliable features of the lower mantle that can be associated with the spin transition effects in the region. Future seismic studies of the lower mantle taking the effects of the spin transition as well as temperature and composition parameters into account are needed to decipher the geophysical consequences of the spin transition of ferropericlase in the region. Our results point to potential consequences of the spin transition for seismic heterogeneities in the lower mantle and also highlight a much more complex picture of the elasticity of ferropericlase in the lower mantle, affecting our understanding of seismology, geochemistry, and geodynamics in the region.

Methods

Single-crystal ferropericlase ($(\text{Mg}_{0.92}\text{Fe}_{0.08})\text{O}$) was synthesized via inter-diffusion of Fe and Mg between a single-crystal periclase and pre-synthesized $(\text{Mg,Fe})\text{O}$ powder in a H_2/CO_2 gas-mixing furnace at the Institute for Study of the Earth's Interior (ISEI) of Okayama University at Misasa. The MgO crystal with a pre-oriented (100) crystallographic plane purchased from the MTI Corporation was cut down to 7 mm in length by 7 mm wide and 0.25 mm thick, and was sandwiched between two layers of compacted polycrystalline $(\text{Mg}_{0.75}\text{Fe}_{0.25})\text{O}$ powder approximately 1 mm thick each. The starting sample assemblage was then placed in a Pt holder into the furnace operating at 1350 °C and 10^{-2} Pa oxygen fugacity for approximately 2 weeks. The synthesized single-crystal ferropericlase was then extracted and polished for further sample analyses. Electron microprobe and X-ray diffraction analyses of the sample at The University of Texas at Austin showed that the sample was chemically homogeneous with the chemical composition of $(\text{Mg}_{0.92}\text{Fe}_{0.08})\text{O}$ and a unit cell parameter of $a = 4.1996$ (4) Å. The (100)-oriented sample was double polished down to approximately 15 µm thick, and cut into squared platelets 50–80 µm in length for high-pressure DAC experiments. The orientation of the platelet was confirmed by the single-crystal X-ray diffraction patterns at ambient and high pressure (Fig. S1B).

High-pressure X-ray diffraction patterns were collected from the single-crystal sample at room temperature in a DAC at the Sector 13-BMD of the GSECARS of the Advanced Photon Source (APS), Argonne National Lab (ANL) (Fig. S1). A pair of diamond anvils with 200 µm culets were used to pre-indent a rhenium gasket with an initial thickness of 250 µm to approximately 25 GPa (or approximately 25 µm thick). Consequently, a hole of 120 µm was drilled in the pre-indented area and used as the sample chamber. A piece of the platelet 50 µm in length was loaded, together with Au powder as the pressure calibrant and Ne as the pressure medium, in a short symmetric DAC. An incident X-ray beam with a wavelength of 0.3344 Å and focused size of 20 µm (FWHM) in diameter was used for the diffraction experiments (Figs S1 and S2). X-ray diffraction patterns of the sample were collected at pressure intervals of 1–3 GPa up to 91 GPa by a MAR CCD by continuously rotating the DAC around the vertical axis of the sample stage by $\pm 15^\circ$ (Fig. S2). The unit cell parameters and their uncertainties for the sample were calculated based on four sets of the diffraction peaks corresponding to {200}, {220}, {400}, and {420} equivalent reflections (Fig. S1). The uncertainties of the unit cell parameters are typically in the order of 0.04% and are approximately 0.08% at the highest pressure of 91 GPa, indicating that the sample remained sufficiently high quality for the X-ray diffraction, BLS, and ISS experiments. Analysis of the XRD patterns of the sample also confirmed that the crystal was indeed oriented in the (100) crystallographic plane within approximately $\pm 1^\circ$ angular uncertainty.

High-pressure BLS and ISS experiments were performed on the single-crystal ferropericlase at up to 96 GPa in a short symmetric DAC in the Mineral Physics Laboratory of The University of Texas at Austin. Ultralow birefringence and microscopically defect-free diamond anvils were selected for these experiments using a petrographic microscope under crossed-polar, because we had observed that the pulsed laser of the ISS system with a 1064 nm wavelength could potentially damage diamond anvils having defects and high strained areas. Similar to the sample preparation in the XRD experiments, a pair of diamond anvils with 200 µm culets was used to pre-indent a rhenium gasket and a hole of 120 µm was drilled and used as the sample chamber. A piece of the platelet 50–70 µm in length was loaded, together with a few ruby spheres as the pressure calibrant and Ne as the pressure medium, in a short symmetric DAC. Two runs were conducted for the BLS and ISS experiments (Table S1). Pressure uncertainties were determined from measured ruby fluorescence spectra before and after the BLS and ISS measurements. BLS spectra of the sample were collected from the (100) platelet along the [100] and [110] crystallographic directions in the transmitted geometry with a pressure interval of 3–5 GPa up to 96 GPa (Figs 1 and 2). The BLS system is equipped with a Coherent Verdi V2 laser operating at 532 nm wavelength and 600 mW laser power, together with a JRS interferometer and an APD detector (Count-10B Photo Counting Module with approximately 5 cps from Laser Components, Inc.). The focused laser beam size at the sample position was approximately 20 µm in diameter while the scattering angle of the BLS system was set at 48° and calibrated against SiO_2 glass and purified water standards. The data collection time was typically 1 hour at pressures below 50 GPa and 2 hours at higher pressures. Analyses of the Brillouin spectra using OriginPro 9.1 software showed strong V_p and V_s peaks with high signal-to-noise ratios at pressures below 20 GPa, while only the V_s peak of the sample was observed at higher pressures as the V_s peak of the diamond anvils saturated the V_p peak of the sample. The measured V_p and V_s velocities

of the sample at pressures below 20 GPa were used to derive the full elastic constants of the crystal and also to cross check the reliability of the ISS results at lower pressures (see further discussion below).

High-pressure ISS spectra were also collected from the single-crystal sample along the [100] and [110] crystallographic directions up to 96 GPa (Fig. 1). The ISS system is a pump-and-probe technique that is equipped with the pump laser with a 1064 nm wavelength and a pulse width of 15 ps and the probe laser with a 532 nm wavelength. The pump laser from Talisker of the Coherent Company was split into two beams which were then recombined at the sample position with a crossing angle of 20.3° and a beam size of 30 μm. The probe laser was delayed by an Aerotech linear stage as long as 20 ns, while the diffracted ISS signals were collected by a photodiode detector. The data collection time for each ISS spectrum was typically 2 hours. Using MATLAB and OriginPro 9.1 softwares, we implemented the Burg method to analyze the time-domain ISS spectra in order to derive the frequency-domain power spectra and the acoustic wave velocities of the sample at high pressures (Fig. 1). The derived V_p values from ISS measurements are consistent with those from the BLS measurements at pressures below 20 GPa, confirming the calibration and reliability of both techniques. The interfacial waves were also observed in most of the ISS experiments, but their signals were much weaker than that of the longitudinal acoustic waves; in fact, the interfacial waves were too weak to be observed in our analyses in some experiments. Given the uncertainties involved in interpretation of the interfacial wave results from ISS experiments, we have only used the V_p from the ISS measurements and the V_s from the BLS experiments, together with the density results from XRD measurements, to derive full elastic constants of the single-crystal ferropericline at high pressures.

References

1. Crowhurst, J., Brown, J., Goncharov, A. & Jacobsen, S. Elasticity of (Mg,Fe)O through the spin transition of iron in the lower mantle. *Science* **319**, 451 (2008).
2. Marquardt, H., Speziale, S., Reichmann, H. J., Frost, D. J. & Schilling, F. R. Single-crystal elasticity of $(\text{Mg}_{0.9}\text{Fe}_{0.1})\text{O}$ to 81 GPa. *Earth Planet. Sci. Lett.* **287**, 345–352 (2009).
3. Antonangeli, D. *et al.* Spin crossover in ferropericline at high pressure: A seismologically transparent transition? *Science* **331**, 64 (2011).
4. Marquardt, H. *et al.* Elastic shear anisotropy of ferropericline in Earth's lower mantle. *Science* **324**, 224 (2009).
5. Lin, J. F., Speziale, S., Mao, Z. & Marquardt, H. Effects of the electronic spin transitions of iron in lower mantle minerals: implications for deep mantle geophysics and geochemistry. *Rev. Geophys.* **51**, 244–275 (2013).
6. Dziewonski, A. M. & Anderson, D. L. Preliminary reference Earth model. *Phys. Earth Planet. Inter.* **25**, 297–356 (1981).
7. Kennett, B., Engdahl, E. & Buland, R. Constraints on seismic velocities in the Earth from traveltimes. *Geophys. J. Int.* **122**, 108–124 (1995).
8. Ritsema, J., Deuss, A., Van Heijst, H. & Woodhouse, J. S40RTS: a degree-40 shear-velocity model for the mantle from new Rayleigh wave dispersion, teleseismic traveltime and normal-mode splitting function measurements. *Geophys. J. Int.* **184**, 1223–1236 (2011).
9. Ringwood, A. E. *Composition and Petrology of the Earth's Mantle*. Vol. 618 (McGraw-Hill New York, 1975).
10. Irifune, T. Absence of an aluminous phase in the upper part of the Earth's lower mantle. *Nature* **370**, 131–133 (1994).
11. Matas, J., Bass, J., Ricard, Y., Mattern, E. & Bukowski, M. On the bulk composition of the lower mantle: predictions and limitations from generalized inversion of radial seismic profiles. *Geophys. J. Int.* **170**, 764–780 (2007).
12. Karato, S. I. & Karki, B. B. Origin of lateral variation of seismic wave velocities and density in the deep mantle. *J. Geophys. Res.-Sol. Ea.* (1978–2012) **106**, 21771–21783 (2001).
13. Kellogg, L. H., Hager, B. H. & van der Hilst, R. D. Compositional stratification in the deep mantle. *Science* **283**, 1881–1884 (1999).
14. Murakami, M., Ohishi, Y., Hirao, N. & Hirose, K. A perovskitic lower mantle inferred from high-pressure, high-temperature sound velocity data. *Nature* **485**, 90–94 (2012).
15. Cammarano, F., Marquardt, H., Speziale, S. & Tackley, P. J. Role of iron-spin transition in ferropericline on seismic interpretation: A broad thermochemical transition in the mid mantle? *Geophys. Res. Lett.* **37**, L03308 (2010).
16. Murakami, M., Hirose, K., Kawamura, K., Sata, N. & Ohishi, Y. Post-perovskite phase transition in MgSiO_3 . *Science* **304**, 855 (2004).
17. Wentzcovitch, R. *et al.* Anomalous compressibility of ferropericline throughout the iron spin cross-over. *Proc. Natl. Acad. Sci. USA* **106**, 8447 (2009).
18. Badro, J. *et al.* Iron partitioning in Earth's mantle: toward a deep lower mantle discontinuity. *Science* **300**, 789–791 (2003).
19. Badro, J. *et al.* Electronic transitions in perovskite: Possible nonconvecting layers in the lower mantle. *Science* **305**, 383–386 (2004).
20. Lin, J. F. *et al.* Spin transition of iron in magnesiowüstite in the Earth's lower mantle. *Nature* **436**, 377–380 (2005).
21. Lin, J. F. *et al.* Electrical conductivity of the lower-mantle ferropericline across the electronic spin transition. *Geophys. Res. Lett.* **34**, 16305 (2007).
22. Tsuchiya, T., Wentzcovitch, R. M., da Silva, C. R. S. & de Gironcoli, S. Spin transition in magnesiowüstite in Earth's lower mantle. *Phys. Rev. Lett.* **96**, 198501 (2006).
23. Mao, Z., Lin, J. F., Liu, J. & Prakapenka, V. B. Thermal equation of state of lower-mantle ferropericline across the spin crossover. *Geophys. Res. Lett.* **38**, L23308 (2011).
24. Wu, Z. & Wentzcovitch, R. M. Spin crossover in ferropericline and velocity heterogeneities in the lower mantle. *Proc. Natl. Acad. Sci. USA* **111**, 10468–10472 (2014).
25. Wu, Z., Justo, J. F. & Wentzcovitch, R. M. Elastic anomalies in a spin-crossover system: ferropericline at lower mantle conditions. *Phys. Rev. Lett.* **110**, 228501 (2013).
26. Brown, J. & Shankland, T. Thermodynamic parameters in the Earth as determined from seismic profiles. *Geophys. J. Roy. Astr. Soc.* **66**, 579–596 (1981).
27. Fei, Y. *et al.* Spin transition and equations of state of (Mg, Fe)O solid solutions. *Geophys. Res. Lett.* **34**, L17307 (2007).
28. Birch, F. Finite strain isotherm and velocities for single-crystal and polycrystalline NaCl at high pressures and 300 K. *J. Geophys. Res.* **83**, 1257–1268 (1978).
29. Mainprice, D., Barruol, G. & Ismail, W. B. The seismic anisotropy of the Earth's mantle: from single crystal to polycrystal. *Geophys. Monog.* **AGU 117**, 237–264 (2000).
30. Lin, J. F. *et al.* Spin transition zone in Earth's lower mantle. *Science* **317**, 1740 (2007).

31. Karki, B., Wentzcovitch, R., De Gironcoli, S. & Baroni, S. First-principles determination of elastic anisotropy and wave velocities of MgO at lower mantle conditions. *Science* **286**, 1705–1707 (1999).
32. Auzende, A. L. *et al.* Element partitioning between magnesium silicate perovskite and ferropericlase: new insights into bulk lower-mantle geochemistry. *Earth Planet. Sci. Lett.* **269**, 164–174 (2008).
33. Sakai, T. *et al.* Fe-Mg partitioning between perovskite and ferropericlase in the lower mantle. *Am. Mineral.* **94**, 921–925 (2009).

Acknowledgements

We greatly appreciate the critical comments and helpful suggestions by S.P. Grand and Z. Mao. We acknowledge J. Liu, Y. Wu, M. Song, C. Prescher, K. Zhuravlev, and V. Prapapenka for their assistance in the high-pressure experiments. J.F. Lin acknowledges support from the Geophysics Program (EAR-1446946) of the U.S. National Science Foundation. The ISS system at UT Austin is funded by the Instrumentation and Facility Program (EAR-1053446) of the US NSF and the Instrumentation Matching Program of the Jackson School of Geosciences, while the Brillouin system was built partially using funds from the CDAC (Carnegie-DOE Alliance Center) and the Jackson School Matching Program. Synchrotron X-ray diffraction work was performed at the GSECARS of the APS, ANL. APS is supported by 263 DOE-BES, under Contract No. DE-AC02-06CH11357. GeoSoilEnviroCARS is supported by the National Science Foundation - Earth Sciences and Department of Energy Geosciences. This work was partly supported from ISEI for long term Joint-Use Research.

Author Contributions

J.F.L. conceived and designed the research. J.F.L., N.T. and T.O. synthesized the starting sample. J.F.L. and X.T. designed and built the ISS system, and X.T. built the analytical programs for analyzing the ISS data. J.Y., X.T. and J.F.L. collected and analyzed the XRD, BLS, and ISS data. J.Y. conducted thermodynamic modelling. J.F.L. and J.Y. wrote the paper, and all authors were involved in the revision of the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Yang, J. *et al.* Elasticity of Ferropericlase across the Spin Crossover in the Earth's Lower Mantle. *Sci. Rep.* **5**, 17188; doi: 10.1038/srep17188 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>