SCIENTIFIC REPORTS

OPEN

A single-nucleotide exon found in *Arabidopsis*

Lei Guo^{1,2} & Chun-Ming Liu¹

Received: 24 July 2015

Accepted: 11 November 2015

Published: 10 December 2015

The presence of introns in gene-coding regions is one of the most mysterious evolutionary inventions in eukaryotic organisms. It has been proposed that, although sequences involved in intron recognition and splicing are mainly located in introns, exonic sequences also contribute to intron splicing. The smallest constitutively spliced exon known so far has 6 nucleotides, and the smallest alternatively spliced exon has 3 nucleotides. Here we report that the *Anaphase Promoting Complex subunit 11* (*APC11*) gene in *Arabidopsis thaliana* carries a constitutive single-nucleotide exon. *In vivo* transcription and translation assays performed using *APC11-Green Fluorescence Protein (GFP)* fusion constructs revealed that intron splicing surrounding the single-nucleotide exon is effective in both *Arabidopsis* and rice. This discovery warrants attention to genome annotations in the future.

Most eukaryotic genes carry protein-coding exons that are separated by non-coding introns^{1,2}. Pre-mRNA splicing is performed by the spliceosome, a large ribonucleoprotein complex comprised of five small nuclear ribonucleoproteins (snRNPs U1, U2, U4, U5 and U6) and a large number of associated proteins^{3,4}. The size of introns ranges from 13 to over 300,000 nucleotides^{5,6}. Sufficient evidence suggest that intronic sequences not only determine the splicing pattern⁷, but also have regulatory functions in gene expression⁸. Although most known regulatory sequences including the conserved GT and AG located at the beginning and the end of introns, respectively, an A at the branch point and a pyrimidine tract in spliceosome-binding and intron splicing are located in introns⁹, exonic sequences play an important role in accurate splicing as well^{10–12}. The average size of exons is approximately 130 nucleotides in vertebrates, and 180 nucleotides in plants¹³. Studies have showed that exons with less than 51 nucleotides may cause exon skipping, and exons that are too small in size may hinder the recognition of adjacent spliceosome binding^{14–17}. However, internal micro-exons with less than 25 nucleotides have been identified in different eukaryotic organisms by sequencing and computational analyses^{18,19}. The smallest naturally available exon that has been experimentally characterized so far has 3 nucleotides^{16,20}. Here we report the identification of a single-nucleotide exon in *Arabidopsis*.

Results

***APC11* cDNA in GenBank is mis-annotated.** *APC11* (*At3g05870*) is a single-copy gene in the genome of *Arabidopsis thaliana*²¹. Current annotation predicts that *APC11* has three exons and two introns, and its coding sequence (CDS) contains 261 nucleotides, producing a polypeptide with 87 amino acids (AAs)²¹. However, sequencing of *APC11* cDNA performed in this study has identified only one CDS with 252 nucleotides (highlighted in red; Fig. 1), encoding a polypeptide with 84 AAs. The discrepancy was partially caused by the inclusion of 10 nucleotides from the first intron to the exon in previous annotation (highlighted in blue; Fig. 1).

Further, alignment of the cDNA obtained with the *APC11* genomic sequence revealed a single-nucleotide A inserted into the cDNA. The mysterious A is not in continuity with the CDS in the genomic region. The insertion is absolutely required for in-frame *APC11* translation. Re-sequencing of the *APC11* genomic DNA extracted from both Col-0 and *Ler* ecotypes confirmed that the genomic sequence available in the GenBank of National Center for Biotechnology Information (NCBI) is correct, while its cDNA annotated is wrong. We therefore speculate that the extra A may originate from a single-nucleotide exon located in the intron between the previously annotated first and second exons. Within the assigned 422-nucleotide intronic sequence we identified a putative A (designated as A333 in Fig. 1), surrounded by GT and AG, located 333 nucleotides after the upstream exon-intron junction. A putative branch point A was detected 44 nucleotides upstream of the A333 (highlighted in purple; Fig. 1).

A333 is a functional single-nucleotide exon. To test whether A333 indeed represents a single-nucleotide exon, six constructs with nucleus-localized *APC11-SV40-GFP* fusion proteins expressed under the control of

¹Key Laboratory of Plant Molecular Physiology, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China. ²Graduate University of the Chinese Academy of Sciences, Beijing 100049, China. Correspondence and requests for materials should be addressed to C.M.L. (email: cmlu@ibcas.ac.cn)

1 ATGAAAGTCAAGATCTTGCgtatccttttgttgc
 35 ttcattctcttcttagcttttttgggggaaagttt
 69 atattaggttaagtgataaatcatcccaggactt
 103 ttgatggtgatcatctgatatgataataaacatc
 137 ttaagtctgtttatcgtgtattagctttaccgag
 171 ttagattcccagcttgatgattcataataacttt
 205 ttgcatcttccattggtttatttaatagactagg
 239 tttgttcaaaatgcttgcatttcgctttgttgca
 273 caagtaaaagaagaaaaacgaaattgttgctctt
 307 gactttctatgctacttctgtcctgcatacccaa
 341 aaaaaaacagA³³³gtaggttcctcgtcccttta
 373 gctttaacacatgagctcttacaatgcggagttt
 407 ctattgttgtaactgattgtttctccctggaata
 441 gATGGCATGCAGTTGCTTCATGGACATGGGATGC
 475 TCAAGACGAAACATGTGGGATATGTCGGATGGCT
 509 TTTGATGGTTGTTGTCTGATTGCAAACCTCCTG
 543 GAGATGATTGCCCTCTAAgtaaagtctctcttctt
 577 tcatgtcttcgcttttatatatgttgcttagaac
 611 acccactatttctcttgccttacggaacattgct
 645 aaacatttacttatggtaataactcaaaacaaaa
 679 ctgcatgccttagaccttatctgttttgggtgtt
 713 tgggtgttgggcagTTTGGGGAGCCTGCAACCAC
 747 GCGTTTCATCTTCACTGTATATTGAAATGGGTGA
 781 ATTTCGACAGACGAGCCAAGCTCATTGCCCAATGTG
 815 CAGAAGAGAATGGCAGTTCAAAGAGTAA

Figure 1. The genomic sequence of *APC11*. The coding region of *APC11*, in which putative exons are highlighted in red and capital, introns are denoted in black and lower case, and the putative branch point “a” is highlighted in purple. A333 is the putative single-nucleotide exon. Conserved intron-exon splicing sequences “gt” and “ag” are underlined and in lower case. Start and stop codons are underlined and in capital. The mis-annotated exonic sequence in GenBank is highlighted in blue.

the cauliflower mosaic virus (CaMV) 35S promoter were made: 1) *gAPC11-nGFP*: the 839-nucleotide *APC11* genomic sequence, with its stop codon deleted, in-frame fused with a SV40-GFP reporter gene; 2) *cAPC11-nGFP*: a 252-nucleotide *APC11* cDNA, with its stop codon deleted, in-frame fused with the same SV40-GFP; 3) *gAPC11(A > T)-nGFP*: the same as *gAPC11-nGFP* except the A333 was substituted by a T, which is expected to produce a cDNA with T333 if the A333 is indeed a single-nucleotide exon; 4) *gAPC11(A > G)-nGFP*: the A333 in *gAPC11-nGFP* was substituted by a G to determine whether nucleotide types affect the splicing; 5) *gAPC11(A > TT)-nGFP*: the A333 in *gAPC11-nGFP* was substituted by TT, which shall cause a TT substitution in the *APC11* cDNA and a frame shift in *APC11* translation, leading to disappearance of GFP fluorescence; and 6) *gAPC11(-A)-nGFP*: A333 in *gAPC11-nGFP* was deleted, which shall produce a cDNA without A333, leading to a frame-shift in *APC11* translation and disappearance in GFP fluorescence (Fig. 2a). These constructs were introduced into *A. thaliana* mesophyll protoplasts individually using a polyethylene glycol (PEG)-mediated transfection²² for *in vivo* transcriptional and translational assays.

cDNAs were prepared from RNAs extracted from protoplasts transfected with different fusion constructs to examine their splicing patterns. Afterwards, *APC11-nGFP* cDNAs were amplified from individual cDNAs by polymerase chain reaction (PCR) using a forward *APC11* primer and a reverse *GFP* primer (Supplementary Table S1), and sequenced. Results obtained showed that, when either *cAPC11-nGFP* or *gAPC11-nGFP* was used, a sequence identical to *APC11* cDNA was produced. Interestingly, substitutions of A333 by T [*gAPC11(A > T)-nGFP*], G [*gAPC11(A > G)-nGFP*] or TT [*gAPC11(A > TT)-nGFP*] led to T, G or TT substitutions in the cDNA, respectively (Fig. 2b). Further, deletion of A333 made in *gAPC11(-A)-nGFP* led to production of a cDNA without the A.

Detections of GFP fluorescence were used to define the translation of different fusion constructs. When examined under a confocal microscope after twelve-hour incubations, nucleus-localized GFP fluorescence was observed in protoplasts transfected with either *cAPC11-nGFP*, *gAPC11-nGFP*, *gAPC11(A > T)-nGFP* or *gAPC11(A > G)-nGFP*, suggesting that in-frame GFP translations were achieved in protoplasts transfected with these constructs. In contrast, no GFP fluorescence was detected when either *gAPC11(A > TT)-nGFP* or *gAPC11(-A)-nGFP* was used (Fig. 2c), indicating that the substitution of the A333 by TT or deletion of the A333 impaired the translation of these fusion constructs. These results confirmed that A333 in the *APC11* is a functional single-nucleotide exon.

Splicing of the single-nucleotide exon is mostly conserved in rice. We then addressed whether the processing capability of the single-nucleotide exon is conserved in rice (*Oryza sativa*, var. Zhonghua 11), a remotely

Figure 2. *In vivo* transcriptional and translational assays in *Arabidopsis* and rice protoplasts. (a) Constructs generated for transient assays. The CaMV 35S promoter was used to drive the expression of APC11 cDNA, genomic or different substitution constructs fused with a nucleus-localized SV40-GFP reporter gene. Boxes in orange, cyan and grey indicate three previously identified exons in APC11. The black lines indicate introns, and the A333 is shown as red vertical bars. (b) Alignment of APC11 cDNA produced in transgenic *Arabidopsis* or rice protoplasts. Identical nucleotides are shaded. (c) Examinations of GFP fluorescence in *Arabidopsis* protoplasts transfected with constructs illustrated in a. Note that GFP signals are detected only in protoplasts transfected with nGFP, cAPC11-nGFP, gAPC11-nGFP, gAPC11(A > T)-nGFP or gAPC11(A > G)-nGFP. Scale bar = 10 μm for all photos in c.

related monocotyledonous species. APC11 in rice has two paralogs, OsAPC11-1 (Os03g0302700) and OsAPC11-2 (Os07g0411101), both of them lack an intron. Protoplasts prepared from 14-day-old etiolated rice seedlings were used to perform *in vitro* transcriptional assay using above-mentioned six constructs (Fig. 2a). Sequencing of APC11-nGFP cDNAs amplified from rice protoplasts showed that, when either cAPC11-nGFP or gAPC11-nGFP was used in transfections, the intact APC11 cDNA produced from the same splicing patterns as those in *Arabidopsis* protoplasts were detected (Fig. 2b). Similarly, T or G substitutions were detected in cDNA isolated from protoplasts transfected with gAPC11(A > T)-nGFP or gAPC11(A > G)-nGFP, respectively (Fig. 2b). A cDNA without A333, and consequently a frame-shift, was detected in protoplasts transfected with gAPC11(-A)-nGFP. These results suggest that protoplasts of rice can splice the single-nucleotide exon accurately and effectively as those from *Arabidopsis*. However, it is interesting to note that, when gAPC11(A > TT)-nGFP was used, the splicing was incorrect. Additional 56 nucleotides from the first intron were incorporated into the cDNA, leading to a frame-shift in the translation of gAPC11(A > TT)-nGFP, suggesting that the substitution of A333 by TT has caused an altered splicing pattern in rice, which was not observed in *Arabidopsis*.

Discussion

Pre-mRNA splicing is essential in gene expression in eukaryotic organisms since most of their genes contain multiple copies of non-coding introns interspersed between exons. Precise removal of introns ensures the accurate production of proteins. Exons in pre-mRNA can be spliced either constitutively or alternatively: the former generates a single splicing product across all cell types and developmental stages in which the gene is expressed, and the latter produces a variety of mRNAs by splicing from the same gene in different arrangements to generate protein diversity^{9,23}. How these intronic sequences are removed effectively and accurately is still largely unknown, given the fact that the sizes of introns and exons varies tremendously^{5,6,13}.

The average size of internal exons in most eukaryotic organisms is from 130 to 180 nucleotides¹³. Although it has been proposed that exons with less than 51 nucleotides may hinder the recognition of adjacent spliceosome

binding, causing exon skipping^{14–17}, micro-exons with less than 25 nucleotides have been identified in different eukaryotic organisms by sequencing and computational analyses^{18,19}. For example, extensive studies have been performed in a 9-nucleotide constitutive micro-exon in the potato invertase gene and a 6-nucleotide constitutive micro-exon from the chicken *cTNT* gene^{24,25}. The potato invertase gene carries an exon with 9 nucleotides. When 8 of these 9 nucleotides were deleted, the artificial 1-nucleotide exon was skipped in 33% transcripts produced. When this 9-nucleotide exon was replaced by a 6-nucleotide exon from the chicken *cTNT* gene, over 50% of the transcripts produced skipped or mis-spliced the exon²⁴. Another recent study in animal and human brains have identified a whole set of genes carrying evolutionally conserved micro-exons, often with the numbers of multiples of three nucleotides, which are involved in modulating interaction domains of neural proteins through alternative splicing²⁰. It is plausible that different regulatory mechanisms are implicated in splicing introns flanking a normal exon or a micro-exon.

Three models have been proposed to explain how pre-mRNA splicing is achieved. The “intron definition” model states that, for introns with moderate sizes, the splicing reaction occurs by pairing of the splice sites at two ends of an intron to remove the introns^{3,7}. The “exon definition” model is proposed to explain the phenomenon that, for short exons separated by a large intervening intron, attaching a 5′ splice site downstream of the second exon in a two-exon splicing substrate greatly enhances the splicing of the upstream intron *in vitro*^{3,16}. A “recursive splicing” model, which is proposed recently to explain the removal of large introns successively in several steps using intronic ratchet points^{26–29}. In this study, we identified a constitutive single-nucleotide exon in *Arabidopsis*. *In vitro* transcriptional and translational assays performed in protoplasts showed that splicing of introns around this exon can be achieved accurately in both *Arabidopsis* and rice. We also demonstrated that nucleotide types, either purine or pyrimidine, have no effect on splicing of introns around the single-nucleotide exon. Given the fact that spliceosomes are very large in size³⁰, it is very unlikely that the exon definition model could be used to explain the splicing of two introns flanking such a single-nucleotide exon. The intron definition model is more plausible, although it is very unlikely that two introns flanking the single-nucleotide exon could be spliced simultaneously. A combined intron definition and recursive splicing model might be applicable to explain the splicing of introns flanking the single-nucleotide exon, to allow two flanking introns to be removed one after another. Consistent with this hypothesis, it has been reported that in the potato invertase gene the splicing of introns surrounding the 9-nucleotide exon occurs recursively in two steps: the second intron was removed before the first one²⁴. Further studies are needed to discriminate these possibilities and to identify regulatory sequences involved in intron splicing around the single-nucleotide exon.

In summary, although how widely such single-nucleotide exons are present in eukaryotic genomes remains to be investigated, the discovery of the functional single-nucleotide exon undoubtedly has significant impact on genome annotation in the future.

Materials and Methods

Plant materials. *Arabidopsis thaliana* plants (ecotypes Col-0 and *Ler*) were grown at 21 °C in a growth room with 16 h of light (100 μmol photons m^{−2}sec^{−1}) per day.

Constructs. *SV40-GFP* was amplified from *pPLV04*³¹. The full-length *APC11* cDNA (*cAPC11*) was amplified from cDNA prepared from Col-0 seedlings using reverse transcription polymerase chain reaction (RT-PCR), and *APC11* genomic DNA (*gAPC11*) was amplified from Col-0 or *Ler* genomic DNA. *SV40-GFP* and either *cAPC11* or *gAPC11* were ligated simultaneously into *p326-cGFP* digested with *Xba*I (NEB, USA) and *Kpn*I (NEB, USA) using a one-step cloning assay³² to produce *p35S:cAPC11-nGFP* or *p35S:gAPC11-nGFP*, respectively. To generate *p35S:nGFP*, *p35S:gAPC11-nGFP* was digested by *Xba*I and ligated with T4 DNA ligase (NEB, USA). For A333 substitutions, point mutations were introduced to *p35S:gAPC11-nGFP* using the primers listed in Supplementary Table S1 to produce *APC11(A > T)*, *APC11(A > G)*, *APC11(A > TT)* or *APC11(−A)*.

Protoplast transfection. For protoplast transient expressions, well-expanded leaves from 4-week-old *Arabidopsis* plants (Col-0) were chosen, and the assays were performed as previously described²². For protoplast transient expression in rice, seeds (*Oryza sativa*, var. Zhonghua 11) were germinated on half-strength MS basal salts medium and cultured in the dark at 26 °C for 10 to 12 days before protoplasts were isolated and assays were performed as in *Arabidopsis* except the Macerozyme R-10 was replaced by Macerozyme RS (Yakult, Japan).

Microscopic analyses. To examine the expression of *GFP* in transfected protoplasts, a confocal laser scanning microscope (FV1000MPE, Olympus, Japan) equipped with 488 nm excitation laser was used.

RNA extraction and RT-PCR. Total RNA was isolated from transfected *Arabidopsis* or rice protoplasts using the Plant Total RNA Purification Kit (GeneMark, China), reverse-transcribed using the FastQuant RT Kit (TIANGEN, China), and sequenced.

References

- Berget, S. M., Moore, C. & Sharp, P. A. Spliced segments at the 5′ terminus of adenovirus 2 late mRNA. *Proc. Natl. Acad. Sci. USA* **74**, 3171–3175 (1977).
- Chow, L. T., Gelinas, R. E., Broker, T. R. & Roberts, R. J. An amazing sequence arrangement at the 5′ ends of adenovirus 2 messenger RNA. *Cell* **12**, 1–8 (1977).
- Will, C. L. & Lührmann, R. Spliceosome structure and function in *The RNA world third edition* (eds Atkins, J. F., Gesteland, R. F. & Cech, T. R.) 369–400 (Cold Spring Harbor Laboratory Press, 2006).
- Hang, J., Wan, R., Yan, C. & Shi, Y. Structural basis of pre-mRNA splicing. *Science* **349**, 1191–1198 (2015).
- Deutsch, M. & Long, M. Intron-exon structures of eukaryotic model organisms. *Nucleic Acids Res.* **27**, 3219–3228 (1999).
- Atambayeva, Sh. A., Khailenko, V. A. & Ivashchenko, A. T. Intron and exon length variation in *Arabidopsis*, rice, nematode, and human. *Mol. Biol. (Moscow)* **42**, 312–320 (2008).

7. Fox-Walsh, K. L. *et al.* The architecture of pre-mRNAs affects mechanisms of splice-site pairing. *Proc. Natl. Acad. Sci. USA* **102**, 16176–16181 (2005).
8. Nott, A., Meislin, S. H. & Moore, M. J. A quantitative analysis of intron effects on mammalian gene expression. *RNA* **9**, 607–617 (2003).
9. Burge, C. B., Tuschl, T. & Sharp, P. A. Splicing of precursors to mRNAs by the spliceosomes in *The RNA world second edition* (eds Gesteland, R. F., Cech, T. R. & Atkins, J. F.) 525–560 (Cold Spring Harbor Laboratory Press, 1999).
10. Furdon, P. J. & Kole, R. The length of the downstream exon and the substitution of specific sequences affect pre-mRNA splicing *in vitro*. *Mol. Cell. Biol.* **8**, 860–866 (1988).
11. Brown, J. W. S. & Simpson, C. G. Splice site selection in plant pre-mRNA splicing. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **49**, 77–95 (1998).
12. Chasin, L. A. Searching for splicing motifs in *Alternative splicing in the postgenomic era* (eds Blencowe, B. & Graveley, B.) 85–106 (Landes Bioscience, 2007).
13. Hawkins, J. D. A survey on intron and exon lengths. *Nucleic Acids Res.* **16**, 9893–9908 (1988).
14. Black, D. L. Does steric interference between splice sites block the splicing of a short c-src neuron-specific exon in non-neuronal cells? *Gene. Dev.* **5**, 389–402 (1991).
15. Dominski, Z. & Kole, R. Selection of splice sites in pre-mRNAs with short internal exons. *Mol. Cell. Biol.* **11**, 6075–6083 (1991).
16. Berget, S. M. Exon recognition in vertebrate splicing. *J. Biol. Chem.* **270**, 2411–2414 (1995).
17. Hwang, D. Y. & Cohen, J. B. U1 small nuclear RNA-promoted exon selection requires a minimal distance between the position of U1 binding and the 3' splice site across the exon. *Mol. Cell. Biol.* **17**, 7099–7107 (1997).
18. Florea, L., Hartzell, G., Zhang, Z., Rubin, G. M. & Miller, W. A computer program for aligning a cDNA sequence with a genomic DNA sequence. *Genome Res.* **8**, 967–974 (1998).
19. Volfovsky, N., Haas, B. J. & Salzberg, S. L. Computational discovery of internal micro-exons. *Genome Res.* **13**, 1216–1221 (2003).
20. Irimia, M. *et al.* A highly conserved program of neuronal microexons is misregulated in autistic brains. *Cell* **159**, 1511–1523 (2014).
21. The Arabidopsis Genome Initiative. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**, 796–815 (2000).
22. Yoo, S. D., Cho, Y. H. & Sheen, J. *Arabidopsis* mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nat. Protoc.* **2**, 1565–1572 (2007).
23. Black, D. L. Mechanisms of alternative pre-messenger RNA splicing. *Ann. Rev. Biochem.* **72**, 291–336 (2003).
24. Simpson, C. G. *et al.* Requirements for mini-exon inclusion in potato invertase mRNAs provides evidence for exon-scanning interactions in plants. *RNA* **6**, 422–433 (2000).
25. Carlo, T., Sterner, D. A. & Berget, S. M. An intron splicing enhancer containing a G-rich repeat facilitates inclusion of a vertebrate micro-exon. *RNA* **2**, 342–353 (1996).
26. Hatton, A. R., Subramaniam, V. & Lopez, A. J. Generation of alternative *Ultrabithorax* isoforms and stepwise removal of a large intron by resplicing at exon-exon junctions. *Mol. Cell* **2**, 787–796 (1998).
27. Burnette, J. M., Miyamoto-Sato, E., Schaub, M. A., Conklin, J. & Lopez, A. J. Subdivision of large introns in *Drosophila* by recursive splicing at nonexonic elements. *Genetics* **170**, 661–674 (2005).
28. Sibley, C. R. *et al.* Recursive splicing in long vertebrate genes. *Nature* **521**, 371–375 (2015).
29. Duff, M. O. *et al.* Genome-wide identification of zero nucleotide recursive splicing in *Drosophila*. *Nature* **521**, 376–379 (2015).
30. Yan, C. *et al.* Structure of a yeast spliceosome at 3.6-angstrom resolution. *Science* **349**, 1182–1191 (2015).
31. De Rybel, B. *et al.* A versatile set of ligation-independent cloning vectors for functional studies in plants. *Plant Physiol.* **156**, 1292–1299 (2011).
32. Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods* **6**, 343–345 (2009).

Acknowledgements

This work was supported by the “Mechanistic dissection of plant embryo and seed development” project (2014CB943400) of the National Basic Research Program of China and the Dutch-China Collaborative Project (31161130531) of NNFC. We thank Jingbo Jin for providing the *p326-cGFP* vector and Dolf Weijers for the *pPLV04* vector.

Author Contributions

C.M.L. designed the research; L.G. performed the experiments; C.M.L. and L.G. wrote the paper. All authors reviewed the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Guo, L. and Liu, C.M. A single-nucleotide exon found in *Arabidopsis*. *Sci. Rep.* **5**, 18087; doi: 10.1038/srep18087 (2015).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>