

SCIENTIFIC REPORTS



OPEN

Amino acid changes in PB2 and HA affect the growth of a recombinant influenza virus expressing a fluorescent reporter protein

Received: 21 September 2015

Accepted: 21 December 2015

Published: 05 February 2016

Hiroaki Katsura¹, Satoshi Fukuyama^{1,2}, Shinji Watanabe^{2,3}, Makoto Ozawa^{4,5}, Gabriele Neumann⁶ & Yoshihiro Kawaoka^{1,2,6,7}

Influenza viruses that express reporter proteins are useful tools, but are often attenuated. Recently, we found that an influenza virus encoding the Venus fluorescent protein acquired two mutations in its PB2 and HA proteins upon mouse adaptation. Here, we demonstrate that the enhanced viral replication and virulence in mice of this Venus-expressing influenza virus are primarily conferred by the PB2-E712D mutation, with only a minor contribution by the HA-T380A mutation.

Recombinant influenza A viruses that express reporter proteins are useful tools for studying the dynamics of influenza virus infection *in vivo*^{1,2}. Building on the strategy of Manicassamy *et al.*³, we recently generated an influenza A/Puerto Rico/8/34 (PR8, H1N1) virus expressing the Venus reporter protein from a fusion construct with the viral NS1 protein⁴. This virus (referred to as wild-type [WT]-Venus-PR8) was attenuated *in vitro* and *in vivo*, but serial passages in mice resulted in a mouse-adapted variant (MA-Venus-PR8) whose replicative ability and virulence in mice were similar to those of wild-type PR8 (WT-PR8) virus⁴. MA-Venus-PR8 differs from WT-Venus-PR8 by two amino acid changes: one in the viral polymerase subunit PB2 (PB2-E712D) and the other in the hemagglutinin (HA) surface glycoprotein (HA-T380A; H1 HA numbering)⁴. Here, we generated single-gene reassortants and assessed the contribution of these mutations to viral growth and pathogenicity to mice.

Results and Discussion

To assess the contributions of PB2-E712D and HA-T380A to the increased replicative ability of MA-Venus-PR8, we used reverse genetics⁵ to generate two single-gene reassortants that possessed the PB2 or HA viral RNA (vRNA) of MA-Venus-PR8 and the remaining vRNA segments from WT-Venus-PR8 (referred to as PB2-Venus-PR8 and HA-Venus-PR8, respectively). We infected Madin-Darby canine kidney (MDCK) cells at a multiplicity of infection (MOI) of 0.001 and determined viral titers in supernatants by using plaque assays (Fig. 1A). As reported previously⁴, MA-Venus-PR8 grew to significantly higher titers than WT-Venus-PR8. The PB2 and HA vRNAs of MA-Venus-PR8 each increased the growth properties of WT-Venus-PR8 virus, although the virus titers did not reach that of MA-Venus-PR8. These data indicate that the PB2-E712D and HA-T380A mutations both contributed to increased virus replication in MDCK cells.

In mice infected with 10^3 – 10^5 plaque-forming units (PFU) of viruses, MA-Venus-PR8 was more virulent than WT-Venus-PR8, as reported previously⁴ (Fig. 1B). The increased virulence of MA-Venus-PR8 compared with WT-Venus-PR8 was primarily conferred by the MA-Venus PB2 vRNA, whereas HA-Venus-PR8 was similar in its virulence to WT-Venus-PR8. These findings were consistent with virus titers in the lungs of mice infected with

¹Division of Virology, Department of Microbiology and Immunology, Institute of Medical Science, University of Tokyo, Tokyo, Japan. ²Exploratory Research for Advanced Technology, Infection-Induced Host Responses Project, Japan Science and Technology Agency, Saitama, Japan. ³Laboratory of Veterinary Microbiology, Department of Veterinary Sciences, University of Miyazaki, Miyazaki, Japan. ⁴Laboratory of Animal Hygiene, Joint Faculty of Veterinary Medicine, Kagoshima University, Kagoshima, Japan. ⁵Transboundary Animal Disease Center, Joint Faculty of Veterinary Medicine, Kagoshima University, Kagoshima, Japan. ⁶Department of Pathobiological Sciences, School of Veterinary Medicine, University of Wisconsin-Madison, Madison, WI, USA. ⁷Department of Special Pathogens, International Research Center for Infectious Diseases, Institute of Medical Science, University of Tokyo, Tokyo, Japan. Correspondence and requests for materials should be addressed to Y.K. (email: kawaoka@ims.u-tokyo.ac.jp)




Figure 1. Viral replication in MDCK cells and virulence in mice. (A) Virus replication in MDCK cells. Results are expressed as the mean titer (\log_{10} PFU/ml) \pm standard deviation. (B) Assessment of virulence in mice. Body weights are shown as the mean \pm standard error. (C) Assessment of virus titers in the lungs of infected mice. Results are expressed as the mean of the titer (\log_{10} PFU/g) \pm standard deviation. Statistical significance was calculated by using the Tukey-Kramer method. Asterisks indicate significant differences from titers in mice infected with WT-Venus-PR8 ($P < 0.05$). ND: Not detected (detection limit, 5 PFU/lung).

10^3 PFU of each virus (Fig. 1C). Hence, the PB2-E712D substitution is primarily responsible for the increased virulence of MA-Venus-PR8 relative to WT-Venus-PR8 in mice.

To demonstrate that the PB2-E712D mutation increased the Venus expression levels, we infected MDCK cells with the indicated viruses at an MOI of 1 and performed confocal microscopy 12 h later (Fig. 2A). As expected, the levels of the NS1-Venus fusion protein were higher in cells infected with MA-Venus-PR8 or PB2-Venus-PR8 than in those infected with WT-Venus-PR8 or HA-Venus-PR8 (Fig. 2A).




Figure 2. The PB2-E712D substitution augments the expression of Venus in infected cells but does not enhance the polymerase activity in minireplicon assays. (A) Venus expression in infected MDCK cells. MDCK cells were infected with each virus at an MOI of 1 and observed 12 h later by confocal microscopy. (B) Polymerase activity in minireplicon assays in human HEK293 cells. Cells were transfected with plasmids encoding the PB1, PA, NP, and wild-type or mutant PB2 proteins, with a plasmid for the expression of the virus-like RNA encoding the firefly luciferase gene under the control of the human RNA polymerase I promoter, and with a control plasmid encoding *Renilla* luciferase. Luciferase activity was measured 48 h later. (C) Polymerase activity in minireplicon assays in canine MDCK cells. Cells were transfected with plasmids encoding the PB1, PA, NP, and wild-type or mutant PB2 proteins, with a plasmid for the expression of the virus-like RNA encoding the firefly luciferase gene under the control of the canine RNA polymerase I promoter, and with a control plasmid encoding *Renilla* luciferase. Luciferase activity was measured 48 h later.




Figure 3. The HA-T380A substitution affects the pH threshold for membrane fusion of HA. (A) The position of the HA-T380A substitution (red) was mapped on the PR8 HA three-dimensional structure⁹. The three-dimensional structure of PR8 HA was obtained from the Protein Data Bank (PDB ID: 1RU7). The fusion peptide of HA is shown in cyan. The image was generated by using Pymol software. (B) The pH threshold of HA-mediated membrane fusion. Representative results of two independent experiments are shown.

Collectively, our data indicate that the PB2-E712D substitution is primarily responsible for the increased replicative ability, Venus expression, and virulence in mice of MA-Venus-PR8 virus. To assess whether the PB2-E712D mutation directly affects the viral polymerase activity, we performed a minireplicon assay in human HEK293 cells^{6,7}. Unexpectedly, the polymerase activity of PB2-E712D was lower than that of the parental PB2 (Fig. 2B). Similar results were obtained with canine MDCK cells⁸ (Fig. 2C). In the context of a minireplicon that measures viral replication and transcription, the PB2-E712D mutation is thus attenuating; in contrast, this mutation enhances viral growth in the context of replicating virus. These findings indicate that the PB2 protein functions not only in viral replication/transcription, but performs additional roles in the viral life cycle.

The HA vRNA of MA-Venus-PR8 did not significantly increase the virulence of WT-Venus-PR8 in mice; however, HA-Venus-PR8 virus grew more efficiently in MDCK cells than WT-Venus-PR8 (Fig. 1A), suggesting a contribution of the HA-T380A mutation to, at least, virus replication in cultured cells. Because the HA-T380A substitution is located on an α -helix in the HA2 subunit⁹ (Fig. 3A), we evaluated its effect on HA membrane-fusion activity by using a polykaryon formation assay¹⁰. The wild-type HA had a threshold for membrane fusion of pH 5.5, whereas the threshold for HA-T380A was pH 5.8 (Fig. 3B), leading to the conformational change in HA at an earlier stage of endosome maturation during influenza virus entry¹¹. Changes in the pH threshold for membrane fusion may affect HA thermostability¹², an effect that we did not observe at 50 °C (data not shown).

In conclusion, the increased replicative ability, Venus expression, and virulence in mice of MA-Venus-PR8 are primarily brought about by the PB2-E712D mutation, with a minor additional contribution by HA-T380A. The glutamic acid residue at PB2-712 is highly conserved among all influenza A viruses; our search of the Influenza Research Database (www.fludb.org) did not uncover a single isolate with an aspartic acid residue at PB2-712. Interestingly, the PB2-E712D mutation does not increase viral replication and transcription in minireplicon assays perhaps because it affects PB2 binding to importins, which are essential for protein import into the nucleus. The PB2 interaction with importin is mediated by the PB2 residues at positions 701¹³ and, to a lesser extent, 714¹⁴. In fact, the PB2-D701N and PB2-S714R mutations facilitate adaptation of avian influenza viruses to mammals by affecting the PB2 interaction with importin, resulting in better exposure of the PB2 nuclear localization signal^{13,14}. A mutation at position 712 (which is located in the three-dimensional structure of the viral polymerase complex next to the residue at position 714¹⁵) may affect the PB2 interaction with importins, potentially reducing the virus's replicative ability. On the other hand, PB2 is known to interact with RIG-I¹⁶, and may also interact with other cellular factors that play roles in innate immune responses^{17,18} or in the assembly and budding of new virions¹⁸. We speculate that the PB2-E712D mutation may affect these steps, resulting in the higher virulence of the PB2-E712D mutant virus *in vivo*.

The NS1 protein is incorporated into progeny virions¹⁹. NS1 functions as an antagonist of host immune responses by interfering with innate immune pathways such as the RIG-I/IPS-1 signaling pathway, resulting in the inhibition of interferon secretion²⁰. Although we have not examined differences in NS1 protein incorporation between WT-Venus-PR8 and mouse-adapted viruses, it is possible that PB2-Venus-PR8 and MA-Venus-PR8 incorporated more NS1 protein because NS1-Venus expression levels were higher in cells infected with these viruses. Since the virion-incorporated NS1 protein could function soon after infection, PB2-Venus-PR8 and MA-Venus-PR8 might suppress the host innate immune responses more efficiently than WT-Venus-PR8, resulting in the higher proliferative ability and virulence of the mouse-adapted viruses.

Our hypothesis that PB2 has additional, as-yet uncharacterized roles in the viral life cycle is supported by two other recent studies^{21,22}. Alternatively, higher NS1 virion-incorporation levels may increase virulence. Although the mechanism is not currently understood, PB2-E712D and HA-T380A (in this study), as well as other mutations found in the MA-Venus-H5N1²³ virus, are useful for generating recombinant fluorescent influenza viruses that replicate efficiently in cultured cells and mice.

Methods

Cells and viruses. Madin-Darby canine kidney (MDCK) cells were maintained in minimum essential medium (MEM) containing 5% of newborn calf serum. Human embryonic kidney 293 (HEK293) cells were maintained in Dulbecco's modified Eagle medium supplemented with 10% fetal calf serum (FCS). PR8 and each Venus-PR8 mutant were generated by using reverse genetics⁵ and were propagated in MDCK cells at 37 °C for 48 h in MEM containing L-(tosylamido-2-phenyl) ethyl chloromethyl ketone (TPCK)-treated trypsin (0.8 µg/ml) and 0.3% bovine serum albumin (BSA).

Growth kinetics of each virus. MDCK cells were infected with each virus at an MOI of 0.001. Supernatants were collected every 12 h and viral titers in the supernatants were determined by means of plaque assay in MDCK cells.

Pathogenicity and replication of viruses in mice. Six-week-old female C57BL/6 mice were intranasally infected with 50 µl of 10³, 10⁴ or 10⁵ PFU of each virus. Six mice per group were monitored for survival and body weight changes for 14 days after infection. Three mice per group were infected with 10³ PFU of each virus and euthanized on the indicated days. Their lungs were collected to determine viral titers by means of plaque assay in MDCK cells. All animal experiments were performed in accordance with the University of Tokyo's Regulations for Animal Care and Use, which were approved by the Animal Experiment Committee of the Institute of Medical Science, the University of Tokyo (approval number PA 10–13).

Minigenome assay. A minigenome assay based on the dual-luciferase system was performed as described previously^{6–8}. Briefly, HEK293 cells were transfected with viral protein expression plasmids for PB1, PA, NP and wild-type PB2 or PB2-E712D, with a plasmid expressing a reporter vRNA encoding the firefly luciferase gene under the control of the human RNA polymerase I promoter [pPolI/NP(0)Fluc(0)], and pRL-null (Promega), which expresses *Renilla* luciferase, as a transfection control. The luciferase activity in the transfected HEK293 cells were measured by using Dual-Glo luciferase assay system (Promega) at 48 h posttransfection. Polymerase activity was calculated by standardization of the firefly luciferase activity to the *Renilla* luciferase activity. For the minigenome assay in canine MDCK cells, we used a plasmid expressing a reporter vRNA encoding the firefly luciferase gene under the control of the canine RNA polymerase I promoter⁸.

Polykaryon formation assay. Polykaryon formation assay was performed as described previously¹⁰ with modification. HEK293 cells were infected with wild-type PR8 or PR8 possessing the HA-T380A mutation in DMEM containing 10% FCS at an MOI of 10. At 18 h post-infection, cells were washed with MEM containing 0.3% BSA and treated with TPCK-treated trypsin (1 µg/ml) in MEM containing 0.3% BSA for 15 min at 37 °C to cleave the HA on the cell surface into HA1 and HA2. Trypsin was inactivated by washing the cells with DMEM containing 10% FCS. To initiate polykaryon formation, cells were exposed to low-pH buffer (145 mM NaCl, 20 mM sodium citrate (pH 6.0–5.4)) for 2 min at 37 °C. Then the low-pH buffer was replaced with DMEM containing 10% FCS and the cells were incubated for 2 h at 37 °C. The cells were then fixed with methanol and stained with Giemsa's solution.

References

- Kittel, C. *et al.* Rescue of influenza virus expressing GFP from the NS1 reading frame. *Virology* **324**, 67–73, doi: 10.1016/j.virol.2004.03.035 (2004).
- Shinya, K., Fujii, Y., Ito, H., Ito, T. & Kawaoka, Y. Characterization of a neuraminidase-deficient influenza A virus as a potential gene delivery vector and a live vaccine. *J Virol* **78**, 3083–3088 (2004).
- Manicassamy, B. *et al.* Analysis of *in vivo* dynamics of influenza virus infection in mice using a GFP reporter virus. *Proc Natl Acad Sci USA* **107**, 11531–11536, doi: 10.1073/pnas.0914994107 (2010).
- Fukuyama, S. *et al.* Multi-spectral fluorescent reporter influenza viruses (Color-flu) as powerful tools for *in vivo* studies. *Nat Commun* **6**, 6600, doi: 10.1038/ncomms7600 (2015).
- Neumann, G. *et al.* Generation of influenza A viruses entirely from cloned cDNAs. *Proc Natl Acad Sci USA* **96**, 9345–9350 (1999).
- Ozawa, M. *et al.* Contributions of two nuclear localization signals of influenza A virus nucleoprotein to viral replication. *J Virol* **81**, 30–41, doi: 10.1128/JVI.01434-06 (2007).
- Yamayoshi, S. *et al.* Virulence-affecting amino acid changes in the PA protein of H7N9 influenza A viruses. *J Virol* **88**, 3127–3134, doi: 10.1128/JVI.03155-13 (2014).
- Murakami, S. *et al.* Establishment of canine RNA polymerase I-driven reverse genetics for influenza A virus: its application for H5N1 vaccine production. *J Virol* **82**, 1605–1609, doi: 10.1128/JVI.01876-07 (2008).
- Gamblin, S. J. *et al.* The structure and receptor binding properties of the 1918 influenza hemagglutinin. *Science* **303**, 1838–1842, doi: 10.1126/science.1093155 (2004).

10. Imai, M. *et al.* Experimental adaptation of an influenza H5 HA confers respiratory droplet transmission to a reassortant H5 HA/H1N1 virus in ferrets. *Nature* **486**, 420–428, doi: 10.1038/nature10831 (2012).
11. Lozach, P. Y., Huotari, J. & Helenius, A. Late-penetrating viruses. *Curr Opin Virol* **1**, 35–43, doi: 10.1016/j.coviro.2011.05.004 (2011).
12. Ruigrok, R. W. *et al.* Conformational changes in the hemagglutinin of influenza virus which accompany heat-induced fusion of virus with liposomes. *Virology* **155**, 484–497 (1986).
13. Tarendeau, F. *et al.* Structure and nuclear import function of the C-terminal domain of influenza virus polymerase PB2 subunit. *Nat Struct Mol Biol* **14**, 229–233, doi: 10.1038/nsmb1212 (2007).
14. Czudai-Matwich, V., Otte, A., Matrosovich, M., Gabriel, G. & Klenk, H. D. PB2 mutations D701N and S714R promote adaptation of an influenza H5N1 virus to a mammalian host. *J Virol* **88**, 8735–8742, doi: 10.1128/JVI.00422-14 (2014).
15. Tarendeau, F. *et al.* Host determinant residue lysine 627 lies on the surface of a discrete, folded domain of influenza virus polymerase PB2 subunit. *PLoS Pathog* **4**, e1000136, doi: 10.1371/journal.ppat.1000136 (2008).
16. Li, W., Chen, H., Sutton, T., Obadan, A. & Perez, D. R. Interactions between the influenza A virus RNA polymerase components and retinoic acid-inducible gene I. *J Virol* **88**, 10432–10447, doi: 10.1128/JVI.01383-14 (2014).
17. Graef, K. M. *et al.* The PB2 subunit of the influenza virus RNA polymerase affects virulence by interacting with the mitochondrial antiviral signaling protein and inhibiting expression of beta interferon. *J Virol* **84**, 8433–8445, doi: 10.1128/JVI.00879-10 (2010).
18. Watanabe, T. *et al.* Influenza virus-host interactome screen as a platform for antiviral drug development. *Cell Host Microbe* **16**, 795–805, doi: 10.1016/j.chom.2014.11.002 (2014).
19. Hutchinson, E. C. *et al.* Conserved and host-specific features of influenza virion architecture. *Nat Commun* **5**, 4816, doi: 10.1038/ncomms5816 (2014).
20. Hale, B. G., Randall, R. E., Ortín, J. & Jackson, D. The multifunctional NS1 protein of influenza A viruses. *J Gen Virol* **89**, 2359–2376, doi: 10.1099/vir.0.2008/004606-0 (2008).
21. Fan, S. *et al.* Novel residues in avian influenza virus PB2 protein affect virulence in mammalian hosts. *Nat Commun* **5**, 5021, doi: 10.1038/ncomms6021 (2014).
22. Taft, A. S. *et al.* Identification of mammalian-adapting mutations in the polymerase complex of an avian H5N1 influenza virus. *Nat Commun* **6**, 7491, doi: 10.1038/ncomms8491 (2015).
23. Zhao, D. *et al.* Molecular Determinants of Virulence and Stability of a Reporter-Expressing H5N1 Influenza A Virus. *J Virol* **89**, 11337–11346, doi: 10.1128/JVI.01886-15 (2015).

Acknowledgements

We thank Takeo Gorai and Dongming Zhao for helpful discussion, Izumi Ishikawa for technical support, and Susan Watson for editing the manuscript. This work was supported by the Japan Initiative for Global Research Network on Infectious Diseases from the Ministry of Education, Culture, Sports, Science, and Technology, Japan and Japan Agency for Medical Research and Development, AMED; by grants-in-aid from the Ministry of Health, Labour and Welfare, Japan; by ERATO and the Strategic Basic Research Programs of the Japan Science and Technology Agency, and by an NIAID-funded Center for Research on Influenza Pathogenesis (CRIP, HHSN266200700010C) to Y.K. H.K. is supported by a Grant-in-Aid from the Japan Society for the Promotion of Science.

Author Contributions

H.K., S.F., M.O. and Y.K. designed the study. H.K. and S.W. performed the experiments. H.K., G.N. and Y.K. wrote the manuscript. All authors reviewed the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Katsura, H. *et al.* Amino acid changes in PB2 and HA affect the growth of a recombinant influenza virus expressing a fluorescent reporter protein. *Sci. Rep.* **6**, 19933; doi: 10.1038/srep19933 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>