SCIENTIFIC REPORTS

Received: 27 October 2015 Accepted: 15 January 2016 Published: 12 February 2016

OPEN Head-on collision and overtaking collision between an envelope solitary wave and a KdV solitary wave in a dusty plasma

Heng Zhang¹, Wen-Shan Duan¹, Xin Qi² & Lei Yang^{2,3}

Head-on collision and overtaking collision between a KdV solitary wave and an envelope solitary wave are first studied in present paper by using Particle-in-cell (PIC) method in a dusty plasma. There are phase shifts of the KdV solitary wave in both head-on collision and the overtaking collision, while no phase shift is found for the envelop solitary wave in any cases. The remarkable difference between head-on collision and the overtaking collision is that the phase shift of KdV solitary wave increases as amplitude of KdV solitary wave increases in head-on collision, while it decreases as amplitude of the KdV solitary wave increases in the overtaking collision. It is found that the maximum amplitude during the collision process is less than sum of two amplitudes of both solitary waves, but is larger than either of the amplitude.

The field of plasma physics is rapidly growing due to its extensive application in space $plasmas^{1-4}$, microwave transmision^{5,6}, fusion plasmas⁷⁻¹⁰ and so on. Plasmas containing dust particles are called the dusty plasmas</sup> which are also extensively studied because of their relevance in many spaces and technological applications¹¹⁻¹³. Meanwhile dusty plasma supports a variety of collective modes and nonlinear coherent structures. The waves in a dusty plasma have been a interesting topic for many years. The existence of dust acoustic wave (DAW) was first theoretically predicted by Rao et al.¹⁴ in an unmagnetized plasma consisting of the inertial dust fluid, Boltzmann distributed electrons and ions. Since then, the DAW has been extensively studied analytically¹⁵⁻¹⁸ and verified by the experiments¹⁹⁻²¹. As well known, this kind of nonlinear wave can be described by either a Korteweg-de-Vries (KdV) equation or a nonlinear Schrödinger equation (NLSE). The KdV equation describes evolution of a unmodulated wave, called KdV soliton. Whereas, the NLSE usually describes the modulated waves. One of the solutions of the NLSE is called envelope soliton. Admin and Shukla et al.²² studied the modulational instability of the dust acoustic waves and the dust-ion-waves. Ghosh et al.²³ studied the effects of the dust charge fluctuations of the low-frequency wave modulation.

The interaction between two KdV solitary waves is a subject of great interest^{24–26}. In general, when two solitary waves propagate in one-dimensional plasmas, they can undergo two kinds of interaction²⁷. One is the overtaking collision when both propagate in the same direction but with different propagation speed which can be investigated by the inverse scattering method²⁸. The other is called the head-on collision when both propagate in oppo-site directions which is usually studied by the extended Poincare-Lighthill-Kuo (PLK) method²⁹⁻³⁵. The head-on collision between two KdV solitary waves have been verified from both the experiments^{36,37} and the numerical simulation by using the PIC simulation method³⁸. Recently, the head-on collision between two envelope solitary waves are studied analytically³⁹. The head-on collision between two envelope solitary waves has also been studied by using the PIC method⁴⁰. Interesting results are found when head-on collision between two envelope solitary waves happen. There is a phase shift after a head-on collision between two KdV solitary waves³⁸, while there is no phase shift between two envelope solitary waves⁴⁰.

¹College of Physics and Electronic Engineering and Joint Laboratory of Atomic and Molecular Physics of Northwest Normal University, Lanzhou 730070, China. ²Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China. ³Department of Physics, Lanzhou University, Lanzhou 730000, China. Correspondence and requests for materials should be addressed to W.-S.D. (email: duanws@126.com) or X.Q. (email: gixin2002@impcas.ac.cn) or L.Y. (email: lyang@impcas.ac.cn)

However, no report is given on collision between a KdV solitary wave and an envelope one. In the present paper we try to discuss this question by using the PIC simulation method. Both the head-on collision and overtaking collision between two kinds of waves will be studied in this paper.

It is found that there are phase shifts for a KdV solitary wave during the collision between KdV solitary wave and envelope solitary wave, which is similar to that between two KdV solitary waves in which there are phase shifts for both colliding solitary waves³⁸. However, there is no phase shift for the envelope solitary wave after the collision.

Results

Analytical solutions of KdV and NLSE by using the perturbation method. The interaction between a KdV solitary wave and an envelope solitary wave in a dusty plasma will be studied in the present paper by using the one dimensional (1D) PIC method in an infinite background plasma. First we give the 1D dimensionless normalized equations of motion for system^{38,40}

$$\frac{\partial n_d}{\partial t} + \frac{\partial}{\partial x}(n_d u_d) = 0, \tag{1}$$

$$\frac{\partial u_d}{\partial t} + u_d \frac{\partial u_d}{\partial x} = \frac{\partial \phi}{\partial x},\tag{2}$$

$$\frac{\partial^2 \phi}{\partial x^2} = 1 + \nu exp(\beta s\phi) - \mu exp(-s\phi), \qquad (3)$$

where $\beta = T_i/T_e$ is the ratio of the ion and electron temperatures, $s = 1/(\mu + \nu\beta)$, μ and ν are the normalized ion and electron number densities, respectively, n_d and u_d refer to the density, the velocity of the dust grains respectively, ϕ is the electrostatic potential. In the simulation, we take $\beta = 0.1$, $\mu = 1.1$ and $\nu = 0.1$.

The spatial coordinate x, the time t, the velocity and the electrostatic potential ϕ are normalized by the Debye length $\lambda_D = (T_{eff}/4\pi Z_d n_{d0} e^2)^{1/2}$, the inverse of effective dust plasma frequency $\omega^{-1} = (m_d/4\pi n_{d0} Z_d^2 e^2)^{1/2}$, the dust acoustic speed $C_d = (Z_d T_{eff}/m_d)^{1/2}$ and T_{eff}/e , respectively, where the effective temperature is defined as $T_{eff} = \frac{T_i T_e}{\mu T_e + \nu T_i}$.

By introducing the appropriate stretched coordinates $\xi = \varepsilon(x - ct)$ and $\tau = \varepsilon^3 t$, we can obtain a KdV equation as follows

$$\frac{\partial \phi}{\partial \tau} + \alpha \phi \frac{\partial \phi}{\partial \xi} + \delta \frac{\partial^3 \phi}{\partial \xi^3} = 0, \tag{4}$$

where $\alpha = -\frac{k_d}{2}(3 + s^2(\delta^2\nu - \mu)), \delta = \frac{k_d^3}{2}$. k_d and c are the wave number and velocity of the KdV solitary wave, respectively, where $k_d = 1.0$. One of the solution of Eq. (4) is $\phi_{kd\nu}(\xi, \tau) = \phi_m \operatorname{sech}^2\left(\frac{\xi - u_0\tau}{D}\right)$, where $\phi_m = \frac{3u_0}{\alpha}$, $D = \sqrt{\frac{2k_d^3}{u_0}}, u_0 = 1.0$. We choose the same parameters as that in ref. 38.

NLSE can also be obtained as follows by using the different transformations⁴⁰

$$i\frac{\partial\phi_{1}^{(1)}}{\partial\tau'} + P\frac{\partial^{2}\phi_{1}^{(1)}}{\partial\xi'^{2}} + Q\left|\phi_{1}^{(1)}\right|^{2}\phi_{1}^{(1)} = 0,$$
(5)

where $\xi' = \varepsilon(x - u_s t)$, $\tau' = \varepsilon^2 t$, u_s is group velocity of envelope solitary wave. P and Q are functions of system parameters. In the simulation, we choose system parameters as follows: wave number of envelope wave $k_n = 0.1$, then we obtain: P = -0.146, $Q = -3.65^{40}$. One of the envelope solitary wave solution of Eq. (5) is: $\phi_1^{(1)} = \pm \sqrt{\frac{2(Pk_n^2 - \omega)}{Q}} \cdot \operatorname{sech} \sqrt{\frac{Pk_n^2 - \omega}{P}} \xi' \cdot e^{i(k_n x - \omega t)}$. If the signs of both ω and u_s are positive, the envelope wave propagates in the positive x direction.

0.1 PIC simulation results of the head-on collision. In simulation, the initial conditions of the envelope solitary wave are given as follows: $\phi_{nls} = \varepsilon_2 \phi_1^{(1)}$, $u_{nls} = \frac{k_n}{\omega} \phi_{nls}$ and $n_{nls} = 1 + \frac{k_n^2}{\omega^2} \phi_{nls}$. ϕ_{nls} , u_{nls} and n_{nls} are electrostatic potential, velocity and density of dust grains corresponding to envelope solitary wave, respectively. The initial conditions of the KdV solitary wave are also chosen from the analytical solution:

 $\phi_{kdv} = -\left(\frac{3\varepsilon_1^2 u_0}{\alpha}\right) \operatorname{sech}^2 \frac{x - \frac{1 + \varepsilon_1^2 u_0}{k_d}}{2\sqrt{\delta/(\varepsilon_1^2 k_d^2 u_0)}}, u_{kdv} = \phi_{kdv} \text{ and } n_{kdv} = 1 + \phi_{kdv}. \phi_{kdv}. u_{kdv} \text{ and } n_{kdv} \text{ are electrostatic potential,}$ velocity and density of dust grains corresponding to KdV solitary wave, respectively.

The dust particles are represented as superparticles, while the ions and the electrons are modeled as Boltzmann distributed background. The area weighting technique is used to deposit the charge on the 2th nearest grid points in order to obtain the spatial distribution of the particles on the mesh. At the start of simulations, 100 superparticles with different weighting parameters are allocated symmetrically in each grid. Weighting parameters and initial velocities of superparticles are carefully chosen to make sure that the density and velocity distribution of dust particles are given as those obtained by the reductive perturbation method. The electric field is determined by the



Figure 1. The evolution of head-on-collision of KdV solitary wave 1 and an envelope solitary wave 2 at the different times where $\varepsilon_1 = 0.002$, $\varepsilon_2 = 0.01$, $\mu = 1.1$, $\nu = 0.1$, $\beta = 0.1$.



Figure 2. The trajectories of the head on collision between an envelope solitary wave 2 and a KdV solitary wave 1, where $\varepsilon_1 = 0.002$, $\varepsilon_2 = 0.01$, $\mu = 1.1$, $\nu = 0.1$, $\beta = 0.1$.

nonlinear Poisson-Boltzmann's equation. The motion equations for superparticles are integrated by a standard leap-frog algorithm.

Initially the KdV solitary wave propagates in the positive *x* direction, while the envelope solitary wave propagates in the negative *x* direction, shown in Fig. 1. Both are far apart. After some time, they interact, collide, and then depart. It seems that the waveform and the propagation speeds of both solitary waves remain unchanged after the collision is over. The colliding process are clearly shown in both Figs 1 and 2.

In Fig. 2 the colliding trajectories are pictured. There are no phase shift of the envelope solitary wave, i.e., the propagation velocity of the envelope solitary wave is a constant. However, the propagation velocity of the KdV solitary wave is not a constant during the collision. During the collision, the propagation velocity of the KdV solitary wave first becomes larger, then smaller and finally become a constant as that of the previous one. Furthermore, a phase delay of KdV solitary wave is observed. The definition of the phase shift is shown in Fig. 2. In order to further understand the dependence of the phase shifts of both solitary waves on their amplitudes, the dependence of phase shifts of the KdV solitary waves on the parameters of both ε_1 and ε_2 are given in Fig. 3. It is noted that the phase shifts of KdV solitary waves only depend on its amplitudes, i.e., ε_1 , but not that of the



Figure 3. The phase delay of KdV solitary wave 1 after collision with envelop solitary wave 2 as a function of amplitudes of both ε_1 and ε_2 .



Figure 4. The dependence of the maximum amplitude A_{max} in colliding process on envelop solitary wave amplitudes A_2 for different amplitudes of KdV solitary wave A_1 , A_2 and A_3 .

.....

envelope wave, i.e., ε_2 . Furthermore, It seems that there is no phase shift for the envelope solitary wave after the collision in any case.

It is observed that a maximum amplitude will be reached during the collision between two solitary waves. It indicates, however, that the maximum value is less than the sum of two amplitudes of both solitary waves, but it is larger than either of the amplitude. In order to understand this situation in detail, the dependence of the maximum amplitudes in the colliding process between two solitary waves on both of their initial amplitudes is shown in Fig. 4. It is noted that the maximum amplitude in colliding process as the amplitudes of both colliding solitary waves increase. It seems that the maximum amplitude is less than the sum of two amplitudes of both the KdV solitary wave and the envelop solitary wave.

PIC simulation results of the overtaking collision. When the overtaking collision between two solitary waves takes place, both solitary waves propagate in the same direction (positive *x* direction). In the simulations, the initial conditions of the envelope solitary wave is: $\phi_{nls} = \varepsilon_2 \phi_1^{(1)}, u_{nls} = -\frac{k_n}{\omega} \phi_{nls}$ and $n_{nls} = 1 - \frac{k_n^2}{\omega^2} \phi_{nls}$. The initial conditions of the KdV solitary wave is: $\phi_{kdv} = -\left(\frac{3\varepsilon_1^2 u_0}{\alpha}\right) \operatorname{sech}^2 \frac{x - \frac{1 + \varepsilon_1^2 u_0}{kd}}{2\sqrt{\delta/(\varepsilon_1^2 k_d^2 u_0)}}, u_{kdv} = \phi_{kdv}$ and $n_{kdv} = 1 + \phi_{kdv}$. Because the propagation velocity of the envelope solitary wave is constant, while for the KdV solitary wave the

Because the propagation velocity of the envelope solitary wave is constant, while for the KdV solitary wave the propagation velocity depends on its amplitude. We choose the large amplitude KdV solitary wave in order that it can overtake the envelope solitary wave in a short period. The trajectories of the overtaking collision between two solitary waves are shown in Fig. 5. It is found that there is no phase shift for the envelope solitary wave after the overtaking collision is over, while there is phase shift for the KdV solitary wave. The dependence of the phase shifts of the KdV solitary waves on the parameter of ε_1 characterizing the amplitude of the KdV solitary wave are given in Fig. 6. It indicates that the phase shift of the KdV solitary wave on the parameter of ε_2 characterizing the amplitude of the envelope solitary wave is given in Fig. 7. It seems that the phase shift of the KdV solitary wave. It is noted that the phase shift of the envelope solitary wave. It is noted that the phase shift of the envelope solitary wave. It is noted that the phase shift of the envelope solitary wave.







Figure 6. The dependence of the phase delay of the KdV solitary wave for the overtaking collision between an envelope solitary wave 2 and a the KdV solitary wave 1 on the amplitude of the KdV solitary wave.



Figure 7. The dependence of the phase delay of the KdV solitary wave for the overtaking collision between an envelope solitary wave 2 and a the KdV solitary wave 1 on the amplitude of the envelope solitary wave.

decreases as the amplitude of the KdV solitary wave increases in the overtaking collision which is different from that in head-on collision.



Figure 8. The summary of a computational cycle of the PIC method.

Discussion

Both the head-on collision and overtaking collision between a KdV solitary wave and an envelope solitary wave are investigated by using PIC method in a dusty plasma. It is found that there are phase delay for KdV solitary wave for either the head on collision or the overtaking collision between a KdV solitary wave and an envelope solitary wave. It is observed that there are no phase delay for the envelop solitary wave in any cases. The remarkable difference between the head-on collision and the overtaking collision is that the phase shift of the KdV solitary wave increases as the amplitude of the KdV solitary wave increases in the head-on collision, while it decreases as the amplitude of the KdV solitary wave increases in the overtaking collision. It is found that the maximum amplitude during the collision process is less than the sum of two amplitudes of both solitary waves, but is larger than either of the amplitude. It is noted that the maximum amplitude in colliding process increases as the amplitudes of both colliding solitary waves increase. Meanwhile, PIC simulation provides a more realistic description of the dynamics of nonlinear dust acoustic waves that can be usefully applied to various low frequency phenomena observed in laboratory as well as space plasmas. The results has potential applications in the instability study of the space plasmas, and the fusion plasmas.

Methods

Numerical experiment is performed by using the one-dimensional PIC simulation method to study interaction between a KdV solitary wave and an envelope solitary wave in a dusty plasma in present work. During the simulation, The dust particles are represented as kinetic particles, while ions and electrons are modeled as Boltzmann distributed background. As well known, the real systems always contain very large amount of particles. In order to make simulations efficient or at least possible, so-called super-particles(SPs) are used. Each SP has a weight factor S specifying the number of real particles contained. Therefore, the equation of motion of the system is the Newton's equation as follows

$$m_j \frac{dv_j}{dt} = q_j E_j, \, \frac{dx}{dt} = v_j, \tag{6}$$

where m_j , q_j , x_j are the mass, charge and position of the *j*th SP, respectively. E_j is the electric field at the position of the *j*th SP. As the dust particles follow their trajectories, they continually exchange information with the background grid. Each dust particle contributes its charge to the corners of its instantaneous host cell. Therefore, the simulation region is divided to contain several grid cells during the PIC simulation. At each time step, the velocities, the positions of SPs are weighted to all the grids to calculate the charge density ρ_g (or electric current density J_g). Once ρ_g is obtained, the Maxwell's equations (electromagnetic model) or Poisson-Boltzmann equation (electrostatic model) will be solved numerically to derive the value of *E* at each grid. In electrostatic model, $B_g = 0$. Then the field imposed on each SP can be worked out and each SP will be driven by electric field according to Eq. (6), which will be solved numerically via the leap-frog algorithm. At last, the new positions and velocities are obtained, the procedure come to repeat until the simulation completed. The flowchart of PIC method is shown in Fig. 8.

In process of head-on collision between the KdV solitary wave and the envelope solitary waves, we assume that there are two opposite propagating solitary waves. the KdV solitary wave is propagating in the positive *x* direction and the envelop wave is in the negative *x* direction. The initial number density and the velocity of the dust particles are: $\phi_{nls} = \varepsilon_2 \sqrt{\frac{2(Pk_n^2 - \omega)}{Q}} \cdot \operatorname{sech} \sqrt{\frac{Pk_n^2 - \omega}{P}} \varepsilon_2 x \cdot \cos k_n (x_1 - x), u_{nls} = \frac{k_n}{\omega} \phi_{nls}$ and $n_{nls} = 1 + \frac{k_n^2}{\omega^2} \phi_{nls}$, where x_1 is the initial phase of the envelope solitary wave. The initial condition of the KdV solitary wave is also chosen from the analytical solution, $\phi_{kdv} = -\left(\frac{3\varepsilon_1^2 u_0}{\alpha}\right)\operatorname{sech}^2 \frac{(x - x_2) - \frac{1 + \varepsilon_1^2 u_0}{k_d}}{2\sqrt{\delta/(\varepsilon_1^2/k_d^2 u_0)}}, u_{kdv} = \phi_{kdv}$ and $n_{kdv} = 1 + \phi_{kdv}$, where x_2 is the initial phase of the KdV solitary wave. In the simulation we choose $x_1 = 2250, x_2 = 725$ which ensure that two solitary waves are far enough. The parameters of two colliding waves are $\varepsilon_1 = 0.002, \varepsilon_2 = 0.01, k_d = 1.0$.

two solitary waves are far enough. The parameters of two colliding waves are $\varepsilon_1 = 0.002$, $\varepsilon_2 = 0.01$, $k_d = 1.0$, $k_n = 0.1$, $\mu = 1.1$, $\nu = 0.1$, $\beta = 0.1$, $x_1 = LX/4$, $x_2 = 3LX/4$, $\Delta x = 0.1$, $\Delta t = \Delta x/100$, $N_x = 10000$, while $\omega = 0.1$, $u_s = -0.985$ and the number of super particles contained per cell is 100, where the negative sign stands for the envelop solitary wave propagating in the negative x direction.

In process of overtaking collision between KdV solitary wave and envelope solitary wave, both solitary waves propagate in same direction (positive x direction). The initial conditions are:

 $\phi_{nls} = \varepsilon_2 \sqrt{\frac{2(Pk_n^2 - \omega)}{Q}} \cdot \operatorname{sech} \sqrt{\frac{Pk_n^2 - \omega}{P}} \varepsilon_2 x \cdot \cos k_n (x - x_3), u_{nls} = -\frac{k_n}{\omega} \phi_{nls} \text{ and } n_{nls} = 1 - \frac{k_n^2}{\omega^2} \phi_{nls}, \text{ where } x_3 \text{ is the initial phase of the envelope solitary wave. The initial condition of the KdV solitary wave is also chosen from the$

analytical solution, $\phi_{kdv} = -\left(\frac{3\varepsilon_1^2 u_0}{\alpha}\right) \operatorname{sech}^2 \frac{(x-x_4) - \frac{1+\varepsilon_1^2 u_0}{k_d}}{2\sqrt{\delta/(\varepsilon_1^2 k_d^2 u_0)}}$, $u_{kdv} = \phi_{kdv}$ and $n_{kdv} = 1 + \phi_{kdv}$, where x_4 is the initial phase of the KdV solitary wave. The parameters of two colliding waves are $\varepsilon_1 = 0.1$, $\varepsilon_2 = 0.025$, $k_d = 1.0$, $k_n = 0.1$,

 $\mu = 1.1, \nu = 0.1, \beta = 0.1, x_3 = LX/2 = 1200, x_4 = LX/4 = 600, \Delta x = 0.3, \Delta t = \Delta x/100, N_x = 8000$, while $\omega = 0.1$, $u_s = 0.985.$

References

- 1. Abid, A. A., Ali, S., Du, J. & Mamun, A. A. Vasyliunas-Cairns distribution function for space plasma species. Phys. Plasmas 22, 084507 (2015).
- 2. Souza, V. M. Enhancement of Space Plasma Images by Complex Wavelets. Braz. J. Phys. 45, 510 (2015).
- 3. Klimushkin, D. Y. & Mager, P. N. Coupled Alfven and drift-mirror modes in non-uniform space plasmas: a gyrokinetic treatment. Plasma Phys. Control. Fusion 54, 015006 (2012).
- 4. Nickeler, D. H. & Wiegelmann, T. Thin current sheets caused by plasma flow gradients in space and astrophysical plasma. Ann. Geophys. 28, 1523 (2010).
- 5. Thoma, C., Rose, D. V., Miller, L. C., Clark, R. E. & Hughes, T. P. Electromagnetic wave propagation through an overdense magnetized collisional plasma layer. J. Appl. Phys. 106, 043301 (2009).
- 6. Kuo, S. P. & Faith, J. Interaction of an electromagnetic wave with a rapidly created spatially periodic plasma. Phys. Rev. E 56, 2143 (1997).
- 7. Joglekar, A. S., Thomas, A. G. R., Fox, W. & Bhattacharjee, A. Magnetic Reconnection in Plasma under Inertial Confinement Fusion Conditions Driven by Heat Flux Effects in Ohm's Law. Phys. Rev. Lett. 112, 105004 (2014).
- 8. Abel, I. G. et al. Multiscale gyrokinetics for rotating tokamak plasmas: fluctuations, transport and energy flows. Rep. Prog. Phys. 76, 116201 (2013)
- 9. Bertelli, N. et al. The influence of the edge density fluctuations on electron cyclotron wave beam propagation in tokamaks. Journal of Physics Conference Series 260, 012002 (2010).
- Kim, J. S. et al. Measurements of nonlinear energy transfer in turbulence in the Tokamak Fusion Test Reactor. Phys. Rev. Lett. 79, 841-844 (1997).
- 11. Chandrappan, J. et al. Target dependent femtosecond laser plasma implantation dynamics in enabling silica for high density erbium doping. Phys. Rev. Lett. 115, 083901 (2015).
- 12. Depresseux, A. et al. Demonstration of a Circularly Polarized Plasma-Based Soft-X-Ray Laser. Phys. Rev. Lett. 79, 841-844 (1997).
- 13. Avinash, K. & Sen, A. Rayleigh-Taylor instability in dusty plasma experiment. Phys. Plasmas 22, 083707 (2015).
- 14. Rao, N. N., Shukla, P. K. & Yu, M. Y. Dust-acoustic waves in dusty plasmas. Plant Space Sci. 38, 543 (1990).
- 15. Kourakis, I. & Shukla, P. K. Nonlinear theory of solitary waves associated with longitudinal particle motion in lattices-Application to longitudinal grain oscillations in a dust crystal. Eur. Phys. J. D 29, 247 (2004).
- 16. Zhdanov, S. K., Samsonov, D. & Morfill, G. E. Anisotropic plasma crystal solitons. Phys. Rev. E 66, 026411 (2002).
- 17. Harvey, P., Durniak, C., Samsonov, D. & Morfill, G. Soliton interaction in a complex plasma. Phys. Rev, E 81, 057401 (2010).
- 18. Bandyopadhyay, P., Prasad, G., Sen, A. & Kaw, P. K. Experimental study of nonlinear dust acoustic solitary waves in a dusty plasma. Phys. Rev. Lett. 101, 065006 (2008).
- Samsonov, D. et al. Dissipative Longitudinal Solitons in a Two-Dimensional Strongly Coupled Complex (Dusty) Plasma. Phys. Rev. Lett. 88, 095004 (2002).
- Bandyopadhyay, P., Prasad, G., Sen, A. & Kaw, P. K. Experimental Study of Nonlinear Dust Acoustic Solitary Waves in a Dusty 20. Plasma. Phys. Rev. Lett. 101, 065006 (2008).
- Heidemann, R., Zhdanov, S., Sutterlin, R., Thomas, H. M. & Morfill, G. E. Dissipative Dark Soliton in a Complex Plasma. Phys. Rev. Lett 102, 135002 (2009)
- 22. Amin, M. R., Morfill, G. E. & Shuklar, P. K. Modulational instability of dust-acoustic and dust-ion-acoustic waves. Phys. Rev. E 58, 5 (1998).
- Ghosh, S., Sarkar, S., Khan, M. & Gupta, M. R. Low-frequency wave modulations in an electronegative dusty plasma in the presence 23 of charge variations. Phys. Rev. E 84, 066401 (2011).
- Carbonaroa, P. Head-on collision of ion-acoustic solitary and shock waves in a two-electron-temperature plasma. Eur. Phys. J. D 66, 24. 302 (2012).
- 25. Huang, G. X. & Velarde, M. G. Head-on collision of two concentric cylindrical ion acoustic solitary waves. Phys. Rev. E 53, 2988 (1996).
- 26. El-Shami, E. F., Moslem, W. M. & Shukla, P. K. Head-on collision of ion-acoustic solitary waves in a Thomas-Fermi plasma containing degenerate electrons and positrons. Phys. Lett. A 374, 290 (2009).
- Su, C. H. & Mirie, R. M. On head-on collision between 2 solitary wavws. J. Fluid Mech. 98, 509 (1980).
- 28. Gardner, C. S., Greener, J. M., Kruskal, M. D. & Miura, R. M. Method for solving korteweg-deveris equation. Phys. Rev. Lett. 19, 1095 (1967)
- 29. Shay, M. A., Drake, J. F. & Dorland, B. Equation free projective integration: A multiscale method applied to a plasma ion acoustic wave. J. Comput. Phys. 226, 571 (2007).
- 30. Israelevich, P. L. & Ofman, L. Hybrid simulation of ion-acoustic waves excitation by non-linear Alfven wave. Adv. Space Res. 48, 25 (2011).
- 31. El-Shamy, E. F., Moslem, W. M. & Shukl, P. K. Head-on collision of ion-acoustic solitary waves in a Thomas-Fermi plasma containing degenerate electrons and positrons. Phys. Lett. A 374, 290 (2009).
- 32. El-Labany, S. K., El-Shamy, E. F., Sabry, R. & Shokry, M. Head-on collision of dust-acoustic solitary waves in an adiabatic hot dusty plasma with external oblique magnetic field and two-temperature ions. Astrophys. Space Sci. 325, 201 (2010).
- Chatterjee, P. et al. Head-on collision of ion acoustic solitary waves in an electron-positron-ion plasma with superthermal electrons. Phys. Plasmas 17, 122314 (2010).
- 34. Ghosh, U. N., Chatterjee, P. & Roychoudhury, R. The effect of q-distributed electrons on the head-on collision of ion acoustic solitary waves. Phys. Plasmas 19, 012113 (2012).
- Ghorui, M. K., Chatterje, P. & Wong, C. S. Head on collision of dust ion acoustic solitary waves in magnetized quantum dusty plasmas. Astrophys. Space Sci. 343, 639 (2013).
- 36. Harvey, P., Durniak, C., Samsonov, D. & Morfill, G. Soliton interaction in a complex plasma. Phys. Rev. E 81, 057401 (2010).
- 37. Sharma, S. K., Boruah, A. & Bailung, H. Head-on collision of dust-acoustic solitons in a strongly coupled dusty plasma. Phys. Rev. E 89,013110 (2014).

- Zhang, J. et al. The study of the Poincare-Lighthill-Kuo method by using the particle-in-cell simulation method in a dusty plasma. Phys. Plasmas 21, 103706 (2014).
- El-Labany, S. K., El-Taibany, W. F., El-Shamy, E. F., El-Depsy, A. & Zedan, N. A. Head-on-collision of modulated dust acoustic waves in strongly coupled dusty plasma. *Phys. Plasmas* 19, 103708 (2012).
- 40. Zhang, H., Qi, X., Duan, W. S. & Yang, L. Envelope solitary waves exist and collide head-on without phase shift in a dusty plasma. *Sci. Rep.* **5**, 14239 (2015).

Acknowledgements

This work is supported by National Science Foundation of China (Grant No. 11547304, 11275156,11505261) and the National Magnetic Confinement Fusion Science Program of China (Grant No. 2014GB104002), the Strategic Priority Research Program of Chinese Academy of Sciences (Grant No. XDA03030100).

Author Contributions

H.Z. wrote the main manuscript, developed algorithm and prepared all figures. X.Q. developed algorithm. W.-S.D. and L.Y. proposed the physical idea, analyzed results and revised the article. All authors review the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhang, H. *et al.* Head-on collision and overtaking collision between an envelope solitary wave and a KdV solitary wave in a dusty plasma. *Sci. Rep.* **6**, 21214; doi: 10.1038/srep21214 (2016).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/