SCIENTIFIC REPORTS

OPEN

Effect of excessive CO₂ on physiological functions in coastal diatom

Feng-Jiao Liu^{1,2}, Shun-Xing Li^{1,†}, Bang-Qin Huang², Feng-Ying Zheng¹ & Xu-Guang Huang¹

Received: 01 July 2015

Accepted: 28 January 2016

Published: 15 February 2016

Rising dissolution of anthropogenic CO₂ in seawater may directly/indirectly cause ocean acidification and desalination. However, little is known about coastal physiological functions sensitivity to these processes. Here we show some links between ocean acidification/desalination and physiological functions in *Thalassiosira weissflogii*. Cell density (CD), protein, chlorophyll *a* (Chl *a*), malonaldehyde (MDA), superoxide dismutase (SOD), and carbonic anhydrase (CAs) were determined for the assessment of algal biomass, nutritional value, photosynthesis and respiration, lipid peroxidation, antioxidant capacity, and carbon sequestration ability. The influence of pH on the algal Chl *a* and MDA were extremely significant ($P < 0.01$). Salinity (S) on cell density and acidity (pH) on protein was significant ($0.01 < P < 0.05$). Additionally, a significant negative-correlation was observed between cell density and CAs. CAs and SOD had negatively correlations with CD, Chl *a*, protein, and MDA under pH or S influence, but positive correlation between themselves. Coastal physiological functions were affected by increasing order was acidification < acidification + desalination < desalination for Chl *a* and protein, desalination < acidification + desalination < acidification for SOD and CAs. Thus, the ongoing excessive CO₂-driven ocean acidification and desalination should be of high attention when assessing the risks of climate change on coastal phytoplankton.

Human induced climate change effects marine environment in the next century and beyond. Future scenarios predict an increasing CO₂ partial pressure (pCO₂) in the atmosphere approximately from 380 to 1,000 ppm until the end of this century^{1–3}. Approximately a quarter of the anthropogenic CO₂ is absorbed by the ocean. This oceanic uptake of CO₂ leads to a change in marine carbonate chemistry resulting in a decrease of seawater pH and carbonate ion concentration, a process commonly called “Ocean Acidification (OA)”⁴. OA is a potential threat to marine ecosystems through its effect on the physiology and ecology of many marine species^{5–7}. OA is also likely to have measurable biological consequences on marine biodiversity and ecosystem functioning, as well as on the provision of ecosystem services^{8–11}.

Climate change results in ocean warming, sea ice coverage reducing (or sea ice melt increasing), global sea-level rising, surface evaporation increasing, and patterns of deep-water ventilation changing, potentially altering future surface ocean carbonate conditions and acidification^{7,12}. Along with warmer air and ocean temperatures, increased sea ice melt and had risen global sea-level can lead to salinity declining¹³. Simultaneously, climate change induced heavy precipitation is expected to reduce surface salinity¹⁴. Where mixing with fresh water runoff from river mouths, seawater salinity can be substantially decline.

The growth and photosynthesis of some seaweeds and seagrasses will be benefit under OA conditions¹⁵. Furthermore, OA could interact with other stressors, such as increasing light, warming water, decreasing oxygen concentration, overfishing, and after eutrophication, showing an additive effect on these stressors^{8,16}. The rising CO₂ combined with light stress may reduce primary production of phytoplankton¹⁷. However, the influence of ocean desalination and its coexistence with acidification on coastal physiological functions is unknown.

Coastal ecosystems account for over a third of the world's ecosystems services, such as nitrogen fixation, phytoplankton abundance and production, bacteria growth rates, food-web type, and the interactions between animals, plants and bacteria^{18–22}. Diatoms are the dominant group of phytoplankton in the modern ocean, specially

¹Fujian Province Key Laboratory of Modern Analytical Science and Separation Technology, Minnan Normal University, Zhangzhou, China, 363000. ²College of the Environment & Ecology, Xiamen University, 361005, China.

[†]Present address: College of Chemistry and Environment, Minnan Normal University, Zhangzhou, China, 363000. Correspondence and requests for materials should be addressed to S.-X.L. (email: shunxing_li@aliyun.com) or B.-Q.H. (email: bqhuang@xmu.edu.cn)

Figure 1. Influence of salinity (S) on algal density. (a) S = Blank; (b) S = 29; (c) S = 30; (d) S = 31; (e) S = 32. Data are mean $\pm$ SD ($n = 3$).

in well-mixed coastal upwelling regions^{23,24}, accounting for approximately 40% of oceanic primary productivity and critical foundation of coastal food web^{25,26}. Coastal physiological functions (including cell density (CD), protein, chlorophyll *a* (Chl *a*), malonaldehyde (MDA), superoxide dismutase (SOD), and carbonic anhydrase (CAs)) have been used in our laboratory for the estimation of algal biomass, nutritional value, photosynthesis and respiration, lipid peroxidation, antioxidant capacity, and carbon sequestration ability, respectively. Algal density controls the stability of food chains and the energy transfer in the food chains of marine systems²⁷. Protein synthesis is one of nitrogen assimilations and its concentration is an indicator of metabolic capacity and nutritional value²⁸. Chlorophyll *a* is an important material for algal photosynthesis and respiration, and its concentration is an important indicator for estimating algal primary productivity²⁸. SOD is one kind of metalloenzyme that can catalyze the dismutation process of superoxide anion into oxygen and hydrogen peroxide. Thus SOD provides protection for the survival of aerobic organisms^{29,30}. MDA, a product of lipid peroxidation, can be used as an indicator of free radical activity and tissue damage³¹. CAs is a zinc-containing metalloenzyme that catalyzes the interconversion of CO_2 and HCO_3^- in many organisms, has been taken as a component of carbon-concentrating mechanisms and plays a role in photosynthetic CO_2 fixation^{32,33}.

Results

Influence of S or/and pH on Cell Density. Cell growth rate was controlled by the speed of cell division. The results were shown in Figs 1 and 2 indicated that cell division speed could be inhibited by desalination but promoted by acidification. When coexistence with S and pH, cell division could be promoted, i.e., cell densities were all higher than blank experiment, but cell growth cycles were shorten significantly, the results were shown in Fig 2. When pH was 7.8 and S was decreased from 32 to 30, cell growth was increased continuously, but cell density was decreased. The overall observed changes was acidification < acidification + desalination < desalination.

Influence of S or/and pH on Phytoplankton Physiological Properties. The influence of S or/and pH on coastal diatom can be seen in Figs 3–7, including Chl *a*, protein, MDA, SOD, and CAs. Figures 3–5 mainly display the effect of S on concentration of Chl *a*, protein, and MDA was clearly obvious when compared with pH. However, the influence trends were reversed as SOD and CAs were observed to be low. The variation trend of protein and MDA affected by S and pH was similar, but that was reverse to CA activity; Influence trend of pH on Chl *a*, SOD, and CAs was also similar.

Coexisted with S and pH, the concentrations of Chl *a* and activities of CAs were decreased with desalination when pH was 7.8 and 8.1, protein was similar when S from 32 to 29 and pH was 8.1. At the same time, the concentrations of MDA and protein and activities of CAs were decreased with S decreased from 31–29 when pH was 7.9, but the influence trends were reversed while S was over the above mentioned range. When pH was 7.9 and 8.0 the influence of desalination on SOD activity was similar.

The influence degree on coastal physiological functions was different, acidification < acidification + desalination < desalination for Chl *a* and protein, desalination < acidification + desalination < acidification for SOD and CAs.

There were significant relationships in the six tested parameters, although these were not always consistent (Table S1). There was a very significant negative correlations between CD and CAs ($r = -0.996$, $P < 0.01$, S; $r = -0.966$, $P < 0.05$, pH). CAs and SOD had negatively correlations with CD, Chl *a*, protein, and MDA under pH or S influence, but positive correlation between themselves ($r = 0.950$, pH; $r = 0.879$, S). Between protein and MDA was also positively correlated ($r = 0.900$, S; $r = 0.870$, pH).

These data were much higher than their corresponding detection limit and within their linear range. The varying of these data could be accepted, shown in Figs 1–7. According to ANOVA analysis as shown in Table S2, the influence of pH on the algal Chl *a* and MDA were extremely significant ($P < 0.01$, $F > F_{\alpha}$), S on CD and pH on protein were significant ($0.01 < P < 0.05$, $F > F_{\alpha}$).

Figure 2. Influence of salinity (S) and acidity (pH) on algal density. Data are mean $\pm$ SD ($n = 3$).

Discussion

Cell growth rate is controlled by the speed of cell division and directly affected by ocean acidification³⁴. Similar to previously published results^{35,36}, our study concurs with these observations, showing that algal density decreased with an increase of acidify (e.g. by decreasing pH). Rising atmospheric CO_2 could increase dissolved CO_2 concentration in seawater, a moderate increase in CO_2 facilitates photosynthetic carbon fixation of some phytoplankton groups³⁷. However, when the acidity is lower than a certain concentration, it will damage cellular structure and physiological functions³⁴. The reduced salinity can cause further osmotic challenge and elevate basal energy demand in a decreased pH environment, leading to lower energy levels available for growth or shell formation in marine organisms^{38,39}. Thus, it can mitigate biological effects for microalgae. High concentrations of inorganic anions and cations from highly saline environments can potentially reduce the permeability of ionic liquid cations through algal cell walls⁴⁰. Therefore, excessive concentrations of salinity can have a harmful effect on algal cell growth. Our results were agreed well with the results by Stedmon *et al.*⁴¹.

Figure 3. (a) Influence of salinity (S) or acidity (pH) on the content of chlorophyll a (Chl a); (b) Influence of S and pH on the content of Chl a. Data are mean $\pm$ SD (n = 3).

Figure 4. (a) Influence of salinity (S) or acidity (pH) on the content of malonaldehyde (MDA); (b) Influence of S and pH on the content of MDA. Data are mean $\pm$ SD (n = 3).

As previously reported that the synthesis process of protein and Chl a is controlled by the activity of NR, GS, G-tRNAs, and NADPH^{27,28}. Ocean acidification may enhance enzyme activity, including NR, GS, G-tRNAs, and NADPH⁴², and then promote the synthesis of protein (including the other enzymes) and chlorophyll a. When pH

Figure 5. (a) Influence of salinity (S) or acidity (pH) on the content of protein; (b) Influence of S and pH on the content of protein. Data are mean $\pm$ SD (n = 3).

Figure 6. (a) Influence of salinity (S) or acidity (pH) on the activity of carbonic anhydrase (CA); (b) Influence of S and pH on the activity of CA. Data are mean $\pm$ SD (n = 3).

was more than a certain concentration, algal cell could induce the overload of hydroxyl radicals, and then inhibited the activities of NR, GS, glutamate-tRNA synthase, and NADPH. Therefore, the influence trends of pH on

Figure 7. (a) Influence of salinity (S) or acidity (pH) on the activity of superoxide dismutase (SOD); (b) Influence of S and pH on the activity of SOD. Data are mean $\pm$ SD ($n = 3$).

the content of protein and chlorophyll *a* were similar and significant ($P < 0.05$, $F > F_{\alpha}$). Our results were in good agreement with the reported influence trends⁴².

SOD activity and MDA content would be changed with the amount of hydroxyl radical (one kind of reactive oxygen species). According to our results, both S and pH could induce the oxidant stress in algal cells. With desalination and acidification, MDA content of algal cells increased, indicating that membrane lipid peroxidation was enhanced. The change in SOD activity of algal cells also confirms the generation of oxidant stress.

CAs abundant enzymes play a key role in carbon-concentrating mechanism of marine phytoplankton. The synthesis process is controlled by an essential element (Zn) in phytoplankton, because of Zn is used as a cofactor in CAs⁴³. The decrease in pH will result in reducing the concentrations of hydroxide (OH^-) in most natural surface waters. Ocean desalination also will reduce the concentration of anion, such as Cl^- and OH^- . When pH in the range from 7.8 to 8.1 and S from 29 to 32, the speciation of Zn included ZnCO_3 , $\text{Zn}(\text{CO}_3)_2^{2-}$, $\text{Zn}(\text{OH})^+$, ZnCl^+ , Zn^{2+} , ZnClOH , ZnCl_2 , and ZnHCO_3^+ , only a few fraction of Zn^{2+} could be existed. Thus, S and pH could inhibit CAs activity through affecting the availability of Zn. CAs expression is very sensitive to variations in $\text{pH}-P_{\text{CO}_2}$ ^{44–46}. Previous studies have also shown a substantial decrease in CAs activity in the diatoms *T. weissflogii* when pH decreases⁴³.

Coastal physiological functions could be affected seriously by the carbon dioxide-associated ocean acidification and desalination. Here fully investigated the coexistence of ocean acidification and desalination affect algal biomass, nutritional value, photosynthesis and respiration, lipid peroxidation, antioxidant capacity, and carbon sequestration ability, respectively. At the same time, desalination can be related to the influence of dilute water mouth, and coexistence of acidification and desalination can also be linked to estuary affected by acid rain. However, there has only laboratory tests and should be further investigated and most importantly on scale-up field experiment for a future angle to risk assessment climate change on coastal environment.

Methods

Seawater Sample. Seawater was collected from the Taiwan strait (22.65 N, 118.82E) shown in Figure S1 (The R Programming Language, 3.0.3), stored at 4°C for about 6 months, and filtered through 0.22 μm acid-washed capsule filters (Pall Supor membrane, Acropak 200) before use. The salinity (S) and acidity (pH) were measured three times using Salinometer Meter and Delta 320-SpH meter (Mettler-Toledo, Greifensee, Switzerland). The Sal and pH were 32.5 and 8.10, respectively.

Phytoplankton culture. Unialgal cultures of *T. weissflogii* were obtained from the State Key Laboratory for Marine Environmental Science, Xiamen University. The cultured growth cycle curve was shown in Figure S2. Exponentially growing cells of *T. weissflogii* cells were filtered and transferred to new medium every 1–2 days and then the cells would adapt to the tested experimental conditions. After 4 transfers, the cells (1×10^4

cells mL⁻¹) were again filtered and added into 5 L of seawater (filtered with 0.22 µm acid-washed capsule filters) in acid-cleaned polycarbonate bottles. Algal cells were illuminated with a light intensity of 140 µmol photons m⁻² s⁻¹ on a 14 hr: 10 hr light: dark cycle and maintained in seawater at various concentrations of S (29, 30, 31, and 32, respectively) and/or pH/ pCO₂ (7.8/770 ppm, predicted CO₂ levels in 2100; 7.9/600 ppm, predicted CO₂ levels in 2060; 8.0/470 ppm, predicted CO₂ levels in 2030; and 8.1/380, current CO₂ levels, respectively)^{8,47} at 19°C. After determination of S and pH in the medium, their values were maintained through compensating addition daily of Na₂CO₃ (0.1 mol/L) and CO₂ for 4 days, respectively, i.e., semi-continuous culture was adopted. Na₂CO₃ and CO₂ were added by drops and gently bubbled with humidified, CO₂-enriched air, respectively. Marine carbonate system considers CO₂ (aq) [the sum of CO₂ and H₂CO₃], HCO₃⁻, CO₃²⁻, H⁺, OH⁻⁴⁸. The additional CO₂ causes re-equilibration of the seawater carbonate system, increasing the concentrations of aqueous CO₂ and bicarbonate ion, HCO₃⁻, while decreasing that of the carbonate ion, CO₃²⁻. Dissolved inorganic carbon (DIC) is the sum of all dissolved inorganic carbon species, which is the main acid-base buffer of seawater³⁴. When addition of Na₂CO₃ and CO₂, the medium was stirring until desirable S and pH were achieved stably. The algal suspensions were stirred at 100 rpm to simulate the current of seawater. Large culture vessels (5 L) were used so as to decrease the thickness of marine phytoplankton suspension to minimize the difference of light illumination between the surface and bottom of marine phytoplankton suspension⁴⁹. These experiments were triplicated ($n = 3$).

Test methods. Cell density was counted microscopically. SOD activity, and the content of protein, Chl *a*, and MDA was determined by monitoring the inhibition of photochemical reduction of nitroblue tetrazolium chloride, coomassie brilliant blue method, spectrophotometric method, and thiobarbituric acid method, respectively, as described in detail previously²⁷. CA activity was measured by an electrometric method as described by Chen and Gao³².

Statistical analysis. Analysis of variance was calculated by using SASPROC MIXED⁵⁰ and *p* value was calculated using two-way ANOVA. The methods of statistical analysis as described in detail previously²⁸. *Post hoc* tests were examined using Tukey's test for two-way ANOVA and Dunnett's Test for multivariate ANOVA⁷.

References

- Kranz, S. A. *et al.* Combined effects of CO₂ and light on the N₂-fixing cyanobacterium Trichodesmium IMS101: physiological responses. *Plant physiol.* **154**, 334–345 (2010).
- Raven, J. *et al.* Ocean acidification due to increasing atmospheric carbon dioxide. *The Royal Society*, 1–68 (2005).
- Raupach, M. R. *et al.* Global and regional drivers of accelerating CO₂ emissions. *Proceed. Natl. Acad. Sci.* **104**, 10288–10293 (2007).
- Land, P. E. *et al.* Salinity from space unlocks satellite-based assessment of ocean acidification. *Environ. Sci. Technol.* **49**, 1987–1994 (2015).
- Kroeker, K. J. *et al.* Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Global Change Biol.* **19**, 1884–1896 (2013).
- Cornwall, C. E. & Hurd, C. L. Experimental design in ocean acidification research: problems and solutions. *ICES J. Mar. Sci.* **118** (2015).
- Gattuso, J. P. *et al.* Contrasting futures for ocean and society from different anthropogenic CO₂ emissions scenarios. *Science* **349**, 4722 (2015).
- Xu, D. *et al.* Effects of CO₂ and seawater acidification on the early stages of *Saccharina saccharina japonica* development. *Environ. Sci. Technol.* **49**, 3548–3556 (2015).
- Hofmann, G. E. *et al.* The effect of ocean acidification on calcifying organisms in marine ecosystems: an organism-to-ecosystem perspective. *Annu. Rev. Ecol. Evol. S.* **41**, 127–147 (2010).
- Fabry, V. J., Seibel, B. A., Feely, R. A. & Orr, J. C. Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES J. Mar. Sci. J. du Conseil* **65**, 414–432 (2008).
- Hall-Spencer, J. M. *et al.* Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature* **454**, 96–99 (2008).
- McNeil, B. I. & Matear, R. J. Southern Ocean acidification: A tipping point at 450-ppm atmospheric CO₂. *Proceed. Natl. Acad. Sci.* **105**, 18860–18864 (2008).
- Massom, R. A. & Stammerjohn, S. E. Antarctic sea ice change and variability – Physical and ecological implications. *Polar Sci.* **4**, 149–186 (2010).
- Groisman, P. Y. *et al.* Changes in the probability of heavy precipitation: important indicators of climatic change. *Clim Change* **42**, 243–283 (1999).
- Zou, D. Effects of elevated atmospheric CO₂ on growth, photosynthesis and nitrogen metabolism in the economic brown seaweed, *Hizikia fusiforme* (Sargassaceae, Phaeophyta). *Aquaculture* **250**, 726–735 (2005).
- Harvey, B. P., Gwynn-Jones, D. & Moore, P. J. Meta-analysis reveals complex marine biological responses to the interactive effects of ocean acidification and warming. *Ecol. Evol.* **3**, 1016–1030 (2013).
- Gao, K. *et al.* Rising CO₂ and increased light exposure synergistically reduce marine primary productivity. *Nat. Clim. change* **2**, 519–523 (2012).
- Sam Dupont, H. P. Get ready for ocean acidification. *Nature* **498**, 429 (2013).
- Powers, J. E. Fisheries: Measuring biodiversity in marine ecosystems. *Nature* **468**, 385–386 (2010).
- Worm, B. *et al.* Impacts of biodiversity loss on ocean ecosystem services. *Science* **314**, 787–790 (2006).
- Smith, V. H. Eutrophication of freshwater and coastal marine ecosystems a global problem. *Environ. Sci. Pollut. Res.* **10**, 126–139 (2003).
- Shaked, Y. & Lis, H. Disassembling iron availability to phytoplankton. *Front Microbiol* **3**, 123 (2012).
- Chao, M., Shen, X., Lun, F., Shen, A. & Yuan, Q. Toxicity of Fuel Oil Water Accommodated Fractions on Two Marine Microalgae, *Skeletonema costatum* and *Chlorella* spp. *Bull. Environ. Contam. Toxicol.* **88**, 712–716 (2012).
- Ken, C. F., Hsiung, T. M., Huang, Z. X., Juang, R. H. & Lin, C. T. Characterization of Fe/Mn-superoxide dismutase from diatom *Thalassiosira weissflogii*: cloning, expression, and property. *J. Agr. Food Chem.* **53**, 1470–1474 (2005).
- Fakness, L. G., Brandvik, P. J. & Sydnes, L. K. Composition of the water accommodated fractions as a function of exposure times and temperatures. *Mar. Pollut. Bull.* **56**, 1746–1754 (2008).
- Rabosky, D. L. & Sorhannus, U. Diversity dynamics of marine planktonic diatoms across the Cenozoic. *Nature* **457**, 183–186 (2009).
- Liu, F. J. *et al.* Risk assessment of nitrate and oxytetracycline addition on coastal ecosystem functions. *Aquat. Toxicol.* **146**, 76–81 (2014).
- Li, S. X., Liu, F. J., Zheng, F. Y., Huang, X. G. & Zuo, Y. G. Risk assessment of nitrate and petroleum-derived hydrocarbon addition on *Contricribia weissflogii* biomass, lifetime, and nutritional value. *J. Hazard. Mater.* **268**, 199–206 (2014).

29. Learman, D. R., Voelker, B. M., Vazquez-Rodriguez, A. I. & Hansel, C. M. Formation of manganese oxides by bacterially generated superoxide. *Nat. Geo.* **4**, 95–98 (2011).
30. Choudhary, M., Jetley, U. K., Abash Khan, M., Zutshi, S. & Fatma, T. Effect of heavy metal stress on proline, malondialdehyde, and superoxide dismutase activity in the cyanobacterium *Spirulina platensis*-S5. *Ecotoxicol. Environ. Saf.* **66**, 204–209 (2007).
31. Jones, M. N. Nitrate reduction by shaking with cadmium alternative to cadmium columns. *Water Res.* **18**, 643–646 (1984).
32. Chen, X. W. & Gao, K. S. Roles of carbonic anhydrase in photosynthesis of *Skeletonema costatum*. *J. Plant Physiol. Mol. Biol.* **30**, 511–516 (2004).
33. Murray, R. & Badger, G. D. P. The role of carbonic anhydrase in photosynthesis. *Annu. Rev. Plant. Physiol. Plant. Mol. Biol.* **45**, 369–392 (1994).
34. Shi, D., Xu, Y. & Morel, F. Effects of the pH/pCO₂ control method on medium chemistry and phytoplankton growth. *Biogeosciences* **6**, 1199–1207 (2009).
35. Shi, D., Kranz, S. A., Kim, J. M. & Morel, F. M. Ocean acidification slows nitrogen fixation and growth in the dominant diazotroph *Trichodesmium* under low-iron conditions. *Proceed. Natl. Acad. Sci.* **109**, 3094–3100 (2012).
36. Tortell, P. D. *et al.* CO₂ sensitivity of Southern Ocean phytoplankton. *Geophys. Res. Lett.* **35**, L04605 (2008).
37. Riebesell, U. Effects of CO₂ enrichment on marine phytoplankton. *J. Oceanogr.* **60**, 719–729 (2004).
38. Ko, G. W. *et al.* Interactive effects of ocean acidification, elevated temperature, and reduced salinity on early-life stages of the Pacific oyster. *Environ. Sci. Technol.* **48**, 10079–10088 (2014).
39. Dickinson, G. H. *et al.* Interactive effects of salinity and elevated CO₂ levels on juvenile eastern oysters, *Crassostrea virginica*. *J. Exp. Biol.* **215**, 29–43 (2012).
40. Samorì, C. *et al.* Effects of imidazolium ionic liquids on growth, photosynthetic efficiency, and cellular components of the diatoms *Skeletonema marinoi* and *Phaeodactylum tricornutum*. *Chem. Res. Toxicol.* **24**, 392–401 (2011).
41. Stedmon, C. A. *et al.* Characteristics of dissolved organic matter in Baltic coastal sea ice: allochthonous or autochthonous origins? *Environ. Sci. Technol.* **41**, 7273–7279 (2007).
42. Rouco, M., Branson O., Lebrato, M. & Iglesias-Rodríguez, M. D. The effect of nitrate and phosphate availability on *Emiliania huxleyi* (NZEH) physiology under different CO₂ scenarios. *Front. Microbiol.* **4**, 155 (2013).
43. Xu, Y., Shi, D., Aristilde, L. & Morel, F. M. The effect of pH on the uptake of zinc and cadmium in marine phytoplankton: Possible role of weak complexes. *Limnol. Oceanogr.* **57**, 293 (2012).
44. Lane, T. W. & Morel, F. M. Regulation of carbonic anhydrase expression by zinc, cobalt, and carbon dioxide in the marine diatom *Thalassiosira weissflogii*. *Plant Physiol.* **123**, 345–352 (2000).
45. Sunda, W. G. & Huntsman, S. A. Effect of CO₂ supply and demand on zinc uptake and growth limitation in a coastal diatom. *Limnol. Oceanogr.* **50**, 1181 (2005).
46. McGinn, P. J. & Morel, F. M. Expression and regulation of carbonic anhydrases in the marine diatom *Thalassiosira pseudonana* and in natural phytoplankton assemblages from Great Bay, New Jersey. *Physiol. Plantarum* **133**, 78–91 (2008).
47. Ellycia Harrould-Kolieb, J. S. Acid Test: can we save our oceans from CO₂? *Oceana* **2**, 1–32 (2009).
48. Gattuso, J. P., Lee, K., Björn, Rost & Schulz, Kai G.. Approaches and tools to manipulate the carbonate chemistry. In: Riebesell U., Fabry V. J., Hansson L. & Gattuso J.-P. (Eds), *Guide to best practices for ocean acidification research and data reporting Luxembourg: Publications Office of the European Union*, 41–52 (2010).
49. Li, S. X., Liu, F. J., Zheng, F. Y., Zuo, Y. G. & Huang, X. G. Effects of nitrate addition and iron speciation on trace element transfer in coastal food webs under phosphate and iron enrichment. *Chemosphere* **91**, 1486–1494 (2013).
50. Littell, R., Milliken, G., Stroup, W. & Wolfinger, R. Cary, NC: SAS Institute Inc. SAS system for mixed models: Some bayesian approaches to mixed models. *Ch.* **13**, 497–524 (1996).

Acknowledgements

This work was supported by the Natural Science Foundation of China (no. 41206096, 21475055, 40506020, and 21175115), the Program for New Century Excellent Talents in University (NCET-110904), Project of Fujian Education Department, China (JA09164), the Science and Technology Committee of Fujian Province, China (2012Y0065), and State Oceanic Administration project (201105021).

Author Contributions

S.X.L. designed of research idea and experimental method, analyzed of experimental data. Experiment and data analysis is done by F.J.L., B.Q.H., F.Y.Z. and X.G.H. All authors co-wrote the paper.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Feng-Jiao, L. *et al.* Effect of excessive CO₂ on physiological functions in coastal diatom. *Sci. Rep.* **6**, 21694; doi: 10.1038/srep21694 (2016).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

SCIENTIFIC REPORTS

OPEN

Corrigendum: Effect of excessive CO₂ on physiological functions in coastal diatom

Feng-Jiao Liu, Shun-Xing Li, Bang-Qin Huang, Feng-Ying Zheng & Xu-Guang Huang

Scientific Reports 6:21694; doi: 10.1038/srep21694; published online 15 February 2016; updated 09 May 2016

The original version of this Article incorrectly listed 'College of Chemistry and Environment, Minnan Normal University, Zhangzhou, China, 363000' as a present address for Bang-Qin Huang rather than Shun-Xing Li.

This error has now been corrected in the PDF and HTML versions of the Article.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>