SCIENTIFIC REPORTS

OPEN

Reconstructing the complex evolutionary history of mobile plasmids in red algal genomes

JunMo Lee¹, Kyeong Mi Kim², Eun Chan Yang³, Kathy Ann Miller⁴, Sung Min Boo⁵, Debashish Bhattacharya⁶ & Hwan Su Yoon¹

Received: 17 July 2015

Accepted: 14 March 2016

Published: 31 March 2016

The integration of foreign DNA into algal and plant plastid genomes is a rare event, with only a few known examples of horizontal gene transfer (HGT). Plasmids, which are well-studied drivers of HGT in prokaryotes, have been reported previously in red algae (Rhodophyta). However, the distribution of these mobile DNA elements and their sites of integration into the plastid (ptDNA), mitochondrial (mtDNA), and nuclear genomes of Rhodophyta remain unknown. Here we reconstructed the complex evolutionary history of plasmid-derived DNAs in red algae. Comparative analysis of 21 rhodophyte ptDNAs, including new genome data for 5 species, turned up 22 plasmid-derived open reading frames (ORFs) that showed syntenic and copy number variation among species, but were conserved within different individuals in three lineages. Several plasmid-derived homologs were found not only in ptDNA but also in mtDNA and in the nuclear genome of green plants, stramenopiles, and rhizarians. Phylogenetic and plasmid-derived ORF analyses showed that the majority of plasmid DNAs originated within red algae, whereas others were derived from cyanobacteria, other bacteria, and viruses. Our results elucidate the evolution of plasmid DNAs in red algae and suggest that they spread as parasitic genetic elements. This hypothesis is consistent with their sporadic distribution within Rhodophyta.

Horizontal gene transfer (HGT) plays a significant role in the evolution of bacterial genomes, promoting environmental adaptation and speciation. Plasmids drive HGT by moving DNA from one genome to another, often between species, in the absence of sexual reproduction^{1–3}. However, the mechanism of eukaryotic HGT is poorly understood, although it is known to occur from prokaryotes to eukaryotes^{4–7}, between different eukaryotes^{8–10}, and from eukaryotes to prokaryotes^{11,12}. A special case of HGT, endosymbiotic gene transfer (EGT), is responsible for massive amounts of intracellular gene movement in eukaryotes. This is an outcome of organellogenesis, whereby 100 s to 1000 s of genes were transferred from the bacterium-derived organelle genomes (i.e., the mitochondrion and plastid) to the host nuclear chromosomes^{13–16}.

Nuclear-encoded, plastid-derived genes have been studied in the glaucophyte alga *Cyanophora paradoxa* Korshikov (6–11%)^{17,18}; the red alga *Cyanidioschyzon merolae* DeLuca, Taddei, & Varano (6–20%)^{19–21}; the green algae *Chlamydomonas reinhardtii* P.C.A. Dangeard (6–14%) and *Ostreococcus tauri* C. Courties & M.-J. Chrétiennot-Dinet (11%)^{20–22}; and in *Arabidopsis thaliana* (L.) Heynh. and other land plants (9–18%)^{20,21,23}. EGT is essentially uni-directional. As a consequence, organelle (e.g., plastid) genomes have been reduced to 100 ~200 Kbp from their original size of several megabases in the cyanobacterial endosymbiont. It is not known, however, if plasmids may have facilitated EGT in algae and plants, thereby contributing significantly to their genome reduction.

HGTs have also been reported between organelle genomes of unrelated organisms. The plastid genome of the common milkweed *Asclepias syriaca* L. contains several mitochondrial genes²⁴, whereas the mitochondrial genome of *Amborella trichopoda* Baill. contains mtDNAs from green algae (including the entire mitochondrial genome in three species), mosses, and other angiosperms^{25–27}. The maize mitochondrial S-1 plasmid was found

¹Department of Biological Sciences, Sungkyunkwan University, Suwon, 16419, Korea. ²Marine Biodiversity Institute of Korea, Seocheon, 325-902, Korea. ³Marine Ecosystem Research Division, Korea Institute of Ocean Science & Technology, Ansan, 15627, Korea. ⁴Herbarium, University of California at Berkeley, 1001 Valley Life Sciences Building 2465, Berkeley, California, 94720-2465, USA. ⁵Department of Biology, Chungnam National University, Daejeon, 34134, Korea. ⁶Department of Ecology, Evolution and Natural Resources and Department of Marine and Coastal Sciences, Rutgers University, New Brunswick, NJ 08901, USA. Correspondence and requests for materials should be addressed to H.S.Y. (email: hsyoon2011@skku.edu)

in the mitochondrial genome of the liverwort *Marchantia polymorpha* L.^{28,29}. Interestingly, these sequences are similar to the mitochondrial *dpo* gene in the red alga *Porphyra* and the golden-brown alga *Ochromonas danica* E. Pringsheim³⁰, suggesting the existence of HGT among different phyla. Plasmid-derived sequences were also reported from mtDNA in the brown alga *Pylaiella littoralis* (L.) Kjellman³¹ and two fungal species, *Agaricus bisporus* (J.E. Lange) Imbach and *Gigaspora rosea* T.H. Nicolson & N.C. Schenck^{32–33}. The plastid genomes of photo-synthetic haptophytes and cryptophytes contain bacterial-derived *rpl36* genes⁷, and the cryptophyte *Rhodomonas salina* D.R.A. Hill & R. Wetherbee has a bacterial *dnaX* gene in its ptDNA³⁴. These examples demonstrated significant HGT between organelles and between organelles and plasmids; however, no such data has been reported for the red algae (Rhodophyta).

In red algae, proteobacterial operons related to leucine biosynthesis (*leuC* and *leuD* subunits) are encoded in the plastid genome of *Gracilaria tenuistipitata* var. *liui* J. Zhang, & B. Xia^{35,36}. This gene cluster (*leuA/B/C/D*) was traced to a plasmid from *Buchnera*, a genus of bacterial endosymbionts in aphids^{37,38}. The ptDNA of the red alga *Pyropia haitanensis* (T. J. Chang & B.F. Zheng) N. Kikuchi & M. Miyata contains plasmid-derived sequences that were discovered in the plasmid of another red alga, *Porphyra pulchra* G.J. Hollenberg^{39,40}. The mtDNA of other red algae, *Gracilaria chilensis* C.J. Bird, J.L. McLachlan, & E.C. de Oliveira, *Gracilariopsis chorda* (E.M. Holmes) Ohmi and *Gracilariopsis lemaneiformis* (Bory de Saint-Vincent) Dawson, Acleto, & Foldvik, contain partial plasmid sequences that have been reported in *Gracilaria robusta* Setchell^{41–43}. However, less is known about the mechanisms of plasmid-derived HGT to the plastid genome.

Plasmids are extrachromosomal genetic materials that are generally referred to as autonomously replicating double-stranded, circular or linear DNA molecules⁴⁴. About 25% of red algal genera contain more than two plasmids per species, and encode open reading frames (ORFs) that are transcriptionally active⁴⁵. Eukaryotic plasmids are widely distributed throughout algae, land plants, fungi, yeast, and other eukaryotes. However, their origins are poorly understood and their functions, including pathogenicity, have been reported only in a few cases^{45–48}. Of 35 red algal species assessed for plasmid sequences, 5 species contain 14 plasmid sequences: *Porphyra pulchra* (five plasmids); *Pyropia tenera* (Kjellman) N. Kikuchi, M. Miyata, M.S. Hwang & H.G. Choi (two plasmids); *Gracilaria chilensis* (three plasmids); *G. robusta* (two plasmids); and *Gracilariopsis lemaneiformis* (two plasmids)^{39,45,49–52}. However, no comprehensive analysis has yet been done to investigate the evolutionary relationship between plasmid DNA and plastid genomes.

To this end, we sequenced five red algal ptDNAs, including two that are plasmid-rich from *Gracilaria chilensis* and *Porphyra pulchra*. We analyzed plasmid-derived sequences from a total of 21 available red algal plastid genomes^{35,36,40,53–61} to elucidate the impact of plasmids over the > 1 billion year evolutionary history of red algae.

Results and discussion

Novel red algal plastid genomes. Five novel plastid genomes were completed using next-generation sequencing (NGS) data from *Gelidium elegans* (1,529 Mbp of total data), *G. vagum* (990 Mbp), *Gracilaria chilensis* (506 Mbp), *Porphyra pulchra* (263 Mbp) and *Sporolithon durum* (3,190 Mbp). The range of average genome coverage from the raw data was 52–445x (Supplementary Table S1). The plastid genome of *P. pulchra* (Supplementary Fig. S1) was the largest (194,175 bp) and had a higher GC-content (33.3%) than that of *S. durum* (191,465 bp, GC = 29.3%, Supplementary Fig. S2), *G. elegans* (174,748 bp, GC = 30.2%, Supplementary Fig. S3), *G. vagum* (179,853 bp, GC = 29.9%, Supplementary Fig. S4) and *Gr. chilensis* (185,637 bp, GC = 29.3%, Supplementary Fig. S5). Basic information about these plastid genomes is summarized in Supplementary Table S2. The plastid genome of *P. pulchra*, similar to those in other bangiophycean species, comprised 207 protein-coding genes, 37 tRNAs and 6 rRNAs; the rRNA operon (*rrs*, *rnl* and *rrf*) was duplicated. Among the florideophycean species, *S. durum* comprised 202 protein-coding genes, 30 tRNAs, 3 rRNAs, 3 rRNAs and 2 introns, as well as several pseudogenes (*dnaB*, *syfB*, *ycf21* and *ycf23*). This genome lacked the *syh* gene and *trnV* tRNA, both of which are present in another member of the Corallinophycidae, *Calliarthron tuberculosum*. These two coralline algae have a unique group II intron in the *chlB* gene³⁶ with intronic *orfs*. *Gelidium vagum* contained 201 protein-coding genes with pseudogenes of *ycf34*; *G. elegans* contained 202 protein-coding genes. These members of the order Gelidiales encode 30 tRNAs, 3 rRNAs and a group II intron in *trnMe* tRNA³⁶. The plastid genome of *Gracilaria chilensis* (order Gracilariales) contained 203 protein-coding genes, 30 tRNAs, 3 rRNAs and a group II intron in *trnMe* tRNA, which had not been found previously in the plastid genomes of *G. salicornia* and *G. tenuistipitata* var. *liui* or in *Grateloupia taiwanensis* (Halymeniales).

The ML tree inferred from the concatenated dataset of 193 plastid protein-coding genes (Supplementary Table S3; Table S4) resolved phylogenetic relationships among red algae (Fig. 1A, Supplementary Fig. S6). The early diverging Cyanidiophyceae was chosen as the outgroup for this phylogeny^{62,63}. The Bangiophyceae and the Florideophyceae grouped together with maximum ML bootstrap support value (MLB, 100%), and each class formed a strongly supported monophyletic clade, as previously reported^{62–65}. Within the Bangiophyceae, *Porphyra pulchra* grouped within *Pyropia* clade (100% MLB) rather than *Porphyra* clade, suggesting a taxonomic revision of *Porphyra pulchra* as *Pyropia pulchra*. Relationships within the Florideophyceae were consistent with previous work^{64–66}. For example, two Corallinophycidae species, *Sporolithon durum* (Sporolithales) and *Calliarthron tuberculosum* (Corallinales) grouped together (100% MLB) and were sister to the rest of florideophycean clades. Within the subclass Rhodymeniophycidae, *Chondrus crispus* (Gigartinales) diverged first, followed by *Gelidium* (Gelidiales), *Grateloupia taiwanensis* (Halymeniales) and *Gracilaria* (Gracilariales). Although internal relationships within the Rhodymeniophycidae were not resolved with the concatenated plastid dataset, we used this ML tree (Fig. 1A) as a reference for inferring the evolution of red algal plasmid DNAs.

Distribution of plasmid-derived genes in red algal ptDNA. We identified 22 plasmid-derived (PD) sequences in nine red algal species when 56 red algal plasmid-encoded proteins were used to query the available 21 red algal plastid genomes (using BLASTx, *e*-value $\leq 1.0e^{-05}$) (GI numbers of the 56 proteins are listed in

Figure 1. Phylogeny of red algae showing the distribution of plasmid-derived DNA. (A) Simplified maximum likelihood (ML) tree topology for red algae based on concatenated 193 protein encoding genes of plastid genomes (see also Supplementary Fig. S6). (B) The plastid genome sizes (kilo base pair, Kbp) are shown beside the taxon names. Colored boxes indicate plastid genes (grey), plasmid-derived regions: *Porphyra pulchra* plasmid Pp6859 *orf4* and *orf5* homolog (green), *P. pulchra* plasmid Pp6427 *orf3* homolog (dark green), *P. pulchra* plasmid Pp6859 *orf6* and *orf7* (orange and bright orange), *P. pulchra* plasmid Pp6427 *orf1* homolog (bright green), *Gracilaria chilensis* plasmid Gch7220 *orf5* homolog (blue), *G. robusta* plasmid Gro4970 *orf5* homolog (cyan), *Gracilariopsis lemaneiformis* plasmid Gle4293 *orfs* homolog (red, pink and bright pink), *Gp. lemaneiformis* plasmid GL3.5 *orfs* homolog (violet and bright violet), and genes that encode the bacterial operon for *leuC* and *leuD* (black). Detailed information about plasmid-derived regions is given in Supplementary Fig. S7. The “p” in *syfB* gene of *Sporolithon durum* designates a pseudogenization of the gene. PCR-confirmed regions in different individuals are marked by asterisk. Syntenic diagrams for the *Cyanidioschyzon*, *Cyanidium*, *Galdieria* and *Porphyridium* are not shown because there was no plasmid-derived DNA.

Table 1). The putative origin, copy number, and distribution in the ptDNAs were different for each ORF (Fig. 1B; Supplementary Table S5). In addition to the previously reported bacterial operon *leuC* and *leuD* gene^{35,36} (two black blocks in Fig. 1B), out of the 22 PD *orfs* (including pseudogenized regions) identified here, six were homologous to *orf4* and *orf5* of the *Porphyra pulchra* plasmid Pp6859 (GI: 11466614; green region in Fig. 1B), five were homologous to the *P. pulchra* plasmid Pp6427 (GI: 11466608) *orf3* (dark green region in Fig. 1B), and two were homologous to the *P. pulchra* plasmid Pp6859 *orf6* (bright orange region in Fig. 1B). The rest of the PD *orfs* were unique to plasmids in their species of origin. Interestingly, six homologous PD sequences from Pp6859 *orf4* and *orf5* (green box in Fig. 1B) were found in four red algal plastid genomes but their copy number and position were not consistent with their phylogenetic relationships. For instance, two copies of the Pp6859 *orf4*-*orf5* homolog were found in *Pyropia haitanensis* among eight *Porphyra*/*Pyropia* species, whereas a single copy was found in each *Gelidium* species, but at different locations. *Sporolithon durum* contained two homologous copies but one was pseudogenized. The sequences homologous to plasmid Pp6427 *orf3* of *P. pulchra*³⁹ were found in the plastid genomes of three *Gracilaria* species and *Grateloupia taiwanensis* (dark-green in Fig. 1B) in addition to that of *P. pulchra*, and were located near ribosomal RNAs and *ycf27* genes. We note that half of the PD *orfs* were positioned near *rRNA* (*rps6*-*rRNA*-*ycf27*-*psbD*, see Fig. 1B), in particular in *Gelidium*, *Grateloupia*, and *Gracilaria*.

We tested whether these PD *orfs* were conserved in populations within a species and in different individuals within a population. To this end, PCR was used to test three populations of *G. elegans* (SKKU18, SKKU22, SKKU28), two individuals of *P. pulchra* selected from a single population (UC1879714 and UC1454976), and three individuals of *S. durum* from a single population (SKKU_SD01, SKKU_SD02, and SKKU_SD03; Supplementary Table S6). All tested PD *orfs* were found in the same position with the same flanking region sequences. Therefore, these PD *orfs* are conserved across different individuals within one species.

For *P. pulchra* and *G. chilensis*, the PD *orfs* found in their ptDNA or their homologs were not detected in the draft genome data but rather only in the plasmid sequence. From draft genome data (NGS), five complete plasmid sequences (103 ~360× average coverage) were recovered from *Porphyra pulchra* (263 Mbp of reads) and three plasmids (401 ~590× average coverage) from *Gracilaria chilensis* (506 Mbp of reads). Thus, NGS data were useful

Species	Plasmid (size)	Gene	GI	Rd	Cy	Bc	Gn	Vr	Rz	St	Remark	Relevant figure
<i>Gracilaria chilensis</i>	GC2 (3827 bp)	Unnamed protein	18477									Fig. 3
		ORF1	11465609									Fig. 3
		ORF2	11465610								-	-
	Gch3937 (3937 bp)	ORF3	11465611									-
		ORF4	11465612									-
		ORF5	11465607									-
		ORF1	11465592									Fig. 3
		ORF2	11465593									-
		ORF3	11465594									-
		ORF4	11465595									Fig. 3
		ORF5	11465596	p								-
		ORF6	11465597									Fig. 3
		ORF7	11465598									Fig. 3
	Gch7220 (7220 bp)	ORF8	11465599									-
		ORF9	11465600									-
		ORF10	11465601									-
		ORF12	11465602									-
		ORF13	11465603									-
		ORF14	11465604									-
		ORF15	11465605									-
		ORF16	11465606									-
		ORF17	11465607									-
<i>Gracilaria robusta</i>	Gro4059 (4059 bp)	ORF1478	11466335									-
		ORF2347	11466334	m,p								Fig. 3
		ORF1	11466337	p								Fig. 3
	Gro4970 (4970 bp)	ORF2	11466338									-
		ORF3	11466339									-
		ORF4	11466340									-
		ORF5	11466341	m,p								-
<i>Gracilariopsis lemaneiformis</i>	GL3.5 (3491 bp)	ORF1	18514	p								-
		ORF2	18515	p							e-value: e^{-03}	Fig. S8
	GLE4293 (4293 bp)	ORF1	11465615	p								Fig. 3
		ORF2	11465616	p								-
		ORF3	11465617	p								-
		ORF4	11465618									-
<i>Porphyra pulchra</i>	Pp6427 (6427 bp)	ORF5	11465619									-
		ORF1	11466609	P								-
		ORF2	11466610								e-value: e^{-03}	Fig. 4
		ORF3	11466611	P								Fig. 3
		ORF4	11466612				p					-
	Pp6859 (6859 bp)	ORF5	11466613									-
		ORF1	11466615									-
		ORF2	11466616									-
		ORF3	11466617									-
		ORF4 & ORF5	11466618-9	p			m				orf4 & orf5	Fig. 2
	Plasmid-1 (2102 bp)	ORF6	11466620	p							e-value: e^{-03}	-
		ORF7	11466621	p								-
		replicase	7108457								viral origin ⁵⁸	Fig. S9
	Plasmid-2 (1899 bp)	replicase	7108459								viral origin ⁵⁸	Fig. S9
	Plasmid-3 (2105 bp)	replicase	7108461								viral origin ⁵⁸	Fig. S9
<i>Pyropia tenera</i>	pPT4 RAP (1808 bp)	Replication associated protein	17980119								m	Fig. S9
	pPT4-NU (1820 bp)	Hypothetical protein	254029131									-
		Putative replicase	254029132								m	Fig. S9

Table 1. Distribution of red algal plasmids and their homologous sequences (BLASTp results, cut-off = e^{-05}). Rd = red algae, Cy = Cyanobacteria, Bc = Bacteria (excluding Cyanobacteria), Gn = green plant lineage (Viridiplantae), Vr = Virus, Rz = Rhizaria, St = Stramenopile. Letters beside the filled circles indicate origins: *m* for the mitochondrial homolog, *p* for the plastid homolog and *g* for the nuclear genome homolog. When the letters *m*, *p*, or *g* are absent, the origin of the gene is unknown.

for identifying plasmid sequences. However, we could not find red algal plasmid sequences in the published complete genome of *Cyanidioschyzon merolae*^{67,68}, *Galdieria sulphuraria*⁶⁹, *Porphyridium purpureum*⁷⁰, *Calliarthron tuberculosum*⁷¹, *Chondrus crispus*⁵⁶, and *Gelidium vagum* (Yoon *et al.* unpublished).

Origin of the plasmid-derived Pp6859 *orf4-orf5* homologs in ptDNA. The origin of plasmid-derived *orfs* was difficult to determine because most plastid-encoded PD *orfs* matched only plasmid *orf* data, except for the following five cases (see, Figs 2–4, S8, S9). A BLAST search against the NCBI database using six homologous plastid genes of the *P. pulchra* plasmid Pp6859 *orf4-orf5* resulted in 26 hypothetical proteins encoded in a bacterial genome, cyanobacterial genomes, cyanobacterial plasmids, and the mitochondrial genome of a liverwort. All homologous sequences of Pp6859 *orf4-orf5* were used to reconstruct the ML phylogeny using RAxML (Fig. 2). In the best tree, red algal plastid PD *orfs* grouped together, including plasmid Pp6859 (98% MLB). It is interesting that plasmid genes of Pp6859 (*P. pulchra*) grouped with pseudogenized plastid genes from *P. haitanensis* (100% MLB), suggesting a possible ORF gene transfer mediated by a plasmid to a plastid genome (see discussion in previous study⁴⁰).

The red algal clade was positioned within cyanobacterial clade Group I (92% MLB) that included hypothetical proteins encoded in the cyanobacterial genome as well as cyanobacterial plasmids (Fig. 2). Group II (72% MLB) contained cyanobacterial species and mitochondrial sequences from the liverwort *Marchantia polymorpha* (combined with two fragmented genes with flanking region data). Moon and Goff³⁹ reported the putative homologous relationship between Pp6859 and the liverwort mitochondrial region. Two cyanobacterial plasmid genes and a hypothetical gene from the Planctomycetes *Zavarzinella formosa* were grouped together (Group III, 100% MLB).

Because only 12 species (16 strains; Fig. 2) out of the 100 cyanobacterial genomes available in NCBI contain a homolog of Pp6859 *orf4-orf5*, it is unlikely to be a core cyanobacterial gene. If this *orf* was inherited from the primary endosymbiosis event, it should be retained in most red algal plastid genomes as well as those of other primary endosymbiotic lineages (i.e., green and glaucophyte algae). However, it is sporadically distributed in only a few species (e.g., *Pyropia*, *Gelidium* and *Sporolithon*) (Fig. 1). We postulate that this *orf* originated from an unknown cyanobacterial species, then spread independently to other cyanobacteria, to a bacterium (*Z. formosa*), to a liverwort (*M. polymorpha*), and to a few red algae.

The cyanobacterium *Crocospaera watsonii* WH8501 contains three copies of this *orf* as a result of gene duplications^{72,73}. However, it is likely that these red algal PD *orfs* originated independently, as a result of plasmid mobility. Alternatively, a red algal species inherited this *orf* from a cyanobacterial genome through the plasmid, after which it was transferred into the plastid genome in random genomic positions (e.g., see Fig. 1B), followed by pseudogenization or complete loss. This plasmid-mediated HGT may have occurred after speciation. For example, two *Gelidium* species both retain PD *orfs*, but they differ in size and genomic position. Similar cases were found in three *Gracilaria* species. If indeed the PD *orfs* were introduced during speciation, the presence and position of PD *orfs* could be used as species-specific markers.

Origin of the plasmid-derived Pp6427 *orfs* homologs in plastid genomes. Pp6427 *orf3* homologs were found in five plastid regions (Fig. 1), nine plasmid *orfs* and a mitochondrial *orf* from seven red algal species. Unlike Pp6859, homologous sequences were not found in any other taxa; therefore, Pp6427 *orf3* homologs are specific to red algae. In the ML tree using 16 homologs (Fig. 3), strong (>95% MLB) plasmid-plastid relationships were recovered, even though all plastid-encoded *orfs* were pseudogenized (see alignment in Fig. 3). For example, the plasmid Pp6427 *orf3* (485 aa) grouped with a short pseudogenized gene (191 aa) in the plastid genome of *P. pulchra* (95% MLB), whereas the plasmid Gro4059 (GI: 11466333) *orf2347* (190 aa) grouped with partial genes from *G. taiwanensis* (101 aa) within a clade of mitochondrial *orf44* from *Palmaria palmata* (Linnaeus) F. Weber & Mohr (88% MLB) (Fig. 3). *Gracilaria chilensis* contained six *orfs* in three plasmids (Gch7220 [GI: 11465591], Gch3937 [GI: 11465608], and GC2 [GI: 18476]); however, those *orfs* did not group with the plastid-encoded homologs that were clustered (99% MLB) with the pseudogenized plastid gene of *G. tenuistipitata* (144 aa) and plasmid Gle4293 (GI: 11465614) *orf1* of *G. lemaneiformis*.

Because the evolutionary trajectories of plasmid and plastid copies are very different (the former presumably functional and therefore subject to purifying selection, but the latter pseudogenized and under relaxed selective constraint), it is difficult to infer evolutionary relationships, since both rates and types of mutation (synonymous versus nonsynonymous) may be very different depending on the genetic background. We think it is likely that the plasmid *orfs* are ancestral because they contain complete *orfs* (405–485 aa), whereas plastids contain pseudogenized genes (up to 190 aa). On the other hand, plastid sequences occur in all the *Gracilaria* clades; the difference may be due to relaxed purifying selection on the shorter, non-functional (pseudogenized) plastid copies. Pp6427 *orf3* homologs were found in the closely related genera *Grateloupia*, *Gracilaria*, and *Gracilariopsis* (multigene phylogeny using mitochondrial genes⁴³), suggesting that an ancestral Pp6427 *orf3* of *P. pulchra* was transferred into the ancestral plastid of these genera and the mitochondrial genome of *Palmaria palmata*. Some plasmid *orfs* were duplicated (e.g., Gch7220 *orfs* and Gch3937 *orf1*) and fragmented (*orf1*, *orf6*, and *orf7*) within a plasmid (Gch7220). Although the origin of the plasmid-derived sequences is unknown, they may have spread into red algal organelle genomes and subsequently undergone relaxed selective constraint.

Two other plasmid *orfs* of Pp6427, *orf2* and *orf4* showed exclusive homology to cyanobacteria and green plants species, respectively. Pp6427 *orf2* was homologous to a putative transcriptional regulator protein (GI: 495464247) from the cyanobacterium *Moorea producens* (*e*-value: $4e^{-08}$) and to other cyanobacterial genes. The red algal plasmid *orf2* was likely transferred from cyanobacteria (Fig. 4; MLB 90% in basal clade). The combined region from Pp6427 *orf4* (Supplementary Table S7, ML tree is not shown) and their flanking regions were homologous to a hypothetical protein (*orf619*) from the plastid genome of *Ettlia pseudoalveolaris* (T.R. Deason & H.C. Bold) J. Komárek (green alga; GI: 725650857; BLASTx result *e*-value: $5e^{-12}$) as well as *orf436* of *Mankyua chejuensis*

Figure 2. Maximum likelihood (ML) tree based on aligned amino acid sequences of homologous regions of *Porphyra pulchra* plasmid Pp6859 *orf4* and *orf5* with 2,000 ML bootstrap replications. Species names are followed by GI, amino acid (aa) length, and location. Colored names indicate cyanobacteria (cyan), bacteria (black), liverwort (bright green) and red algae (red). Locations of the sequences are genome (black), plasmid (green), mitochondria (orange) and plasmid (yellowish brown). Some *orfs* and pseudogenized or non-coding regions were combined and aligned with sampled taxon sequences (Supplementary Table S3; Table S4; Table S7). The clades of the ML tree are divided into three groups based on species composition. Group I includes cyanobacterial plasmids and genomes with red algal plasmid and plastid regions. Group II includes cyanobacterial genomes and mitochondrial regions of liverwort, *Marchantia polymorpha*. Group III includes cyanobacterial plasmids and a bacterial (*Zavarzinella formosa*) genome.

B.Y. Sun, M.H. Kim & C.H. Kim (fern; GI: 727397314; BLASTx result e -value: $7e^{-05}$). Therefore, *orfs* encoded in the plasmid Pp6427 originated from various sources, and some *orfs* were subsequently transferred to the red algal plastid and mitochondrial genomes. Both plasmid Pp6859 *orf4-orf5* and Pp6427 *orf2* were homologous to cyanobacterial *orfs*, including those from several common species, *Calothrix* sp. 336/3, *Moorea producens* 3L, and *Rivularia* sp. PCC7116. Thus, these two plasmids may have served as reservoirs for *orfs* from different sources that eventually were delivered to organelles.

Bacterial and viral origins of red algal plasmid ORFs. Bacterial or viral sequences were detected by a BLASTp search of the NCBI (nr) database using 22 PD red algal plastid *orf* queries (Table S5). The homologous sequence of *Gracilariopsis lemaneiformis* plasmid GL3.5 *orf2* in the *Grateloupia taiwanensis* plastid genome showed a close phylogenetic relationship with bacterial and viral sequences (Supplementary Fig. S8). This red algal clade was positioned within the bacterial clades (100% MLB), suggesting the bacterial origin of the GL3.5 *orf2* homologs. It was, however, unclear whether this plasmid-related sequence was transferred from bacteria directly or by a virus-mediated process, because the clade showed a sister relationship to the viral clade but with weak statistical support (48% MLB).

The ML tree based on red algal plasmid-encoded replicase genes (i.e., *Pyropia tenera*, GI: 17980119, 254029132; *P. pulchra*, GI: 7108457, 7108459, 7108461) and homologous sequences from a BLASTp search suggest a viral origin of these plasmid *orfs* (see Supplementary Fig. S9). Five red algal plasmid *orfs*, mitochondrial *orf98* from *Phytophthora sojae* (stramenopile; GI: 145932354), and nuclear-encoded genes from *Reticulomyxa filosa* (rhizaria; GI: 569382219, 569382219) grouped together with diverse circular virus DNA (52% MLB), as well as with obligate parasitic bacteria (i.e., onion yellow phytoplasma) (78% MLB). A BLASTp search (cutoff e -value $\leq 1.0e^{-05}$, see Table 1) recognized eight sequences from the nuclear genome of *Nicotiana tomentosiformis* (green plant; GI: 697190580, 1587991, 697190578, 697159806, 697175541, 697190576, 697140845, 697149473) that were distantly related to the red algal plasmid clade. The replicase gene from the *P. pulchra* plasmid (GI: 7108457) was reported as a geminivirus-related sequence because it share five conserved motifs and phylogenetic affinities^{51,78}.

Virus-derived plasmid genes (i.e., GL3.5 *orf2*, three replicase genes in *P. pulchra* plasmids, and two replicase genes in *Py. tenera* plasmids) were detected in both eukaryotic nuclear and organellar genomes. These were different from non-viral-derived red algal plasmid homolog sequences that were found only in organelle genomes (Table 1). It is likely that virus-derived plasmid genes could be transferred to the eukaryotic nuclear genome more easily than could non-viral plasmid genes.

Figure 3. Maximum likelihood (ML) tree based on aligned amino acid sequences of homologous regions to *Porphyra pulchra* plasmid Pp6427 orf3 with 2,000 ML bootstrap replications. Species names are followed by source, GI, amino acid (aa) length, and location. Some orfs and pseudogenes or non-coding regions were combined and aligned with sampled taxon sequences (Supplementary Table S7). Location of sequences is indicated by color: plasmid (underlined black), plastid (green) and mitochondria (orange). Synteny is shown with the schematic alignment on the right of the tree based on major regions of homology.

Remnant DNA replication domain in plasmid-derived plastid genes. Plasmids are composed of three essential domains for replication, segregation and conjugation with additional accessory genes^{76,77}. From the alignment of the Pp6859 orf4-orf5 homologs with the size range of 104 ~1,242 amino acid sequences, the functional domain was detected by a conserved domain database search⁷⁹. One distinct domain is the DNA polymerase type-B family catalytic domain (POLBc) superfamily. Nine amino acid sequences were identical in this domain (aligned 142 aa), including highly conserved active sites (R-K-ND motif) and metal binding sites (DG motif) (see Fig. 5). The DNA polymerase type-B family consists of an editing active site and excision region for DNA replication (562 ~3,425 aa in size) that has been reported in a wide range of organisms, including Archaea, Bacteria, eukaryotes, bacteriophages and viruses⁸⁰⁻⁹¹. Although the POLBc motif was generally conserved in nine major subfamilies⁸⁵, we found differences in the catalytic domain of the Pp6859 orf4-orf5 homologs. These unique domains were represented in the ML tree that was reconstructed using homolog regions of the domain (aligned 222 aa) from the public POLBc superfamily database (Supplementary Table S8; Fig. 6). The ML tree showed that all POLBc domains in the Pp6859 orf4-orf5 homolog were grouped into a clade (100% MLB), but the clade did not belong to any other known POLBc subfamilies. This novel POLBc domain might contribute to the insertion of plasmid orfs into the red algal plastid genome.

Conclusions

Plasmids have long been recognized as mobile elements but their origins in red algae remained unclear. Using a comprehensive database of 21 plastid genomes that included five novel red algal ptDNAs, we found evidence for the spread of plasmid DNA into plastid and mitochondrial genomes. There is currently insufficient nuclear genome data from species that contain plasmid-derived DNA to determine whether this compartment is also a major target for integration (Fig. 7). The distribution of plasmid-derived orfs showed a species-specific pattern, consistent with the evolution of a mobile genetic element. Because organelles are inherited maternally, foreign genetic DNA can be rapidly fixed in a population. Consistent with this idea, individual members of three lineages (i.e., *Porphyra pulchra*, *Sporolithon durum*, and *Gelidium elegans*) all showed plasmid DNA retention, although these orfs were absent or located in different genomic positions in closely related sister species (e.g., eight *Porphyra/Pyropia* species, *Sporolithon-Calliarthron*, two *Gelidium* species, see Fig. 1). It is known that the distribution of transposable elements can show variation within a single cyanobacterial species^{72,73,92}. Therefore, plasmids may be regarded as analogous to transposable elements^{76,77,93-96}, with mobility and loss contributing to variation in gain/loss among closely related genomes. For instance, Halary *et al.*⁹⁷ demonstrated that plasmids are key vectors of genetic exchange between bacterial chromosomes on the basis of network analysis using sequences including phage, plasmid and environmental viral genomes.

It should be noted that we were originally interested in testing the idea whether plasmids may have facilitated EGT in algae and thereby played a key role in their genome evolution. Analysis of the available data, however, suggests that plasmids are better thought of as parasitic elements (e.g., group II introns in red algal ptDNA⁹⁸) that spread plasmid-derived DNA regions. As “mobile gene cassettes”^{97,98} it nonetheless remains possible that these selfish elements can mediate gene transfer between foreign DNA and organelles. As the databases of available

Figure 4. Maximum likelihood (ML) tree based on aligned amino acid sequences of homologous genes of *Porphyra pulchra* plasmid Pp6427 orf2 (red) with 2,000 ML bootstrap replications. Blue indicates that the sequences found in the plasmid genome.

organelle and nuclear genome data increase, plasmid involvement in recent instances of EGT may become apparent.

In summary, one of the major challenges in the field of microbial eukaryote genome evolution is to understand how genes move across the tree of life. Species such as *Galdieria sulphuraria* encode at least 5% foreign genes, many of which are clearly of adaptive value⁶⁹. The halotolerant green alga *Picochlorum* SE3 has acquired at least 24 genes of bacterial provenance, putatively to deal with abiotic stress⁹⁹. Plasmids, viruses, symbionts, and pathogens likely all play a role in the HGT process in protists. Therefore, the search for “smoking guns” of recent transfer will continue to fascinate biologists who seek to show that highways of gene sharing¹⁰⁰, common in prokaryotes, are drivers of evolution in eukaryotic microbes.

Methods

Sample preparation, genome sequencing, and assembly. Thalli of the red algal species *Gelidium elegans* Kützinger, *G. vagum* Okamura, *Gracilaria chilensis* C.J. Bird, J.L. McLachlan, & E.C. de Oliveira, and *Sporolithon durum* (Foslie) R. Townsend & W. Woelkerling were collected from nature and immediately dried in silica-gel. Tissue samples of *Porphyra pulchra* were taken from herbarium specimens collected in 1970 and housed at the University Herbarium, University of California at Berkeley (UC). Detailed information about the samples is shown in Table S6 in the Supplementary Information. Genomic DNA was extracted using the DNeasy Plant Mini Kit (Qiagen, Hilden, Germany). Next-Generation Sequencing (NGS) was carried out using Ion Torrent PGM platform (Life Technologies, San Francisco, California, USA). The Ion Xpress Plus gDNA Fragment Library Kit (Life Technologies) was used for 200 bp-sized or 400 bp-sized sequencing library preparation. Genome sequencing was done with the Ion PGM Template OT2 200 or 400 Kit and Ion PGM Sequencing 200 or 400 Kit (Life Technologies, San Francisco, California, USA).

The raw NGS reads were assembled using the CLC Genomics Workbench 5.5.1 (CLC bio, Aarhus, Denmark) and the MIRA assembler that was incorporated in the Ion Server. Contigs of plastid genes were sorted by customized Python scripts with local BLAST searches. Sorted contigs were re-assembled to construct consensus plastid genomes. A draft plastid genome was confirmed by the read-mapping method using CLC Genomics Workbench 5.5.1. Gaps were filled by PCR to generate intact genomes.

Gene annotation and plasmid-derived ORFs search. Putative ORFs in the five novel genomes were predicted using ORF Finder in Geneious 6.1.6¹⁰¹ and annotated based on BLASTx searches (e -value $\leq 1.0e^{-05}$) with codon table 11 (Bacterial, Archaeal and Plant Plastid Code). Ribosomal RNAs and transfer RNAs were predicted using the RNAmmer 1.2 Server¹⁰² and ARAGORN programs¹⁰³. Group II intron and RNase P were searched using the program RNAweasel (<http://megasun.bch.umontreal.ca/cgi-bin/RNAweasel/RNAweaselInterface.pl>). Plasmid-derived sequences were searched by BLASTx (e -value $\leq 1.0e^{-05}$) using 56 proteins encoded in 14 red algal plasmids (Supplementary Table S9) derived from all available red algal ptDNAs. We also searched for plasmid-derived sequences in nuclear genome data. Here 56 plasmid-encoded genes were searched in the complete nuclear genomes of *Cyanidioschyzon merolae*^{67,68}, *Galdieria sulphuraria* (Galdieri) Merola⁶⁹, *Porphyridium purpureum* (Bory) K.M. Drew & R. Ross⁷⁰, *Calliarthron tuberculatum*

Figure 5. Domain search of *Porphyra pulchra* plasmid Pp6859 orf4-orf5 homolog. (A) Cladogram of the best maximum likelihood tree, Fig. 2. (B) Schematic of the DNA polymerase type-B family catalytic domain (POLBc) superfamily related regions (rectangular boundary) in the hypothetical homologous gene (green bars). Active and metal binding sites on the POLBc superfamily are shown below the alignment. In the domain of *Porphyra pulchra* (*), serine (S*) substitutes for glycine (G) in metal binding sites. (C) The composition of amino acids in alignment. Representative color of amino acid follows the Chemistry color scheme in WebLogo with probability-based size differences. The nine red arrowheads indicate conserved amino acids (100%) in the alignment. Blanks in the alignment indicate extremely low contribution by the hypothetical protein in *Fischerella* sp. (GI: 652333536) and in *Scytonema tolypothrichoides* (GI: 750920257).

Figure 6. Maximum likelihood (ML) tree based on aligned amino acid sequences of DNA polymerase type-B family catalytic domain (POLBc) superfamily with 2,000 ML bootstrap replications. A public POLBc database was used from the conserved domain database (CDD) (Supplementary Table S8). Each cluster indicates a subfamily of POLBc superfamily. The novel POLBc subfamily clade comprises the POLBc-related partial domains of the hypothetical gene in this study. Eight public domain data are not identified to a specific subfamily in the POLBc superfamily (bright orange; GI: 16081956, 48477200, 15789745, 124362982, 48425269, 48425257, 6435552, 147921691). The DNA polymerase type-II B (Pol II B; bright green) subfamily is not monophyletic. Most subfamilies of POLBc are monophyletic; however, some inter-clade relationships differ slightly from the public POLBc superfamily database (CDD cl00145).

(Postels & Ruprecht) E.Y. Dawson⁷¹, *Chondrus crispus* Stackhouse⁵⁶ and the 5 novel red algal draft genomes. Reported plasmid sequences from *Gracilaria chilensis* (GI: 11465591, 11465608 and 18476) and *Porphyra pulchra* (GI: 11466614, 11466608, 7108456, 7108458 and 7108460) were used as reference sequences for the read-mapping method for NGS data. To check consistency within individuals and populations, plasmid-derived sequences were determined from three individuals of *Gelidium elegans* from three different sites in Korea (SKKU18, SKKU22 and SKKU28), two individuals of *Porphyra pulchra* from Moss Beach, CA, USA (UC1454976 and UC1879714), and three individuals of *Sporolithon durum* from Army Bay, Whangaparaoa, NZ (SKKU_SD01, SKKU_SD02 and SKKU_SD03) using PCR with custom primer pairs (Supplementary Table S10).

Figure 7. The spread of plasmid DNA in eukaryote genomes. The schematic cell includes the nucleus (dotted line circle), plastid (green circle), mitochondrion (orange circle) and plasmid (black circle) DNA. The plasmid-derived regions are indicated as red boxes in the genomes. The flow of plasmid DNA is indicated by the arrows. Plasmid-mediated HGT in plastid genomes are divided into two types: with plasmid and without plasmid (Figs 2, 3 and S8). Organisms with and without plasmid DNA are listed below as red algae (red taxa), green lineage (green taxa), stramenopiles (brown taxa), and rhizarians (violet taxa). Plastid genomes of *Porphyrta pulchra* and *Gracilaria chilensis* include plasmid-derived homologs in both their plastid and plasmid genomes. The other red algae include plasmid-derived homologs only in the plastid genome. Mitochondrial HGT is found in red algae, the green lineage and stramenopiles (Figs 2, 3 and S9). Plasmid-mediated transfer to the nuclear genome is found only in *Nicotiana tomentosiformis* (plants) and *Reticulomyxa filosa* (rhizarian), with both regions related to viruses (Supplementary Fig. S9).

Phylogenetic analysis of red algal plasmid-derived genes in plastid genome. Plastid-coding genes from 21 taxa (16 reference genomes and our five new genomes) were extracted and sorted by customized Python scripts with local BLAST searches. To identify the independent loss of plastid genes, each gene set was manually analyzed. A selection of 193 plastid-coding genes (e.g., homologous genes present in at least 16 different taxa) and plasmid-derived sequences were aligned using MAFFT 7.110¹⁰⁴. All aligned plastid genes were concatenated for multigene phylogenetic analysis. Based on the alignment, fragmented plasmid-derived *orfs* were combined (Supplementary Table S7). To reconstruct the phylogenetic tree, an evolutionary model was selected using Modeltest implemented in MEGA 6.0¹⁰⁵. Maximum likelihood (ML) tree search and ML bootstrap analysis were done using RaxML 8.0.0 with 2000 replications¹⁰⁶ with the PROT + GAMMA + LG4MF model of evolution.

Domain prediction and phylogenetic analysis. Protein domain prediction was done using the conserved domain database CDD⁷⁹. Predicted domain sequences were aligned and represented using Weblogo¹⁰⁷. Aligned sequences of DNA polymerase type-B family catalytic domain (POLBc) subfamilies (POLBc_B1, POLBc_B2, POLBc_alpha, POLBc_delta, POLBc_zeta, POLBc_epsilon, POLBc_B3, POLBc_Pol_II, POLBc_Pol_II_B and unspecified POLBc domain (Supplementary Table S8) were used to find the inter-subfamily relationship based on the RAXML phylogeny.

References

- Ochman, H., Lawrence, J. G. & Groisman, E. A. Lateral gene transfer and the nature of bacterial innovation. *Nature* **405**, 299–304 (2000).
- Lawrence, J. G. Gene transfer in bacteria: speciation without species? *Theor. Popul. Biol.* **61**, 449–460 (2002).
- Wiedenbeck, J. & Cohan, F. M. Origins of bacterial diversity through horizontal genetic transfer and adaptation to new ecological niches. *FEMS Microbiol. Rev.* **35**, 957–976 (2011).
- Salzberg, S. L., White, O., Peterson, J. & Eisen, J. A. Microbial genes in the human genome: lateral transfer or gene loss? *Science* **292**, 1903–1906 (2001).
- Archibald, J. M., Rogers, M. B., Toop, M., Ishida, K. & Keeling, P. J. Lateral gene transfer and the evolution of plastid-targeted proteins in the secondary plastid-containing alga *Bigeloviella natans*. *Proc. Natl. Acad. Sci. USA* **100**, 7678–7683 (2003).
- Hall, C., Brachat, S. & Dietrich, F. S. Contribution of horizontal gene transfer to the evolution of *Saccharomyces cerevisiae*. *Eukaryot. Cell* **4**, 1102–1115 (2005).
- Rice, D. W. & Palmer, J. D. An exceptional horizontal gene transfer in plastids: gene replacement by a distant bacterial paralog and evidence that haptophyte and cryptophyte plastids are sisters. *BMC Biol.* **4**, 31 (2006).
- Richards, T. A., Dacks, J. B., Jenkinson, J. M., Thornton, C. R. & Talbot, N. J. Evolution of filamentous plant pathogens: gene exchange across eukaryotic kingdoms. *Curr. Biol.* **16**, 1857–1864 (2006).
- Novo, M. *et al.* Eukaryote-to-eukaryote gene transfer events revealed by the genome sequence of the wine yeast *Saccharomyces cerevisiae* EC1118. *Proc. Natl. Acad. Sci. USA* **106**, 16333–16338 (2009).
- Richards, T. A. *et al.* Phylogenomic analysis demonstrates a pattern of rare and ancient horizontal gene transfer between plants and fungi. *Plant Cell* **21**, 4897–1911 (2009).
- Guljamow, A. *et al.* Horizontal gene transfer of two cytoskeletal elements from a eukaryote to a cyanobacterium. *Curr. Biol.* **17**, 757–759 (2007).
- Rogers, M. B., Patron, N. J. & Keeling, P. J. Horizontal transfer of a eukaryotic plastid-targeted protein gene to cyanobacteria. *BMC Biol.* **5**, 26 (2007).
- Bhattacharya, D., Yoon, H. S. & Hackett, J. D. Photosynthetic eukaryotes unite: endosymbiosis connects the dots. *BioEssays* **26**, 50–60 (2004).
- Timmis, J. N., Ayliffe, M. A., Huang, C. Y. & Martin, W. Endosymbiotic gene transfer: organelle genomes forge eukaryotic chromosomes. *Genetics* **5**, 123–135 (2004).
- Richardson, A. O. & Palmer, J. D. Horizontal gene transfer in plants. *J. Exp. Bot.* **58**, 1–9 (2007).
- Keeling, P. J. & Palmer, J. D. Horizontal gene transfer in eukaryotic evolution. *Nat. Rev. Genet.* **9**, 605–618 (2008).
- Reyes-Prieto, A., Hackett, J. D., Soares, M. B., Bonaldo, M. F. & Bhattacharya, D. Cyanobacterial contribution to algal nuclear genomes is primarily limited to plastid functions. *Curr. Biol.* **16**, 2320–2325 (2006).
- Price, D. C. *et al.* *Cyanophora paradoxa* genome elucidates origin of photosynthesis in algae and plants. *Science* **335**, 843–847 (2012).
- Sato, N., Ishikawa, M., Fujiwara, M. & Sonoike, K. Mass identification of chloroplast proteins of endosymbiont origin by phylogenetic profiling based on organism-optimized homologous protein groups. *Genome Inform.* **16**, 56–68 (2005).
- Deusch, O. *et al.* Genes of cyanobacterial origin in plant nuclear genomes point to a heterocyst-forming plastid ancestor. *Mol. Biol. Evol.* **25**, 748–461 (2008).
- Dagan, T. *et al.* Genomes of Stigonematalean cyanobacteria (subsection V) and the evolution of oxygenic photosynthesis from prokaryotes to plastids. *Genome Biol. Evol.* **5**, 31–44 (2013).
- Moustafa, A. & Bhattacharya, D. PhyloSort: a user-friendly phylogenetic sorting tool and its application to estimating the cyanobacterial contribution to the nuclear genome of *Chlamydomonas*. *BMC Evol. Biol.* **8**, 6 (2008).
- Martin, W. *et al.* Evolutionary analysis of *Arabidopsis*, cyanobacterial, and chloroplast genomes reveals plastid phylogeny and thousands of cyanobacterial genes in the nucleus. *Proc. Natl. Acad. Sci. USA* **99**, 12246–12251 (2002).
- Straub, S. C. K., Cronn, R. C., Edwards, C., Fishbein, M. & Liston, A. Horizontal transfer of DNA from the mitochondrial to the plastid genome and its subsequent evolution in milkweeds (Apocynaceae). *Genome Biol. Evol.* **5**, 1872–1885 (2013).
- Koulintchenko, M., Konstantinov, Y. & Dietrich, A. Plant mitochondria actively import DNA via the permeability transition pore complex. *EMBO J.* **22**, 1245–1254 (2003).
- Sheahan, M. B., McCurdy, D. W. & Rose, R. J. Mitochondria as a connected population: ensuring continuity of the mitochondrial genome during plant cell dedifferentiation through massive mitochondrial fusion. *Plant J.* **44**, 744–755 (2005).
- Rice, D. W. *et al.* Horizontal transfer of entire genomes via mitochondrial fusion in the angiosperm *Amborella*. *Science* **342**, 1468–1473 (2013).
- Paillard, M., Sederoff, R. R. & Levings, III C. S. Nucleotide sequence of the S-1 mitochondrial DNA from the S cytoplasm of maize. *EMBO J.* **4**, 1125–1128 (1985).
- Weber, B., Börner, T. & Weihe, A. Remnants of a DNA polymerase gene in the mitochondrial DNA of *Marchantia polymorpha*. *Curr. Genet.* **27**, 488–490 (1995).
- Burger, G., Saint-Louis, D., Gray, M. W. & Lang, B. F. Complete sequence of the mitochondrial DNA of the red alga *Porphyra purpurea*: cyanobacterial introns and shared ancestry of red and green algae. *Plant Cell* **11**, 1675–1694 (1999).
- Rousvoal, S., Oudot, M. P., Fontaine, J. M., Kloareg, B. & Loiseaux-de Goër, S. Witnessing the evolution of transcription in mitochondria: the mitochondrial genome of the primitive brown alga *Pylaiella littoralis* (L.) Kjellm. Encodes a T7-like RNA polymerase. *J. Mol. Biol.* **277**, 1047–1057 (1998).
- Férandon, C., Xu, J. & Barroso, G. The 135kbp mitochondrial genome of *Agaricus bisporus* is the largest known eukaryotic reservoir of group I introns and plasmid-derived sequences. *Fungal Genet. Biol.* **55**, 58–91 (2013).
- Nadimi, M., Beaudet, D., Forget, L., Hijri, M. & Lang, B. F. Group I intron-mediated trans-splicing in mitochondria of *Gigaspora rosea* and a robust phylogenetic affiliation of arbuscular mycorrhizal fungi with Mortierellales. *Mol. Biol. Evol.* **29**, 2199–2210 (2012).
- Khan, H. *et al.* Plastid genome sequence of the cryptophyte alga *Rhodomonas salina* CCMP1319: lateral transfer of putative DNA replication machinery and a test of chromist plastid phylogeny. *Mol. Biol. Evol.* **24**, 1832–1842 (2007).
- Hagopian, J. C., Reis, M., Kitajima, J. P., Bhattacharya, D. & de Oliveira, M. C. Comparative analysis of the complete plastid genome sequence of the red alga *Gracilaria tenuistipitata* var. *liui* provides insights into the evolution of rhodoplasts and their relationship to other plastid. *J. Mol. Evol.* **59**, 464–447 (2004).
- Janoušková, J. *et al.* Evolution of red algal plastid genome: ancient architectures, introns, horizontal gene transfer, and taxonomic utility of plastid markers. *PLoS ONE* **8**, e59001 (2013).
- Baumann, L., Baumann, P., Moran, N. A., Sandwtröm, J. & Thao, M. L. Genetic characterization of plasmids containing genes encoding enzymes of leucine biosynthesis in endosymbionts (*Buchnera*) of aphids. *J. Mol. Evol.* **48**, 77–85 (1999).
- Sabater-Muñoz, B., van Ham, R. C. H. J., Moya, A., Silva, F. J. & Latorre, A. Evolution of the leucine gene cluster in *Buchnera aphidicola*: insights from chromosomal versions of the cluster. *J. Bacteriol.* **186**, 2646–2654 (2004).
- Moon, D. A. & Goff, L. J. Molecular characterization of two large DNA plasmids in the red alga *Porphyra pulchra*. *Curr. Genet.* **32**, 132–138 (1997).
- Wang, L. *et al.* Complete sequence and analysis of plastid genomes of two economically important red algae: *Pyropia haitanensis* and *Pyropia yezoensis*. *PLoS ONE* **8**, e65902 (2013).

41. Zhang, L. *et al.* Complete sequences of the mitochondrial DNA of the wild *Gracilariopsis lemaneiformis* and two mutagenic cultivated breeds (Gracilariaceae, Rhodophyta). *PLoS ONE* **7**, e40241 (2012).
42. Yang, E. C., Kim, K. M., Kim, S. Y. & Yoon, H. S. Complete mitochondrial genome of agar-producing red alga *Gracilariopsis chorda* (Gracilariaceae). *Mitochondrial DNA* **25**, 339–341 (2014).
43. Lee, J.-M., Boo, S. M., Mansilla, A. & Yoon, H. S. Unique repeat and plasmid sequences in the mitochondrial genome of *Gracilaria chilensis* (Gracilariaceae, Rhodophyta). *Phycologia* **54**, 20–23 (2015).
44. Brown, G. G. & Finnegan, P. M. RNA plasmids. *Int. Rev. Cytol.* **117**, 1–56 (1989).
45. Goff, L. J. & Coleman, A. W. Red algal plasmids. *Curr. Genet.* **18**, 557–565 (1990).
46. Esser, K. *et al.* in *Heidelberg Science Library, Plasmids of eukaryotes: Fundamentals and applications* (eds Esser, K *et al.*) 1–124 (Springer, 1986).
47. Meinhardt, F., Kempken, F., Kämper, J. & Esser, K. Linear plasmids among eukaryotes: fundamentals and application. *Curr. Genet.* **17**, 89–95 (1990).
48. Klassen, R. & Meinhardt, F. in *Microbial Linear Plasmids*, Vol. 7 (eds Meinhardt, F. & Klassen, R.) *Linear protein-primed replicating plasmids in eukaryotic microbes*, 188–226 (Springer, 2007).
49. Villemur, R. Circular plasmid DNAs from the red alga *Gracilaria chilensis*. *Curr. Genet.* **18**, 251–257 (1990).
50. Villemur, R. The DNA sequence and structural organization of the GC2 plasmid from the red alga *Gracilaria chilensis*. *Plant Mol. Biol.* **15**, 237–243 (1990).
51. Moon, D. A. & Goff, L. J. Small plasmids in *Porphyra pulchra* are related to flowering plant geminiviruses. *J. Phycol.* **32**, 33–34 (1996).
52. Choi, H. S., Choi, K. H., Kim, T. H., Lee, C. H. & Rhew, T. H. Characterization of natural plasmid and construction of putative transformation vector using the plasmid in the Korean red alga, *Porphyra tenera*. *Algae* **16**, 287–294 (2001).
53. Reith, M. & Munholland, J. Complete nucleotide sequence of the *Porphyra purpurea* chloroplast genome. *Plant Mol. Biol. Rep.* **13**, 333–335 (1995).
54. Glöckner, G., Rosenthal, A. & Valentin, K. The structure and gene repertoire of an ancient red algal plastid genome. *J. Mol. Evol.* **51**, 382–390 (2000).
55. Ohta, N. *et al.* Complete sequence and analysis of the plastid genome of the unicellular red alga *Cyanidioschyzon merolae*. *DNA Res.* **10**, 67–77 (2003).
56. Collén, J. *et al.* Genome structure and metabolic features in the red seaweed *Chondrus crispus* shed light on evolution of the Archaeplastida. *Proc. Natl. Acad. Sci. USA* **110**, 5247–5252 (2013).
57. DePriest, M. S., Bhattacharya, D. & López-Bautista, J. M. The plastid genome of the red macroalga *Grateloupia taiwanensis* (Halymeniaceae). *PLoS ONE* **8**, e68246 (2013).
58. Campbell, M. A., Presting, G., Bennett, M. S. & Sherwood, A. R. Highly conserved organellar genomes in the Gracilariaceae as inferred using new data from the Hawaiian invasive alga *Gracilaria salicornia* (Rhodophyta). *Phycologia* **53**, 109–116 (2014).
59. Hughey, J. R. *et al.* Minimally destructive sampling of type specimens of *Pyropia* (Bangiales, Rhodophyta) recovers complete plastid and mitochondrial genomes. *Sci. Rep.* **4**, 5113 (2014).
60. Jain, K. *et al.* Extreme features of the *Galdieria sulphuraria* organellar genomes: a consequence of polyextremophily? *Genome Biol. Evol.* **7**, 367–380 (2015).
61. Tajima, N. *et al.* Analysis of the complete plastid genome of the unicellular red alga *Porphyridium purpureum*. *J. Plant Res.* **127**, 389–397 (2014).
62. Yoon, H. S., Hackett, J. D. & Bhattacharya, D. A single origin of the peridinin- and fucoxanthin-containing plastids in dinoflagellates through tertiary endosymbiosis. *Proc. Natl. Acad. Sci. USA* **99**, 11724–11729 (2002).
63. Yoon, H. S., Hackett, J. D., Ciniglia, C., Pinto, G. & Bhattacharya, D. A molecular timeline for the origin of photosynthetic eukaryotes. *Mol. Biol. Evol.* **21**, 809–818 (2004).
64. Le Gall, L. & Saunders, G. W. A nuclear phylogeny of the Florideophyceae (Rhodophyta) inferred from combined EF2, small subunit and large subunit ribosomal DNA: establishing the new red algal subclass Corallinophycidae. *Mol. Phylogenet. Evol.* **43**, 1118–1130 (2007).
65. Verbruggen, H. *et al.* Data mining approach identifies research priorities and data requirements for resolving the red algal tree of life. *BMC Evol. Biol.* **10**, 16 (2010).
66. Yang, E. C. *et al.* Highly conserved mitochondrial genomes among multicellular red algae of the Florideophyceae. *Genome Biol. Evol.* **7**, 2394–2406 (2015).
67. Matsuzaki, M. *et al.* Genome sequence of the ultrasmall unicellular red alga *Cyanidioschyzon merolae* 10D. *Nature* **428**, 653–657 (2004).
68. Nozaki, H. *et al.* 100%-complete sequence reveals unusually simple genomic features in the hot-spring red alga *Cyanidioschyzon merolae*. *BMC Biol.* **5**, 28 (2007).
69. Schönknecht, G. *et al.* Gene transfer from bacteria and archaea facilitated evolution of an extremophilic eukaryote. *Science* **339**, 1207–1210 (2013).
70. Bhattacharya, D. *et al.* Genome of the red alga *Porphyridium purpureum*. *Nat. Commun.* **4**, 1941 (2013).
71. Chan, C. X. *et al.* Red and green algal monophyly and extensive gene sharing found in a rich repertoire of red algal genes. *Curr. Biol.* **21**, 328–333 (2011).
72. Bench, S. R., Ilikchyan, I. N., Tripp, H. J. & Zehr, J. P. Two strains of *Crocospaera watsonii* with highly conserved genomes are distinguished by strain-specific features. *Front. Microbiol.* **2**, 261 (2011).
73. Bench, S. R. *et al.* Whole genome comparison of six *Crocospaera watsonii* strains with differing phenotypes. *J. Phycol.* **49**, 786–801 (2013).
74. Palmer, K. M., Turner, S. L. & Young, J. P. W. Sequence diversity of the plasmid replication gene *repC* in the Rhizobiaceae. *Plasmid* **44**, 209–219 (2000).
75. Sørensen, S. J., Bailey, M., Hansen, L. H., Kroer, N. & Wuertz, S. Studying plasmid horizontal transfer *in situ*: a critical review. *Nat. Rev. Microbiol.* **3**, 700–710 (2005).
76. Norman, A., Hansen, L. H. & Sørensen, S. J. Conjugative plasmids: vessels of the communal gene pool. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **364**, 2275–2289 (2009).
77. Harrison, E. & Brockhurst, M. A. Plasmid-mediated horizontal gene transfer is a coevolutionary process. *Trends Microbiol.* **20**, 262–267 (2012).
78. Vadivukarasi, T., Girish, K. R. & Usha, R. Sequence and recombination analyses of the geminivirus replication initiator protein. *J. Biosci.* **32**, 17–29 (2007).
79. Aron, M. B. *et al.* CDD: a conserved domain database for the functional annotation of proteins. *Nucleic Acids Res.* **39**, D225–229 (2011).
80. Braithwaite, D. K. & Ito, J. Compilation, alignment, and phylogenetic relationships of DNA polymerases. *Nucleic Acids Res.* **21**, 787–802 (1993).
81. Hopfner, K. *et al.* Crystal structure of a thermostable type B DNA polymerase from *Thermococcus gorgonarius*. *Proc. Natl. Acad. Sci. USA* **96**, 3600–3605 (1999).
82. Shamoo, Y. & Steitz, T. A. Building a replisome from interacting pieces: sliding clamp complexed to a peptide from DNA polymerase and a polymerase editing complex. *Cell* **99**, 155–166 (1999).

83. Iwai, T., Kurosawa, N., Itoh, Y. H., Kimura, N. & Horiuchi, T. Sequence analysis of three family B DNA polymerases from the thermoacidophilic crenarchaeon *Sulfurisphaera ohwakuensis*. *DNA Res.* **7**, 243–251 (2000).
84. Rodriguez, A. C., Park, H., Mao, C. & Beese, L. S. Crystal structure of a pol α family DNA polymerase from the hyperthermophilic archaeon *Thermococcus* sp. 9° N-7. *J. Mol. Biol.* **299**, 447–462 (2000).
85. Albà, M. M. 2001. Replicative DNA polymerases. *Genome Biol.* **2**, 3002.1–3002.4 (2001).
86. Burgers, P. M. *et al.* Eukaryotic DNA polymerases: proposal for a revised nomenclature. *J. Biol. Chem.* **276**, 43487–43490 (2001).
87. Hashimoto, H. *et al.* Crystal structure of DNA polymerase from hyperthermophilic archaeon *Pyrococcus kodakaraensis* KOD1. *J. Mol. Biol.* **306**, 469–477 (2001).
88. Filée, J., Forterre, P., Sen-Lin, T. & Laurent, J. Evolution of DNA polymerase families: evidences for multiple gene exchange between cellular and viral proteins. *J. Mol. Evol.* **54**, 763–773 (2002).
89. Freisinger, E., Grollman, A. P., Miller, H. & Kisker, C. Lesion (in) tolerance reveals insights into DNA replication fidelity. *EMBO J.* **23**, 1494–1505 (2004).
90. Koonin, E. V. Temporal order of evolution of DNA replication systems inferred by comparison of cellular and viral DNA polymerases. *Biol. Direct* **1**, 39 (2006).
91. Liu, S. *et al.* Crystal structure of the herpes simplex virus 1 DNA polymerase. *J. Biol. Chem.* **281**, 18193–18200 (2006).
92. Barrick, J. E. *et al.* Genome evolution and adaptation in a long-term experiment with *Escherichia coli*. *Nature* **461**, 1243–1247 (2009).
93. Thorsted, P. B. *et al.* Complete sequence of the IncPb plasmid R751: implications for evolution and organisation of the IncP backbone. *J. Mol. Biol.* **282**, 969–990 (1998).
94. Rawlings, D. E. & Tietze, E. Comparative biology of IncQ and IncQ-like plasmids. *Microbiol. Mol. Biol. Rev.* **65**, 481–496 (2001).
95. Haines, A. S., Jones, K., Cheung, M. & Thomas, C. M. The IncP-6 plasmid Rms149 consists of a small mobilizable backbone with multiple large insertions. *J. Bacteriol.* **187**, 4728–4738 (2005).
96. Thomas, C. M. & Nielsen, K. M. Mechanisms of, and barriers to, horizontal gene transfer between bacteria. *Nature Rev. Microbiol.* **3**, 711–721 (2005).
97. Halary, S., Leigh, J. W., Cheaib, B., Lopez, P. & Baptiste, E. Network analyses structure genetic diversity in independent genetic worlds. *Proc. Natl. Acad. Sci. USA* **107**, 127–132 (2010).
98. Perrineau, M. M., Price, D. C., Mohr, G. & Bhattacharya, D. Recent mobility of plastid encoded group II introns and twintrons in five strains of the unicellular red alga *Porphyridium*. *PeerJ* **3**, e1017 (2015).
99. Foflonker, F. *et al.* Genome of the halotolerant green alga *Picochlorum* sp. reveals strategies for thriving under fluctuating environmental conditions. *Environ. Microbiol.* **17**, 412–426 (2015).
100. Beiko, R. G., Harlow, T. J. & Ragan, M. A. Highways of gene sharing in prokaryotes. *Proc. Natl. Acad. Sci. USA* **102**, 14332–14337 (2005).
101. Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–1649 (2012).
102. Lagesen, K. *et al.* RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res.* **35**, 3100–3108 (2007).
103. Laslett, D. & Canback, B. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res.* **32**, 11–16 (2004).
104. Katoh, K. & Toh, H. Recent developments in the MAFFT multiple sequence alignment program. *Brief. Bioinform.* **9**, 286–298 (2008).
105. Tamura, K., Stecher, G., Peterson, D., Filipski, A. & Kumar, S. MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Mol. Biol. Evol.* **30**, 2725–2729 (2013).
106. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).
107. Crooks, G. E., Hon, G., Chandonia, J. & Brenner, S. E. WebLogo: a sequence logo generator. *Genome Res.* **14**, 1188–1190 (2004).

Acknowledgements

We thank the two anonymous reviewers for their constructive comments, which helped improve the manuscript. This study was supported by the Global Ph.D. Fellowship (GPF) Program from the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2013–034853) of Korea to J.L., NRF (2013–0699 and MEST: 2014R1A2A2A01003588) and Marine Biotechnology Program (PJT200620) funded by Ministry of Oceans and Fisheries, Korea to ECY, SMB and HSY, Korean RDA Next-generation BioGreen21 (PJ011121) to HSY.

Author Contributions

J.L., K.M.K., E.C.Y., K.A.M., S.M.B. and H.S.Y. collected samples from field and K.A.M. contributed tissue samples of *Porphyra pulchra* housed in the University Herbarium, University of California at Berkeley (UC). J.L., K.M.K. and E.C.Y. produced the genome sequencing. J.L. lead the plastid genome assembly, genome prediction, and wrote the manuscript. H.S.Y., S.M.B. and D.B. contributed to designing the work, interpreting the results, and editing the manuscript. All authors discussed the results and commented on the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Lee, J.M. *et al.* Reconstructing the complex evolutionary history of mobile plasmids in red algal genomes. *Sci. Rep.* **6**, 23744; doi: 10.1038/srep23744 (2016).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>