SCIENTIFIC REPORTS

OPEN

Goos-Hänchen shift of partially coherent light fields in epsilon-near-zero metamaterials

Ziauddin^{1,2}, You-Lin Chuang¹, Sajid Qamar² & Ray-Kuang Lee¹

Received: 25 January 2016

Accepted: 03 May 2016

Published: 23 May 2016

The Goos-Hänchen (GH) shifts in the reflected light are investigated both for *p* and *s* polarized partial coherent light beams incident on epsilon-near-zero (ENZ) metamaterials. In contrary to the coherent counterparts, the magnitude of GH shift becomes non-zero for *p* polarized partial coherent light beam; while GH shift can be relatively large with a small degree of spatial coherence for *s* polarized partial coherent beam. Dependence on the beam width and the permittivity of ENZ metamaterials is also revealed for partial coherent light fields. Our results on the GH shifts provide a direction on the applications for partial coherent light sources in ENZ metamaterials.

Goos-Hänchen (GH) shift reveals the discrepancy between geometric optics and wave optics, when a light beam undergoes a small lateral displacement from the interface of two media¹. There are a variety of interesting applications on the GH shift. For example, in optical heterodyne sensors, GH shift is used to measure beam angle, refractive index, displacement, temperature, and film thickness². The GH shift has also been used to characterize the permittivity and permeability of different materials in optical microscopy and lithography^{3,4}.

In the past decades, the GH shift has been studied extensively in various interfaces including absorptive dielectrics^{5–7}, metals^{8,9} and photonics^{10,11}. In addition to these natural materials, recently, a lot of interest has been emerged for zero index metamaterials (ZIMs) due to its stimulating electromagnetic features^{12–15}. Even though metamaterials cannot be found in nature, they could be manufactured very easily through plasmonic materials¹⁶ or doped semiconductors¹⁷. In particular, the GH shift in the reflected light is studied for *s* or *p* polarized light incident on epsilon-zero-index (ENZ) metamaterials¹⁸. Here, ENZ metamaterial is a kind of ZIMs. It is found that when a linear polarized light beam incident from air onto ENZ metamaterials, there is no GH shift for *p* polarized light beam; while a constant GH shift remains for *s* polarized light beam¹⁸.

As the coherent effect plays a key role in every investigation on wave phenomena^{19–24}, the influence from partial coherent light fields on the GH shifts raised a debate^{25–28}. This contradiction has been revealed thoroughly by Zubairy and coworkers for the first time²⁹, then by Ziauddin and coworkers in considering a partial coherent light beam incident on a medium³⁰. With a partial coherent light field, a counter-intuitive result on the GH shift with a strong dependence on spatial coherence is revealed^{29,30}.

In this Report, we consider a *p* or *s* polarized partial coherent light field incident on ENZ metamaterials and study the GH shift from the reflection. We show the influence of spatial coherence and beam width of partial coherent light on the GH shift for both *p* and *s* polarized lights. In contrary to the coherent counterparts, a totally different scenario for GH shift happens with partially coherent light fields. For *p* polarized partial coherent light beam, the magnitude of GH shift becomes non-zero with a small degree of spatial coherence; while for *s* polarized partial coherent beam, a relatively large GH shift can also be achieved in low spatial coherence. We also reveal the dependence on the beam width and the permittivity of ENZ metamaterials, both for *p* and *s* partial coherent light fields. By combining ENZ metamaterials and partial coherent light, our results provide a direction for highly precise measurement on GH shifts.

Results

Fundamental concept. We consider a partial coherent light field (*p* or *s* polarized) that incidents from air on ENZ metamaterials, which have the permeability $\mu_2 = 1$ and permittivity $\epsilon_2 \approx 0$, respectively. The partial coherent light makes an angle θ with respect to the *z*-axis, as depicted in Fig. 1. Based on Mercer's mode expansion, the electric field for *m* th-order mode in a partial coherent light at *z* = 0 can have the form^{30,31}

¹Institute of Photonics and Technologies, National Tsing-Hua University, Hsinchu 300, Taiwan. ²Department of Physics, COMSATS institute of information technology, Islamabad, Pakistan. Correspondence and requests for materials should be addressed to Z. (email: ziauddin@comsats.edu.pk) or R.-K.L. (email: rklee@ee.nthu.edu.tw)

Figure 1. Schematics of a partial coherent light incident on ENZ metamaterials, with an angle θ to the z -axis.

$$E_m^i(z, y) = \frac{1}{\sqrt{2\pi}} \int E_m(k_y - k_{y_0}) e^{i(k_z z + k_y y)} dk_y, \quad (1)$$

where

$$E_m(k_y - k_{y_0}) = \frac{1}{(2c\pi)^{1/4}} \times \frac{(-i)^m}{\sqrt{2^m m!}} \times e^{\frac{(k_y - k_{y_0})^2}{4c}} \times H_m\left(\frac{k_y - k_{y_0}}{\sqrt{2c}}\right), \quad (2)$$

being the angular spectrum that can be calculated using a Gaussian Schell-model (GSM) beam. For the normalized eigen-function in GSM light beams, one has³¹

$$E_m(y) = (2c/\pi)^{1/4} \times \frac{1}{\sqrt{2^m m!}} \times H_m[y\sqrt{2c}] e^{-cy^2}. \quad (3)$$

The corresponding angular spectrum $E_m(k_y)$ can be calculated by taking Fourier transform of Eq. (3). Here, $E_m(k_y)$ is replaced by $E_m(k_y - k_{y_0})$, and k_y is the y -component of the wave vector k , $k_{y_0} = k \sin \theta$, and θ is the incident angle. In Eqs (2–3), H_m is the Hermite polynomials and $c = [a^2 + 2ab]^{1/2}$, which can be calculated using the eigenvalues $\beta_m = A^2 [\pi/(a + b + c)]^{1/2} [b/(a + b + c)]^m$ of GSM beams with $a = (4w_s^2)^{-1}$, $b = (2w_g^2)^{-1}$. Furthermore, w_g and w_s are the spectral coherence and beam width of partial coherent light, respectively. We have the expression for $c = (q^2 + 4)^{1/2} / (4qw_s^2 \sec^2 \theta)$ with $w_s \rightarrow w_s \sec \theta$ and $w_g \rightarrow w_g \sec \theta$. To describe a partially coherent light field, the parameter $q = w_g/w_s$ measures the degree of the coherence in a GSM beam, which is also denoted as the spatial coherence.

For the reflected partial coherent light in m th-order mode, we have^{29,30}

$$E_m^r(y) = \frac{1}{\sqrt{2\pi}} \int r_{s,p}(k_y) E_m(k_y - k_{y_0}) e^{ik_y y} dk_y, \quad (4)$$

where $r_s(k_y)$ and $r_p(k_y)$ are the reflection coefficients.

For s and p polarized light, the corresponding reflection coefficients can be calculated as¹⁸:

$$r_s(k_y) = \frac{k_z - i\alpha}{k_z + i\alpha}, \quad (5)$$

$$r_p(k_y) = \frac{\epsilon_2 k_z - i\alpha}{\epsilon_2 k_z + i\alpha}, \quad (6)$$

with $k_z = k \cos \theta$ and $\alpha = \sqrt{k_y^2 - k\epsilon_2}$. Then, the compact expression for GH shifts in the reflected light in the m th-order mode can be derived analytically by the following expression²⁹

$$S_m^r = \frac{\int y |E_m^r(y)|^2 dy}{\int |E_m^r(y)|^2 dy}. \quad (7)$$

Effect of spatial coherence on GH shift. It has been reported earlier for a coherent light beam incident on lossless ENZ metamaterials, there is no GH shift for p polarized light; while for s polarized light the GH shift

Figure 2. (a) The GH shifts versus spatial coherence (q) for p polarized light. (b) The GH shifts versus spatial coherence (q) for s polarized light, when $w_s = 125\lambda$ and $\theta = 0.12$ rad.

is a constant value for different incident angles¹⁸. In the following, we proceed the study on the GH shifts in the reflected light for p or s polarization, but with partially coherent light beams. From the studies of spatial coherence on negative and positive GH shifts in the reflected light^{29,30}, the amplitude of GH shift becomes larger for small values of spatial coherence (q) and decreases as the value of the spatial coherence increases. It is remarked that the incident light beam becomes more and more coherent with a larger value of q and becomes incoherent with a small value of q . In using Eq. (7) to analyze the behavior of GH shift for different number of modes of p and s polarized partial coherent light, each mode of partial coherent light is perfectly coherent.

Now, to see the influence of spatial coherence on the GH shift, we start with a p polarized light incident on ENZ metamaterials. In Fig. 2(a), we plot the corresponding GH shifts in the reflected light versus spatial coherence q for $m = 0$ th- and $m = 8$ th-orders, respectively, with a fixed incident angle $\theta = 0.12$ rad. To give a clear representation, first, we concentrate our attention to study the effect of spatial coherence over the GH shift in the reflected light by considering a lossless ENZ metamaterial with the permittivity $\varepsilon_2 = 10^{-3}$. For the lossless ENZ metamaterial, the amplitude of GH shift in the reflected light is high for small values of q , and approaches to zero when q increases. This clearly shows that the amplitude of the GH shift is not zero for partial coherent light (small q) and approaches to zero when light becomes coherent (large q), see the black and green curves for different modes of partial coherent light in Fig. 2(a). As reported earlier in ref. 18 that there is no GH shift in the reflected light for p polarization because they have used a coherent light beam. This is also shown in our analysis that the GH shift in the reflected light for p polarized coherent light approaches to zero as the coherence of the incident beam increases.

Next, we consider the loss effect in the materials, *i.e.*, by setting $\text{Im}[\varepsilon_2] = 0.01$. As shown in Fig. 2(a), the corresponding GH shifts in the reflected light versus the coherence parameter q for p polarization are shown in the red and blue curves for the $m = 0$ and 8 -th modes, respectively. By introducing absorption in a system, the GH shift in the reflected light becomes negative. This is due to the fact that when a p polarized light reflected from an absorptive medium, a phase changes abruptly at the Brewster angle and induces a negative GH shift⁶. Similar behavior of the GH shift in the reflected light is also reported for absorptive ENZ metamaterials in ref. 18.

Furthermore, we consider a s polarized beam incident on ENZ metamaterials and study the influence of spatial coherence on the GH shift in the reflected light for both lossless and absorptive media. For lossless ENZ metamaterials, positive GH shifts are revealed for a fixed incident angle $\theta = 0.12$ rad by considering $m = 0$ and $m = 8$, see the dashed black and green curves in Fig. 2(b). The amplitude of GH shift is larger for small values of spatial coherence and decreases with an increment in the spatial coherence. In the limit of a perfect coherent light field, the amplitude of GH shifts remains as a constant value. To give a clear illustration for such a constant GH shift in the reflected light for coherent light fields, we show the enlarged plot for the GH shifts ranging from $\log_{10}[q] = 0$ to $\log_{10}[q] = 1.7$ in the inset of Fig. 2(b). The GH shifts in the reflected light in the inset clearly shows that the amplitude remains constant when the value of q increases. The corresponding constant amplitude of GH shifts for the coherent light region are $S_0^r = 0.01\lambda$ (black curve) and $S_8^r = 0.063\lambda$ (green curve), respec-

Figure 3. (a) The GH shifts versus beam width (w_s) for p polarized light. (b) The GH shifts versus beam width (w_s) for s polarized light, when $q = 0.01$ and $\theta = 0.12$ rad.

tively. Here, the amplitude of GH shifts in the reflected light is quite small because we consider a large beam width to suppress the distortion of light beam. In contrary to the coherent incident light field, we also reveal that the GH shift in the reflected light does not remain constant for partial coherent light at all the incident angles θ . When loss is considered in ENZ metamaterials, a similar behavior on the modified GH shifts in the reflected light can be seen for the dashed red and blue curves in Fig. 2(b).

Effect of beam widths. It is known that the beam width also plays a key role in the observation of GH shifts with a Gaussian beam^{22,24}, as well as a partial coherent light beam^{30,31}. By increasing the beam width, the effect of partial coherence on the GH shift decreases and vice versa. Following this concept, here, we consider a spatial coherence $q = 0.01$ for p polarized partial coherent light incident on ENZ metamaterials. For lossless ENZ metamaterials, the GH shift in the reflected light for different modes of partial coherent light beam is depicted in Fig. 3(a). The black and green curves in Fig. 3(a) show that the amplitude of GH shift varies with the beam width of partial coherent light, *i.e.*, the amplitude is larger for small values of beam widths and decreases to zero as the beam width increases. Based on these results, one can safely conclude that for small values of beam widths the amplitude of the GH shift is not equal to zero, but approaches to zero for a larger beam width in a p polarized light beam.

As an absorptive ENZ metamaterial is considered, in Fig. 3(a). The red and blue curves represent negative GH shifts in the reflected light for p polarized light beam, which varies with the beam width, too. These curves of negative GH shifts demonstrate similar behavior as those shown in Fig. 2(a). As for s polarized beams, shown in Fig. 3(b), the GH shift in the reflected light is depicted for different modes of partial coherent light field versus beam width w_s . Again, we have similar behavior on the GH shift in the reflected light versus the beam width, as those shown in Fig. 2(b) for different spatial coherence q . In comparison of Figs 2 and 3, we notice that the influence of spatial coherence q and beam width w_s have the same role on the GH shifts in the reflected light of ENZ metamaterials.

Effect of $\text{Re}[\epsilon_2]$. By varying the $\text{Re}[\epsilon_2]$ (permittivity of metamaterials), it is known that the GH shifts in the reflected light are a little bit affected both for p and s polarized light¹⁸. Here, we examine the influence of $\text{Re}[\epsilon_2]$ on the GH shift in the reflected light for partially coherent light fields. We consider different values of spatial coherence q and study the GH shift for p and s polarized light, as a function of real part of metamaterial permittivity, see Fig. 4. For different values of q , the GH shift in the reflected light for p polarized light is shown in Fig. 4(a). There is no GH shift when $\text{Re}[\epsilon_2] = 0$ for different spatial coherence q , but for $\text{Re}[\epsilon_2] \approx 0$ the amplitude of GH shift is non-zero for partial coherent light. Furthermore, the magnitude of the GH shift is large for small values of spatial coherence and decreases as the spatial coherence increases.

Figure 4. (a) The GH shifts versus permittivity of ENZ metamaterials (ϵ_2) for p polarized light. (b) The GH shifts versus permittivity of ENZ metamaterials (ϵ_2) for s polarized light, when $m = 0$ and $\theta = 0.12$ rad.

The comparison between the black (for $q = 0.015$) and blue (for $q = 10$) curves in Fig. 4(a), clearly shows that the magnitude of GH shift is not zero for partial coherent light (p polarized) in ENZ metamaterials. Again, the magnitude of the GH shift approaches to zero in ENZ metamaterials when the light becomes perfectly coherent, *i.e.*, at $q = 10$, see the blue curve in Fig. 4(a). It is also noticed that the magnitude of GH shift for partial coherent light (p polarized) increases more quickly with the increment in $\text{Re}[\epsilon_2]$, as compared to a coherent light beam (p polarized), see the black and blue curves in Fig. 4(a). Similarly, for s polarized light beams, as shown in Fig. 4(b), the GH shift is not zero for $\text{Re}[\epsilon_2] = 0$ and the magnitude increases for ENZ metamaterials when $\text{Re}[\epsilon_2]$ increases. The magnitude of GH shift for s polarized partially coherent light is larger than that for a coherent light beam; see the black and blue curves in Fig. 4(b).

Discussion

In this work, we considered a partial coherent light field incident on ENZ metamaterial with an angle θ to the z -axis. The corresponding GH shifts in the reflected light both for p and s polarized beams are studied, with the influence from spatial coherence, beam width, and permittivity of metamaterials. We show that for lossless ENZ metamaterials, the magnitude of GH shift is non-zero for p polarized partially coherent light beams; while it approaches to zero as the degree of spatial coherence increases. With small values of spatial coherence and beam width, the GH shifts both for p and s polarized beams increase first, and then decrease for higher values of spatial coherence and beam width. As in most of practical implementations, such as the x-ray beams^{32,33}, the coherent light sources are usually not available. With the introduction of partially coherent beams, our results on the GH shifts may provide a direction on the applications for partial coherent light sources in ENZ metamaterials.

References

- Goos, F. & Hänchen, H. Ein neuer und fundamentaler Versuch zur Totalreflexion. *Ann. Phys.* **436**, 333–346 (1947).
- Hashimoto, T. & Yoshino, T. Optical heterodyne sensor using the Goos-Hänchen shift. *Opt. Lett.* **14**, 913 (1989).
- Hu, X., Huang, Zhang, Y. W., Qing, D.-K. & Peng, J. Opposite Goos-Hänchen shifts for transverse-electric and transverse-magnetic beams at the interface associated with single-negative materials. *Opt. Lett.* **30**, 899 (2005).
- de Fornel, F. “*Evanescent Waves*” (Springer, Berlin, 2001).
- Wild, W. J. & Giles, C. L. Goos-Hänchen shifts from absorbing media. *Phys. Rev. A* **25**, 2099 (1982).
- Lai, H. M. & Chan, S. W. Large and negative Goos-Hänchen shift near the Brewster dip on reflection from weakly absorbing media. *Opt. Lett.* **27**, 9 (2002).
- Wang, L. G., Chen, H. & Zhu, S. Y. Large negative Goos-Hänchen shift from a weakly absorbing dielectric slab. *Opt. Lett.* **30**, 2936 (2005).
- Leung, P. T., Chen, C. W. & Chiang, H. P. Large negative Goos-Hänchen shift at metal surfaces. *Opt. Commun.* **276**, 206 (2007).
- Merano, M. *et al.* Observation of Goos-Hänchen shifts in metallic reflection. *Opt. Express* **15**, 15928 (2007).
- He, J., Yi, J. & He, S. Giant negative Goos-Hänchen shifts for a photonic crystal with a negative effective index. *Opt. Express* **14**, 3024 (2006).

11. Soboleva, I. V., Moskalenko, V. V. & Fedyanin, A. A. Giant Goos-Hänchen effect and Fano resonance at photonic crystal surfaces. *Phys. Rev. Lett.* **108**, 123901 (2012).
12. Silveirinha, M. & Engheta, N. Tunneling of electromagnetic energy through subwavelength channels and bends using ϵ -near-zero materials. *Phys. Rev. Lett.* **97**, 157403 (2006).
13. Liu, R. *et al.* Experimental demonstration of electromagnetic tunneling through an epsilon-near-zero metamaterial at microwave frequencies. *Phys. Rev. Lett.* **100**, 023903 (2008).
14. Nguyen, V. C., Chen, L. & Halterman, K. Total transmission and total reflection by zero index metamaterials with defects. *Phys. Rev. Lett.* **105**, 233908 (2010).
15. Huang, X. *et al.* Dirac cones induced by accidental degeneracy in photonic crystals and zero-refractive-index materials. *Nat. Mater.* **10**, 582 (2011).
16. Wes, P. R. *et al.* Searching for better plasmonic materials. *Laser Photon. Rev.* **4**, 6 (2010).
17. Adams, D. C. *et al.* Funneling light through a subwavelength aperture with epsilon-near-zero materials. *Phys. Rev. Lett.* **107**, 133901 (2011).
18. Xu, Y., Chan, C. T. & Chen, H. Goos-Hänchen effect in epsilon-near-zero-metamaterials. *Sci. Reports* **5**, 8681 (2015).
19. Wang, L. G., Ikram, M. & Zubairy, M. S. Control of the Goos-Hänchen shift of a light beam via a coherent driving field. *Phys. Rev. A* **77**, 023811 (2008).
20. Ziauddin, Qamar, S. & Zubairy, M. S. Coherent control of the Goos-Hänchen shift. *Phys. Rev. A* **81**, 023821 (2010).
21. Ziauddin & Qamar, S. Gain-assisted control of the Goos-Hänchen shift. *Phys. Rev. A* **84**, 053844 (2011).
22. Ziauddin & Qamar, S. Control of the Goos-Hänchen shift using a duplicated two-level atomic medium. *Phys. Rev. A* **85**, 055804 (2012).
23. Ziauddin, Qamar, S. & Abbas, M. Amplitude control of the Goos-Hänchen shift via Kerr nonlinearity. *Laser Phys. Lett.* **11**, 015201 (2014).
24. Ziauddin & Qamar, S. Effect of width of incident Gaussian beam on the longitudinal shifts and distortion in the reflected beam. *Opt. Commun.* **319**, 1 (2014).
25. Aiello, A. & Woerdman, J. P. Role of spatial coherence in Goos-Hänchen and Imbert-Fedorov shifts: reply to comment. *Opt. Lett.* **36**, 3151 (2011).
26. Löffler, W., Aiello, A. & Woerdman, J. P. Spatial coherence and optical beam shifts. *Phys. Rev. Lett.* **109**, 213901 (2012).
27. Merano, M., Umbriaco, G. & Mistura, G. Observation of nonspecular effects for Gaussian Schell-model light beam. *Phys. Rev. A* **86**, 033842 (2012).
28. Wang, L.-Q., Wang, L.-G., Zhu, S.-Y. & Zubairy, M. S. The influence of spatial coherence on the Goos-Hänchen shift at total internal reflection. *J. Phys. B* **41**, 055401 (2008).
29. Wang, L.-G., Zhu, S. Y. & Zubairy, M. S. Goos-Hänchen shifts of partially coherent light field. *Phys. Rev. Lett.* **111**, 223901 (2013).
30. Ziauddin, Chuang, Y.-L. & Lee, R.-K. Negative and positive Goos-Hänchen shifts of partially coherent light fields. *Phys. Rev. A* **91**, 013803 (2015).
31. Mandel, L. & Wolf, E. *“Optical Coherence and Quantum Optics”* (Cambridge University Press, Cambridge, England, 1995).
32. Abbey, B. *et al.* *Adv. Phys.* **59**, 1 (2010).
33. Tamasaku, K. & Ishikawa, T. *Acta Crystallogr. Sect. A* **58**, 408 (2002).

Acknowledgements

This work is supported in part by Ministry of Science and Technology, Taiwan, under contract No. 101-2628-M-007-003-MY4.

Author Contributions

Z. and R.-K.L. conceived the idea. Y.-L.C. and S.Q. helped in the calculations, numerical simulations and analysis. R.-K.L. supervised the whole project. All the authors contributed in writing the manuscript.

Additional Information

Competing financial interests: The authors announce no competing financial interests.

How to cite this article: Ziauddin *et al.* Goos-Hänchen shift of partially coherent light fields in epsilon-near-zero metamaterials. *Sci. Rep.* **6**, 26504; doi: 10.1038/srep26504 (2016).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>