

SCIENTIFIC REPORTS



OPEN

Vitamin k3 inhibits protein aggregation: Implication in the treatment of amyloid diseases

Parvez Alam¹, Sumit Kumar Chaturvedi¹, Mohammad Khursheed Siddiqi¹, Ravi Kant Rajpoot², Mohd Rehan Ajmal¹, Masihuz Zaman¹ & Rizwan Hasan Khan¹

Received: 09 February 2016

Accepted: 29 April 2016

Published: 27 May 2016

Protein misfolding and aggregation have been associated with several human diseases such as Alzheimer's, Parkinson's and familial amyloid polyneuropathy etc. In this study, anti-fibrillation activity of vitamin k3 and its effect on the kinetics of amyloid formation of hen egg white lysozyme (HEWL) and A β -42 peptide were investigated. Here, in combination with Thioflavin T (ThT) fluorescence assay, circular dichroism (CD), transmission electron microscopy and cell cytotoxicity assay, we demonstrated that vitamin k3 significantly inhibits fibril formation as well as the inhibitory effect is dose dependent manner. Our experimental studies inferred that vitamin k3 exert its neuro protective effect against amyloid induced cytotoxicity through concerted pathway, modifying the aggregation formation towards formation of nontoxic aggregates. Molecular docking demonstrated that vitamin k3 mediated inhibition of HEWL and A β -42 fibrillogenesis may be initiated by interacting with proteolytic resistant and aggregation prone regions respectively. This work would provide an insight into the mechanism of protein aggregation inhibition by vitamin k3; pave the way for discovery of other small molecules that may exert similar effect against amyloid formation and its associated neurodegenerative diseases.

Amyloid fibrillation of proteins and peptides are responsible for wide number of diseases in humans such as Alzheimer's, Parkinson's, Huntington's, and Diabetes II etc^{1,2}. Despite the unrelated amino acid sequences and tertiary structure, proteins can unfold and assemble to form amyloid fibrils with similar ultrastructure and biochemical properties³. Amyloid fibrils do have long unbranched shape, enriched beta sheet content, cytotoxic nature, high surface hydrophobicity and specific tinctorial properties⁴. Although, the propensity to form amyloid fibrils is modulated by polypeptide sequence and varies from sequence to sequence⁵.

HEWL is a small size, 129 amino acid residues long protein that has been used as a model to study the mechanism of amyloid formation and inhibition⁶. Further, HEWL is homologous to human lysozyme which is responsible for non-neuropathic systemic amyloidosis resulting in deposition of amyloids in liver and kidney in humans. Alzheimer's diseases is one of the most common form of dementia, fibrillar aggregates of amyloid beta peptides of 39–43 amino acids is the main factor responsible for the disease. The amyloid cascade hypothesis presumes that amyloid aggregates self-assembled from misfolded A β peptides, affect the structure and function of neuronal cells as well as stimulate apoptosis that is responsible for synaptic dysfunction and neurodegeneration⁷.

Currently, many methodologies have been developed to find suitable cure for amyloid associated diseases and is still progressing. Small molecules/drugs which can stabilize the native state of protein retard the fibrillation process or reverse the misfolding process can serve as promising agent against debilitating diseases⁸. Many compounds such as polyphenols, various drugs, flavonoids, vitamins and some metal complexes are known to inhibit the aggregation process of proteins either *in vitro* or *in vivo*^{9–12}. Recently numbers of reports are available advocating the anti-aggregation effect of nanomaterials^{13–16}. Varieties of vitamins including vitamin A, vitamin E, vitamin D and vitamin k attenuated the aggregation process in proteins and also delayed the onset of dementia^{17–21}. In order to get further insight into the mechanism of vitamins mediated aggregation inhibition of proteins; we evaluated the effect of vitamin k3, a synthetic analogue of vitamin k on fibrillation of HEWL and A β -42 peptide.

¹Molecular Biophysics and Biophysical Chemistry Group, Interdisciplinary Biotechnology Unit, Aligarh Muslim University, Aligarh-202002, India. ²Recombinant Gene Product Group, International Centre for Genetic Engineering and Biotechnology, Aruna Asaf Ali Marg, New Delhi-110067, India. Correspondence and requests for materials should be addressed to R.H.K. (email:rizwanhkh@hotmail.com or rizwanhkh@1@gmail.com)




Figure 1. (a) ThT fluorescence intensity of HEWL (15 μM) at 485 nm in presence of varying concentration of vitamin k3 (0–200 μM). Samples were incubated at 65 °C for 120 hours. (b) ThT fluorescence kinetics of HEWL in absence and presence of vitamin k3 (50 and 100 μM) (c) ThT fluorescence spectra of HEWL incubated at 65 °C over 120 hours in absence and presence of vitamin k3 (50 and 100 μM) (d) Congo red binding absorption spectra of HEWL (15 μM) in the absence and presence of vitamin k3. Samples were incubated with two different concentration of vitamin k3 (50 and 100 μM) at 65 °C for 120 hours. Experimental data represent the average ± s.d (n = 3).

Vitamin k3, an anti-inflammatory and anti-cancer agent, is employed for post translational modification of proteins including blood clotting factors. Similar behaviour to vitamin k makes it a good choice as a health supplement and component of multivitamin drugs^{22–24}.

In this study, mechanism of action of vitamin k3 has been investigated on HEWL and Aβ-42 amyloid fibrillogenesis process by exploiting various complementary techniques. To assess the effect of vitamin k3 on the growth phase of amyloid, aggregation kinetics study was performed by ThT binding assay. Changes in secondary structure were examined by CD spectroscopy. Hydrodynamic radii were monitored by using dynamic light scattering measurements. Morphology of aggregates was studied by using transmission electron microscopy. Interaction mode of vitamin k3 with HEWL and Aβ-42 was studied using molecular docking. Further, cell viability (MTT) assay was accomplished by using SH-SY5Y cell lines.

Results and Discussion

ThT Binding Assay. Thioflavin T (ThT) fluorescence assay was carried out to quantify amyloid formation of HEWL aggregates and examining the impact of vitamin k3 on rate of fibril formation. ThT fluorescence has been widely used to detect cross β-sheet structure of amyloids²⁵. The growth of amyloid formation of HEWL was monitored and characterized by ThT. On interaction with cross-β sheet structure, ThT results in significant rise in fluorescence intensity^{26,27}. The inhibitory effect of vitamin k3 on fibrillation process was concentration dependent (Fig. 1a). Kinetics of amyloid formation in the absence and presence of vitamin k3 (50 μM and 100 μM) is shown in Fig. 1b. As shown in Fig. 1b, the aggregation kinetics of HEWL exhibit a typical sigmoidal appearance containing a lag phase associated with nucleation, a fast growth phase linked to the elongation and propagation of fibrils, and a final stationary phase²⁸. Further, these results are in agreement with the nucleation dependent polymerisation for amyloidogenic proteins. However, simultaneous incubation of HEWL with 50 and 100 μM vitamins k3 significantly attenuated the ThT fluorescence intensity throughout the time frame of the experiment (Fig. 1b). ThT fluorescence spectra of HEWL and HEWL in combination with vitamin k3 after 120 hours incubation at 65 °C are shown in Fig. 1c. It can be seen from the figure that ThT fluorescence intensity of HEWL was reduced from 222 to 96 and 50 in presence of 50 and 100 μM of vitamin k3 respectively. Further addition of vitamin k3 to preformed HEWL amyloid fibrils did not lead to decrease in ThT fluorescence intensity, it suggests that vitamin k3 cannot reverse the fibril formation but can only inhibit its formation. These results revealed that vitamin k3 inhibits fibrillation from very beginning of log phase as observed in the case of EGCG which inhibits fibrillogenesis of Chicken Cystatin²⁹.




Figure 2. (a) ANS fluorescence intensity of HEWL (15 μM) at 480 nm incubated at 65 $^{\circ}\text{C}$ over 120 hours in absence and presence of vitamin k3 (50 and 100 μM). *Statistically different from the HEWL $p \leq 0.01$. (b) Far-UV CD spectra of HEWL (15 μM) at 25 $^{\circ}\text{C}$ in absence and presence of vitamin k3. Inset (b) shows HEWL (15 μM) incubated at 65 $^{\circ}\text{C}$ over a period of 120 hours in the absence and presence of vitamin k3 (50 and 100 μM).

Congo red binding assay. To corroborate the findings of ThT assay, the inhibition of HEWL aggregation was also confirmed by Congo red binding assay. A red shift in absorbance maxima of Congo red absorption spectrum from 490 nm to 540 nm indicates presence of increased cross β -sheet rich structure³⁰. HEWL samples incubated for 120 hours at 65 $^{\circ}\text{C}$ showed marked increase in absorbance accompanied with red shift in comparison to Congo red alone, signifying the presence of amyloid fibrils as represented in Fig. 1d. We observed that this increased peak shift in HEWL was prevented in presence of vitamin k3 and more pronounced effect was observed in presence of 100 μM ³¹. These results also in accord with preceding ThT fluorescence results suggesting that fibril formation is inversely related to the concentration of vitamin k3.

Surface hydrophobicity modulation by vitamin k3. The impacts of vitamin k3 on surface hydrophobicity of HEWL were examined based on ANS fluorescence emission. ANS is widely used to characterise the protein folding intermediate states and to detect the presence of hydrophobic patches on the surface of protein/amyloids³². Interaction of ANS with solvent exposed hydrophobic patches on the protein leads to significant increase in fluorescence intensity³³. The surface hydrophobicity of HEWL at 25 $^{\circ}\text{C}$ with or without vitamin k3 was found to be very low because in native protein hydrophobic patches are hidden inside in compactly folded structure (data not shown). In contrast, a marked increase in ANS fluorescence intensity was observed in HEWL upon incubation at 65 $^{\circ}\text{C}$ for 120 hours (Fig. 2a)³⁴. Conversely, low ANS fluorescence intensity was observed in HEWL-vitamin k3 (50 and 100 μM) co-incubated samples. This suggests that exposure of hydrophobic regions is significantly decreased in presence of vitamin k3. The reduced ANS fluorescence intensity observed in presence of vitamin k3 which indicates less exposure of hydrophobic core to solvent, suggested that vitamin k3 could stabilise the whole conformation of HEWL even though it was subjected to at high temperature. It is believed that non covalent bonds stabilize the core structure of all amyloid fibrils³⁵. Vitamin k3 might have interacted via non covalent interaction with amino acid residues and interfere with the formation of HEWL fibrils.

Conformational transition in HEWL during fibrillation. CD is well known spectroscopic technique to elucidate the secondary structure of proteins³⁶. The conversion of alpha helix or random coil to beta sheet is hallmark of amyloid formation³⁷. The CD spectra of HEWL obtained prior to incubation exhibited the typical profile of predominantly alpha helical conformation and in presence of vitamin k3, increased in alpha helical content from 32 to 41% was observed at 25 $^{\circ}\text{C}$ (Fig. 2b)³⁸. As was expected, the fibrillation of HEWL while incubating at 65 $^{\circ}\text{C}$ for 120 hours resulted in decreased of alpha helical fraction and concomitant increase in beta sheet fraction (inset Fig. 2b and supplementary Table 1). Although we observed single minima at around 218 nm in HEWL- vitamin k3 co-incubated samples but the negative CD value at this point was reduced. To calculate the secondary structure composition of each CD spectra, k2d3 software was used for calculating various secondary structure components (alpha helix, beta sheet and random coil)³⁹. The anticipated decrease in beta sheet content was observed in presence of vitamin k3 in HEWL. These results suggest that structural transition in HEWL was mitigated upon co-incubation with vitamin k3⁴⁰.

Dynamic light scattering measurements. In order to determine the size of protein aggregates formed in the absence and presence of vitamin k3, dynamic light scattering measurements were performed. Results are summarised in Fig. 3. It is quite apparent from figure that hydrodynamic radii (R_h) of HEWL at 25 $^{\circ}\text{C}$ decreases slightly (2 nm to 1.8 nm) in presence of vitamin k3. Decrease in R_h might be attributed to disturbance of solvation sphere around protein suggesting that vitamin k3 stabilize the native state of protein which is in consistent with our CD data. The R_h value of HEWL increases with respect to time upon incubation at 65 $^{\circ}\text{C}$ but in presence of vitamin k3, concentration dependent decrement in R_h was observed. HEWL incubated for 120 hours shows R_h




Figure 3. DLS pattern of HEWL in absence and presence of vitamin k3 at 25 °C (a) HEWL (b) HEWL + 50 μ M vitamin k3 (c) HEWL + 100 μ M vitamin k3 and after incubation at 65 °C over a period of 120 hours (d) HEWL (e) HEWL + 50 μ M vitamin k3 and (f) HEWL + 100 μ M vitamin k3.




Figure 4. Transmission Electron Microscopic images (a) HEWL at 25 °C (b) HEWL incubated at 65 °C over 120 hours (c) HEWL + 50 μ M vitamin k3 incubated at 65 °C over 120 hours (d) HEWL + 100 μ M vitamin k3 incubated at 65 °C over 120 hours.

value around 400 nm but in presence of vitamin k3 this value decrease to 40–60 nm. This decrease in R_h value might be attributed to the stabilization of protein in presence of vitamin k3.

Transmission electron microscopy. To further investigate the efficacy of vitamin k3 against HEWL fibril formation, TEM analysis was performed. As shown in Fig. 4, HEWL sample alone incubated for 120 hours




Figure 5. MTT reduction assay for cell cytotoxicity of 120-hours aged HEWL amyloid fibrils in absence and presence of vitamin k3 (50 and 100 μM) on SH-SY5Y cell lines. Control represents the cells without exposed to HEWL fibrils. *Statistically different from the control group, $p \leq 0.01$ and #statistically different from the HEWL, $p \leq 0.01$.

formed large, branched fibrils which are characteristic feature of amyloids⁴¹. Samples of HEWL in presence of vitamin k3 (50 and 100 μM) showed no fibrillar aggregate. This result reveals that vitamin k3 inhibits aggregation process of HEWL in concentration dependent manner. The possible mechanism for vitamin k3 mediated reduction in HEWL aggregation may be coercing of the protein in its native state³⁴. This observation is similar to previous studies on the effect of osmolytes on insulin fibrillation⁴².

Vitamin k3 reduces amyloid inflicted cytotoxicity. To investigate whether HEWL amyloids are cytotoxic or not and if this cytotoxicity could be relieved by adding vitamin k3, we employed SH-SY5Y (human neuroblastoma) cell line for cell viability assay at 24 hours. When vitamin k3 (50 and 100 μM) added to SH-SY5Y cell line showed no cytotoxic effect (data not shown). Further vitamin k3 added to preformed amyloids have no protective effect on cytotoxicity, ruling out other protective mechanism apart from aggregation inhibition (data not shown). SH-SY5Y cell lines were treated with 120 hours aged HEWL amyloids and their effect on cell viability was checked and data is represented in Fig. 5. It is inferred from the figure that cell viability decreases in presence of 120 hours aged HEWL amyloid and reaches up to 38%³⁴. Further, it is clear from the cytotoxicity profile that HEWL incubated with vitamin k3 enhance cell viability, even in presence of 120 hour aged amyloid. Cell viability was rescued to 73% and 80% in presence of 50 μM and 100 μM vitamin k3 respectively. Cell toxicity of HEWL may be attributed to the formation of amyloids which disrupts the cell membrane. In presence of vitamin k3, gain in cell viability was observed. This confirmed that increase in cell viability was due to anti amyloidogenic behaviour of vitamin k3⁴³. These results suggest that the non fibrillar aggregates formed in the presence of vitamin k3 are less toxic to neuronal cells therefore related compounds may have therapeutics intervention against amyloids diseases⁴⁴.

Binding mode of vitamin k3 to HEWL. Based on the spectroscopic and electron microscopy data it can be inferred that vitamin k3 inhibits amyloid formation of HEWL. Information on putative binding site and mode of interaction between HEWL and vitamin k3 was obtained from molecular docking using Autodock 4.2 program⁴⁵. Molecular docking results are summarises in supplementary Figure S1 and supplementary Table 2. It is clear from the figure that vitamin k3 interact with HEWL with the following amino acid residues Lys¹, Phe³, Glu⁷, Ala¹⁰, Ala¹¹, Arg¹⁴, His¹⁵, Thr⁴⁰, Gln⁴¹, Leu⁸⁴, Ser⁸⁵, Ser⁸⁶, Asp⁸⁷ and Ile⁸⁸. The main interaction forces between HEWL and vitamin k3 are hydrophobic interaction. It is clear from these results that vitamin k3 interact with proteolytically resistant region (32–108). Interaction with this region might be responsible for anti amyloidogenic behavior. In spite of no interaction with amyloid prone region of HEWL vitamin k3 inhibits aggregation similar to inhibitory action of nanobody on aggregation of human lysozyme³⁴. As generally acknowledged that protein aggregate mostly through hydrogen bonding, aromatic and hydrophobic interaction, disrupting of such interactions by vitamin k3 may thus inhibits amyloid formation⁴⁶. The value of Gibbs free energy for the best pose was $-3.98 \text{ kcal mol}^{-1}$, that suggests the formation of a stable protein–ligand complex⁴⁷. Our results implied that the inhibition of HEWL fibrillogenesis can be attributed mainly to the binding of vitamin k3 to HEWL proteolytically resistant region.

Vitamin k3 inhibits amyloid fibrillation of A β peptide. Inhibition of A β -42 fibrillation was also examined in presence of vitamin k3 (50 and 100 μM). Figure 6a,b represented the ThT kinetics of A β -42 aggregation and ThT fluorescence intensity of 70 hours aged amyloids respectively, in the absence and presence of vitamin k3. A β -42 aggregates gave strong ThT emission, producing a typical sigmoidal curve that reaches the plateau stage after almost 50 hours incubation. As predicted this observed behavior was opposite to that we observed when A β -42 incubated in presence of vitamin k3. Under these conditions the kinetic of A β -42 aggregation was delayed and initiated only after 25 hours and 40 hours of incubation in presence of 50 and 100 μM of vitamin k3 respectively. Addition of vitamin k3 significantly inhibited the aggregation of A β -42 which is in well agreement with previous reports delineating the effect of small molecules on aggregation⁴⁸.




Figure 6. (a) ThT fluorescence kinetics measurement of A β -42 aggregation when incubated at 37 °C over a period of 70 hours in absence and in presence of vitamin k3 (50 and 100 μ M) (b) ThT fluorescence spectra of A β -42 after incubation at 37 °C for 70 hours in absence and in presence of vitamin k3 (50 and 100 μ M). Experimental data represent the average \pm s.d (n = 3).

Congo red binding assay was used as an indicator of extended β -pleated sheet structures and A β -42 aggregation. Congo red absorption spectra of A β -42 in absence and presence of vitamin k3 is shown in supplementary Figure S2a. Protein solution containing amyloid fibrils exhibited an increase in Congo red absorption accompanied with a red shift of the spectral maximum. Analogous to ThT fluorescence results, co-incubation of A β -42 with vitamin k3 resulted in both reduction of absorbance intensity and also blue shift in wavelength maximum.

ANS has long been used as fluorescent molecular probe for examining hydrophobicity of proteins. ANS binding was used to study the changes in hydrophobicity in A β -42 upon incubation in absence and presence of vitamin k3. The ANS fluorescence maxima in absence and presence of vitamin k3 are shown in supplementary Figure S2b. It is clear from the figure that A β -42 alone shows very high ANS binding suggesting exposure of hydrophobic patches in contrast the ANS binding was significantly reduced in presence of vitamin k3. Similar observation was made by Ghaghghaei *et al.* in the protective role of crocin in A β -42 fibril formation⁴⁹.

In order to detect the change in the secondary structure, CD spectroscopy in the far UV region (200–250) was carried out for A β -42 in absence and presence of vitamin k3. CD spectra of native A β -42 displayed characteristic spectra of random coil structure of the peptide (supplementary Figure S3) but almost no structural change was observed in the peptide at 0 hour in the presence of vitamin k3 (data not shown). As shown in figure 70 hour of incubation at 37 °C a negative peak at around 218 nm was observed suggesting the structural transition from random coil to beta sheet in the peptide. But the structural conversion from random coil to beta sheet was retarded by vitamin k3 manifested by decrease negative band at around 218 nm as compared to native A β -42. These results indicate that vitamin k3 inhibits A β -42 aggregation by decreasing the beta sheet formation. Similar results were observed by Dai *et al.* for Chitosan Oligosaccharides inhibition of A β -42 aggregation⁵⁰.

Dynamic light scattering measurements was employed to get qualitative estimation of the size of A β -42 aggregates. A β -42 peptide without incubation has hydrodynamic radii (R_h) of 4 nm as shown in supplementary Figure S4 and almost no change in R_h was observed in presence of vitamin k3 (date not shown)⁵¹. But after incubation of A β -42 for 70 hour at 37 °C results in increase in hydrodynamic radii to around 550 nm. In contrast when A β -42 was incubated with vitamin k3 the R_h value was found to be between to 35–55 nm (Fig. S4). This result further indicates that vitamin k3 inhibits A β -42 aggregation.

TEM was performed to determine the morphology of aggregates formed. It is clear from the Fig. 7 that control A β -42 samples shows long branched fibrils which are characteristic feature of amyloids⁵². As compared to control samples in presence of vitamin k3 short and sparsely populated non fibrillar aggregate were formed. Based on TEM images, our results thus indicate that incubation of A β -42 with vitamin k3 inhibit A β -42 aggregation and significantly alter the morphology of A β fibrillar aggregates.

Vitamin k3 reduce amyloid beta inflicted cell cytotoxicity. Cultured human neuroblastoma cells (SH-SY5Y) were treated with an aliquot of A β -42 with and without vitamin k3 and cell viability was assessed by MTT assay, which measures metabolic activity. As shown in Fig. 8, A β -42 amyloids significantly reduce cell viability approximately up to 28% in SH-SY5Y cells. However A β -42 co-incubated with vitamin k3 (50 and 100 μ M) showed higher level of viability suggesting its chemo preventive role against A β -42 aggregates induced cytotoxicity. It has been reported previously that exposure of neurons to β - amyloid induces degeneration and cell death involving apoptotic pathway that may contribute to the neuronal loss associated with Alzheimer disease⁵³. Similar mechanism may involve in cell degeneration in the present study after exposure to A β -42 amyloid and our data suggest that vitamin k3 prevented this cellular degeneration. Previous works suggest that β - amyloid induced cytotoxicity




Figure 7. Transmission Electron Microscopic images of (a) A β -42 at 25°C (b) A β -42 incubated for 70 hours at 37°C (c) A β -42 + 50 μ M vitamin k3 incubated for 70 hours at 37°C (d) A β -42 + 100 μ M vitamin k3 incubated for 70 hours at 37°C.




Figure 8. SH-SY5Y cell viability after being exposed to 70 hours aged A β -42 fibrils formed in the absence and presence of (50 and 100 μ M) vitamin k3. *Statistically different from the control group, $p \leq 0.01$ and #statistically different from the A β -42, $p \leq 0.01$.

could be mediated by oxidative stress and free radicals. It is known that vitamin k3 has free radical scavenging and anti-oxidant activity and this could inhibit the A β -42 amyloid induced cell death⁵⁴.

Binding mode of vitamin k3 to A β -42 peptide. Molecular docking study was performed to get insight into the type of interactions involved between vitamin k3 and A β -42 that are responsible for aggregation inhibition. Docking results are summarized in supplementary Figure S5 and supplementary Table 3. Figure S5b shows that vitamin k3 interacts with six amino acids residues of A β -42 including Ser⁸, Glu¹¹, Val¹², Gln¹⁵, Lys¹⁶ and Phe¹⁹. Vitamin k3 interact with A β -42 with Ser⁸, Glu¹¹, Val¹², Gln¹⁵ and Phe¹⁹ via hydrophobic interactions. Further hydrogen bonding is involved between Lys¹⁶ of A β -42 and vitamin k3. From these results it can be inferred that Val¹², Gln¹⁵, Lys¹⁶ are common residues that are generally involved in the interaction of other inhibitors like Myricetin, EGCG, Curcumin and Pyrazinamide which suggests that vitamin k3 may follow the same mechanism of action against amyloidogenesis^{55–57}. Recently, Sinha *et al.* reported that the molecular tweezer (CLR01) a Lys-specific synthetic compound prevented cytotoxicity and oligomerization of A β -42 through non-covalent interaction and their study advocates the crucial role of Lys16 and Lys28 in A β -42 aggregation⁵⁸.

Conclusion

To conclude, current study reports that vitamin k3 is a potent therapeutic molecule that inhibits the progression of HEWL and A β -42 aggregation process by exploration of various biophysical and imaging tools including ThT fluorescence, ANS binding, CD measurements, DLS and TEM. Vitamin k3 inhibits the aggregation process starting from very beginning of nucleation phase. The non fibrillar aggregates formed in the presence of vitamin k3 contain less β -sheet structures as obtained from CD results. Sparsely populated and smaller sized aggregates were formed in presence of vitamin k3 as revealed by TEM and DLS results. Further, vitamin k3 which seems to be therapeutic via conventional biophysical and imaging techniques also imparts beneficial effects in reducing cytotoxicity in human neuronal cell line. Hydrophobic interactions are key forces involved between HEWL and vitamin k3 whereas in case of A β -42 vitamin k3 complex both hydrogen bonding as well as hydrophobic interactions are involved, implicated to play a dominating role in the inhibiting activity. The future projections of this study are to investigate the modulatory role of not only vitamin k3 but also its related molecules in disease progression in systemic amyloidosis and Alzheimer disease mice models.

Methods

Materials. Hen egg white lysozyme (HEWL), A β -42 peptide, Thioflavin T (ThT), 1-anilino 8 naphthalene sulphate (ANS) and vitamin k3 were procured from Sigma Aldrich, India. All other reagents used were of analytical grade.

pH measurements. pH measurements were carried out using Mettler Toledo Seven Easy pH meter (model S20) which was routinely calibrated with standard buffers. The experiments were performed at the 20 mM pH 7.4 sodium phosphate buffer. All preparations used in the experiments were filtered through 0.45 μ m Millipore Millex-HV PVDF filter.

Sample preparation. A stock solution of HEWL was made in 20 mM phosphate buffer pH 7.4 and extensively dialyzed against the same buffer and concentration was determined using a UV-visible spectrophotometer (Perkin Elmer Lambda 25) $E_{cm}^{1\%} = 12.4$ at 280 nm. For preparation of amyloid, 100 μ M of HEWL was used and samples were incubated at 65 °C in the presence of 50 mM NaCl for 120 hours in a circulating shaking water bath. Aliquots were taken from the stock at different time intervals. The A β -42 was prepared as described in literature⁵⁹. For preparation amyloid 100 μ M of A β -42 was kept at 37 °C and incubated for 70 hours under constant stirring in absence and presence of 50 and 100 μ M of vitamin k3. For further analysis, aliquots were taken from each set at different time intervals.

ThT fluorescence spectroscopy measurements. ThT fluorescence assay was performed with Shimadzu fluorescence spectrophotometer (RF-5301 PC). ThT stock solution was prepared in double distilled water and filtered with 0.2 micron millipore filter. HEWL samples incubated with or without vitamin k3, from each set samples were withdrawn at definite interval of time and mixed with ThT to accomplish final protein and dye concentration of 15 μ M. A β -42 samples incubated with or without vitamin k3, from each set samples were withdrawn at definite interval of time and mixed with ThT to accomplish final protein and dye concentration of 30 μ M. Samples were incubated in dark for 30 minutes. The ThT was excited at 440 nm and spectra were recorded from 450 to 600 nm. The excitation and emission slit widths were set at 5 and 10 nm, respectively. Sodium phosphate buffer 20 mM (pH 7.4) was used for dilution and spectra were corrected from respective blanks. All measurements were performed in triplicates. All curves were fitted as we described previously⁴³.

Congo red binding assay. Congo red was dissolved in a 20 mM phosphate buffer (pH 7.4) consisting of 50 mM NaCl and filtered through 0.45 μ m membrane filter. The concentration was determined using ϵ_M 45,000 $M^{-1} cm^{-1}$ at 498 nm. The HEWL and A β -42 concentration were fixed at 15 and 30 μ M respectively. CR and protein were mixed at a molar ratio of 1:1 in the absence and presence of vitamin k3 and kept for 30 minutes. The absorbance spectra (400–650 nm) of the samples were recorded with a UV-Visible spectrophotometer (Perkin Elmer Lambda 25) in a 1 cm path length cuvette.

ANS fluorescence measurements. HEWL samples (15 μ M) in absence and presence of vitamin k3 were mixed with 50 fold molar excess of ANS and then mixtures were kept in dark for 30 minutes at room temperature. A β -42 (30 μ M) sample in absence and presence of vitamin k3 was mixed with 50 fold molar excess of ANS and then mixtures were kept in dark for 30 minutes at room temperature. ANS fluorescence intensities were recorded with excitation at 380 nm and emission between 400 to 600 nm on a fluorescence spectrophotometer (RF-5301 PC). Excitation and emission slit width were set at 5 nm. All measurements were performed in triplicates.

Far-UV circular dichroism measurements. The circular dichroic measurements were performed on a JASCO spectropolarimeter (J-815) with a thermostatically controlled cell holder attached to a peltier with multitech water circulator. The experiments were carried out with HEWL (15 μ M) and A β -42 (30 μ M) in absence and presence of vitamin k3. Spectra were scanned in the range of 200–250 nm in a cuvette of 0.1 cm path length with scanning speed of 100 nm/min. Each spectrum was an average of three scans.

Dynamic light scattering (DLS) measurements. The change in aggregation behaviour of HEWL and A β -42 in the presence of varying concentration of vitamin k3 were determined using DLS. The hydrodynamic radii (R_h), measurements were done using a protein concentration of 10 μ M at 830 nm on a DynaPro-TC-04 dynamic light scattering instrument (Protein Solutions, Wyatt Technology, Santa Barbara, CA) equipped with a temperature controlled microsampler. All solutions were filtered through a 0.22 μ M pore sized micro filter (Whatman International, Maidstone, UK). The measured hydrodynamic radius (R_h) was the average of 50 measurements. The mean R_h and polydispersity (Pd) were estimated, on the basis of an autocorrelation analysis of scattered light intensity based on the translational diffusion coefficient, from the Stokes–Einstein equation⁶⁰

$$R_h = \frac{kT}{6\pi\eta D_W^{25^\circ C}} \quad (1)$$

where R_h is the hydrodynamic radius (nm), k is the Boltzmann's constant, T is the absolute temperature (K), η is the viscosity of water and $D_W^{25^\circ C}$ is the translational diffusion coefficient. All the samples were incubated for 120 hour at 65 $^\circ$ C prior to measurements.

Transmission electron microscopy (TEM). TEM images were taken on Philips CM-10 transmission electron microscope operating at an accelerating voltage of 200 kV. The amyloid fibril formation was assessed by applying 6 μ L of HEWL (15 μ M) and A β -42 (30 μ M) in absence and presence of vitamin k3 on 200-mesh copper grid covered by carbon-stabilized formvar film. Excess of fluid was removed after 2 min and the grids were then negatively stained with 2% (w/v) uranyl acetate. Images were viewed at 10,000X.

Cell culture. SH-SY5Y (human neuroblastoma cell line) cells were cultured in DMEM medium in humidified 5% (v/v) CO₂/air at 37 $^\circ$ C in 10% (v/v) (fetal bovine serum) FBS and 100 U/ml penicillin.

Cell viability assay. MTT (3, (4, 5-dimethylthiazol-2-yl) 2, 5-diphenyltetrazolium bromide) reduction assay was used to measure the cell viabilities of SH-SY5Y. MTT, in the presence of viable cells reduce to form blue formazon crystals, toxicity leads to inhibition of formazon production²⁶. For the MTT reduction assays, sample solutions of HEWL and A β -42 in absence and presence of vitamin k3 were added to the SH-SY5Y cells in the 96-well plates. Cells were seeded at 5,000 cells/well on 96-well plates and incubated for 24 hours before the treatment. The HEWL and A β -42 sample solutions (incubated over a period of 120 hours and 70 hours at 65 $^\circ$ C and 37 $^\circ$ C respectively) were incubated with SH-SY5Y cells for 24 hours, and MTT reduction was performed. MTT was added to the culture medium to yield a final concentration of 0.5 mg/ml and incubated for 4 hours at 37 $^\circ$ C in CO₂ incubator then removed supernatant carefully, 200 μ L of DMSO was added and mixed. After 20 hours of incubation in a humidified CO₂ incubator, the absorbance at 585 nm was read using a Micro plate absorbance reader (Bio-Rad Instruments, iMarkTM). Cell viability was compared to control cells without prior exposure to the fibril solutions.

Molecular docking study of HEWL-vitamin k3 interaction. The molecular docking study was performed using Autodock 4.2 and Autodock tools (ADT) using Lamarckian genetic algorithm. The crystal structure of HEWL (PDB id: 2LYZ) and A β -42 (PDB id: 1IYT) were obtained from Brookhaven Protein Data Bank. Three dimensional structure of vitamin k3 (CID: 4055) was obtained from PubChem. Water molecules, ions were removed and all hydrogen atoms were added. Then partial Kollman charges were assigned to protein. The protein was set to be rigid and there was no consideration of solvent molecules on docking. The grid size was set to be 116, 116 and 116 along X, Y and Z axes with 0.522 Å grid spacing for HEWL- vitamin k3. But in case of A β -42- vitamin k3 grid size was set to be 60, 60 and 60 along X, Y and Z axes with 0.426 Å grid spacing. Auto dock parameters used were GA population size: 150 and maximum number of energy evolutions: 250,000. 10 best solution based on docking score was retained for further analysis, Discovery studio 3.5 were used for visualization and for the identification of residues involved in binding.

Statistical analysis. All data were presented as mean \pm standard deviation from 3 independent determinations. The statistical analysis was made by performing one-way ANOVA for 3 independent determinations. Significance of results was determined as $p \leq 0.01$, unless otherwise stated.

References

- Huang, Y. & Mucke, L. Alzheimer mechanisms and therapeutic strategies. *Cell* **148**, 1204–1222 (2012).
- Stout, J. C. *et al.* Neurocognitive signs in prodromal Huntington disease. *Neuropsychology* **25**, 1 (2011).
- DuBay, K. F. *et al.* Prediction of the absolute aggregation rates of amyloidogenic polypeptide chains. *Journal of molecular biology* **341**, 1317–1326 (2004).
- Knowles, T. P. J., Vendruscolo, M. & Dobson, C. M. The amyloid state and its association with protein misfolding diseases. *Nature Reviews Molecular Cell Biology* **15**, 384–396 (2014).
- Marshall, K. E. *et al.* Hydrophobic, aromatic, and electrostatic interactions play a central role in amyloid fibril formation and stability. *Biochemistry* **50**, 2061–2071 (2011).
- Swaminathan, R., Ravi, V. K., Kumar, S., Kumar, M. V. S. & Chandra, N. Lysozyme: a model protein for amyloid research. *Adv Protein Chem Struct Biol* **84**, 63–111 (2011).

7. Karran, E., Mercken, M. & De Strooper, B. The amyloid cascade hypothesis for Alzheimer's disease: an appraisal for the development of the therapeutics. *Nature Reviews Drug Discovery* **10**, 698–712 (2011).
8. Soldi, G., Plakoutsi, G., Taddei, N. & Chiti, F. Stabilization of a native protein mediated by ligand binding inhibits amyloid formation independently of the aggregation pathway. *Journal of medicinal chemistry* **49**, 6057–6064 (2006).
9. Jayamani, J. & Shanmugam, G. Gallic acid, one of the components in many plant tissues, is a potential inhibitor for insulin amyloid fibril formation. *European journal of medicinal chemistry* **85**, 352–358 (2014).
10. Xu, S. *et al.* Rosiglitazone prevents the memory deficits induced by amyloid-beta oligomers via inhibition of inflammatory responses. *Neuroscience letters* **578**, 7–11 (2014).
11. Takasaki, J. *et al.* Vitamin A has anti-oligomerization effects on amyloid-beta *in vitro*. *Journal of Alzheimer's Disease* **27**, 271–280 (2011).
12. Valensin, D., Gabbiani, C. & Messori, L. Metal compounds as inhibitors of Beta-amyloid aggregation. Perspectives for an innovative metalloterapeutics on Alzheimer's disease. *Coordination Chemistry Reviews* **256**, 2357–2366 (2012).
13. Nedumpully-Govindan, P., Yang, Y., Andorfer, R., Cao, W. & Ding, F. Promotion or Inhibition of IAPP Aggregation by Zinc Coordination Depends on Its Relative Concentration. **54**, 7335–7344 (2015).
14. Nedumpully-Govindan, P. *et al.* Stabilizing Off-pathway Oligomers by Polyphenol Nanoassemblies for IAPP Aggregation Inhibition. *Scientific reports* **6**, 19463 (2016).
15. Nedumpully-Govindan, P. *et al.* Graphene oxide inhibits hIAPP amyloid fibrillation and toxicity in insulin-producing NIT-1 cells. *Physical Chemistry Chemical Physics* **18**, 94–100 (2016).
16. Mirsadeghi, S., Shanehazzadeh, S., Atyabi, F. & Dinarvand, R. Effect of PEGylated superparamagnetic iron oxide nanoparticles (SPIONs) under magnetic field on amyloid beta fibrillation process. *Materials Science and Engineering: C* **59**, 390–397 (2016).
17. Tayeb, H. O., Yang, H. D., Price, B. H. & Tarazi, F. I. Pharmacotherapies for Alzheimer's disease: beyond cholinesterase inhibitors. *Pharmacology & therapeutics* **134**, 8–25 (2012).
18. Lee, V., Rekhi, E., Hoh Kam, J. & Jeffery, G. Vitamin D rejuvenates aging eyes by reducing inflammation, clearing amyloid beta and improving visual function. *Neurobiology of aging* **33**, 2382–2389 (2012).
19. Annweiler, C. d., Brugg, B., Peyrin, J.-M., Bartha, R. & Beauchet, O. Combination of memantine and vitamin D prevents axon degeneration induced by amyloid-beta and glutamate. *Neurobiology of aging* **35**, 331–335 (2014).
20. Huy, P. D. Q. *et al.* In silico and *in vitro* characterization of anti-amyloidogenic activity of vitamin K3 analogues for Alzheimer's disease. *Biochimica et Biophysica Acta (BBA)-General Subjects* **1830**, 2960–2969 (2013).
21. Giraldo, E., Lloret, A., Fuchsberger, T. & Vina, J. Aβ and tau toxicities in Alzheimer's are linked via oxidative stress-induced p38 activation. Protective role of vitamin E. in *Redox Biology* Vol. 2, 873–877 (2014).
22. Truong, J. T. & Booth, S. L. Emerging issues in vitamin K research. *Journal of Evidence-Based Complementary & Alternative Medicine* **16**, 73–79 (2011).
23. McGuire, K., Jamison, J. M., Gilloteaux, J. & Summers, J. L. Synergistic Antitumor Activity of Vitamins C and K3 on Human Bladder Cancer Cell Lines. *Journal of Cancer Therapy* **4**, 7–19 (2013).
24. Checker, R. *et al.* Vitamin K3 suppressed inflammatory and immune responses in a redox-dependent manner. *Free radical research* **45**, 975–985 (2011).
25. Chaturvedi, S. K., Khan, J. M., Siddiqi, M. K., Alam, P. & Khan, R. H. Comparative insight into surfactants mediated amyloidogenesis of lysozyme. *International journal of biological macromolecules* **83**, 315–325 (2016).
26. Khurana, R. *et al.* Mechanism of thioflavin T binding to amyloid fibrils. *Journal of structural biology* **151**, 229–238 (2005).
27. Biancalana, M. & Koide, S. Molecular mechanism of Thioflavin-T binding to amyloid fibrils. *Biochimica et Biophysica Acta (BBA)-Proteins and Proteomics* **1804**, 1405–1412 (2010).
28. Marino, L. *et al.* Ortho-methylated 3-hydroxypyridines hinder hen egg-white lysozyme fibrillogenesis. *Scientific reports* **5**, 22052 (2015).
29. Wang, N. *et al.* Epigallocatechin-3-gallate Inhibits Fibrillogenesis of Chicken Cystatin. *Journal of agricultural and food chemistry* **63**, 1347–1351 (2015).
30. Khan, J. M. *et al.* Protonation favors aggregation of lysozyme with SDS. *Soft matter* **10**, 2591–2599 (2014).
31. Ahsan, N., Mishra, S., Jain, M. K., Suroli, A. & Gupta, S. Curcumin Pyrazole and its derivative (N-(3-Nitrophenyl)pyrazole) Curcumin inhibit aggregation, disrupt fibrils and modulate toxicity of Wild type and Mutant alpha-Synuclein. *Scientific reports* **5**, 9862 (2015).
32. Alam, P., Rabbani, G., Badr, G., Badr, B. M. & Khan, R. H. The Surfactant-Induced Conformational and Activity Alterations in *Rhizopus niveus* Lipase. *Cell biochemistry and biophysics* **71**, 1199–1206 (2015).
33. Haq, S. K., Rasheedi, S. & Khan, R. H. Characterization of a partially folded intermediate of stem bromelain at low pH. *European Journal of Biochemistry* **269**, 47–52 (2002).
34. He, J. *et al.* Myricetin Prevents Fibrillogenesis of Hen Egg White Lysozyme. *Journal of agricultural and food chemistry* **62**, 9442–9449 (2014).
35. Zhou, W. *et al.* At low concentrations, 3, 4-dihydroxyphenylacetic acid (DOPAC) binds non-covalently to alpha-synuclein and prevents its fibrillation. *Journal of molecular biology* **388**, 597–610 (2009).
36. Alam, P. *et al.* Biophysical and molecular docking insight into the interaction of cytosine beta-D arabinofuranoside with human serum albumin. *Journal of Luminescence* **164**, 123–130 (2015).
37. Khan, J. M. *et al.* SDS can be utilized as an amyloid inducer: a case study on diverse proteins. *PLoS one* **7**, e29694 (2012).
38. Shariatizi, S., Meratan, A. A., Ghasemi, A. & Nemat-Gorgani, M. Inhibition of amyloid fibrillation and cytotoxicity of lysozyme fibrillation products by polyphenols. *International journal of biological macromolecules* **80**, 95–106 (2015).
39. Louis-Jeune, C., Andrade-Navarro, M. A. & Perez-Iratxeta, C. Prediction of protein secondary structure from circular dichroism using theoretically derived spectra. *Proteins: Structure, Function, and Bioinformatics* **80**, 374–381 (2012).
40. Ghosh, S. *et al.* Copper (II) directs formation of toxic amorphous aggregates resulting in inhibition of hen egg white lysozyme fibrillation under alkaline salt-mediated conditions. *Journal of Biomolecular Structure and Dynamics* **33**, 991–1007 (2015).
41. Jayaram, D. T., Shankar, B. H. & Ramaiah, D. Effective Amyloid Defibrillation by Polyhydroxyl-Substituted Squaraine Dyes. *Chemistry- An Asian Journal* **10**, 2689–2694 (2015).
42. Choudhary, S., Kishore, N. & Hosur, R. V. Inhibition of insulin fibrillation by osmolytes: Mechanistic Insights. *Scientific reports* **5**, 17599 (2015).
43. Chaturvedi, S. K. *et al.* Biophysical insight into the anti-amyloidogenic behavior of taurine. *International journal of biological macromolecules* **80**, 375–384 (2015).
44. Hayden, E. Y. *et al.* Heme stabilization of alpha-Synuclein oligomers during amyloid fibril formation. *Biochemistry* **54**, 4599–4610 (2015).
45. Dal Piaz, F. *et al.* Natural iminosugar (+)-lentiginosine inhibits ATPase and chaperone activity of hsp90. **7**, 1–7 (2012).
46. Wu, C., Lei, H., Wang, Z., Zhang, W. & Duan, Y. Phenol red interacts with the protofibril-like oligomers of an amyloidogenic hexapeptide NFGAIL through both hydrophobic and aromatic contacts. *Biophysical journal* **91**, 3664–3672 (2006).
47. Alam, P., Abdelhameed, A. S., Rajpoot, R. K. & Khan, R. H. Interplay of multiple interaction forces: Binding of tyrosine kinase inhibitor nintedanib with human serum albumin. *Journal of Photochemistry and Photobiology B: Biology* **157**, 70–76 (2016).
48. Zhang, J. *et al.* Epigallocatechin-3-gallate (EGCG)-Stabilized Selenium Nanoparticles Coated with Tet-1 Peptide To Reduce Amyloid-beta Aggregation and Cytotoxicity. *ACS applied materials & interfaces* **6**, 8475–8487 (2014).

49. Ghahghaei, A., Bathaie, S., Kheirkhah, H. & Bahraminejad, E. The protective effect of crocin on the amyloid fibril formation of Amyloid beta-42 peptide *in vitro*. *Cellular and Molecular Biology Letters* **18**, 328–339 (2013).
50. Dai, X. *et al.* Chitosan Oligosaccharides Inhibit/Disaggregate Fibrils and Attenuate Amyloid beta-Mediated Neurotoxicity. *International journal of molecular sciences* **16**, 10526–10536 (2015).
51. Peng, J., Xiong, Y., Lin, Z., Sun, L. & Weng, J. Few-layer bismuth selenides exfoliated by hemin inhibit amyloid-beta 1–42 fibril formation. *Scientific reports* **5**, 10171 (2015).
52. Thapa, A., Jett, S. D. & Chi, E. Y. Curcumin Attenuates Amyloid beta Aggregate Toxicity and Modulates Amyloid beta Aggregation Pathway. *ACS chemical neuroscience* **7**, 56–58 (2015).
53. Loo, D. T. *et al.* Apoptosis is induced by beta-amyloid in cultured central nervous system neurons. *Proceedings of the National Academy of Sciences* **90**, 7951–7955 (1993).
54. Okayasu, H., Ishihara, M., Satoh, K. & Sakagami, H. Cytotoxic activity of vitamins K1, K2 and K3 against human oral tumor cell lines. *Anticancer research* **21**, 2387–2392 (2000).
55. Churches, Q. I. *et al.* Naturally occurring polyphenolic inhibitors of amyloid beta aggregation. *Bioorganic & medicinal chemistry letters* **24**, 3108–3112 (2014).
56. Chaturvedi, S. K. *et al.* Unraveling Comparative Anti-Amyloidogenic Behavior of Pyrazinamide and D-Cycloserine: A Mechanistic Biophysical Insight. *PLoS one* **10**, e0136528 (2015).
57. Rao, P. P. N., Mohamed, T., Teckwani, K. & Tin, G. Curcumin Binding to Beta Amyloid: A Computational Study. *Chemical biology & drug design* **130**, 813–820 (2015).
58. Sinha, S. *et al.* Lysine-specific molecular tweezers are broad-spectrum inhibitors of assembly and toxicity of amyloid proteins. *Journal of the American Chemical Society* **133**, 16958–16969 (2011).
59. Du, W.-J. *et al.* Brazilin inhibits amyloid [bgr]-protein fibrillogenesis, remodels amyloid fibrils and reduces amyloid cytotoxicity. *Scientific reports* **5**, 7992 (2015).
60. Edward, J. T. Molecular volumes and the Stokes-Einstein equation. *Journal of Chemical Education* **47**, 261 (1970).

Acknowledgements

Facilities provided by Interdisciplinary Biotechnology Unit, Aligarh Muslim University, Aligarh are gratefully acknowledged. P A and SKC are thankful to Council of Scientific and Industrial Research (CSIR), New Delhi, India for providing financial assistance. MKS is thankful to Department of Biotechnology (DBT), New Delhi, India for providing fellowship. We thank Mohd Akram (Max Planck Institute of Medical Research, Heidelberg, Germany) and Manu Shree (Indian Institute of technology, Mandi, India) for language editing.

Author Contributions

Designed the studies: P.A., S.K.C. and R.H.K. Undertook the experimental work: P.A., S.K.C., M.K.S., R.K.R., M.R.A., M.Z. and R.H.K. Analysed the data P.A., S.K.C., M.K.S. and R.H.K. Contributed to figure and manuscript preparation: P.A., S.K.C. and R.H.K.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Alam, P. *et al.* Vitamin k3 inhibits protein aggregation: Implication in the treatment of amyloid diseases. *Sci. Rep.* **6**, 26759; doi: 10.1038/srep26759 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>