

SCIENTIFIC REPORTS

OPEN

New Highly Oxygenated Germacranolides from *Carpesium divaricatum* and their Cytotoxic Activity

Received: 08 January 2016

Accepted: 13 May 2016

Published: 06 June 2016

Tao Zhang¹, Jin-Guang Si^{1,2}, Qiu-Bo Zhang¹, Gang Ding¹ & Zhong-Mei Zou¹

Eight highly oxygenated germacranolides (1–8) including four new ones (2–5) were isolated from the whole plant of *Carpesium divaricatum*. The planar structures and relative configurations of the new compounds were determined by NMR experiment and HRESIMS data. The absolute configuration of **1** was established by circular dichroism (CD) method and X-ray diffraction, and the stereochemistry of the new compounds 2–5 were determined by similar CD spectra with **1**. Compound **2** is the first hydroperoxyl germacrane from the genus *Carpesium*. The ¹³C NMR data of **1**, NMR data of 6–7, and their absolute configurations were reported for the first time. Two new compounds (**2** and **4**) and two known compounds (**6** and **8**) exhibited potent cytotoxicity against human cervical cancer (HeLa) cells, superior to that of the positive control doxorubicin.

Carpesium divaricatum Sieb. et Zucc, belonging to the genus *Carpesium* (Asteraceae), is widely distributed in China, traditionally used for the treatment of fevers, colds, bruises, and inflammatory diseases^{1–5}. The constituents of this plant have been previously investigated and shown to contain a number of sesquiterpenoids^{6–9}. Previous investigations indicate that sesquiterpene lactones possessing an α -methylene- γ -lactone moiety in the structure have cytotoxic activity to human cancer cells^{9–14}. Recently, *Carpesium* plants have attracted much attention due to eleven isolated sesquiterpene lactone dimers with novel skeletons displaying significantly cytotoxic activity^{15–18}. The parent nucleus of the germacrane contains a special ten member ring with different post-modification to produce diverse structural features. A survey of the literature has shown that a large number of germacranolides are isolated from the genus *Carpesium*, but their absolute configurations have rarely been reported^{16–8,12–13,19–23}.

In our ongoing search for new/novel and bioactive products from the medicinal plants in China, four new (2–5) and four known (1 and 6–8) germacranolides were isolated from the whole plant of *C. divaricatum*. In this paper, the structural elucidation including absolute configuration and bioactive evaluation of these compounds were present.

Results and Discussion

Compound **1** (Fig. 1) was identified as incaspitolide A (**1**)²⁴, by comparison of its MS, ¹H NMR data, as well as optical rotation data with reported data. However, its ¹³C NMR data have not been reported and absolute configuration has not been determined. The ¹³C NMR data was assigned by ¹H-¹H COSY and HSQC spectral data. The CD spectrum (Fig. 2) of **1** exhibited two positive Cotton effects at near 252 nm (α -methylene- γ -lactone region) and 294 nm (ketone n, π^* region), supporting 6S, 7S configuration¹². Fortunately, a suitable crystal was obtained for X-ray diffraction to confirm the absolute configuration. The X-ray crystallographic analysis [flack parameter: -0.02 (10)] established unambiguously the absolute configuration of **1** to be 4S, 5R, 6S, 7S, 8R and 10R (Fig. 3). Herein, the ¹³C NMR data and absolute configuration of **1** were reported for the first time.

Compound **2** was obtained as white needles. The molecular formula was assigned as C₂₄H₃₄O₁₀ on the basis of the positive-ion HRESIMS peak at m/z 505.2036 [M + Na]⁺, together with its ¹H and ¹³C NMR data (Tables 1 and 2). Its IR spectrum showed hydroxyl (3400 cm⁻¹) and carbonyl (1775 and 1729 cm⁻¹) absorptions. The ¹H and ¹³C NMR spectra of **2** showed an α -methylene- γ -lactone at δ_{H} 6.27 (1H, d, J = 2.0 Hz, Ha-13) and 5.96

¹Institute of Medicinal Plant Development, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100193, P.R. China. ²School of Pharmacy, Henan University of Traditional Chinese Medicine, Zhengzhou 450046, P.R. China. Correspondence and requests for materials should be addressed to Z.-M.Z. (email: zzmou@implad.ac.cn)




Figure 1. chemical structures of compounds 1–8.

(1H, d, $J = 2.0$ Hz, Hb-13), δ_C 134.7 (C-11), 127.4 (C-13) and 170.6 (C-12); three carbonyl carbons at δ_C 213.5 (C-9), 175.0 (C-1') and 167.1 (C-1''); an oxygenated quaternary carbons at δ_C 73.9 (C-4); five methines including three oxygenated ones at δ_H 4.75 (1H, d, $J = 6.5$ Hz, H-5), 4.63 (1H, dd, $J = 6.5, 2.0$ Hz, H-6), 3.87 (1H, dd, $J = 11.5, 2.0$ Hz, H-7), 4.92 (1H, d, $J = 11.0$ Hz, H-8) and 3.27 (1H, m, H-10), δ_C 79.4 (C-5), 73.3 (C-6), 46.3 (C-7), 79.8 (C-8), and 42.7 (C-10); and two methyl groups at δ_H 1.01 (3H, d, $J = 7.0$ Hz, CH₃-14), 1.19 (3H, s, CH₃-15). These signals (¹H and ¹³C NMR data) of **2** were similar to those of **8**¹² (Table S8.2, Supplementary




Figure 2. CD spectra of compounds 1–5.

Information), except for the ester residue at C-5. The singlet signals of 3'-Me and 4'-Me together with chemical shift difference of C-2' (δ 34.9 \rightarrow 84.8) implied that C-2' was an oxygenated quaternary carbon compared with **8**. Considering the chemical shift value of C-2' (δ 84.8) and molecular formula of **2** confirmed that a hydroperoxy moiety was attached at C-2'^{25–27}, which was further confirmed by the HRESIMS with fragment peaks at m/z 345.1645 [$M + 1 - H_2O - HOiBu - OOH$]⁺ and 245.1102 [$M + 1 - H_2O - HOiBu - OOH - HOAng$]⁺. The HMBC correlations from both of H₃-3' (δ_H 1.46, 3H, s) and H₃-4' (δ_H 1.42, 3H, s) to C-1' (δ_C 175.0) and C-2' (δ_C 84.8) allowed a reasonable connection of the hydroperoxy moiety to C-2'. The ¹H-¹H COSY spectrum (Fig. 4) showed two partial structure sequences for **2**: CH₂(3)CH₂(2)CH₂(1)CH(10)CH₃(14) and CH(5)CH(6)CH(7)CH(8). The C-C interconnectivity of all fragments was established from the HMBC spectrum (Fig. 4) as correlations of H-15 with C-3 and C-5, H-14 with C-1 and C-9, H-13 with C-7 and C-12, H-8 with C-1' (ester carbonyl of angeloyloxy group), and H-5 with C-1' (ester carbonyl of 2'-hydroperoxy-isobutyryloxy group). On the basis of these data, the planar structure of **2** was established.




Figure 3. X-ray ORTEP drawing of 1.

No.	1 ^a	2 ^a	3 ^a	4 ^a	5 ^a	6 ^a	7 ^a
1	1.68 o ^b , 1.24 m	1.69 o, 1.24 m	1.69 o, 1.24 m	1.72 m, 1.24 m	2.37 m, 2.18 m	1.69 o, 1.24 m	1.71 o, 1.23 m
2	1.54 m, 1.44 m	1.57 m, 1.46 m	1.56 m, 1.45 m	1.58 m, 1.45 m	5.98 m	1.55 m, 1.45 m	1.54 m, 1.45 m
3	1.68 o, 1.68 o	1.69 o, 1.69 o	1.69 o, 1.69 o	1.69 o, 1.69 o	5.63 br d (17.0)	1.69 o, 1.69 o	1.71 o, 1.71 o
5	4.67 d (6.0)	4.75 d (6.5)	4.70 d (6.5)	4.69 d (6.0)	4.68 d (8.5)	4.68 d (6.0)	4.69 d (6.0)
6	4.58 dd (6.5, 2.0)	4.63 dd (6.5, 2.0)	4.62 dd (6.5, 1.5)	4.62 dd (6.0, 2.0)	4.40 br d (8.5)	4.59 dd (6.5, 2.0)	4.60 dd (6.0, 1.5)
7	3.79 dd (11.5, 2.0)	3.87 dd (11.5, 2.0)	3.87 dd (11.5, 1.5)	3.88 dd (11.5, 2.0)	3.64 dd (10.0, 1.5)	3.78 dd (11.5, 2.0)	3.81 dd (11.5, 1.5)
8	4.76 d (11.5)	4.92 d (11.0)	4.91 d (11.5)	4.92 d (11.0)	4.74 d (10.5)	4.84 d (11.5)	4.80 d (11.5)
10	3.24 m	3.27 m	3.27 m	3.29 m	3.43 m	3.24 m	3.25 m
13a	6.30 d (1.5)	6.27 d (2.0)	6.27 d (2.0)	6.27 d (1.5)	6.27 d (1.5)	6.29 d (2.0)	6.32 d (2.0)
13b	6.03 d (1.5)	5.96 d (2.0)	5.97 d (2.0)	5.98 d (1.5)	5.88 d (1.5)	6.03 d (2.0)	6.05 d (2.0)
14	1.00 d (6.5)	1.01 d (7.0)	1.01 d (7.0)	1.00 d (6.5)	1.07 d (6.5)	1.03 d (6.5)	1.04 d (7.0)
15	1.13 s	1.19 s	1.15 s	1.15 s	1.24 s	1.14 s	1.15 s
2'	2.69 o			2.68 m	2.67 m	2.67 m	2.68 m
3'	1.18 d (7.0)	1.46 s	1.45 s	1.19 d (7.0)	1.20 d (7.0)	1.19 d (7.0)	1.19 d (7.0)
4'	1.17 d (7.0)	1.42 s	1.41 s	1.17 d (7.0)	1.17 d (7.0)	1.17 d (7.0)	1.17 d (7.0)
2''	2.69 o					2.36 o, 2.36 o	2.53 m
3''	1.22 d (7.0)	6.30 q (6.5)	6.31 q (6.5)	6.31 dq (3.0, 1.0), 5.83 dq (3.0, 1.0)	6.28 q (6.5)	2.09 m	1.71 m, 1.54 o
4''	1.15 d (7.0)	2.00 br s	2.00 br s	1.99 br s	1.95 br s	0.98 d (6.5)	1.19 d (7.0)
5''		1.98 dq (6.5, 1.5)	1.99 dq (6.5, 1.5)		1.97 dq (6.5, 1.5)	0.97 d (7.0)	0.94 t (9.0)

Table 1. ¹H NMR spectral data for compounds 1–7 (*J* in Hz within parentheses). ^aMeasured at 500 MHz in CD₃OD. ^bOverlapped with other signals.

No.	1 ^a	2 ^a	3 ^a	4 ^a	5 ^a	6 ^a	7 ^a
1	22.8	22.8	22.8	22.9	36.5	22.9	22.7
2	37.8	37.6	37.5	37.7	131.0	37.7	37.8
3	34.5	34.5	34.5	34.4	130.9	34.5	34.7
4	73.8	73.9	74.0	73.8	75.5	73.8	73.8
5	78.6	79.4	79.3	78.7	78.5	78.6	78.4
6	73.2	73.3	73.7	73.1	73.4	73.2	73.3
7	46.2	46.3	46.2	46.3	46.0	46.2	46.2
8	80.0	79.8	79.8	80.3	79.9	80.0	79.8
9	213.5	213.5	213.5	213.3	210.7	213.5	213.5
10	42.4	42.7	42.7	42.7	44.4	42.5	42.3
11	134.7	134.7	134.7	134.9	134.9	134.8	134.6
12	170.7	170.6	170.6	170.7	170.8	170.7	170.7
13	127.6	127.4	127.4	127.4	127.4	127.6	127.7
14	20.9	20.9	20.9	20.8	18.6	21.0	21.0
15	24.8	24.8	24.8	24.8	25.9	24.8	24.9
1'	178.7	175.0	177.2	178.7	178.6	178.7	178.7
2'	34.9	84.8	73.2	34.9	35.1	34.9	34.9
3'	19.4	23.1	27.2	19.2	19.3	19.2	19.2
4'	19.2	22.7	27.7	19.2	19.4	19.2	19.2
1''	177.0	167.1	167.1	166.9	167.4	173.0	176.6
2''	35.1	127.6	127.6	136.8	127.6	43.6	42.2
3''	19.2	143.1	143.1	128.2	142.8	26.5	27.7
4''	19.1	20.7	20.7	18.4	20.6	22.7	16.8
5''		16.1	16.1		16.1	22.7	12.0

Table 2. ¹³C NMR spectral data for compounds 1–7. ^aMeasured at 125MHz in CD₃OD.

The relative configuration of **2** was determined by analysis of ROESY data. The key NOE correlations of H-8/H-6, H-7/H-10, H-7/H-5 and H-5/H₃-15 indicated that **2** had the same relative configuration as **1**. The CD spectrum of **2** showed two positive Cotton effects at near 252 and 294 nm, which closely resembled those of **1**. Similar ROESY and CD data of **2** and **1** assigned the absolute configuration of **2** as 4*S*, 5*R*, 6*S*, 7*S*, 8*R* and 10*R*. Thus, the structure of compound **2** was defined as shown, named divarolide A.

Compounds **3–4** possessed molecular formulas of C₂₄H₃₄O₉ and C₂₃H₃₂O₈, from their HRESIMS at *m/z* 489.2108 [M + Na]⁺, and *m/z* 459.1971 [M + Na]⁺, respectively. The ¹H and ¹³C NMR data of **3–4** were similar to those of incaspitolide A (**1**)²⁴, except that the 2'-hydroxy-isobutyryloxy group at C-5 and the angeloyloxy group at C-8 in **3** were observed in place of two isobutyryloxy groups in **1**, and an isobutyryloxy group at C-8 in **1** was replaced by the 2-methylacryloyl group in **4**, respectively. These observations were confirmed by analyses of relevant ¹H-¹H COSY, HSQC and HMBC data (Fig. 4). The relative configurations of **3–4** were determined to be the same as that of **1** by comparison of ROESY data for relevant protons. Similar CD data of **3–4** and **1** revealed the same absolute configurations of **3–4** as that of **1**. Thus, the structures of compounds **3–4** were established as shown, named divarolide B and divarolide C, respectively.

The molecular formula of compound **5** was assigned as C₂₄H₃₂O₈ by HRESIMS (471.1988 [M + Na]⁺). A comparison of the NMR data of **5** with those of **8** suggested that both of them had the same substituted groups at C-5 and C-8, but that the two mutually coupled methylene units (C-2–C-3) in **8** were oxidized to an olefin moiety in **5**. The C-2/C-3 double bond was assigned *E*-geometry on the basis of the large coupling constant observed for olefinic protons (17.0 Hz). The H-¹H COSY, HSQC and HMBC spectra (Fig. 4) of **5** confirmed this observation, leading to the assignment of its planar structure. The relative and absolute configurations of **5** were deduced to be the same as those of **1**, on the basis of similar ROESY and CD data. Thus, the structure of compound **5** was elucidated as shown, named divarolide D.

Compounds **6–7** shared the same molecular formula C₂₄H₃₆O₈, established from their HRESIMS at *m/z* 475.2317 [M + Na]⁺ and *m/z* 475.2305 [M + Na]⁺. The ¹H and ¹³C NMR data of **6–7** showed a great similarity with those of **1**, except for the ester residues at C-8. The isobutyryloxy group at C-8 in **1** was placed by a 3-methylbutyryloxy group in **6** and the 2-methylbutyryloxy group of **7**, respectively. Compounds **6–7** have been reported as a mixture from *Inula cuspidata*²⁴. Actually, the exact linkage sites of the substituted groups have not been determined in the previous report and the authors speculate the mixture may contain two pairs of mixtures (incaspitolide B and C). Although the isolation of **6–7** is a huge challenge as they are highly oxygenated and similar, both of them were separated successfully in the present paper. Similarly, their relative and absolute configurations were determined as same as those of **1** by comparison of the ROESY and CD data. Thus, the structures of compounds **6–7** were established as shown, named incaspitolide B₁ and incaspitolide B₂, respectively.

Compound **8** was a known analogue of **1–7**, identified as (4*S*, 5*R*, 6*S*, 7*S*, 8*R*, 10*R*)-8-angeloyloxy-4-hydroxy-5-isobutyryloxy-9-oxo-germacran-7, 12-olide, by comparison of its MS, NMR and optical rotation data with reported data¹².




Figure 4. Key ^1H - ^1H COSY and HMBC correlations of compounds 2–5.

Compounds **1**, **2**, **4** and **6–8** were obtained in sufficient amounts to be evaluated for their cytotoxic activity against human cervical cancer (HeLa), hepatocellular cancer (Hep G2), stomach cancer (MGC-803), and lung cancer (A549) cell lines. All evaluated compounds exhibited strong cytotoxicity against HeLa (IC_{50} values of 4.36, 0.83, 1.18, 0.57, 3.58 and 1.70 μM), Hep G2 (IC_{50} values of 6.41, 8.40, 14.20, 18.10, 9.55 and 8.28 μM), and MGC-803 (IC_{50} values of 4.63, 4.48, 2.93, 3.49, 4.63 and 2.70 μM) cell lines, but only compounds **2**, **4**, **6** and **8** had IC_{50} values of 0.83, 1.18, 0.57 and 1.70 μM against HeLa cell lines, superior to that of the positive control doxorubicin (IC_{50} value 2.21 μM). Besides, new compound **2** also displayed strong cytotoxicity against A549 with IC_{50} value of 8.93 μM (the positive control doxorubicin showed IC_{50} value of 4.18 μM).

In conclusion, eight highly oxygenated germacranolides including four new ones (**2–5**) were isolated from the whole plant of *C. divaricatum*. To the best of our knowledge, this is the first report of hydroperoxyl germacranolide from the genus *Carpesium*. New compounds **2** and **4**, as well as known compounds **6** and **8**, exhibited potent cytotoxicity against HeLa cell lines, superior to that of the positive control doxorubicin. These findings are an important addition to the present knowledge on the structurally diverse and biologically important germacranolide family.

Methods

General Experimental Procedures. Optical rotations were measured on a Perkin-Elmer 241 polarimeter (Perkin-Elmer, Waltham, MA, USA) and UV spectra were recorded on Shimadzu UV-2501 PC (Shimadzu, Kyoto, Japan). IR data were recorded using a Shimadzu FTIR-8400S spectrophotometer (Shimadzu, Kyoto, Japan). ^1H and ^{13}C -NMR data were acquired with Bruker 500 instruments (Bruker, Rheinstetten, Germany) using the solvent signals (CD_3OD : δ_{H} 4.87/ δ_{C} 49.0 ppm;) as references. HRESIMS data were acquired using Q-TOF analyzer in SYNAPT HDMS system (Waters, Milford, MA, USA). CD spectra were recorded on a JASCO J-815 Spectropolarimeter (Jasco, Tokyo, Japan).

X-ray diffraction data were collected on the Agilent GEMINITME instrument (CrysAlisPro software, Version 1.171.35.11; Agilent, Santa Clara, CA, USA). HPLC was performed using Waters 2535 system (Waters, Milford, MA, USA) with the following components: preparative column, a Daisogel-C₁₈-100A (10 μm , 30 \times 250 mm, ChuangXinTongHeng Sci.&Tech., Beijing, China) and a YMC-Pack ODS-A column (5 μm , 10 \times 250 mm, YMC, Kyoto, Japan); and detector, Waters 2489 UV. Sephadex LH-20 (40–70 μm , Pharmacia Biotech AB, Uppsala, Sweden), silica gel (60–100, 100–200, and 200–300 mesh) and silica gel GF254 sheets (0.20–0.25 mm) (Qingdao Marine Chemical Plant, Qingdao, China) were used for column chromatography and TLC, respectively. TLC spots were visualized under UV light and by dipping into 5% H_2SO_4 in EtOH followed by heating.

Plant Material. The whole plant of *C. divaricatum* were collected from EnShi, Sichuan province of China, in August, 2013. They were identified by Prof. Ben-Gang Zhang of Institute of Medicinal Plant Development. A voucher specimen (No. 20130828) was deposited in the Institute of Medicinal Plant Development, Chinese Academy of Medical Sciences and Peking Union Medical College (CAMS & PUMC), China.

Extraction and Isolation. The air-dried plants (9 kg) were extracted three times (7 days each time) with EtOH–H₂O (95:5) at room temperature. The combined extract was concentrated under reduced pressure to furnish a dark brown residue (570 g), which was suspended in H₂O and partitioned in turn with petroleum ether (bp 60–90 $^{\circ}\text{C}$), EtOAc, and *n*-BuOH. The EtOAc extract (207 g) was separated chromatographically on silica gel column (60–100 mesh, 16 \times 20 cm) with a gradient mixture of CH_2Cl_2 –MeOH (100:1, 60:1, 30:1, 15:1, and 6:1) as eluent. Five fractions (fraction A–E) were collected according to thin layer chromatography (TLC) analysis. Fraction A (CH_2Cl_2 –MeOH, 100:1, 140 g) was separated by silica gel column chromatography (CC) (100–200 mesh, 16 \times 20 cm) with petroleum ether–acetone (50:1, 25:1, 20:1, 15:1, 12:1, 10:1, 7:1, 5:1, 3:1 and 1:1) as eluent to give fraction A₁–A₁₁. Fraction A₇ (petroleum ether–acetone, 7:1, 8 g) was separated by Sephadex LH-20 CC

compounds	IC ₅₀ (μM)			
	HeLa	Hep G2	MGC-803	A549
1	4.36 ± 0.12	6.41 ± 0.23	4.63 ± 0.25	21.7 ± 1.41
2	0.83 ± 0.09	8.40 ± 0.84	4.48 ± 1.01	8.93 ± 1.73
4	1.18 ± 0.14	14.20 ± 0.21	2.93 ± 0.57	27.8 ± 2.34
6	0.57 ± 0.06	18.10 ± 0.72	3.49 ± 0.40	20.7 ± 0.84
7	3.58 ± 0.21	9.55 ± 0.27	4.63 ± 0.59	22.7 ± 0.94
8	1.70 ± 0.24	8.28 ± 0.64	2.70 ± 0.65	16.3 ± 1.41
doxorubicin	2.21 ± 0.18	5.40 ± 0.80	0.74 ± 0.05	4.18 ± 0.52

Table 3. Cytotoxicity of Compounds 1, 2, 4 and 6–8. Values were mean ± SD. Doxorubicin, positive control. Cell lines: HeLa: cervical cancer, Hep G2: hepatocellular cancer, MGC-803: stomach cancer, and A549: lung cancer.

(5 × 200 cm, MeOH) to give Fr.A₇S₁–Fr.A₇S₃. Fraction A₇S₂ (MeOH–H₂O, 5 g) was purified using preparative HPLC (Daisogel–C₁₈–100A, 10 μm; 250 × 30 mm; 20 mL/min, 70% MeOH in H₂O) to yield **8** (3.9 g) and a mixture of **1–7** (800 mg). The mixture of **1–7** (800 mg) was further purified using semipreparative HPLC (60–90% MeOH in H₂O for 40 min; 40–80% MeCN in H₂O for 40 min) to yield **1** (100 mg), **2** (5 mg), **3** (3.5 mg), **4** (8 mg), **5** (2.6 mg), **6** (30 mg) and **7** (25 mg).

Incasitolide A (**1**): white needles (CH₃OH), [α]_D²⁰ +57.7 (*c* 0.25, CHCl₃); UV (MeOH) λ_{max}(logε): 207 (4.35) nm, IR (neat) ν_{max}: 3544, 1776, 1746, 1720, 1666 cm⁻¹; CD (MeOH) 256 (Δε +0.09), 310 (Δε +0.22), 207 (Δε -0.44) nm; HRESIMS (pos.): *m/z* 461.2154 [M + Na]⁺ (calcd for C₂₃H₃₄O₈Na, 461.2151); ¹H NMR data see Table 1, ¹³C NMR data see Table 2.

Divarolide A (**2**): white needles (CH₃OH), [α]_D²⁰ +35.7 (*c* 0.2, CHCl₃); UV (MeOH) λ_{max}(logε): 214 (3.99) nm, IR (neat) ν_{max}: 3400, 1775, 1729, 1645 cm⁻¹; CD (MeOH) 250 (Δε +0.06), 310 (Δε +0.11), 209 (Δε -0.17) nm; HRESIMS (pos.): *m/z* 505.2036 [M + Na]⁺ (calcd for C₂₄H₃₄O₁₀Na, 505.2050); ¹H NMR data see Table 1, ¹³C NMR data see Table 2.

Divarolide B (**3**): white needles (CH₃OH), [α]_D²⁰ +121.0 (*c* 0.11, CHCl₃); UV (MeOH) λ_{max}(logε): 213 (4.22) nm, IR (neat) ν_{max}: 3500, 1768, 1726, 1643 cm⁻¹; CD (MeOH) 249 (Δε +0.12), 309 (Δε +0.21); HRESIMS (pos.): *m/z* 489.2108 [M + Na]⁺ (calcd for C₂₄H₃₄O₉Na, 489.2101); ¹H NMR data see Table 1, ¹³C NMR data see Table 2.

Divarolide C (**4**): white needles (CH₃OH), [α]_D²⁰ +48.7 (*c* 0.10, CHCl₃); UV (MeOH) λ_{max}(logε): 208 (3.93) nm, IR (neat) ν_{max}: 3508, 1776, 1728, 1665, 1636 cm⁻¹; CD (MeOH) 220 (Δε +0.07), 254 (Δε +0.08), 310 (Δε +0.15) nm; HRESIMS (pos.): *m/z* 459.1971 [M + Na]⁺ (calcd for C₂₃H₃₂O₈Na, 459.1995); ¹H NMR data see Table 1, ¹³C NMR data see Table 2.

Divarolide D (**5**): white needles (CH₃OH), [α]_D²⁰ +18.9 (*c* 0.25, CHCl₃); UV (MeOH) λ_{max}(logε): 210 (4.03) nm, IR (neat) ν_{max}: 3514, 1771, 1728, 1665, 1646 cm⁻¹; CD (MeOH) 255 (Δε +0.04), 309 (Δε +0.07), 205 (Δε -0.10) nm; HRESIMS (pos.): *m/z* 471.1988 [M + Na]⁺ (calcd for C₂₄H₃₂O₈Na, 471.1995); ¹H NMR data see Table 1, ¹³C NMR data see Table 2.

Incasitolide B₁ (**6**): white needles (CH₃OH), [α]_D²⁰ +123.1 (*c* 0.15, CHCl₃); UV (MeOH) λ_{max}(logε): 210 (3.83) nm, IR (neat) ν_{max}: 3529, 1777, 1746, 1721, 1665 cm⁻¹; CD (MeOH) 256 (Δε +0.08), 310 (Δε +0.19), 207 (Δε -0.39) nm; HRESIMS (pos.): *m/z* 475.2317 [M + Na]⁺ (calcd for C₂₄H₃₆O₈Na, 475.2308); ¹H NMR data see Table 1, ¹³C NMR data see Table 2.

Incasitolide B₂ (**7**): white needles (CH₃OH), [α]_D²⁰ +31.2 (*c* 0.18, CHCl₃); UV (MeOH) λ_{max}(logε): 202 (3.77) nm, IR (neat) ν_{max}: 3532, 1781, 1746, 1719, 1667 cm⁻¹; CD (MeOH) 257 (Δε +0.06), 310 (Δε +0.17), 206 (Δε -0.34) nm; HRESIMS (pos.): *m/z* 475.2305 [M + Na]⁺ (calcd for C₂₄H₃₆O₈Na, 475.2308); ¹H NMR data see Table 1, ¹³C NMR data see Table 2.

X-ray crystal structure analysis. X-ray diffraction data were collected on the Agilent GEMINITME instrument (CrysAlisPro software, Version 1.171.35.11), with enhanced Cu Kα radiation (λ = 1.54184 Å). The structure was solved by direct methods and refined by full-matrix least-squares techniques (SHELXL-97). All non-hydrogen atoms were refined with anisotropic thermal parameters. Hydrogen atoms were located by geometrical calculations and from positions in the electron density maps. Crystallographic data (excluding structure factors) for **1** in this paper has been deposited with the Cambridge Crystallographic Data Centre (deposition number CCDC 1441395). Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (fax: +44 12 23336033 or e-mail: deposit@ccdc.cam.ac.uk).

A colorless triclinic crystal (0.50 × 0.50 × 0.40 mm) of **1** was obtained from CH₂Cl₂–MeOH (3:1). Crystal data: 3C₂₃H₃₄O₈·2H₂O (*M* = 459.86), *T* = 105.5 K, triclinic, space group *P*₁, *a* = 9.7574(3) Å, *b* = 10.9450(5) Å, *c* = 18.3652(8) Å, α = 102.359(4)°, β = 99.183(3)°, γ = 101.095(3)°, *V* = 1838.35(13) Å³, *Z* = 3, ρ = 1.246 mg/mm³, μ(Cu Kα) = 0.781 mm⁻¹, measured reflections = 24294, unique reflections = 12313 (*R*_{int} = 0.0202), largest difference peak/hole = 1.010/−0.331 e Å⁻³, and flack parameter = −0.02(10). The final *R* indexes [*I* > 2σ(*I*)] were *R*₁ = 0.0435, and *wR*₂ = 0.1176. The final *R* indexes (all data) were *R*₁ = 0.0448, and *wR*₂ = 0.1191. The goodness of fit on *F*² was 1.046.

Cytotoxicity assays. The assay was run in triplicate. In a 96-well plate, each well was plated with 2 × 10⁴ cells. After cell attachment overnight, the medium was removed, and each well was treated with 100 μL of medium

containing 0.1% DMSO or different concentrations of the test compounds and the positive control doxorubicin. The plate was incubated for 4 days at 37 °C in a humidified, 5% CO₂ atmosphere. Cytotoxicity was determined using a modified 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) colorimetric assay²⁸. After addition of 10 μL MTT solution (5 mg/mL), cells were incubated at 37 °C for 4 h. After adding 150 μL DMSO, cells were shaken to mix thoroughly. The absorbance of each well was measured at 490 nm in a Multiscan photometer. The IC₅₀ values were calculated by SPSS software and listed in Table 3.

References

1. Editorial board of Chinese Materia Medica. *Chinese Materia Medica*, Vol. 21 (eds Hu, X. M. *et al.*) 7761 (Shanghai Science & Technology Press, Shanghai (1999)).
2. Kim, E. J. *et al.* Suppression by a sesquiterpene lactone from *Carpesium divaricatum* of inducible nitric oxide synthase by inhibiting nuclear factor-κB activation. *Biochem. Pharmacol.* **61**, 903–910 (2001).
3. Zee, O. P. *et al.* A new cytotoxic acyclic diterpene from *Carpesium divaricatum*. *Arch. Pharm. Res.* **22**, 225–227 (1999).
4. Zee, O. P., Kim, D. K. & Lee, K. R. Thymol derivatives from *Carpesium divaricatum*. *Arch. Pharm. Res.* **21**, 618–620 (1998).
5. Chung, I. M. *et al.* Antiplasmodial activity of isolated compounds from *Carpesium divaricatum*. *Phytother. Res.* **24**, 451–453 (2010).
6. Kim, D. K. *et al.* Four new cytotoxic germacranolides from *Carpesium divaricatum*. *J. Nat. Prod.* **60**, 1199–1202 (1997).
7. Maruyama, M. Sesquiterpene lactones from *Carpesium divaricatum*. *Phytochemistry*. **29**, 547–550 (1990).
8. Kim, D. K., Lee, K. R. & Zee, O. P. Sesquiterpene lactones from *Carpesium divaricatum*. *Phytochemistry*. **46**, 1245–1247 (1997).
9. Xie, W. D. *et al.* Sesquiterpenoids from *Carpesium divaricatum* and their cytotoxic activity. *Fitoterapia*. **83**, 1351–1355 (2012).
10. Zhang, J. P. *et al.* The genus *Carpesium*: A review of its ethnopharmacology, phytochemistry and pharmacology. *J. Ethnopharmacol.* **163**, 173–191 (2015).
11. Yang, C., Yuan, C. S. & Jia, Z. J. Xanthanolides, germacranolides, and other constituents from *Carpesium longifolium*. *J. Nat. Prod.* **66**, 1554–1557 (2003).
12. Gao, X., Lin, C. J. & Jia, Z. J. Cytotoxic germacranolides and acyclic diterpenoids from the seeds of *Carpesium triste*. *J. Nat. Prod.* **70**, 830–834 (2007).
13. Lin, Y. L. & Ou, J. C. Napalolides A–D, four new sesquiterpene lactones from *Carpesium nepalense*. *J. Nat. Prod.* **59**, 991–993 (1996).
14. Li, X. W. *et al.* Antiproliferative and apoptotic sesquiterpene lactones from *Carpesium faberi*. *Bio. Med. Chem. Lett.* **21**, 366–372 (2011).
15. Yang, Y. X. *et al.* Carpedilactones A–D, four new isomeric sesquiterpene lactone dimers with potent cytotoxicity from *Carpesium faberi*. *Org. Lett.* **16**, 4216–4219 (2014).
16. Wu, J. W. *et al.* Dicarabrones A and B, a pair of new epimers dimerized from sesquiterpene lactones via a [3 + 2] cycloaddition from *Carpesium abrotanoides*. *Org. Lett.* **17**, 1656–1659 (2015).
17. Wang, J. F. *et al.* Dicarabrol, a new dimeric sesquiterpene from *Carpesium abrotanoides* L. *Bioorg. Med. Chem. Lett.* **25**, 4082–4084 (2015).
18. Xu, X. K. *et al.* Four new isomeric sesquiterpene lactone dimers from *Carpesium faberi*. *Tetrahedron Lett.* **56**, 6381–6384 (2015).
19. Lee, H. J. *et al.* A germacranolide sesquiterpene lactone suppressed inducible nitric oxide synthase by downregulating NF-κB activity. *Can. J. Physiol. Pharmacol.* **89**, 232–237 (2011).
20. Moon, H. I. & Zee, O. Antiproliferative effect from sesquiterpene lactones of *Carpesium rosulatum* MIQ consumed in South Korea on the five human cancer cell Lines. *Rec. Nat. Prod.* **4**, 3149–3155 (2010).
21. Moon, H. I. & Zee, O. Sesquiterpene lactones from *Carpesium rosulatum* with potential cytotoxicity against five human cancer cell lines. *Hum. Exp. Toxicol.* **30**, 1083–1087 (2010).
22. Kim, M. R. *et al.* Sesquiterpene lactones from *Carpesium triste* var. *manshuricum*. *Phytochemistry*. **52**, 113–115 (1999).
23. Kim, M. R. *et al.* Cytotoxic germacranolide sesquiterpene lactones from *Carpesium triste* var. *manshuricum*. *Arch. Pharm. Res.* **30**, 556–560 (2007).
24. Bohlmann, F., Singh, P. & Jakupovic, J. Further ineupatorolide-like germacranolides from *Inula cuspidata*. *Phytochemistry*. **21**, 157–160 (1982).
25. Hegazy, M. E. *et al.* Sesquiterpenes from an Egyptian herbal medicine, *Pulicaria undulate*, with inhibitory effects on nitric oxide production in RAW264.7 macrophage cells. *Chem. Pharm. Bull.* **60**, 363–370 (2012).
26. Rustaiyan, A. & Faramarzi, S. Sesquiterpene lactones from *Serratula latifolia*. *Phytochemistry*. **27**, 479–481 (1988).
27. Hegazy, M. E. *et al.* Rare hydroperoxyl guaianolide sesquiterpenes from *Pulicaria undulata*. *Phytochem. Lett.* **12**, 177–181 (2015).
28. Wang, B. J., Won, S. J., Yu, Z. R. & Su, C. L. Free radical scavenging and apoptotic effects of *Cordyceps sinensis* fractionated by supercritical carbon dioxide. *Food Chem. Toxicol.* **43**, 543–552 (2005).

Acknowledgements

This work was financially supported by the Chinese National S&T Special Project on Major New Drug Innovation (2013ZX09508104), the China Postdoctoral Science Foundation (2014M560915), Peking Union Medical College (PUMC) Postdoctoral Science Foundation, and Program for Innovative Research Team in Institute of Medicinal Plant Development (IMPLAD).

Author Contributions

Z.-M.Z. designed the study; T.Z. performed the experiments with the help of J.-G.S., Q.-B.Z. and G.D. The manuscript was prepared by T.Z. and Z.-M.Z. All authors discussed the results and their interpretation and commented on the manuscript at all stages.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhang, T. *et al.* New Highly Oxygenated Germacranolides from *Carpesium divaricatum* and their Cytotoxic Activity. *Sci. Rep.* **6**, 27237; doi: 10.1038/srep27237 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>