

SCIENTIFIC REPORTS



OPEN

A two-year field measurement of methane and nitrous oxide fluxes from rice paddies under contrasting climate conditions

Received: 08 February 2016

Accepted: 02 June 2016

Published: 20 June 2016

Huifeng Sun^{1,2}, Sheng Zhou^{1,2}, Zishi Fu^{1,2}, Guifa Chen^{1,2}, Guoyan Zou^{1,2} & Xiangfu Song^{1,2}

The effects of three irrigation levels (traditional normal amount of irrigation [NA100%], 70%, and 30% of the normal amount [NA70% and NA30%]) and two rice varieties (*Oryza sativa* L. Huayou14 and Hanyou8) on CH₄ and N₂O emissions were investigated over two years under contrasting climate conditions (a 'warm and dry' season in 2013 and a normal season in 2014). Hanyou8 was developed as a drought-resistant variety. The mean seasonal air temperature in 2013 was 2.3 °C higher than in 2014, while the amount of precipitation from transplanting to the grain-filling stage in 2013 was only 36% of that in 2014. CH₄ emission rose by 93–161%, but rice grain yield fell by 7–13% in 2013, compared to 2014 under the NA100% conditions. Surface standing water depths (SSWD) were higher in Hanyou8 than in Huayou14 due to the lower water demand by Hanyou8. A reduction in the amount of irrigation water applied can effectively reduce the CH₄ emissions regardless of the rice variety and climate condition. However, less irrigation during the 'warm and dry' season greatly decreased Huayou14 grain yield, but had little impact on Hanyou8. In contrast, N₂O emission depended more on fertilization and SSWD than on rice variety.

Modern changes in global climate, including greater atmospheric concentrations of greenhouse gases (CO₂, CH₄, N₂O, etc.), temperature shifts, higher frequency of extreme events, such as drought or heavy rainfall, and heat waves, are directly or indirectly associated with the combustion of fossil fuels, land use change and other human activities that have occurred since the Industrial Revolution¹. Such changes have greatly affected terrestrial ecosystems. For example, extreme events could lead to negative and even disastrous effects on plants².

Rice (*Oryza sativa* L.) is the most widely consumed staple food crop in the world, and is particularly important in Asia. The food safety and national security of some countries depends on its production. Rice cultivation is also a significant source of greenhouse gas emissions, primarily methane (CH₄) and nitrous oxide (N₂O). Annual CH₄ emissions from rice fields have been estimated to account for about 5–19% of global CH₄ emissions, and agricultural N₂O emissions increased by nearly 17% between 1990 and 2005, and now account for 60% of global anthropogenic N₂O emissions³. The flooded environment, created during rice cultivation, provides anaerobic conditions that favor CH₄ production by methanogens. The resulting CH₄ can be oxidized by methanotrophs under aerobic conditions (e.g., in the rhizosphere and at the soil-water interface) and is finally emitted to the atmosphere through soil- or water-atmosphere interfaces and by the rice plant aerenchyma⁴. Nitrogen fertilization and water management (e.g. alternating wetting and drying) facilitates N₂O emission via the processes of nitrification and/or denitrification in rice paddies⁴, hence, when applied appropriately, fertilizer and management interventions can play important roles in effectively controlling CH₄ and N₂O emissions during rice cultivation^{5–9}. However, rice varieties differ significantly in terms of total CH₄ and N₂O emissions^{10–13}. Selecting a rice variety that results in low CH₄ and N₂O emissions may therefore be a promising way to mitigate greenhouse gas emissions from rice paddies.

Drought is a serious limiting factor on crop production and the most damaging stressor in modern agriculture^{14,15}. Rice consumes 70–90% of the total amount of irrigation water used in agriculture^{16,17}, and as roughly a half of the world's 158 million ha of rice land is paddy rice¹⁸, the production of this crop is very susceptible to

¹Eco-Environmental Protection Research Institute, Shanghai Academy of Agricultural Sciences, Shanghai 201403, China. ²Shanghai Engineering Research Center of Low-carbon Agriculture (SERCLA), Shanghai 201415, China. Correspondence and requests for materials should be addressed to S.Z. (email: zhous@outlook.com)

Rice season	Treatments	Mean seasonal air temperature (°C)	Mean seasonal soil temperature (°C)	Seasonal total precipitation (mm)	Precipitation from June to September (mm)	Amount of irrigation (mm)	Water input from June to September (mm)
2013 ('warm and dry')	NA100%	26.7	27.8	492.1	271.8	611.7	883.5
	NA70%	26.7	27.7	492.1	271.8	428.3	700.1
	NA30%	26.7	28.0	492.1	271.8	185.0	456.8
2014 (normal)	NA100%	24.4	24.7	762.7	762.7	116.7	879.4
	NA70%	24.4	24.7	762.7	762.7	81.7	844.4
	NA30%	24.4	24.8	762.7	762.7	35.0	797.7

Table 1. Mean seasonal air and soil temperatures, total precipitation, and amount of irrigation water in the 2013 and 2014 rice growing seasons. NA100% = normal amount of irrigation, NA70% = 70% of normal irrigation, NA30% = 30% of normal irrigation.

water stress¹⁹. A series of agricultural practices, such as the use of saturated soil cultures and alternate wetting and drying, are recommended to reduce water input from irrigation and to enhance water use efficiency in rice cultivation. These practices could greatly reduce the water input required. However, they also lower rice grain yields to a certain extent¹⁸. Breeding new varieties of rice with drought tolerance may be an effective way of sustainably addressing the water scarcity issue²⁰. These new varieties are called Water-saving and Drought-resistance Rice (WDR), and are characterized by having a similar yield potential and grain quality as wild type varieties, but require less water (50% less water use). In water-limited environments, they show higher drought resistance and minimize yield loss²⁰. However, to our knowledge, little is known about the CH₄ and N₂O emissions from rice paddies by WDR under different climate conditions.

The annual mean surface air temperature for China over the past 97 years has experienced a warming of 0.79 °C, with a warming rate of 0.08 °C/10a, which is slightly larger than the global or northern hemispheric average as given by IPCC Third Assessment Report²¹. In addition, in the experimental region, the mean air temperature during the rice growing season in the 1990s was 0.2 °C higher than that in the 1960–80s. However, the mean seasonal air temperature between 2001 and 2008 increased by 1.1 °C relative to that in the 1960–80s²². Climate models project that the increase in global surface air temperatures may exceed 1.5 °C by the end of the 21st century relative to the average between 1850 and 1900²³. Higher temperatures are known to influence water, ion, and organic solute movement across plant membranes, which interferes with photosynthesis and respiration. Excessive temperatures can reduce plant leaf photosynthesis and decrease the allocation of dry matter to the shoots and roots²⁴. Therefore, high temperatures may have caused rice grain yield reductions in many rice growing areas^{25–27}. Generally, weather patterns can be highly variable. For example, there may be little precipitation in one rice growing season and more in another. To satisfy the rice plant water demand, large amounts of water are used to irrigate rice fields in a low precipitation season, but less may be needed in a high precipitation season. Although some studies have demonstrated that CH₄ and N₂O emissions differed among rice growing seasons, little explanation has been provided for the variation^{6,9}. Investigating the mechanisms underlying the differences in CH₄ and N₂O emissions under different climate conditions (air temperature and precipitation) is important when accurately assessing the total greenhouse gas emission caused by rice cultivation under global warming scenarios. Some process-based models (e.g. the Denitrification and Decomposition (DNDC) model) have been used to estimate the effects of changes in air temperature and precipitation on CH₄ and N₂O emissions from arable soils^{28,29}. However, these results are not strongly supported by field experimental data. The field experiments examining the effects of climate change on CH₄ and N₂O emissions provide important information that may help improve long-term predictions with those models³⁰.

In a previous study, we found that CH₄ emissions fell significantly when less irrigation water was applied to rice paddies over two growing seasons³¹. In this study, the effects of two rice varieties (a WDR versus a common rice variety) and irrigation management technique on CH₄ and N₂O emissions from rice paddies were investigated over two growing seasons under contrasting climate conditions (a 'warm and dry' season and a normal season). In addition, the impacts of climate condition and irrigation management on rice grain yield and equivalent CO₂ (CO₂-eq) emission were also evaluated.

Results and Discussion

Irrigation management and SSWD dynamics under contrasting climate conditions. The total precipitation was 492.1 and 762.7 mm over the whole rice cultivation periods of 2013 and 2014, respectively (Table 1). In the 'warm and dry' rice season of 2013, only 271.8 mm (~55.2% of the total) of precipitation occurred from June to September, which is the period of active rice plant growth and development. In contrast, total precipitation in the normal rice season of 2014 occurred between June and September. Additionally, the total amount of evaporation was 591.6 and 481.6 mm in 2013 and 2014, respectively. Therefore, to satisfy the plant demand for water during growth and development, irrigation was applied up to 12 times in 2013 to keep the water table in the plots similar to traditional irrigation management (Fig. 1a). However, irrigation was applied only three times in 2014 (Fig. 1b). The total amount of irrigation in the NA100% plot was 611.7 mm in 2013 compared to only 116.7 mm in 2014 (Table 1). Consequently, the effective water inputs (precipitation + irrigation) from June to September for the NA100% plots were similar for the 'warm and dry' season (883.5 mm) and the normal season (879.4 mm). In contrast, the total water inputs in the NA70% and NA30% plots from June to September in 2013 were only 83% and 57% of those in 2014 (Table 1).




Figure 1. Seasonal changes in daily mean air temperature, daily precipitation, and soil temperature at 5 cm depth in the 2013 (a) and 2014 (b) rice growing seasons. Soil temperature values were obtained by averaging the values from all treatments during gas sampling events.

Similar trends in SSWD were observed for all treatments in both seasons when the rice plants were small before MD (Fig. 2). The maximum SSWD reached similar values (approximately 6–7 cm). However, there were considerable differences in SSWD after MD between the rice varieties in 2013 (Fig. 2a,b). Under the same water inputs (precipitation + irrigation), the SSWDs of Hanyou8 in the NA100% and NA70% plots were significantly higher than those of Huayou14. Hanyou8 was developed as a drought-resistant variety and requires less water to maintain growth and development²⁰. The average evapotranspiration rate for Hanyou8 was 5.4 mm day⁻¹, a significantly lower rate than for Huayou14 (6.2 mm day⁻¹), which probably resulted in higher SSWDs in the NA100% and NA70% plots when the WDR (Hanyou8) variety was grown compared to the common rice variety. However, the SSWD in the NA30% plots for both varieties was almost zero after MD in 2013 due to the reduced water input (Fig. 2a,b). In contrast, Hanyou8 mean SSWD was not significantly different to Huayou14 after MD in 2014. Furthermore, the SSWDs in the NA30% plots were similar to NA70% for both varieties in 2014 (Fig. 2c,d). Irrigation management appeared to have a lower effect on SSWD at the different irrigation levels due to higher precipitation in 2014.

CH₄ emissions under contrasting climate conditions. The CH₄ flux from all plots increased from the beginning, and peaked for the first time at 25–30 days after transplanting, then rapidly decreased to zero due to MD. Thereafter, CH₄ fluxes from the NA100% and NA70% plots increased, peaked for a second time at 80–100 days after transplanting, and decreased afterwards until the end of the seasons. However, CH₄ fluxes from the NA30% plot were very low after MD (Fig. 3). Consequently, the NA100% plots had the largest CH₄ emissions for both rice varieties (Table 2). The average total CH₄ emissions from the NA100% plots was 122 kg CH₄ ha⁻¹ (the range was from 47.8 to 252 kg CH₄ ha⁻¹ in this study), which was within the range of 5.4–275 kg CH₄ ha⁻¹ range recorded in numerous other studies where paddies were subject to traditional water management regimes (i.e. NA100% condition) and a N fertilizer application rate of 200–250 kg N ha⁻¹, which is typical for the Yangtze River delta zone^{7,9,32–35}.

A significant difference in total CH₄ emissions was observed among irrigation levels ($P < 0.01$, Table 2). The NA70% and NA30% plots respectively reduced total CH₄ emissions by 31–53% and 32–77% in both seasons compared to the NA100% plots. Similarly, Hou *et al.* reported that controlled irrigation (NA30%) mitigated CH₄ emissions by 82% relative to traditional irrigation (NA100%)³³. Wang *et al.* found that intermittent irrigation (corresponding to NA70%) and constant moisture (corresponding to NA30%) reduced CH₄ emissions by 25% and 58%, respectively, compared to the continuous flooding condition (NA100%)³⁶. Reductions in irrigation water to the rice paddies led to a lower SSWD and even no standing water above the surface. This increased oxygen penetration into the soil and led to soil organic C being oxidized to CO₂ instead of CH₄, which ultimately suppressed CH₄ emissions. In this study, reducing the amount of irrigation was found to mitigate CH₄ emissions not only under normal climate condition but also in the ‘warm and dry’ season.

There was a significant difference in CH₄ emissions between Huayou14 and Hanyou8 ($P < 0.05$, Table 2). The CH₄ emissions from Huayou14 were always lower than from Hanyou8, especially in the NA100% and NA70%




Figure 2. Seasonal variation in surface standing water depth (SSWD) in the 2013 (a) Huayou14; (b) Hanyou8 and 2014 (c) Huayou14; (d) Hanyou8 rice growing seasons. Abbreviations: NA100% = normal amount of irrigation, NA70 = 70% of normal irrigation, NA30 = 30% of normal irrigation. Gray belts represent the mid-season drainage periods. Error bars represent standard errors of the means.




Figure 3. Seasonal variation in CH_4 fluxes in the 2013 (a) Huayou14; (b) Hanyou8 and 2014 (c) Huayou14; (d) Hanyou8 rice growing seasons. Abbreviations: NA100% = normal amount of irrigation, NA70% = 70% of normal irrigation, NA30% = 30% of normal irrigation, BF = base fertilizer, TF = tillering fertilizer, HF = heading fertilizer. Gray belts represent the periods of mid-season drainage. Arrows denote the fertilization date, and error bars represent standard errors of the means.

Treatments	CH ₄ emission (kg CH ₄ ha ⁻¹)		N ₂ O emission (kg N ₂ O ha ⁻¹)	
	2013	2014	2013	2014
H14-NA100%	92.3 ± 18.8	47.8 ± 14.7	0.6 ± 0.2	1.8 ± 0.4
H14-NA70%	63.4 ± 19.9	30.6 ± 6.0	1.5 ± 0.5	2.2 ± 0.8
H14-NA30%	44.9 ± 3.2	32.6 ± 4.9	0.8 ± 0.4	2.7 ± 0.6
H8-NA100%	252.2 ± 108.2	96.5 ± 31.4	0.2 ± 0.3	2.0 ± 0.8
H8-NA70%	117.8 ± 17.8	47.9 ± 16.7	0.4 ± 0.3	2.6 ± 0.2
H8-NA30%	58.7 ± 13.3	29.7 ± 6.2	0.5 ± 1.0	8.9 ± 6.4
Analysis of variance				
IM	**		ns	
RV	*		ns	
Year	**		*	
IM × RV	ns		ns	
IM × Year	ns		ns	
RV × Year	ns		ns	
IM × RV × Year	ns		ns	

Table 2. Seasonal total CH₄ and N₂O emissions in the 2013 and 2014 rice growing seasons. Numbers in the table represent means ± standard errors. H8 = Hanyou8, H14 = Huayou14, NA100% = normal amount of irrigation, NA70% = 70% of normal irrigation, NA30% = 30% of normal irrigation, IM = Irrigation management, RV = Rice Variety. * $P < 0.05$, ** $P < 0.01$, ns = not significance at 0.05 level.

plots. That may be related to the relatively lower SSWD in the Huayou14 plots compared to the Hanyou8 plots, which used less water to maintain growth and development and led to higher SSWDs (Fig. 2). The higher SSWD in the Hanyou8 plots could decrease soil Eh by 54 mV (averaged across the values for the NA100% and NA70% plots), and this facilitated CH₄ production. Additionally, it has been well documented that the difference in CH₄ emissions among rice varieties is correlated with the total amount of root exudate-C (e.g. sugars and organic acids)^{37,38}. Similarly, in this study, some organic C materials, such as soluble sugar and proline, were observed higher in Hanyou8 than in Huayou14 (unpublished data), which may enhance CH₄ production by changing the quantity and quality of the root exudates³⁹. However, when the amount of irrigation water was reduced, CH₄ emission was mitigated more in the Hanyou8 plots (average reduction of 52% and 73% by NA70% and NA30%, respectively, compared to NA100%) than that in the Huayou14 plots (average reduction of 34% and 42% by NA70% and NA30%, respectively, compared to NA100%) (Table 2).

CH₄ emissions were significantly higher in the ‘warm and dry’ season of 2013 than in the normal season of 2014 ($P < 0.01$, Table 2). It is difficult to control climate conditions in field experiments, so there is little data available from field experiments but model simulations have been used to compare CH₄ and N₂O under contrasting climate conditions. By employing the DNDC biogeochemical model to simulate greenhouse gas emissions in Chinese rice growing fields between 1971 and 2010, Tian *et al.* found that CH₄ emission was enhanced during the second 20 years (1991–2010) compared to the first 20 years (1971–1990) due to a 0.5 °C increase in air temperature²⁹. In this study, the CH₄ emissions from the NA100% plots by Huayou14 and Hanyou8 increased by 93% and 161% in the ‘warm and dry’ season of 2013, respectively, compared to the normal season of 2014. Our results partially confirmed the reliability of the model and they improved the accuracy and efficiency of the model when attempting to predict the effects of climate conditions on CH₄ emissions in the future. There were several possible reasons for the relatively higher CH₄ emissions in the ‘warm and dry’ season. Firstly, although there was less precipitation in 2013, large amounts of irrigation water were added to the plots to meet rice growth demand and made the SSWD values higher than or similar to those in 2014. Secondly, the higher air/soil temperature could potentially enhance CH₄ production and emission^{40,41}. Under higher air/soil temperature conditions, more root exudates are released into the soil and the potential availability of C for methanogens is enhanced^{40,42}. For example, Tokida *et al.* found that soil warming enhanced rice root decay and provided more substrates for CH₄ production⁴³. Higher air temperatures also enhanced paddy soil organic C mineralization⁴⁴ and potentially provided available C for methanogens. Moreover, the higher soil temperature around the rice roots may accelerate the CH₄ transport process through rice plants⁴⁵. Dijkstra *et al.* reviewed more than 100 rice paddy field studies that had investigated the response of CH₄ and N₂O emissions to elevated air/soil temperatures and reported that elevated temperature enhanced CH₄ emissions in 73 studies, but depressed emissions in 41 studies³⁰. These studies generally elevated the air/soil temperatures by using OTC (Open Top Chamber) or heating cables buried in the soil, or used infrared heaters installed above the canopy. However, they did not take the variations in precipitation into consideration. It is well known that the relative humidity of the atmosphere can have a negative relationship with the transpiration rate at certain temperatures. The lower relative humidity, caused by reduced precipitation and higher air temperatures in 2013, may have increased rice plant transpiration and allowed more CH₄-rich water from underground to migrate aboveground, which would eventually increase the release of CH₄ through micropores in the leaf sheaths.

N₂O emissions under contrasting climate conditions. As shown in Fig. 4, clear N₂O fluxes peaks were detected mainly in the NA30% or NA70% plots for both varieties. However, at all the irrigation levels, no significant differences in N₂O emissions were observed between Hanyou8 and Huayou14 in either season. Most of the




Figure 4. Seasonal variation in N_2O fluxes in the 2013 (a) Huayou14; (b) Hanyou8 and 2014 (c) Huayou14; (d) Hanyou8 rice growing seasons. Abbreviations: NA100% = normal amount of irrigation, NA70% = 70% of normal irrigation, NA30% = 30% of normal irrigation, BF = base fertilizer, TF = tillering fertilizer, HF = heading fertilizer. Gray belts represent the periods of mid-season drainage. Arrows denote the fertilization date and error bars represent standard errors of the means.

N_2O flux peaks were detected after fertilization, regardless of the rice variety. The influence of fertilization on N_2O emissions was relatively transitory and vigorous (Fig. 4). The average total N_2O emission was $1.2 \text{ kg N}_2\text{O ha}^{-1}$ and ranged from 0.2 to $2.0 \text{ kg N}_2\text{O ha}^{-1}$ in the NA100% plots, which was consistent with the values (0.4 – $5.3 \text{ kg N}_2\text{O ha}^{-1}$) recorded in many previous studies from the same region on paddies subjected to similar water management regimes and N fertilizer application rates^{9,32–35,46}.

The NA70% and NA30% plots increased total N_2O emissions by 22–146% and 26–338%, respectively, compared to the NA100% plots. However, no significant differences in total N_2O emissions were detected amongst the irrigation treatments (Table 2). Hou *et al.* also found that N_2O emissions under controlled irrigation (NA30%) increased by 135% relative to traditional irrigation (NA100%)³³. Reductions in irrigation water to the rice paddies frequently subject the soil to alternating wet/dry conditions, which stimulates N_2O producers activity and increases N_2O emissions⁴. The increased N_2O emissions in the NA70% and NA30% plots after fertilization, relative to the NA100% plot, are probably due to the abundant, newly-added N^6 and the suitable soil moisture conditions³³. Finally, there was no significant difference in total N_2O emissions between Huayou14 and Hanyou8 (Table 2).

There were larger and more N_2O flux peaks in the normal season of 2014 than in the ‘warm and dry’ season of 2013. Similarly, Tian *et al.* also suggested that an increase in precipitation could enhance N_2O emissions after the DNDC model simulation²⁹. Alternative wet and dry soil conditions caused by frequent precipitation would enhance N_2O production through nitrification and denitrification in 2014⁴⁷. However, in this study, the SSWD when fertilizer was applied may play an important role in controlling N_2O emissions. The averaged SSWD over all treatments was higher after fertilization in the ‘warm and dry’ season of 2013 (5.3 cm and 1.9 cm after base and heading fertilization, respectively) than in the normal season of 2014 (2.6 cm and 1.0 cm after base and heading fertilization, respectively) (Fig. 2). In a freshwater marsh, Yang *et al.* also found that a lower water table position (−11 to 0 cm) enhanced N_2O emissions relative to higher water tables (+2 to +14 cm)⁴⁸. Similarly, in a northern boreal fen located in north-western Finland, Lohila *et al.* reported that the highest N_2O fluxes occurred when the SSWD was about 4 cm, whereas atmospheric N_2O was consumed when the SSWD was 15 cm⁴⁹. When the SSWD is lower, less N_2O dissolves into the surface standing water and more N_2O is probably quickly released into the atmosphere. In contrast, higher SSWD could restrict the availability of oxygen and therefore favor the formation of molecular nitrogen (N_2) instead of N_2O . Hence, in this study, N_2O emission might be more related to fertilization and SSWD rather than rice variety and high air temperature.

Rice grain yield and equivalent CO_2 emissions ($\text{CO}_2\text{-eq}$) under contrasting climate conditions. In the NA100% plots, the average rice yield was 9.3 t ha^{-1} with a range of 8.7 – 10.1 t ha^{-1} (Table 3), which was higher than that recorded in other studies (4.8 – 9.3 t ha^{-1}) conducted in the same region^{7,9,32–35,46}. The potentially

Treatments	Rice grain yield ($t\ ha^{-1}$)		CO ₂ -eq emission (kg CO ₂ -eq t^{-1})	
	2013	2014	2013	2014
H14-NA100%	8.8 ± 0.1	10.1 ± 0.2	283.1 ± 48.2	169.7 ± 25.2
H14-NA70%	8.3 ± 0.2	10.0 ± 0.2	246.4 ± 73.9	142.2 ± 39.2
H14-NA30%	7.9 ± 0.0	9.8 ± 0.1	170.2 ± 9.0	165.8 ± 25.8
H8-NA100%	8.7 ± 0.1	9.4 ± 0.1	735.7 ± 324.8	322.8 ± 69.3
H8-NA70%	8.6 ± 0.3	9.3 ± 0.1	352.9 ± 30.7	211.5 ± 45.8
H8-NA30%	8.2 ± 0.4	9.3 ± 0.1	194.1 ± 58.1	371.9 ± 201.7
Analysis of variance				
IM	**		ns	
RV	*		*	
Year	***		ns	
IM × RV	ns		ns	
IM × Year	ns		ns	
RV × Year	***		ns	
IM × RV × Year	ns		ns	

Table 3. Rice grain yield and CO₂-eq emissions in the 2013 and 2014 rice growing seasons. Numbers in the table represent means ± standard errors. H8 = Hanyou8, H14 = Huayou14, NA100% = normal amount of irrigation, NA70% = 70% of normal irrigation, NA30% = 30% of normal irrigation, IM = Irrigation management, RV = Rice Variety. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, ns = not significant at 0.05 level.

negative effect of less precipitation on rice grain yield may be offset by irrigation in the ‘warm and dry’ season. However, rice grain yield significantly decreased with the reduction in the amount of irrigation ($P < 0.01$, Table 3). In the ‘warm and dry’ season, the Huayou14 yield loss was 10% in the NA30% plots compared to the NA100% plots, which was greater than the yield loss recorded for Hanyou8 (6%). This indicates that Hanyou8 is more drought-resistant with regards to rice grain yield, which agrees with the results reported by Luo²⁰.

Recently, Kim *et al.* used the CERES-Rice 4.0 crop simulation model to investigate the effects of climate change on rice grain yield in the temperate climate regions under the East Asian monsoon system and suggested that the air temperature increases could lead to a significant decrease in rice grain yield by 22.1–35.0%⁵⁰. In this study, rice grain yield was significantly lower in the ‘warm and dry’ season (2013) than in the normal season (2014) ($P < 0.001$, Table 3). The yield decreased by 13–19% for Huayou14 and 7–12% for Hanyou8 in 2013 compared to the yield in 2014 and this was probably due to higher air temperature and reduced precipitation. In another case study conducted in the same region, Liu *et al.* reported that although the average rice grain yield increased by 46% from the 1980 s to 2000 s due to the soil improvement, rice variety updating and agricultural management advances, the climate conditions (e.g. higher air temperature, less precipitation) had a negative effect on rice grain yield⁵¹. Generally, air temperature enrichment in rice growing seasons could shorten rice development stages and reduce grain yield^{52–54}. Elevated air temperature can result in rice grain yield loss, mainly through reduced photosynthesis caused by chloroplast damage, spikelet sterility caused by decreased pollen production, and increased energy consumption caused by higher respiration demand²⁵. Moreover, there was a significant interaction between rice variety and year ($P < 0.001$, Table 3).

The average CO₂-eq emissions in the NA100% plots was 378 kg CO₂-eq t^{-1} with a range of 170 to 736 kg CO₂-eq t^{-1} (Table 3), which was within the 33–557 kg CO₂-eq t^{-1} range reported in previous studies conducted in the same region^{9,32–35}. In most treatments, the detected CO₂-eq emissions were higher in the ‘warm and dry’ season than in the normal season. However, the difference between the two seasons was not significant (Table 3). Higher CH₄ emissions and lower rice grain yields from the treatments led to greater CO₂-eq emissions in the ‘warm and dry’ season relative to the normal season. Interestingly, there was a significant difference in CO₂-eq emissions between Huayou14 and Hanyou8 ($P < 0.05$, Table 3). This suggests that the CO₂-eq emissions vary considerably with rice growing season or rice variety. Furthermore, CO₂-eq emissions from the NA70% and NA30% plots of both rice varieties (excluding H8-NA30% in the 2014 normal season) were potentially depressed when compared to the NA100% plot (Table 3). The decreases in CH₄ emissions from the NA70% and NA30% plots were the main cause of the effective depression in CO₂-eq emissions, especially during the ‘warm and dry’ season. However, in the normal season, due to the higher N₂O emissions, reductions in the amount of irrigation had little effect on CO₂-eq emissions.

In conclusion, CH₄ and N₂O emissions strongly differed according to rice variety and irrigation management between the two rice growing seasons under contrasting climate conditions. The amount of irrigation water was significantly higher in the ‘warm and dry’ season than that in the normal season. Although the same amount of irrigation water was applied to the two rice varieties, the SSWDs in the plots planted with Hanyou8 were higher than Huayou14, due to lower water demand from Hanyou8. The CH₄ emissions by Huayou14 and Hanyou8 increased 93% and 161% in the ‘warm and dry’ season (2013), respectively, compared with that in normal season (2014). Moreover, the CH₄ emissions from Hanyou8 were higher than from Huayou14 in both seasons. Reducing the amount of irrigation water can effectively reduce the CH₄ emissions, regardless of the rice variety and climate conditions. However, less irrigation during the ‘warm and dry’ season greatly decreased the Huayou14 grain yield, but had little impact on Hanyou8 yield. In contrast, compared to the effect of rice variety, N₂O emission depended

more on fertilization and surface standing water depth when the fertilizer was applied. Under the global warming scenarios, feasible reductions in the amount of irrigation water applied and the suitable selection of rice varieties would be a promising way to mitigate greenhouse gas emissions as well as maintain rice grain yield.

Materials and Methods

Study site and experimental design. The study was conducted in an experimental field at the Shanghai Engineering Research Center of Low-carbon Agriculture, which is a part of the Zhuanghang Experimental Station ($30^{\circ}53'N$, $121^{\circ}23'E$), and is located in the Yangtze River delta zone in east China. A rice-wheat cropping rotation system is the typical practice in this area. The paddy field soil was plowed to a depth of ~ 15 cm, and its chemical and physical properties were as follows: soil organic C (SOC) $13.7\text{ g}\cdot\text{kg}^{-1}$, total N $1.4\text{ g}\cdot\text{kg}^{-1}$, bulk density $1.4\text{ g}\cdot\text{cm}^{-3}$, and pH (H_2O) 7.6.

Each experimental plot was 60 m^2 and an impermeable membrane was buried vertically in the soil around each plot at 1.1 m depth to prevent lateral seepage between the experimental plots. Then a concrete wall (30 cm width \times 60 cm height) was built around the experimental plots. It was half buried into the soil over the impermeable membrane. A centrifugal pump (SW100-160, $100\text{ m}^3\cdot\text{h}^{-1}$, Shanghai Sanxing Supply and Drainage Equipment Co., Ltd., Shanghai, China) and polyethylene pipes were used to transport river water to each plot for irrigation. A meteorological station was established nearby in 2012, which provided data about the air/soil temperature, dry/wet precipitation, evaporation, solar radiation, wind speed/direction etc.

Two rice varieties (*Oryza sativa* L. Huayou14 and Hanyou8) and three types of irrigation management were employed in this study. Each treatment was replicated three times, resulting in a total of 18 plots in this experiment (i.e., 2 rice varieties \times 3 irrigation levels \times 3 replicates). Huayou14 is a high-yielding hybrid that is often cultivated by local farmers and Hanyou8 was recently developed by the Shanghai Agrobiological Gene Center for its water-saving and drought-resistant traits. The three types of irrigation management applied were normal amount of traditional irrigation management (NA100%), 70% of normal (NA70%), and 30% of normal (NA30%). The performance of NA100% was consistent with conventional irrigation practice for meeting rice growth demand. The other two types were applied proportionately during every irrigation event when little or no surface-standing water was observed in the NA100% plot for Huayou14. The irrigation was mainly carried out between 30 and 100 days after transplanting.

Climate conditions and agricultural practices. Seasonal changes in daily mean air temperature, daily precipitation, and soil temperature at 5 cm depth in the 2013 and 2014 rice growing seasons are shown in Fig. 1. The mean seasonal air temperature was 26.7°C and ranged from 14.8 to 33.5°C during the 2013 rice growing season (Table 1 and Fig. 1a), whereas it was 24.4°C (ranging from 14.7 to 30.2°C) in 2014 (Table 1 and Fig. 1b). The mean seasonal air temperature (24.4°C) in 2014 was similar to the normal value (24.7°C) reported by Su *et al.* in this region⁵⁵. The mean seasonal air temperature in 2013 was 2.3°C higher than that in 2014. Seasonal variations in soil temperature were similar to the daily mean air temperature (Fig. 1). Total precipitation in the 2013 and 2014 seasons was 492.1 and 762.7 mm, respectively (Table 1). In this region, annual precipitation is about 1200 mm, and about 60% of precipitation occurs between May and September⁵⁶. In the 2013 season, precipitation between June and September (i.e., a duration from transplanting to the grain-filling stage) was only 271.8 mm, and this season was considered as a ‘warm and dry’ season. In contrast, in 2014, the total precipitation between June and September was similar to the average for the area, and this season was regarded as normal season.

Rice plants were transplanted at a density of 20 hills per m^2 on June 14/16 and harvested on October 21/22 in 2013 and 2014, respectively. The N fertilizer application rate was $225\text{ kg}\cdot\text{ha}^{-1}$, which was applied at a ratio of 5:3:2 (w/w/w) as base, tillering and heading applications, respectively. The base fertilizer was applied in the form of a compound fertilizer at 1–2 days before transplanting. The tillering and heading fertilizers were applied in the form of urea at about 1 week and 7 weeks after transplanting, respectively. Phosphorous (P_2O_5) fertilizer was applied as a base, compound fertilizer at a rate of $112.5\text{ kg}\cdot\text{ha}^{-1}$, and 44% potassium (K_2O) fertilizer was applied as a base, compound fertilizer at a rate of $255\text{ kg}\cdot\text{ha}^{-1}$. The remaining of potassium (K_2O) fertilizer was applied as a heading fertilizer in the form of commercial potassium chloride (KCl).

There were 12 and three irrigation events in the 2013 and 2014 seasons, respectively. Mid-season drainage (MD) is a conventional agricultural practice during the rice growing season. The MD began on July 19/21 and finished on July 29/August 4 in 2013 and 2014, respectively.

Measurements. The samples used to determine CH_4 and N_2O concentrations were taken using a static transparent chamber consisting of a plexiglass base frame ($50\text{ cm length} \times 40\text{ cm width} \times 20\text{ cm height}$) and a plexiglass lid ($50\text{ cm} \times 40\text{ cm} \times 50\text{ cm}$) equipped with a battery-driven 12 V fan at the center of its inner top. Other plexiglass frames were used to extend the lid height, by 20 , 40 , or 60 cm depending on the height of rice plants. The base frames were inserted approximately 15 cm into the soil, and four hills of rice plants were transplanted. One base frame was placed in each plot. Four gas samples were collected from each chamber at 6-min intervals using an auto gas sampler attached to four aluminum foil gas bags (1 L, Dalian Delin Gas Packing Co., Ltd., Dalian, China) at each sampling time. The auto gas sampler was composed of a 12 V rechargeable battery (NP7-12, YUASA Battery (Guangdong) Co., Ltd, Guangzhou, China), a gas pump (FAY4002, 2 L min^{-1} , Chengdu Qihai E&M Manufacturing Co., Ltd., Chengdu, China), a box containing a circuit board (Nanjing Weina Electronic Co., Ltd., Nanjing, China), and a series of compact direct-operated 2-port solenoid valves (VDW23-6G-1, SMC Pneumatics Ltd., Tokyo, Japan).

The gas samples from all the plots were collected between 08:00 and 10:00 and immediately taken to the laboratory. The concentrations of CH_4 and N_2O were determined by a gas chromatograph (7820 A, Agilent Technologies, Inc., Santa Clara, CA, USA) equipped with a flame ionization detector and an electron capture

detector, respectively. The sampling frequency was once a week. However, whenever there was a fertilizer application, an MD, or irrigation after MD, a higher sampling frequency (once every 2 days) was used and daily sampling lasted for one week. CH_4 and N_2O flux were calculated by examining the linear increases of CH_4 and N_2O concentrations in the headspace of the chambers over time. The seasonal total CH_4 and N_2O emissions from all plots were calculated directly from the fluxes.

The surface-standing water depth (SSWD) was measured directly using a ruler after each gas-sampling event. Soil Eh (Oxidation-Reduction Potential) was detected at 5 cm soil depth with a pH/ NO_3^- /Eh meter (PRN-41, DKK-TOA Co., Tokyo, Japan). Soil temperature was monitored at 5 cm soil depth using a moisture meter (HH2, Delta-T Devices Ltd, Cambridge, UK) during each gas-sampling event around the base frames. At the end of each rice growing season, the plants in each plot were manually harvested. The dry weight of the rice grains was determined using an oven at 75 °C. Finally, the rice grain yield of each plot was calculated on a rice grain dry weight basis using the equation:

$$RY = DW / (1 - 14.5\%)$$

where RY is the rice grain yield ($\text{t}\cdot\text{ha}^{-1}$), DW is the dry weight of rice grains ($\text{t}\cdot\text{ha}^{-1}$), and 14.5% was used as the standard moisture content for storage of the rice varieties used in this study.

The equivalent CO_2 (CO_2 -eq) emission for total CH_4 and N_2O emissions (greenhouse gas intensity) was calculated on a rice grain yield basis using the equation:

$$T_{\text{CO}_2-\text{eq}} = (T_{\text{CH}_4} \times 25 + T_{\text{N}_2\text{O}} \times 298) / RY$$

where $T_{\text{CO}_2-\text{eq}}$ is the total amount of equivalent CO_2 emission ($\text{kg CO}_2\text{-eq}\cdot\text{t}^{-1}$), T_{CH_4} is the total amount of CH_4 emission ($\text{kg}\cdot\text{ha}^{-1}$), $T_{\text{N}_2\text{O}}$ is the total amount of N_2O emission ($\text{kg}\cdot\text{ha}^{-1}$), 25 and 298 are the multiples of GWP (global warming potential) for CH_4 and N_2O versus CO_2 over 100 years³, and RY is the rice grain yield ($\text{t}\cdot\text{ha}^{-1}$).

Statistical analysis. The impacts of the three parameters (irrigation management, rice variety, and year) on CH_4 and N_2O emissions from rice paddies were examined. Their effects were analyzed using the general linear model for analysis of variance along with the least significant difference test. The significance level for both tests was 5%. SPSS 20.0 statistical software (IBM Co., New York, USA) was used to conduct the analysis. The figures were prepared using SigmaPlot 12.5 software (Systat Software Inc., San Jose, CA, USA).

References

- Le Treut, H. *et al.* Historical overview of climate change. In: Climate Change 2007: The physical Science Basis. *Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (eds. Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M. & Miller, H. L.) 93–127 (Cambridge University Press, Cambridge and New York, 2007).
- De Boeck, H. J., Dreesen, F. E., Janssens, I. A. & Nijs, I. Whole-system responses of experimental plant communities to climate extremes imposed in different seasons. *New Phytol.* **189**, 806–817 (2011).
- Forster, P. *et al.* Changes in atmospheric constituents and in radiative forcing. In: Climate Change 2007: The physical Science Basis. *Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (eds. Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M. & Miller, H. L.) 129–234. (Cambridge University Press, Cambridge and New York, 2007)
- Hou, A. *et al.* Methane and nitrous oxide emissions from a rice field in relation to soil redox and microbiological process. *Soil Sci. Soc. Am. J.* **60**, 2180–2186 (2000).
- Cai, Z. *et al.* Methane and nitrous oxide emissions from rice paddy fields as affected by nitrogen fertilizers and water management. *Plant Soil*, **196**, 7–14 (1997).
- Zou, J. *et al.* A 3-year field measurement of methane and nitrous oxide emissions from rice paddies in China: Effects of water regime, crop residue, and fertilizer application. *Global Biogeochem. Cy.* **19**, GB2021 (2005).
- Dong, H. *et al.* Effect of ammonium-based, non-sulfate fertilizers on CH_4 emissions from a paddy field with a typical Chinese water management regime. *Atmos. Environ.* **45**, 1095–1101 (2011).
- Li, X. *et al.* Effect of timing and duration of midseason aeration on CH_4 and N_2O emissions from irrigated lowland rice paddies in China. *Nutr. Cycl. Agroecosyst.* **91**, 293–305 (2011).
- Yao, Z. *et al.* A 3-year record of N_2O and CH_4 emissions from a sandy loam paddy during rice seasons as affected by different nitrogen application rates. *Agr. Ecosyst. Environ.* **152**, 1–9 (2012).
- Liou, R., Huang, S. & Lin, C. Methane emission from fields with differences in nitrogen fertilizers and rice varieties in Taiwan paddy soils. *Chemosphere* **50**, 237–246 (2003).
- Kerdchoechuen, O. Methane emission in four rice varieties as related to sugars and organic acids of roots and root exudates and biomass yield. *Agr. Ecosyst. Environ.* **108**, 155–163 (2005).
- Gogoi, B. & Baruah, K. K. Nitrous oxide emissions from fields with different wheat and rice varieties. *Pedosphere* **22**, 112–121 (2012).
- Riya, S. *et al.* CH_4 and N_2O emissions from different varieties of forage rice (*Oryza sativa* L.) treating liquid cattle waste. *Sci. Total Environ.* **419**, 178–186 (2012).
- Passioura, J. B. The drought environment: physical, biological and agricultural perspectives. *J. Exp. Bot.* **58**, 113–117 (2007).
- Majeed, A. *et al.* Physiology and productivity of rice crop influenced by drought stress induced at different developmental stages. *Afr. J. Biotech.* **10**, 5121–5136 (2011).
- Bhuiyan, S. L. Water management in relation to crop production: case study on rice. *Outlook Agr.* **21**, 293–299 (1992).
- Zhang, Q. Strategies for developing green super rice. *P. Natl Acad. Sci. USA.* **104**, 16404–16409 (2007).
- Bouman, B. A. M., Lampayan, R. M. & Tuong, T. P. *Water management in irrigated rice: coping with water scarcity*. International Rice Research Institute, Los Baños, Philippines, 54p (2007).
- Tao, H. *et al.* Growth and yield formation for rice (*Oryza sativa* L.) in the water-saving ground cover rice production system (GCRPS). *Field Crops Res.* **95**, 1–12 (2006).
- Luo, L. Breeding for water-saving and drought-resistance rice (WDR) in China. *J. Exp. Bot.* **61**, 3509–3517 (2010).
- Tang, G. & Ren, G. Reanalysis of surface air temperature change of the last 100 years over China. *Clim. Environ. Res.* **10**(4), 791–798 (2005). (in Chinese with English abstract).

22. Gu, P. & Jin, Q. Influence of climate warming on single-crop late rice yield in Fengxian District of Shanghai. *Acta Agr. Shanghai* **25**(3), 115–119 (2009). (in Chinese with English abstract).
23. IPCC (Intergovernmental Panel on Climate Change). Summary for policymakers. In: *Climate Change 2013: The Physical Science Basis. Contribution of working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds. Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V. & Midgley, P. M.) 3–29 (Cambridge University Press, Cambridge and New York, 2013).
24. Krishnan, P., Ramakrishnan, B., Reddy, K. R. & Reddy, V. R. High-temperature effects on rice growth, yield, and grain quality. *Adv. Agron.* **111**, 87–206 (2011).
25. Wassmann, R. *et al.* Climate change affecting rice production: The physiological and agronomic basis for possible adaptation strategies. *Adv. Agron.* **101**, 59–122 (2009).
26. Nagarajan, S. *et al.* Local climate affects growth, yield and grain quality of aromatic and non-aromatic rice in northwestern India. *Agr. Ecosyst. Environ.* **138**, 274–281 (2010).
27. Welch, J. R. *et al.* Rice yields in tropical/subtropical Asia exhibit large but opposing sensitivities to minimum and maximum temperatures. *P. Natl Acad. Sci. USA.* **107**, 14562–14567 (2010).
28. Syp, A. *et al.* Modelling impact of climate change and management practices on greenhouse gas emissions from arable soils. *Pol. J. Environ. Stud.* **20**, 1593–1602 (2011).
29. Tian, Z. *et al.* China's rice field greenhouse gas emission under climate change based on DNDC model simulation. *Chin. J. Appl. Ecol.* **26**, 793–799 (2015). (in Chinese with English abstract).
30. Dijkstra, F. A. *et al.* Effects of elevated carbon dioxide and increased temperature on methane and nitrous oxide fluxes: evidence from field experiments. *Front. Ecol. Environ.* **10**, 520–527 (2012).
31. Sun, H. *et al.* CH₄ emission in response to Water-saving and Drought-resistance Rice (WDR) and common rice varieties under different irrigation managements. *Water Air Soil Poll.* **227**, 47 (2016).
32. Liu, S. *et al.* Methane and nitrous oxide emissions from direct-seeded and seedling-transplanted rice paddies in southeast China. *Plant Soil* **374**, 285–297 (2014).
33. Hou, H. *et al.* Seasonal variations of CH₄ and N₂O emissions in response to water management of paddy fields located in Southeast China. *Chemosphere* **89**, 884–892 (2012).
34. Zhang, X. *et al.* Comparison of greenhouse gas emissions from rice paddy fields under different nitrogen fertilization loads in Chongming Island, Eastern China. *Sci. Total Environ.* **472**, 381–388 (2014).
35. Zou, J. *et al.* Sewage irrigation increased methane and nitrous oxide emissions from rice paddies in southeast China. *Agr. Ecosyst. Environ.* **129**, 516–522 (2009).
36. Wang, B. *et al.* Methane emissions from ricefields as affected by organic amendment, water regime, crop establishment, and rice cultivar. *Environ. Mon. Ass.* **57**, 213–228 (1999).
37. Aulakh, M. S., Wassmann, R., Bueno, C. & Rennenberg, H. Impact of root exudates of different cultivars and plant development stages of rice (*Oryza sativa* L.) on methane production in a paddy soil. *Plant Soil* **230**, 77–86 (2001).
38. Kerdchoechuen, O. Methane emission in four rice varieties as related to sugars and organic acids of roots and root exudates and biomass yield. *Agr. Ecosyst. Environ.* **108**, 155–163 (2005).
39. Holzapfel-Pschorn, A., Conrad, R. & Seiler, W. Effects of vegetation on the emission of methane by submerged paddy soil. *Plant Soil* **92**, 223–233 (1986).
40. Allen, L. H. Jr. *et al.* Methane emissions of rice increased by elevated carbon dioxide and temperature. *J. Environ. Q.* **32**, 1978–1991 (2003).
41. Tokida, T. *et al.* Effects of free-air CO₂ enrichment (FACE) and soil warming on CH₄ emission from rice paddy field: impact assessment and stoichiometric evaluation. *Biogeosciences* **7**, 2639–2653 (2010).
42. Denier van der Gon, H. A. C. *et al.* Optimizing grain yields reduces CH₄ emissions from rice paddy fields. *P. Natl Acad. Sci. USA.* **99**, 12021–12024 (2002).
43. Tokida *et al.* Methane and soil CO₂ production from current-season photosynthates in a rice paddy exposed to elevated CO₂ concentration and soil temperature. *Global Change Biol.* **17**, 3327–3337 (2011).
44. Ren *et al.* Effects of temperature on organic carbon mineralization in paddy soils with different clay content. *Chin. J. Appl. Ecol.* **18**, 2245–2250 (2007). (in Chinese with English abstract).
45. Hosono, T. & Nouchi, I. The dependence of methane transport in rice plants on the root zone temperature. *Plant Soil* **191**, 233–240 (1997).
46. Liu, S., Qin, Y., Zou, J. & Liu, Q. Effects of water regime during rice-growing season on annual direct N₂O emission in a paddy rice-winter wheat rotation system in southeast China. *Sci. Total Environ.* **408**, 906–913 (2010).
47. Chen, W. *et al.* Effects of increasing precipitation and nitrogen deposition on CH₄ and N₂O fluxes and ecosystem respiration in a degraded steppe in Inner Mongolia, China. *Geoderma* **192**, 335–340 (2013).
48. Yang, J. *et al.* Effect of water table level on CO₂, CH₄ and N₂O emissions in a freshwater marsh of Northeast China. *Soil Biol. Biochem.* **61**, 52–60 (2013).
49. Lohila *et al.* Responses of N₂O fluxes to temperature, water table and N deposition in a northern boreal fen. *Eur. J. Soil Sci.* **61**, 651–661 (2010).
50. Kim, H. Y., Ko, J., Kang, S. & Tenhunen, J. Impacts of climate change on paddy rice yield in a temperate climate. *Global Change Biol.* **19**, 548–562 (2013).
51. Liu, L. *et al.* Impacts of climate changes, soil nutrients, variety types and management practices on rice yield in East China: A case study in the Taihu region. *Field Crops Res.* **149**, 40–48 (2013).
52. Sheehy, J. E., Mitchell, P. L. & Ferrer, A. B. Decline in rice grain yields with temperature: models and correlations can give different estimates. *Field Crops Res.* **98**, 151–156 (2006).
53. Liu, L., Wang, E., Zhu, Y. & Tang, L. Contrasting effects of warming and autonomous breeding on single-rice productivity in China. *Agr. Ecosyst. Environ.* **149**, 20–29 (2012).
54. Jalota, S. K., Kaur, H., Kaur, S. & Vashisht, B. B. Impact of climate change scenarios on yield, water and nitrogen-balance and -use efficiency of rice-wheat cropping system. *Agr. Water Manag.* **116**, 29–38 (2013).
55. Su, R. *et al.* Analysis on the single-crop late rice yield in Fengxian District as influenced by meteorological factors in 2013. *Acta Agr. Shanghai* **31**(3), 105–107 (2015). (in Chinese with English abstract).
56. Sun, H. *et al.* Effects of rice cultivars on CH₄ and N₂O emissions from rice fields. *J Agro-Environ. Sci.* **34**, 1595–1602 (2015). (in Chinese with English abstract).

Acknowledgements

This study was financed by National Natural Science Foundation of China (No. 41375157), National Key Technology Support Program of the Ministry of Science and Technology of China (No. 2013BAD11B02) and the Shanghai Municipal Agriculture Commission (Hu Nong Ke Tui Zi (2013) No. 1-1). The language in this paper has been re-edited and polished by International Science Editing.

Author Contributions

X.S., G.Z. and S.Z. conceived and participated the experimental design. H.S., Z.F. and G.C. carried out the experiments. H.S. and S.Z. wrote the main manuscript text and H.S. prepared all the figures and tables. All authors reviewed the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Sun, H. *et al.* A two-year field measurement of methane and nitrous oxide fluxes from rice paddies under contrasting climate conditions. *Sci. Rep.* **6**, 28255; doi: 10.1038/srep28255 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>