

SCIENTIFIC REPORTS



OPEN

Distributed lag effects and vulnerable groups of floods on bacillary dysentery in Huaihua, China

Received: 06 January 2016

Accepted: 17 June 2016

Published: 18 July 2016

Zhi-Dong Liu^{1,2,*}, Jing Li^{1,2,*}, Ying Zhang³, Guo-Yong Ding⁴, Xin Xu^{1,2}, Lu Gao^{1,2}, Xue-Na Liu^{1,2}, Qi-Yong Liu^{2,5} & Bao-Fa Jiang^{1,2}

Understanding the potential links between floods and bacillary dysentery in China is important to develop appropriate intervention programs after floods. This study aimed to explore the distributed lag effects of floods on bacillary dysentery and to identify the vulnerable groups in Huaihua, China. Weekly number of bacillary dysentery cases from 2005–2011 were obtained during flood season. Flood data and meteorological data over the same period were obtained from the China Meteorological Data Sharing Service System. To examine the distributed lag effects, a generalized linear mixed model combined with a distributed lag non-linear model were developed to assess the relationship between floods and bacillary dysentery. A total of 3,709 cases of bacillary dysentery were notified over the study period. The effects of floods on bacillary dysentery continued for approximately 3 weeks with a cumulative risk ratio equal to 1.52 (95% CI: 1.08–2.12). The risks of bacillary dysentery were higher in females, farmers and people aged 15–64 years old. This study suggests floods have increased the risk of bacillary dysentery with 3 weeks' effects, especially for the vulnerable groups identified. Public health programs should be taken to prevent and control a potential risk of bacillary dysentery after floods.

Floods are the most common type of natural disaster globally. On average, floods and other hydrological events had accounted for over 50% of the natural disasters between 2001 and 2010 in the world¹. Floods are expected to increase in frequency and intensity due to rising sea levels and more frequent and extreme precipitation events². Hunan, located in the basin of Yangtze River, is a flood-prone province in China. Persistent and heavy rainfall caused several floods from 2005–2011 in Huaihua City of Hunan Province³.

Bacillary dysentery, a diarrheal disease caused by ingestion of water or food contaminated by different species of *Shigella* bacteria, remains a major public health problem in some developing countries⁴. In China, 237,930 new cases of bacillary dysentery were reported in 2011, which ranked fourth among the national notifiable diseases from the Chinese National Notifiable Disease Report. Hunan is one of the most seriously affected provinces in the Yangtze River Region. Huaihua city, which located in the west of Hunan, had 5,414 cases from 2005–2011⁵.

The association between floods and bacillary dysentery is not clear. Some epidemiologic evidence suggests that floods are positively associated with diarrheal diseases, such as dysentery, cholera, and paratyphoid fever^{6,7}. During the flood in the Midwestern United States in 2001, Wade *et al.* found 1.29 times increase in the incidence of gastrointestinal symptoms⁸. A significant positive association between floods and diarrhea was reported by Heller *et al.* in Brazil⁹. A study from Henan of China revealed that floods were significantly associated with dysentery¹⁰. However, a study from Mozambique revealed that there was no outbreak of dysentery after the flooding in 2000¹¹. Another study also found no clear evidence of excess mortality or diarrhea risk during or after flooding

¹Department of Epidemiology, School of Public Health, Shandong University, Jinan, Shandong Province, People's Republic of China. ²Shandong University Climate Change and Health Center, Jinan, Shandong Province, People's Republic of China. ³School of Public Health, China Studies Centre, The University of Sydney, New South Wales, Australia. ⁴Department of Epidemiology, School of Public Health, Taishan Medical College, Taian, Shandong Province, People's Republic of China. ⁵State Key Laboratory for Infectious Disease Prevention and Control, National Institute for Communicable Disease Control and Prevention, China CDC, Beijing 102206, P. R. China. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to B. F. J. (email: bjiang@sdu.edu.cn)

Category	Mean \pm SD	Min	P25	Median	P75	Max
Gender						
Males	11.33 \pm 6.11	1	7	10	15	30
Females	8.49 \pm 4.58	0	5	8	11	30
Age (years)						
0–4	6.16 \pm 3.49	0	3	6	8	19
5–14	2.42 \pm 2.31	0	1	2	3	15
15–64	8.87 \pm 5.17	0	5	8	12	23
65+	2.37 \pm 1.90	0	1	2	3	11
Occupation						
Farmers	7.50 \pm 15.51	0	4	7	10	24
Students	2.33 \pm 2.60	0	1	2	3	22
Children	6.58 \pm 3.69	0	4	6	8	21
Works	1.83 \pm 1.57	0	1	2	3	7
Others	1.58 \pm 1.54	0	0	1	2	7
Total	19.62 \pm 9.38	2	12	19	25	50

Table 1. Description of weekly bacillary dysentery cases from April to September in Huaihua, China, 2005–2011.




Figure 1. The weekly distribution of bacillary dysentery cases during study period from 2005 to 2011 in Huaihua, China. Weeks with flood is indicated as a dashed line.

after controlling for pre-flood rate differences and seasonality¹². More research is needed to elucidate the potential risk of bacillary dysentery related to floods. Research linking floods to bacillary dysentery in China is limited. The effect of the floods on bacillary dysentery in Huaihua remains unknown. This study aimed to investigate the distributed lag effects on bacillary dysentery related to floods and identify the vulnerable groups. Results will contribute to a better understanding of the health impacts of floods and provide more evidence to support decision-making for the prevention and control of bacillary dysentery after floods.

Results

Description of the disease and meteorological data. A total of 3,709 cases of bacillary dysentery were notified in the study area over the study period. Table 1 presents the descriptive statistics for weekly bacillary dysentery cases in different categories from 2005–2011 in the flood season of Huaihua. There were more cases in male living in rural areas. Most of the bacillary dysentery cases were children and farmers. Patients aged 0–4 and 15–64 accounted for approximately 76% of all bacillary dysentery cases. Figures 1 and 2 show the time-series distribution of weekly bacillary dysentery cases and weekly mean temperature, weekly mean relative humidity and weekly cumulative precipitation in the flood season from 2005–2011. The average number of weekly bacillary dysentery cases was 20 (range: 2–50). The average values of weekly mean temperature, weekly mean relative humidity and weekly cumulative precipitation were 24 °C (range: 10–31 °C), 73% (range: 58–84%) and 28 mm (range: 0–153 mm), respectively. Fourteen flood events were recorded during the study period.

Association between floods and bacillary dysentery. The risk ratios (RRs) of floods on the risk of bacillary dysentery from the DLNM model were presented in Table 2. Results showed that bacillary dysentery was associated with floods significantly at lag 1 (RR = 1.32, 95% CI: 1.12–1.56), but there was no significant association between bacillary dysentery and floods at other lag periods. The cumulative effects of floods on bacillary dysentery were presented in Fig. 3. After controlling for precipitation, temperature, relative humidity, seasonality and long-term trends, floods were associated with bacillary dysentery significantly with a cumulative RR value at lag 0–2 weeks equal to 1.52 (95% CI: 1.08–2.12). The effect of floods on bacillary dysentery at lag 1 was similar for females (RR = 1.32, 95% CI: 1.06–1.65) and males (RR = 1.30, 95% CI: 1.06–1.61). But the cumulative effects




Figure 2. Weekly mean temperature (A) weekly mean relative humidity (B) and weekly cumulative precipitation (C) during study period from 2005–2011 in Huaihua. Weeks with flood is indicated as a dashed line in (C).

at lag 0–2 weeks were significant in females (RR = 1.65, 95% CI: 1.06–2.56), while not in males (RR = 1.35, 95% CI: 0.88–2.07). The most vulnerable age group was 15–64 years old group (RR at lag 1 = 1.39, 95% CI: 1.12–1.72). Farmers appeared to be more vulnerable than workers, children, students and other occupations.

Sensitivity analyses. Sensitivity analyses were conducted to check whether our coefficient estimates were robust. The effects changed little when using the full-year data instead of the data in flood season (Supplementary Figure S1). When changing df (2–8) for precipitation, relative humidity, time and week of year, we found that the effect estimates at lag 1 period did not change substantially (Supplementary Figure S2). Similar effects of floods on bacillary dysentery were observed when using different models (e.g. single lag model, unconstrained DLNM model, and constrained DLNM model) (Supplementary Figure S3). Supplementary Figure S4 shows that model residuals were independent over time with a normal distribution. Supplementary Figure S5 is the ACF and PACF plots of residuals, which showed that there was no apparent autocorrelation of model residuals.

Discussion

In recent years, bacillary dysentery has been recognized as a significant infectious disease related to climate change. Our study has quantified the lagged and cumulative effects of floods on the risk of bacillary dysentery in Huaihua, China using a distributed lag non-linear model. After controlling for the meteorological factors,

Category	Lag0	Lag1	Lag2	Lag3	Lag4	Lag0–2
Total	1.09(0.88,1.35)	1.32(1.12,1.56)*	1.06(0.89,1.25)	0.85(0.71,1.02)	1.02(0.86,1.20)	1.52(1.08,2.12)*
Gender						
Males	1.00(0.75,1.31)	1.30(1.06,1.61)*	1.04(0.84,1.29)	0.84(0.67,1.05)	1.04(0.85,1.27)	1.35(0.88,2.07)
Females	1.18(0.89,1.55)	1.32(1.06,1.65)*	1.06(0.85,1.33)	0.88(0.69,1.11)	0.96(0.77,1.20)	1.65(1.06,2.56)*
Age (years)						
0–4	1.00(0.74,1.37)	1.21(0.96,1.53)	0.99(0.78,1.26)	0.85(0.66,1.09)	0.82(0.64,1.05)	1.20(0.75,1.94)
5–14	1.17(0.73,1.88)	1.31(0.85,2.01)	1.07(0.72,1.60)	0.66(0.42,1.04)	1.08(0.75,1.57)	1.64(0.74,3.64)
15–64	1.03(0.78,1.37)	1.39(1.12,1.72)*	1.02(0.81,1.27)	0.84(0.66,1.05)	1.08(0.88,1.34)	1.46(0.94,2.27)
65+	0.98(0.55,1.73)	1.22(0.79,1.87)	1.35(0.90,2.01)	1.20(0.79,1.82)	1.09(0.73,1.63)	1.61(0.68,3.78)
Occupation						
Farmers	0.85(0.60,1.20)	1.42(1.11,1.82)*	1.02(0.79,1.32)	0.80(0.60,1.05)	1.07(0.83,1.37)	1.23(0.73,2.08)
Students	1.58(0.99,2.52)	1.17(0.73,1.87)	1.07(0.69,1.66)	0.71(0.44,1.14)	1.05(0.71,1.57)	1.99(0.86,4.61)
Children	0.98(0.72,1.33)	1.21(0.96,1.53)	0.97(0.77,1.23)	0.88(0.69,1.12)	0.87(0.69,1.10)	1.15(0.72,1.84)
Works	0.92(0.57,1.50)	1.07(0.72,1.58)	1.26(0.89,1.79)	0.87(0.58,1.28)	0.90(0.62,1.30)	1.25(0.58,2.67)
Others	1.59(0.91,2.81)	1.52(0.98,2.36)	1.30(0.83,2.05)	1.18(0.77,1.80)	1.28(0.84,1.97)	3.16(1.29,7.77)*

Table 2. The risk ratios of floods on bacillary dysentery at various lag weeks from the DLNM models in Huaihua, China. * $p < 0.05$.




Figure 3. The cumulative effects of floods on bacillary dysentery in Huaihua, China from 2005 to 2011.

seasonality and long-term trend, results indicate that floods may play an important role in the epidemic of bacillary dysentery. Although this study is based on Huaihua city only, the real impact of bacillary dysentery due to floods might be much greater, given the large population at risk and frequent floods in China. From 2000–2010 floods has affected more than 1.7 billion people in China. The direct economic losses caused by floods was approximately 1.39 trillion yuan. People at risk of floods may increase due to the rapid urbanization in China¹³. The results from this study might be applicable to most city in south and east China, because climates in those places were similar with that in Hunan.

An increased risk of diarrheal disease following floods has been reported all over the world. During the 1993 flooding in Brazil, the flooding was significantly associated with diarrhea⁹. A German study also showed that the major risk factor for diarrhea was contacting with floodwater¹⁴. A study from Anhui of China indicated floods were significantly associated with an increased risk of infectious diarrhea¹⁵. However, limited number of studies have examined the effects of floods on bacillary dysentery^{10,16}.

In our study, results of the DLNM show that floods were associated with an increased risk of bacillary dysentery with 3 weeks' effects after adjustment for meteorological factors, seasonality and long-term trend. However, the underlying mechanisms by which floods influences the bacillary dysentery are not yet clear. The probability of ingesting water or food contaminated by *Shigella* is likely to increase during floods. A study in Pakistan indicated that twenty percent of the drinking water samples collected during flood period were contaminated with *Shigella* and other enter pathogens including *Vibrio cholera*, *Salmonella*, *Staphylococcus aureus* and others¹⁷. Other studies also showed that contamination of drinking water was associated with water-borne disease outbreaks such as dysentery, cholera, hepatitis A, typhoid fever, and other gastrointestinal diseases after floods^{18,19}.

Continuous precipitation during floods can mobilize pathogens and transport them into the aquatic environment, increasing the microbiological agents on surface water¹⁹. Floods can also destroy the sewage systems and waste-disposal systems, washing contaminants into drinking water. These may cause the local water quality seriously deteriorated and lead to a lack of clean water and food supply. As a result, the transmission of enteric pathogens and communicable diseases may increase during floods²⁰. A study in Bangladesh also showed the flooding-induced breakdown of sanitary conditions is likely the principal mediator of the effect of climate on the infectious disease²¹.

Our study also found that the effect of floods on bacillary dysentery in the group of 15–64 was significant. This may be because that people aged 15–64 years old participated in more relief work and engaged the reconstruction work more frequently than the other groups, leading to a higher exposure in the flood period. Farmers appeared




Figure 4. Location of the study area in Hunan province, China. The map was generated using ArcGIS 10.2 (Environmental Systems Research Institute, Redlands California, America) Available at: <http://www.esri.com/software/arcgis/arcgis-for-desktop>.

to be more vulnerable than workers, children, students and other occupations. Kirsch *et al.* also demonstrates significantly worse impact and a slower recovery for rural area after the flood²². A possible reason is that rural areas usually have poor sanitation and medical conditions. Farmers usually do not have access to a flush toilet and probably not to clean water after floods, which may increase the risk of bacillary dysentery. Davis *et al.* found that rural communities are faced with a myriad of health care disparities, each posing as a barrier to timely response and complete recovery from a disaster, including insufficient public health infrastructure and disproportionate access to adequate medical care²³.

It is not clear whether the difference between genders was caused by different behaviors to flood responses. A possible explanation is that during floods females may not only participated in the reconstruction work, but also prepare the food and water, care for the elderly and children. Psychological stress from increased responsibilities may cause fatigue and increased vulnerability to diseases during and after floods. Lowe *et al.* also found that the psychological and physiological health effects of floods appear disproportionately borne by females, elderly and children during floods²⁴.

Urban areas around the world have expanded rapidly in recent years²⁵. Nearly half of the urban expansion is projected to take place in Asia, especially in China which was highly prone to flooding from rivers and coastal surge²⁶. Rapid urbanization in China has an adverse impact on urban hydrological processes, particularly in increasing the urban flood risks¹³. A study in Indonesia showed that urban expansion drives large increases in flood risk²⁷. In the rapidly urbanized China cities, urban floods may cause health risks such as breaks out of infectious diseases by causing sewer water overflow and flushing foul water to public area, especially for the immigrant rural workers with poor sanitation and health services.

Limitations of our study should be acknowledged. Firstly, some confounding factors, such as the degree of flood disaster and different population immune levels could not be included in our study. Secondly, we could only obtain the weekly disease data while the daily data may be a better choice for the analysis of lag effects. Thirdly, we only focused on one city. The results might not be generalizable to other areas, particularly for those places with different climates.

In conclusion, this study provides evidence that floods may play an important role in the epidemic of bacillary dysentery in study area. People aged 15–64 years old, females, and farmers appeared to be more vulnerable than the others. Our findings have significant implications for developing local strategies to prevent and reduce bacillary dysentery given more floods have been predicted in the future due to climate change.

Methods

Study area and period. The study was conducted in Huaihua a city located along the Yaun River, a tributary of the Yangtze River, in the Hunan province between latitudes 25°54' and 29°00'N and longitudes 108°48' and 111°06'E (Fig. 4). The city is generally characterized by a subtropical humid monsoon climate with an annual average temperature of 16.4 °C and an annual average rainfall of 1600 mm. Huaihua has an area of 27,564 square kilometers and a population of 4.74 million in 2010. Given the seasonal distribution of floods and bacillary dysentery in Huaihua, periods between April and September (i.e. the flood season) from 2005–2011 were chosen as the main study periods.

Data collection and management. *Disease surveillance data.* Weekly number of bacillary dysentery cases from 2005–2011 were obtained from the National Notifiable Disease Surveillance System (NDSS). The definition of bacillary dysentery, according to the NSDD, is a group of the human diseases that are caused by *Shigellae*, accompanied by fever, abdominal pain, tenesmus and bloody or mucus stool as the typical clinical presentation. In this study, all bacillary dysentery cases were defined based on the diagnostic criteria and principles of management for dysentery (GB 16002–1995), issued by the Ministry of Health of the People's Republic of China²⁸. Only the cases confirmed both clinically and by laboratory tests, including microscopic examination and biochemical identification, were included in our study. In China, bacillary dysentery is a statutory notifiable category B infectious disease. According to the National Communicable Disease Control Act, physicians in hospitals must report every case of bacillary dysentery to the local health authority. The local health authority must report these cases to the next level of the organization within 24 h²⁹. The Direct Network Report system for infectious diseases has been established and applied well since Jan 1, 2004 in China³⁰. Therefore, it is believed that the degree of compliance with disease notification over the study period was consistent.

Floods and meteorological data. Floods is defined as an extreme climate event with flooding and geological hazards such as debris flow, landslide after local or regional heavy rain process, it must fulfill at least one of the following criteria: (1) Ten or more people reported killed. (2) 50000 or more hectares of farmland reported damaged. (3) Cause a direct economic loss of 100 or more million Chinese yuan. Flood data was extracted from the Yearbooks of Meteorological Disasters in China, which recorded the occurrence time, number of deaths, damaged areas, and economic loss of floods³.

Weekly meteorological data over the same period were obtained from the China Meteorological Data Sharing Service System (<http://cdc.cma.gov.cn/>). The meteorological variables included weekly cumulative precipitation, weekly average temperature, and weekly average relative humidity.

Statistical analysis. A generalized linear mixed model (GLMM) combined with a distributed lag non-linear model (DLNM) was applied to quantify the distributed lag effects of floods on bacillary dysentery, with weekly counts of bacillary dysentery as the dependent variable and floods as the independent variable adjusted for potential confounders. A quasi-Poisson regression was used to deal with the over dispersion of Poisson distribution. Generalized linear mixed model combined with distributed lag non-linear model is widely used in time series studies examining temperature, air pollution and health outcomes^{31–33}. To control for confounder, in study design, time series analysis of diseases and extreme weather events such as heat wave or extreme precipitation event was usually conducted in a period when most extreme event occurred^{34,35}. For our study, we chose the flood season (April–September) as our study period because almost all floods occurred in this period. The natural cubic spline used in GLMM is a flexible and effective technique for adjustment for nonlinear confounding effects of seasonality, long-term trends, and weather variables. The main advantage of DLNM is that it allows the model to describe the lag structure of exposure–response relationships, which in turn provides an estimate of the cumulative effect and delayed effect³⁶.

Potential confounders. To control for any long term trend, we used a natural cubic spline with three degrees of freedom (df) for time. We used a natural cubic spline with four df for week of year (woy) to control for any seasonal trend³⁷. Previous studies have reported that temperature, precipitation and humidity, which linked to the replication, persistence, and transmission of pathogens in the environment, were associated with diarrheal diseases^{38–40}. Therefore, we used a smooth function of natural cubic spline with three df in DLNM for mean temperature, cumulative precipitation and relative humidity to adjust for potential effects of these meteorological factors.

Lags. Due to the delayed environmental transport of pathogens and delayed onset of clinical symptoms, morbidity of bacillary dysentery was expected to peak several days after the occurrence of floods. For example, a study from China reported floods can significantly increase the risk of dysentery within one month¹⁰. Thus, the association in our model was explored across a 4-week lag. Models which allowed for lagged exposure effects can be divide into the single lag model, unconstrained DLNM model and the constrained DLNM model. The selection of our final model was based on the distribution of lagged effects, not the index of model fitting. We chose the constrained DLNM model for two reasons. Firstly, all the lagged predictors can be simultaneously entered in the model. Secondly, after imposing some constraints on the effect estimates for the different lags, collinearity was significantly reduced. Therefore, fewer parameters need to be estimated, and associations at individual lags could be estimated with a greater precision⁴¹.

The main analysis was conducted with a distributed lag model to evaluate the lag effects and cumulative effects of floods on bacillary dysentery. Distributed lag models, widely utilized in air pollution studies and temperature studies, provide a systematic way to investigate the distribution of effects over time^{31,42,43}. We constrained model coefficients using the lag number to fit a natural cubic spline function to reduce collinearity³⁶. This model can estimate the delayed and cumulative effect of floods on the morbidity of bacillary dysentery over the entire lag period simultaneously (Model 1):

Model 1:

$$\log[E(Y_t)] = \beta + \sum_{p=0}^4 \alpha_p \text{Floods}_{t-p} + \sum_{q=0}^4 \gamma_q \text{Temperature}_{t-q} + ns_1(\text{Precipitation}, 3) + ns_2(\text{Humidity}, 4) + ns_3(\text{Woy}, 4) + ns_4(\text{Time}, 3) + \text{Lag}(\text{res}, 1)$$

where Y_t denoted the weekly number of bacillary dysentery cases at time t . *Floods* were a categorical variable including non-flooded or flooded weeks and represented by 0 and 1, respectively. The α_p was the effect estimate of the floods p days before the day of illness. The γ_q was the effect estimate of temperature q days before the day of illness. The $ns_1(\text{Precipitation}, 3)$, $ns_2(\text{Humidity}, 3)$, $ns_3(\text{Woy}, 4)$, and $ns_4(\text{Time}, 3)$ were natural cubic splines of weekly cumulative precipitation, weekly average relative humidity, week of year, and time, respectively, which were designed to control the effects of meteorological factor, seasonality, and long-term trend. The *Lag (res, 1)* was the first-order lagged variable of the model residual error designed to control the autocorrelation.

Sensitivity analysis. As effect estimates vary with different choices of model selections and parameters specification, we conducted the following sensitivity analyses:

(1) using the full-year data instead of the data in flood season; (2) varying the df (2–8) for week of year to adjust for seasonality; (3) varying the df (2–8) for time to adjust for long-term trend; (4) changing the df (2–8) for relative humidity; (5) using different methods to evaluate the estimates, single lag model, unconstrained DLNM model, and constrained DLNM model (using the lag number to fit a polynomial function)⁴¹.

The level of statistical significance was set at 0.05 (two-tailed). Analyses were conducted using “dlnm” package³⁷ in R 3.1.3 (R Foundation for Statistical Computing, Vienna, Austria).

References

- Guha-Sapir, D., Vos, F., Below, R. & Ponslerre, S. Annual disaster statistical review 2010: The Numbers and Trends. Brussels: CRED. (2011). Available at: http://www.Cred.be/Sites/Default/Files/Adsr_2010.Pdf. (Accessed: 17th April 2016).
- Parry, M. L. *Climate Change: impacts, adaptation and vulnerability: contribution of Working Group II to the fourth assessment report of the Intergovernmental Panel On Climate Change*. Cambridge University Press, Cambridge, United Kingdom (2007).
- Song, L. *The Yearbook of meteorological disasters in China* [in Chinese]. China Meteorological Press, Beijing, China (2012).
- Kosek, M., Bern, C. & Guerrant, R. L. The global burden of diarrhoeal disease, as estimated from studies published between 1992 and 2000. *Bull World Health Organ.* **81**, 197–204 (2003).
- Data Center for Public Health. *National notifiable infectious disease database* [in Chinese]. Available at: http://www.phsciencedata.cn/Share/ky_sjml.jsp. (Accessed: 17th April 2016).
- Ahern, M., Kovats, R. S., Wilkinson, P., Few, R. & Matthies, F. Global health impacts of floods: epidemiologic evidence. *Epidemiol Rev.* **27**, 36–46 (2005).
- Alderman, K., Turner, L. R. & Tong, S. Floods and human health: a systematic review. *Environ Int.* **47**, 37–47 (2012).
- Wade, T. J. *et al.* Did a severe flood in the Midwest cause an increase in the incidence of gastrointestinal symptoms? *Am J Epidemiol.* **159**, 398–405 (2004).
- Heller, L., Colosimo, E. A. & Antunes, C. M. Environmental sanitation conditions and health impact: a case-control study. *Rev Soc Bras Med Trop.* **36**, 41–50 (2003).
- Ni, W. *et al.* Effects of the floods on dysentery in north central region of Henan Province, China from 2004 to 2009. *J Infection.* **69**, 430–439 (2014).
- Kondo, H. *et al.* Post-flood–infectious diseases in Mozambique. *Prehosp Disaster Med.* **17**, 126–133 (2002).
- Milojevic, A. *et al.* Health effects of flooding in rural Bangladesh. *Epidemiology.* **23**, 107–115 (2012).
- Chen, Y., Zhou, H., Zhang, H., Du G. & Zhou, J. Urban flood risk warning under rapid urbanization. *Environ Res.* **139**, 3–10 (2015).
- Schnitzler, J., Benzler, J., Altmann, D., Mucke, I. & Krause, G. Survey on the population's needs and the public health response during floods in Germany 2002. *J Public Health Manag Pract.* **13**, 461–464 (2007).
- Ding, G. *et al.* Quantitative analysis of burden of infectious diarrhea associated with floods in northwest of anhui province, china: a mixed method evaluation. *PLoS One.* **8**, e65112 (2013).
- Ni, W., Ding, G., Li, Y., Li, H. & Jiang, B. Impacts of floods on dysentery in Xinxiang city, China, during 2004–2010: a time-series Poisson analysis. *Glob Health Action.* **7**, 23904 (2014).
- Farhat, A. K., Javed, A. & Riaz, U. & Sultan, A. Bacteriological quality assessment of drinking water available at the flood affected areas of Peshawar. *Toxicological & Environmental Chemistry.* **95**, 1448–1454 (2013).
- McMichael, A. J., Woodruff, R. E. & Hales, S. Climate change and human health: present and future risks. *Lancet.* **367**, 859–869 (2006).
- Patz, J. A., Vavrus, S. J., Uejio, C. K. & McLellan, S. L. Climate change and waterborne disease risk in the Great Lakes region of the US. *Am J Prev Med.* **35**, 451–458 (2008).
- CDC. Morbidity and mortality associated with Hurricane Floyd–North Carolina, September–October 1999. *MMWR Morb Mortal Wkly Rep.* **49**, 369–372 (2000).
- Reiner, R. J. *et al.* Highly localized sensitivity to climate forcing drives endemic cholera in a megacity. *Proc Natl Acad Sci USA.* **109**, 2033–2036 (2012).
- Kirsch, T. D., Wadhvani, C., Sauer, L., Doocy, S. & Catlett, C. Impact of the 2010 Pakistan floods on rural and urban populations at six months. *PLoS Curr.* **4**, e4f-e212f (2012).
- Davis, J. R., Wilson, S., Brock-Martin, A., Glover, S. & Svendsen, E. R. The impact of disasters on populations with health and health care disparities. *Disaster Med Public Health Prep.* **4**, 30–38 (2010).
- Lowe, D., Ebi, K. L. & Forsberg, B. Factors increasing vulnerability to health effects before, during and after floods. *Int J Environ Res Public Health.* **10**, 7015–7067 (2013).
- Seto, K. C., Fragkias, M., Guneralp, B. & Reilly, M. K. A meta-analysis of global urban land expansion. *PLoS One.* **6**, e23777 (2011).
- Jongman, B., Ward, P. J. & Aerts, J. C. J. H. Global exposure to river and coastal flooding: Long term trends and changes. *Global Environmental Change.* **22**, 823–835 (2012).
- Muis, S., Guneralp, B., Jongman, B., Aerts, J. C. & Ward, P. J. Flood risk and adaptation strategies under climate change and urban expansion: A probabilistic analysis using global data. *Sci Total Environ.* **538**, 445–457 (2015).
- State Bureau of Technical Supervision of China. Ministry of Health of the People's Republic of China. GB 16002–1995. Diagnostic criteria and principles of management of bacillary and amebic dysentery [in Chinese]. Standards Press of China, (1995).
- Ministry of Health of China, Emergency events and regulation of the notifiable disease surveillance system [in Chinese]. 2006. Available at: http://www.gov.cn/zwqk/2006-09/08/content_382018.htm (Accessed: 17th April 2016).
- Wu, X., Fu, Z., Deng, G. & Zhou, J. Analysis on the operational status of Direct Network Report system for infectious diseases [in Chinese]. *Applied Prev Med.* **19**, 26–28 (2013).
- Gasparrini, A. & Armstrong, B. The impact of heat waves on mortality. *Epidemiology.* **22**, 68–73 (2011).
- Welty, L. J. & Zeger, S. L. Are the acute effects of particulate matter on mortality in the National Morbidity, Mortality, and Air Pollution Study the result of inadequate control for weather and season? A sensitivity analysis using flexible distributed lag models. *Am J Epidemiol.* **162**, 80–88 (2005).
- Zeng, W. *et al.* The effect of heat waves on mortality and effect modifiers in four communities of Guangdong Province, China. *Sci Total Environ.* **482–483**, 214–221 (2014).
- Cheng, J. *et al.* Associations between extreme precipitation and childhood hand, foot and mouth disease in urban and rural areas in Hefei, China. *Sci Total Environ.* **497–498**, 484–490 (2014).
- Son, J. Y., Lee, J. T., Anderson, G. B. & Bell, M. L. The impact of heat waves on mortality in seven major cities in Korea. *Environ Health Perspect.* **120**, 566–571 (2012).
- Gasparrini, A., Armstrong, B. & Kenward, M. G. Distributed lag non-linear models. *Stat Med.* **29**, 2224–2234 (2010).
- Gasparrini, A. Distributed Lag Linear and Non-Linear Models in R: The Package dlnm. *J Stat Softw.* **43**, 1–20 (2011).
- Chou, W. C. *et al.* Modeling the impact of climate variability on diarrhea-associated diseases in Taiwan (1996–2007). *Sci Total Environ.* **409**, 43–51 (2010).
- Naumova, E. N. *et al.* Seasonality in six enterically transmitted diseases and ambient temperature. *Epidemiol Infect.* **135**, 281–292 (2007).
- Onozuka, D., Hashizume, M. & Hagihara, A. Effects of weather variability on infectious gastroenteritis. *Epidemiol Infect.* **138**, 236–243 (2010).
- Bhaskaran, K., Gasparrini, A., Hajat, S., Smeeth, L. & Armstrong, B. Time series regression studies in environmental epidemiology. *Int J Epidemiol.* **42**, 1187–1195 (2013).
- Schwartz, J. The distributed lag between air pollution and daily deaths. *Epidemiology.* **11**, 320–326 (2000).
- Zanobetti, A., Wand, M. P., Schwartz, J. & Ryan, L. M. Generalized additive distributed lag models: quantifying mortality displacement. *Biostatistics.* **1**, 279–292 (2000).

Acknowledgements

This study was supported by the National Basic Research Program of China (973 Program) (Grant No. 2012CB955502) and Natural Science Foundation of Shandong Province for the Joint Specific Program (Grant No. ZR2015HL100).

Author Contributions

Z.-D.L., J.L. and B.-F.J. initiated the study. Q.-Y.L. and B.-F.J. collected the data. Z.-D.L. cleaned the data and performed statistical analysis. Z.-D.L., J.L. and B.-F.J. drafted the manuscript. Y.Z., G.-Y.D., X.X., L.G. and X.-N.L. revised the manuscript. All authors read and approved the final manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Liu, Z.-D. *et al.* Distributed lag effects and vulnerable groups of floods on bacillary dysentery in Huaihua, China. *Sci. Rep.* **6**, 29456; doi: 10.1038/srep29456 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>