SCIENTIFIC REPORTS

OPEN

Relationships between functional diversity and aboveground biomass production in the Northern Tibetan alpine grasslands

Received: 10 March 2016
Accepted: 06 September 2016
Published: 26 September 2016

Juntao Zhu¹, Lin Jiang² & Yangjian Zhang^{1,3}

Functional diversity, the extent of functional differences among species in a community, drives biodiversity–ecosystem function (BEF) relationships. Here, four species traits and aboveground biomass production (ABP) were considered. We used two community-wide measures of plant functional composition, (1) community weighted means of trait values (CWM) and (2) functional trait diversity based on Rao's quadratic diversity (FD_Q) to evaluate the effects of functional diversity on the ABP in the Northern Tibetan alpine grasslands. Both species and functional diversity were positively related to the ABP. Functional trait composition had a larger predictive power for the ABP than species diversity and FD_Q , indicating a primary dependence of ecosystem property on the identity of dominant species in our study system. Multivariate functional diversity was ineffective in predicting ecosystem function due to the trade-offs among different traits or traits selection criterions. Our study contributes to a better understanding of the mechanisms driving the BEF relationships in stressed ecosystems, and especially emphasizes that abiotic and biotic factors affect the BEF relationships in alpine grasslands.

The rapid decline of global biodiversity has motivated considerable research directed towards understanding its potential consequences for ecosystem functioning¹. While much work has focused on species diversity as an important dimension of biodiversity², it has been increasingly recognized that functional diversity, the extent of functional differences among species in a community³, not the taxonomic richness, ultimately drives biodiversity–ecosystem functioning (BEF) relationships^{4–6}. There is now extensive evidence that species traits are better at capturing the functional characteristics of a community and that the use of species traits in ecology significantly contributes to achieving a predictive framework for ecosystem functioning^{7–10}.

Effects of functional traits on ecosystem properties have been quantified by two conceptually different approaches¹¹. On the one hand, community-weighted means of trait values (CWM) are calculated as mean trait values weighted by species relative abundances in a community¹², and are consequently closely related to the “mass ratio hypothesis”¹³, which proposes that ecosystem processes are strongly influenced by the functional traits of dominant species in a community. The CWM is therefore also linked to the sampling or selection effects associated with the greater chance of including highly productive species in more diverse communities¹⁴. On the other hand, a number of continuous measures have been developed which assess functional trait diversity of a community by quantifying the distribution of trait values among species¹⁵. Rao's quadratic diversity FD_Q is the sum of pairwise functional distances between species weighted by their relative abundances. It reaches a maximum value when functionally different species, i.e. those with large trait differences, reach similarly high abundances¹⁶. Defined as such, FD_Q is related to facilitation and/or complementary resource use among species¹⁷.

Recent studies have shown that, CWM^{18,19}, FD_Q ^{20,21}, or a combination of CWM and FD_Q ^{22–24}, can explain variation in ecosystem functioning. However, these results were mainly obtained by using experimentally created assemblages^{25,26}, a method being criticized on the artificiality of the communities created^{27–29}. In all, we still lack knowledge about relationships between functional diversity and ecosystem functioning for natural ecosystems,

¹Lhasa Plateau Ecosystem Research Station, Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing, 100101, China. ²School of Biology, Georgia Institute of Technology, Atlanta, GA, 30332, USA. ³CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing 100101, China. Correspondence and requests for materials should be addressed to J.Z. (email: zhujt@igsnr.ac.cn)

Figure 1. Distribution of the 63 study sites along the Northern Tibetan Plateau Alpine Grassland Transect. The national and county boundary datasets in shapfile format were downloaded from the National Geomatics Center of China (<http://www.ngcc.cn/>) freely. This figure was exported in a JPEG format from ArcGIS 9.3 software (<http://www.esri.com/software/arcgis/index.html>).

raising the question of whether the various species diversity-productivity relationships found in nature³⁰ would apply to functional diversity.

Ecosystems are subject to natural environmental conditions with temporal and spatial variations, such as temperature, precipitation and nutrient availability³¹, as well as to influences from other species and human activities^{25,32}. These effects vary in their frequencies and intensities, including regularly recurring variations to which organisms living in a given environment are more or less adapted, and episodic, catastrophic disturbances that lead to extensive mortality and local species extinction³³. Currently, a few studies^{34–36} have been done in BEF relationships under these stressful conditions. The small number of studies on this topic have reported contrasting results, ranging from clearly positive to no or in some circumstances even negative effects of diversity on ecosystem functioning under environmental stress³⁷.

In the Northern Tibetan plateau, there are three main natural vegetation types, alpine meadow, alpine steppe and alpine desert steppe³⁸, and most of plants are perennial herbs. The vegetation in this area is exposed to extreme environmental conditions, including intense radiation, strong winds, low temperatures, low soil nutrients and drought stress^{38,39}. The northern Tibetan ecosystem is an ideal site to evaluate relationships between functional diversity and ecosystem functioning for natural ecosystems under stressful conditions. In this paper, we selected four species traits, plant height (*H*), plant coverage (*C*), leaf mass per area (*LMA*), leaf dry matter content (*LDMC*), and aboveground biomass production (*ABP*). Firstly, we tested the hypotheses that functional diversity could explain more variation in ecosystem functioning than species diversity. Secondly, we related aboveground biomass production and trait-based indices to assess whether *CWM*, i.e. functional identity of dominant species, or *FD_Q*, i.e. functional dissimilarity among species, are better predictors for ecosystem functioning.

Materials and Methods

Study sites. The Northern Tibetan Plateau (locally named Changtang) is located in the hinterland of the Qinghai–Tibet Plateau (29°53′–36°32′ N; 78°41′–92°16′ E), covering an area of 597,000 km² (Fig. 1). We set up a west–east alpine grassland transect (the Northern Tibetan Plateau Alpine Grassland Transect, NTPAGT) in May 2009⁴⁰. The NTPAGT covers longitudes from 79.71 to 92.03° E and latitudes from 30.50 to 33.45° N, and was approximately 1,200 km long and 400 km wide³⁸. The mean annual precipitation (MAP) decreases from east of 550 mm to west of 60 mm and mean annual temperature (MAT) ranges from −2.3 °C to 1.2 °C (a more sophisticated description of the environmental settings see the climate diagrams in Supplement, Figs S1 and S2). The elevation of the sample sites ranges from 4374 to 4953 m. The growing season in this region usually begins in May and ends in September, with 65 to 85% of precipitation occurring during this period. The zonal alpine grassland types and their aboveground biomass production follow the general climate pattern⁴¹.

The NTPAGT traverses three main natural vegetation types: alpine meadow, alpine steppe and alpine desert steppe³⁸. Vegetation in alpine meadow is dominated by *Kobresia pygmaea*, associated with *Potentilla saundersiana*, *Potentilla cuneata*, *Stipa purpurea* and *Festuca coelestis*. The alpine meadow is located in the eastern Changtang, where an alpine semi-humid climate dominates. Alpine steppe, dominated by *S. purpurea*, *Stipa capillacea* and

Stipa subsessiliflora var. *basiplumosa*, associated with *Kobresia humilis*, *Carex moorcroftii*, *Leontopodium nanum*, *Oxytropis microphylla*. The alpine steppe is widely distributed in the middle Changtang, where an alpine semi-arid climate dominates. Alpine desert steppe (dominated by *S. purpurea*, *Ceratoides lateens* and *Stipa glareosa*) is scattered across the western and northwestern Changtang, where the climate is an alpine arid type⁴⁰.

Species abundance, aboveground biomass production and leaf traits. Field surveys were conducted during late July to early August in 2011 and 2012. Sixty-two fenced sites (310 plant quadrats) were surveyed one time across the growing season along the transect (Fig. 1). There were 18, 28 and 16 sites in the alpine meadow, alpine steppe and alpine desert steppe system, respectively. We recorded geographical coordinates, elevation, and vegetation type for each site. Five 1 m × 1 m quadrats were laid out randomly within each 100 m × 100 m site and all vascular plant species were recorded. The number of species and individual plants (genets and ramets), individual plant height, coverage and weight, and canopy coverage were measured in each quadrat. We harvested the aboveground biomass down to the soil surface and sorted the individuals by species. The major sampled species reached peak coverage usually during the field period (late July to early August). Therefore, the aboveground biomass in this region can serve as a surrogate for ANPP⁴². Aboveground biomass was weighed after removing dead parts and being oven-dried at 65 °C for 72 hours to a constant weight.

According to the standard measurement methods of plant traits⁴³, we selected four plant traits associated with productivity: plant height (*H*), plant coverage (*C*), leaf mass per area (*LMA*) and leaf dry matter content (*LDMC*). Five dominant species in each quadrat were measured, with the biomass of these species accounting for more than 80% of the total biomass. Twenty intact leaves were randomly selected for each species. We scanned the leaf area, and oven-dried leaves at 65 °C for 72 hours to a constant weight. Leaf area was calculated using Sigmascan 4.1.

Functional diversity and Shannon-Weaver index. The community-weighted mean trait values (*CWM*) for each trait were calculated for each plant quadrat (*n* = 310), following Garnier *et al.*¹⁸:

$$CWM = \sum_{i=1}^S p_i \times t_i \quad (1)$$

where *S* is the number of species in a community, *p_i* is the relative abundance of species *i* and the biomass for each species as the relative abundance, *t_i* is the species-specific trait value.

The Rao's functional diversity index (*FD_Q*) was calculated using the Excel-macro developed by Lepš *et al.*⁴⁴ according to the equation:

$$FD_Q = \sum_{i=1}^S \sum_{j=1}^S p_i \times p_j \times d_{ij} \quad (2)$$

where *S* is the number of species in a community, *p_i* and *p_j* are the relative abundances of species *i* and *j*, and *d_{ij}* is the trait distance between species *i* and *j* in a community.

The Shannon-Weaver index (*H*) was calculated for each community using the equation (Shannon & Weaver 1949)⁴⁵:

$$H = - \sum_{i=1}^S p_i \times \ln(p_i) \quad (3)$$

where *S* is the number of species in a community, *p_i* is the relative abundances of species.

Single- and multi-trait indices contained unique information about functional composition of the communities, and both are likely to have a place in predicting variations in ecosystem functions under different scenarios⁴⁶. So we also calculated multi-trait functional diversity as functional dispersion (*FD_{is}*). In multidimensional trait space, *FD_{is}* is the mean distance of each species, weighted by its relative abundances, to the centroid of all species in a community⁴⁷. *CWM*, *FD_Q* for each trait individually and *FD_{is}* for all traits in combination were calculated using the *FD* package⁴⁸ in R version 3.1.1 (R Development Core Team 2014).

Statistical analysis. Data on species traits, species diversity and aboveground biomass production were log transformed prior to analysis to meet the assumptions of normality. A general linear regression analyses was used to examine relationships between diversity and aboveground biomass production. According to the major plant composition, we divided our 310 sample plots into five plant functional groups, including (group I, 57 plots) *Stipa purpurea* - *Stipa subsessiliflora* - *Carex moorcroftii*; (group II, 63 plots) *Stipa purpurea* - *Artemisia duthreuil-de-rhinsi*; (group III, 70 plots) *Stipa purpurea* - *Carex moorcroftii*; (group IV, 60 plots) *Kobresia pygmaea* - *Stipa purpurea* - *Leontopodium ochroleucum*; (group V, 60 plots) *Kobresia pygmaea* - *Potentilla saundersiana*, respectively (Table 1). The mean annual precipitation (MAP) of five groups is 60–150, 150–260, 280–360, 360–440, and 440–550 mm, respectively. We assess the BEF relationships for each plant functional group. Multiple stepwise regression method was used to determine the major traits affecting ecosystem functions. All statistical analyses were performed using the R statistical package version 3.1.1 (R Core Team 2014).

Results

For the Northern Tibetan alpine grasslands, aboveground biomass production (*ABP*) significantly increased with species richness (*SR*) and Shannon-Weaver index (*H*) (*P* < 0.001, Fig. 2). The *ABP* significantly increased with community-weighted means of plant coverage (*CWM_C*), leaf mass per area (*CWM_{LMA}*) and leaf dry matter content (*CWM_{LDMC}*) (*P* < 0.001), but significantly decreased with community-weighted means of plant height

Groups	Plant composition	Grassland types	Altitude (m)	Sample number	Longitude (°)	Latitude (°)
Group I	<i>Stipa purpurea</i> - <i>Stipa subessiflora</i> - <i>Carex moorcroftii</i>	Alpine desert steppe	4374–4814	57	80.38–84.05	32.08–32.51
Group II	<i>Stipa purpurea</i> - <i>Artemisia duthreuil-de-rhinsi</i>	Degraded alpine steppe	4435–4953	63	84.26–87.09	31.75–32.26
Group III	<i>Stipa purpurea</i> - <i>Carex moorcroftii</i>	Alpine steppe	4533–4803	70	86.80–90.08	31.50–32.18
Group IV	<i>Kobresia pygmaea</i> - <i>Stipa purpurea</i> - <i>Leontopodium ochroleucum</i>	Degraded alpine meadow	4548–4788	60	89.35–91.72	31.36–31.86
Group V	<i>Kobresia pygmaea</i> - <i>Potentilla saundersiana</i>	Alpine meadow	4537–4788	60	91.72–92.01	31.22–31.94

Table 1. According to the major plant composition, we divided 310 sample plots into five plant functional groups.

Figure 2. Relationships between species richness (A), Shannon-Weaver index (B) and aboveground biomass production. Lines show the fitted lg–lg relationships. n is the sample size (number of plots). Asterisks indicate significance: ** $P < 0.001$; * $P < 0.01$; NS = not significant.

(CWM_H) ($P < 0.001$, Fig. 3). CWM_{LMA} , CWM_C and CWM_{LDMC} had a higher explanatory power on the ABP than CWM_H (Fig. 3). Multiple regression analyses showed that the ABP mainly depended on CWM_{LMA} , CWM_C and CWM_{LDMC} ($P < 0.001$, $R^2 = 0.82$, Table 2). CWM explained a larger proportion of variation in the ABP than did SR and H , except for CWM_H (Figs 2 and 3).

The ABP significantly decreased with Rao's functional diversity index of plant height (FD_{QH}) ($P < 0.001$, Fig. 4A), but significantly increased with Rao's index of plant coverage (FD_{QC}) ($P < 0.001$, Fig. 4B). However, FD_Q did not explain a larger amount of variation in the ABP (20% and 11%) than SR and H (26% and 12% of explained variation respectively). No significant relationships were found between Rao's index of leaf mass per area (FD_{QLMA}), leaf dry matter content (FD_{QLDMC}) and the ABP (Fig. 4C,D). When all traits were considered together, no significant relationships between functional diversity and ecosystem functions were identified (Fig. 5).

For different plant functional groups, the ABP significantly increased with SR in group I, II and IV ($P < 0.001$), but no significant relationships were found in group III and V (Fig. S3). The ABP in group II significantly increased with H ($P < 0.001$), but no significant relationships were found in other groups (Fig. S4). The ABP significantly increased with CWM_H in group I ($P < 0.001$), but significantly decreased in group III and IV ($P < 0.001$) and no significant relationships in group II and IV (Fig. S5). The ABP significantly increased with CWM_C in group I, III and IV ($P < 0.001$), but no significant relationships in group II and V (Fig. S6). The ABP significantly increased with CWM_{LMA} in group II, III, IV and V ($P < 0.001$), but no significant relationships in group I (Fig. S7). The ABP significantly increased with CWM_{LDMC} in all five groups ($P < 0.001$, Fig. S8). No significant relationships between FD_Q and the ABP were identified for most of the groups (Figs S9–12). For group I, II and V, CWM_{LDMC} had a higher explanatory power on the ABP than other three traits (Fig. S8A,B and E). For group III and IV, CWM_{LMA} explained a larger proportion of variation in the ABP than other three traits (Fig. S7C,D). Multiple regression analyses showed that the ABP in group I, II and V mainly depended on CWM_{LDMC} and CWM_C , and the ABP in group III and IV mainly depended on CWM_{LMA} and CWM_C (Table 2).

Figure 3. Relationships between community-weighted means of four traits (CWM_H , CWM_C , CWM_{LMA} and CWM_{LDMC} ; (A–D) and aboveground biomass production. Lines show the fitted lg–lg relationships. CWM_H , CWM_C , CWM_{LMA} and CWM_{LDMC} indicate community-weighted means of plant height, plant coverage, leaf mass per area and leaf dry matter content, respectively. n is the sample size (number of plots). Asterisks indicate significance levels: ** $P < 0.001$; * $P < 0.01$; NS = not significant.

Groups	Regression equations	R^2	P
Total	$ABP = -0.346 + 0.609CWM_{LMA} + 0.565CWM_C + 0.503CWM_{LDMC}$	0.82	$P < 0.001$
Group I	$ABP = 0.887 + 0.615CWM_{LDMC} + 0.459CWM_C$	0.73	$P < 0.001$
Group II	$ABP = 1.715 + 1.084CWM_{LDMC}$	0.79	$P < 0.001$
Group III	$ABP = -1.651 + 1.077CWM_{LMA} + 0.506CWM_C$	0.76	$P < 0.001$
Group IV	$ABP = -1.557 + 1.013CWM_{LMA} + 0.534CWM_C$	0.75	$P < 0.001$
Group V	$ABP = 1.722 + 0.903CWM_{LDMC} + 0.274CWM_C$	0.76	$P < 0.001$

Table 2. Stepwise regression equations of community aboveground biomass production (ABP) and species diversity, functional diversity. CWM_C , CWM_{LMA} and CWM_{LDMC} , indicated community weighted means of plant coverage, leaf mass per area, leaf dry matter content, respectively. Group I, II, III, IV, V, indicate five plant functional groups, respectively.

Discussion

It is widely reported that species diversity affects ecosystem functioning². Consistent with this general pattern, we found a positive relationship between species richness (SR), Shannon-Weaver index (H) and aboveground biomass production (ABP). Our study also demonstrated that community-weighted mean values (CWM) of species traits explain a larger proportion of variation in the ABP than SR and H , thereby extending the results of previous related studies^{4–6} to the alpine grassland ecosystem. Collectively, we further evidenced that functional diversity, not the taxonomic richness, ultimately drives BEF relationships³. Alpine habitats are well known for their severe physical living conditions⁴⁹, where plants are confronted by low temperatures, excessive radiation, strong winds, low soil nutrients, unstable substrates and short growing seasons^{38,39}. In accord with previous findings^{34,35}, our study indicated that both species diversity and functional diversity were positively related to ecosystem property, and evidenced positive effects of biodiversity on ecosystem functioning under stressful conditions.

We examined relationships between functional diversity based on CWM, FD_Q and ecosystem properties in natural alpine grasslands. Our results showed that the variation of ABP is better explained by CWM, indicating a

Figure 4. Relationships between Rao's functional diversity index of four traits (FD_{QH} , FD_{QC} , FD_{QLMA} and FD_{QLDMC} ; (A–D) and aboveground biomass production. Lines show the fitted lg–lg relationships. FD_{QH} , FD_{QC} , FD_{QLMA} and FD_{QLDMC} indicate Rao's functional diversity index of plant height, plant coverage, leaf mass per area and leaf dry matter content, respectively. n is the sample size (number of plots). Asterisks indicate significance level: ** $P < 0.001$; * $P < 0.01$; NS = not significant.

Figure 5. Relationships between multivariate functional diversity of plant height, plant coverage, leaf mass per area, leaf dry matter content and aboveground biomass production. n is the sample size (number of plots). Asterisks indicate significance level: ** $P < 0.001$; * $P < 0.01$; NS = not significant.

primary dependence of ecosystem properties on the identity of dominant species and their functional traits following the 'mass ratio hypothesis'^{13,19} in alpine grasslands. Indeed, functional identity attributed to the selection effect¹⁴, usually expressed as CWM for species traits, have been demonstrated to be a key predictor of ecosystem functioning at different scales^{18,22}. Regardless of the varying dominant species in each grassland type and functional group, coverage and biomass of the dominant species account for 40–80% of the total in each community (sample plots data), which lends further support on the above conclusion.

Beyond functional identity, functional diversity (FD_Q) has been shown to be associated with ecosystem functioning^{25,50}, indicating the importance of niche complementarity in facilitating ecosystem processes⁵¹. However, our analyses did not find that FD_Q explain a larger amount of variation in the ABP than SR and H , suggesting a lack of the operation of the complementarity effects in our study system. In our study area, soil moisture and nutrients are in shortage for belowground plant parts⁵², which may have led to strong competitive interactions for below-ground resources (nitrogen and water), resulting in weakened complementary resource use among plant species.

For the alpine grasslands, species traits, e.g., leaf mass per area (LMA), plant coverage (C) and leaf dry matter content ($LDMC$) significantly contribute to achieving a predictive framework for ecosystem functioning^{7–10}. Specifically, our analyses identified LMA , which is related to resource acquisition and plant growth strategy^{53,54}, as a key functional predictor of ecosystem functioning. Our results were in accordance with previous studies suggesting that LMA can be used to predict productivity and carbon storage^{11,55}. The leaf economic spectrum framework predicts that low LMA should promote productivity⁷. However, alpine grasslands had their maximum ABP at high LMA . Plants with high LMA could be adapted to low temperatures and high irradiance, and tend to have thick, leathery leaves^{56,57}. Although plant height has been used to predict root/shoot ratio in alpine grasslands⁵⁸, our results found that there was no significant relationship between plant height and ecosystem functioning.

For different plant functional groups, such as group II and IV, community-weighted means of leaf dry matter content (CWM_{LDMC}) and leaf mass per area (CWM_{LMA}), plant coverage (CWM_C) significantly contribute to achieving a predictive framework for the ABP in group II (degraded alpine steppe) and group IV (degraded alpine meadow), respectively. Hence, we should select the dominant species with higher leaf dry matter content ($LDMC$) to maintain high productivity for restoring degenerative alpine steppe. For the degraded alpine meadow, the dominant species with higher leaf mass per area (LMA) and plant coverage (C) should be selected to maintain high productivity.

When all traits were considered together (multivariate functional diversity), no significant relationships between functional diversity and ecosystem functions were identified in this study. This result supports that in a complex landscape with multiple environmental gradients, for example our alpine grassland transect, including three vegetation types and multiple gradients (precipitation, elevation, and soil nutrients, etc.), variation in a single trait can explain more variation in ecosystem functioning than functional diversity calculated based on multiple traits⁴⁶. In addition, plant traits are positively (e.g. leaf mass per area) or negatively (e.g. plant height) related to the ABP , that is trade-offs among different traits could render diverse communities less capable of providing multiple functional diversity in our study. Therefore, the trade-offs among different traits or traits selection criterions must be considered when multivariate functional diversity is used to predict ecosystem function.

References

- Loreau, M. *et al.* Biodiversity and ecosystem functioning: current knowledge and future challenges. *Science* **294**, 804–808 (2001).
- Tilman, D. & Snell-Rood, E. C. Diversity breeds complementarity. *Nature* **515**, 44–5 (2014).
- Tilman, D. Functional diversity. In: *Encyclopedia of Biodiversity* (ed. Levin, S. A.). Academic Press, San Diego, CA, pp. 109–120 (2001).
- Cadotte, M. W. *et al.* Using phylogenetic, functional and trait diversity to understand patterns of plant community productivity. *PLoS ONE* **4**, e5695 (2009).
- Flynn, D. F. B. *et al.* Loss of functional diversity under land use intensification across multiple taxa. *Ecol. Lett.* **12**, 22–33 (2009).
- Milcu, A. *et al.* Functionally and phylogenetically diverse plant communities key to soil biota. *Ecology* **94**, 1878–1885 (2013).
- Wright, I. J. *et al.* The world-wide leaf economics spectrum. *Nature* **428**, 821–827 (2004).
- Violle, C. *et al.* Let the concept of trait be functional. *Oikos* **116**, 882–892 (2007).
- Reiss, J. *et al.* Emerging horizons in biodiversity and ecosystem functioning research. *Trends Ecol. Evol.* **24**, 505–514 (2009).
- Osnas, J. L. *et al.* Global leaf trait relationships: mass, area, and the leaf economics spectrum. *Science* **340**, 741–744 (2013).
- Roscher, C. *et al.* Using plant functional traits to explain diversity–productivity relationships. *PLoS ONE* **7**, e36760 (2012).
- Shipley, B. Vile, D. & Garnier, E. From plant traits to plant communities: a statistical mechanistic approach to biodiversity. *Science* **314**, 812–814 (2006).
- Grime, J. P. Benefits of plant diversity to ecosystems: immediate, filter and founder effects. *J. Ecol.* **86**, 902–910 (1998).
- Huston, M. A. Hidden treatments in ecological experiments: re-evaluating the ecosystem function of biodiversity. *Oecologia* **110**, 449–460 (1997).
- Petchev, O. L. & Gaston, K. J. Functional diversity: back to basics and looking forward. *Ecol. Lett.* **9**, 741–758 (2006).
- Mouchet, M. A. *et al.* Functional diversity measures: an overview of their redundancy and their ability to discriminate community assembly rules. *Funct. Ecol.* **24**, 867–876 (2010).
- Hooper, D. U. *et al.* Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecol. Monogr.* **75**, 3–35 (2005).
- Garnier, E. *et al.* Plant functional markers capture ecosystem properties during secondary succession. *Ecology* **85**, 2630–2637 (2004).
- Hillebrand, H. Bennet, D. M. & Cadotte, M. W. Consequences of dominance: a review of evenness effects on local and regional ecosystem processes. *Ecology* **89**, 1510–1520 (2008).
- Petchev, O. L. Hector, A. & Gaston, K. J. How do different measures of functional diversity perform? *Ecology* **85**, 847–857 (2004).
- Wacker, L. *et al.* Diversity effects in early- and mid-successional species pools along a nitrogen gradient. *Ecology* **90**, 637–648 (2009).
- Mokany, K. Ash, J. & Roxburgh, S. Functional identity is more important than diversity in influencing ecosystem processes in a temperate native grassland. *J. Ecol.* **96**, 884–893 (2008).
- Mouillot, D. *et al.* Functional structure of biological communities predicts ecosystem multifunctionality. *PLoS ONE* **6**, e17476 (2011).
- Paloma, R. B. *et al.* Diversity increases carbon storage and tree productivity in Spanish forests. *Global Ecol. Biogeogr.* **23**, 311–322 (2014).
- Tilman, D. Distinguishing between the effects of species diversity and species composition. *Oikos* **80**, 185 (1997).
- Hector, A. *et al.* Plant diversity and productivity experiments in European grasslands. *Science* **286**, 1123–1127 (1999).
- Grime, J. P. *et al.* Integrated screening validates primary axes of specialisation in plants. *Oikos* **79**, 259–281 (1997).
- Huston, M. A. *et al.* No consistent effect of plant diversity on productivity. *Science* **289**, 1255a (2000).
- Jiang, L. Wan, S. Q. & Li, L. H. Species diversity and productivity: why do results of diversity-manipulation experiments differ from natural patterns? *J. Ecol.* **97**, 603–608 (2009).

30. Mittelbach, G. G. *et al.* What is the observed relationship between species richness and productivity? *Ecology* **82**, 2381–2396 (2001).
31. Steudel, B. *et al.* Biodiversity effects on ecosystem functioning change along environmental stress gradients. *Ecol. Lett.* **15**, 1397–1405 (2012).
32. Chapin, F. S. *et al.* Consequences of changing biodiversity. *Nature* **405**, 234–242 (2000).
33. Lugo, A. E. Visible and invisible effects of hurricanes on forest ecosystems: an international review. *Austral Ecol.* **33**, 368–398 (2008).
34. Ma, W. H. *et al.* Environmental factors covary with plant diversity-productivity relationships among Chinese grassland sites. *Global Ecol. Biogeogr.* **19**, 233–243 (2010).
35. Sun, J. *et al.* Meta-analysis of relationships between environmental factors and aboveground biomass in the alpine grassland on the Tibetan Plateau. *Biogeosciences* **10**, 1707–1715 (2013).
36. Caldeira, M. C. *et al.* Species richness, temporal variability and resistance of biomass production in a Mediterranean grassland. *Oikos* **110**, 115–123 (2005).
37. Wang, Z. *et al.* Causes for the unimodal pattern of biomass and productivity in alpine grasslands along a large altitudinal gradient in semi-arid regions. *J. Veg. Sci.* **24**, 189–201 (2013).
38. Li, X. J. *et al.* Root biomass distribution in alpine ecosystems of the northern Tibetan Plateau. *Environ. Earth Sci.* **64**, 1911–1919 (2011).
39. Körner, C. The alpine life zone under global change. *Gayana Bot.* **57**, 1–17 (2000).
40. Wu, J. S. Shen, Z. X. & Zhang, X. Z. Precipitation and species composition primarily determine the diversity-productivity relationship of alpine grasslands on the Northern Tibetan Plateau. *Alpine Bot.* **124**, 13–25 (2014).
41. Zhong, L. *et al.* Assessment of vegetation dynamics and their response to variations in precipitation and temperature in the Tibetan Plateau. *Climatic Change* **103**, 519–535 (2010).
42. Ma, W. L. *et al.* Changes in individual plant traits and biomass allocation in alpine meadow with elevation variation on the Qinghai-Tibetan plateau. *Sci. China Life Sci.* **53**, 1142–1151 (2010).
43. Cornelissen, J. H. C. *et al.* A handbook of protocols for standardized and easy measurements of plant functional traits worldwide. *Ann. Bot.-London.* **51**, 335–380 (2003).
44. Lepš, J. *et al.* Quantifying and interpreting functional diversity of natural communities: practical considerations matter. *Preslia* **78**, 481–501 (2006).
45. Shannon, C. E. & Weaver, W. *The Mathematical Theory of Communication*. Urbana, Illinois, University of Illinois Press (1949).
46. Butterfield, B. J. & Suding, K. N. Single-trait functional indices outperform multi-trait indices in linking environmental gradients and ecosystem services in a complex landscape. *J. Ecol.* **101**, 9–17 (2013).
47. Spasojevic, M. J. & Suding, K. N. Inferring community assembly mechanisms from functional diversity patterns: the importance of multiple assembly processes. *J. Ecol.* **100**, 652–661 (2012).
48. Laliberté, E. & Legendre, P. A distance-based framework for measuring functional diversity from multiple traits. *Ecology* **91**, 299–305 (2010).
49. Cavieres, L. A. *et al.* Nurse effect of *Bolax gummifera* cushion plants in the alpine vegetation of the Chilean Patagonian Andes. *J. Veg. Sci.* **13**, 547–554 (2002).
50. Scherer-Lorenzen, M. Functional diversity affects decomposition processes in experimental grasslands. *Funct. Ecol.* **22**, 547–555 (2008).
51. Fargione, J. *et al.* From selection to complementarity: shifts in the causes of biodiversity-productivity relationships in a long-term biodiversity experiment. *P. Roy. Soc. Lond. B. Bio.* **274**, 871–876 (2007).
52. Zhu, J. T. *et al.* Below-ground competition drives the self-thinning process of *Stipa purpurea* populations in northern Tibet. *J. Veg. Sci.* **26**, 166–174 (2015).
53. Paquette, A. & Messier, C. The effect of biodiversity on tree productivity: from temperate to boreal forests. *Global Ecol. Biogeogr.* **20**, 170–180 (2011).
54. Swenson, N. G. *et al.* The biogeography and filtering of woody plant functional diversity in North and South America. *Global Ecol. Biogeogr.* **21**, 798–808 (2012).
55. Ruiz-Benito, P. *et al.* Diversity increases carbon storage and tree productivity in Spanish forests. *Global Ecol. Biogeogr.* **23**, 311–322 (2014).
56. Fonseca, C. R. *et al.* Shifts in trait combinations along rainfall and phosphorus gradients. *J. Ecol.* **88**, 964–977 (2000).
57. Wright, I. J. *et al.* Modulation of leaf economic traits and trait relationships by climate. *Global Ecol. Biogeogr.* **14**, 411–421 (2005).
58. Li, Y. H. Luo, T. X. & Lu, Q. Plant height as a simple predictor of the root to shoot ratio: Evidence from alpine grasslands on the Tibetan Plateau. *J. Veg. Sci.* **19**, 245–252 (2008).

Acknowledgements

We are grateful to Dr. Yanbin Jiang, Jian Tao, Xiaoyong Yuan, and Yongqi Huang for offering their help on field investigation in the Northern Tibetan Plateau. This research is supported by the 973 Program of the Ministry of Science and Technology of China (2013CB956302); National Natural Science Foundation of China (41571195); West Light Foundation of the Chinese Academy of Sciences; the Construction Service Program for Cultivating Unique Institute of the Chinese Academy of Sciences (TSYSJ05); and State Key Laboratory of Resources and Environmental Information System.

Author Contributions

J.Z. carried out this research, collected field data, drafted the manuscript, carried out data analysis, and participated in the design of the study; Y.Z. conceived of the study, designed the study, coordinated the study and L.J. helped draft the manuscript; All authors gave final approval for publication.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhu, J. *et al.* Relationships between functional diversity and aboveground biomass production in the Northern Tibetan alpine grasslands. *Sci. Rep.* **6**, 34105; doi: 10.1038/srep34105 (2016).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>