SCIENTIFIC REPORTS

OPEN

The complex evolutionary history of the tympanic middle ear in frogs and toads (Anura)

Martín O. Pereyra^{1,*}, Molly C. Womack^{2,*}, J. Sebastián Barrionuevo¹, Boris L. Blotto^{1,3}, Diego Baldo⁴, Mariane Targino³, Jhon Jairo Ospina-Sarria³, Juan M. Guayasamin^{5,6}, Luis A. Coloma^{7,8}, Kim L. Hoke², Taran Grant³ & Julián Faivovich^{1,9}

Received: 13 May 2016

Accepted: 30 August 2016

Published: 28 September 2016

Most anurans possess a tympanic middle ear (TME) that transmits sound waves to the inner ear; however, numerous species lack some or all TME components. To understand the evolution of these structures, we undertook a comprehensive assessment of their occurrence across anurans and performed ancestral character state reconstructions. Our analysis indicates that the TME was completely lost at least 38 independent times in Anura. The inferred evolutionary history of the TME is exceptionally complex in true toads (Bufonidae), where it was lost in the most recent common ancestor, preceding a radiation of >150 earless species. Following that initial loss, independent regains of some or all TME structures were inferred within two minor clades and in a radiation of >400 species. The reappearance of the TME in the latter clade was followed by at least 10 losses of the entire TME. The many losses and gains of the TME in anurans is unparalleled among tetrapods. Our results show that anurans, and especially bufonid toads, are an excellent model to study the behavioural correlates of earlessness, extratympanic sound pathways, and the genetic and developmental mechanisms that underlie the morphogenesis of TME structures.

The function of audition in frogs and toads (Anura) is primarily the perception of airborne sounds, including those involved in social communication¹. Thus, hearing in anurans is thought to be a key trait for survival and reproduction. In most anurans, perception of airborne sounds is enabled by a tympanic middle ear (TME) composed minimally of a tympanic membrane, middle ear cavity, and middle ear bone (=columella, columella auris, stapes, plectrum) that conducts sound waves from the environment to the inner ear where they are transduced into electrical signals via hair cells^{1–4}.

Among other tetrapods, a TME is absent in caecilians and salamanders^{1,5} but present in amniotes. Nevertheless, although the TME is primitively present in all extant amniotes, it is not homologous across amniote lineages, having evolved independently at least five times in turtles, lepidosaurs, archosaurs, an extinct lineage of parareptiles, and the synapsid ancestor of mammals^{6–10}. TME losses are extremely rare in amniotes. All mammals, turtles, and archosaurs possess a complete TME, and even the amphisbaenians, snakes, and lizards that have lost the tympanic membrane and middle ear cavity retain a columella¹¹, the sole exceptions being the pygopod lizard *Aprasia repens*¹² and possibly the snakes *Atractaspis* and *Xenocalamus*¹³. In contrast, loss is widespread

¹Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”- CONICET, Buenos Aires, C1405DJR, Argentina.

²Department of Biology, Colorado State University, Fort Collins, CO 80523, USA. ³Departamento de Zoología, Instituto de Biociências, Universidade de São Paulo, São Paulo, SP 05508-090, Brazil. ⁴Laboratorio de Genética Evolutiva, Instituto de Biología Subtropical (CONICET-UNaM), Facultad de Ciencias Exactas Químicas y Naturales, Universidad Nacional de Misiones, Posadas, N3300LQF, Argentina. ⁵Centro de Investigación de la Biodiversidad y Cambio Climático (BioCamb), Ingeniería en Biodiversidad y Cambio Climático, Facultad de Medio Ambiente, Universidad Tecnológica Indoamérica, Diego de Robles y Vía Interoceánica, 17-1200-841, Quito, EC170103, Ecuador.

⁶Colegio de Ciencias Biológicas y Ambientales COCIBA, Laboratorio de Biología Evolutiva, Universidad San Francisco de Quito, Campus Cumbayá, Quito, 170901, Ecuador. ⁷Centro Jambatu de Investigación y Conservación de Anfibios, Fundación Otonga, Giovanni Farina 566 y Baltra, San Rafael, Quito, Ecuador. ⁸Universidad Regional Amazónica Ikiam, Muyuna, Tena, Ecuador. ⁹Departamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, C1428EGA, Argentina. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to T.G. (email: taran.grant@ib.usp.br) or J.F. (email: julian@macn.gov.ar)

among anurans, with at least a few species of several families lacking the entire TME, a condition referred to as “earlessness”¹⁴.

Earlessness is especially common in the true toad family Bufonidae, in which the TME is completely lacking in more than 200 species. Bufonidae is one of the most diversified groups of amphibians, comprising more than 580 species in 51 genera naturally distributed over numerous ecoregions of the Americas, Africa, and Eurasia¹⁵. Several authors have noted the reduction and loss of TME structures (e.g.,^{16–20}) or morphological variations in middle ear structure (e.g.,^{1,21–23}) in bufonids, but the phylogenetic distribution of earlessness has never been studied either within Bufonidae or across Anura.

As such, the goals of this study are to explore the sequences of gains and losses of the TME evolution across all anurans and evaluate in a phylogenetic framework the patterns of diversification of TME in Bufonidae.

Materials and Methods

Tympanic middle ear morphology and data collection. Although many variations in the structure of the anuran auditory system exist, a generalized model can be described^{1–3}: the lateral-most portion of this system is composed of a highly differentiated disc of thin, non-glandular skin, termed the tympanic membrane. The rim of the tympanic membrane is attached to a cartilaginous tympanic ring, the tympanic annulus. The middle ear cavity is a diverticulum of the pharynx that opens ventrally to the buccal cavity via the Eustachian tubes. The columella contacts the tympanic membrane laterally and the otic capsule medially and is divided into three portions^{1,24,25}: (1) the pars externa plectri or extracolumella, a cartilaginous structure that contacts the tympanic membrane and often presents a slim, flattened strip of cartilage called the ascending process or pars ascendens plectri, that extends anterodorsally to contact the crista parotica of the prootic, (2) the pars media plectri or columellar shaft, an ossified, rod-shaped portion with a dilated medial end, and (3) the pars interna plectri, a mainly cartilaginous structure that is continuous with the pars media and extends posteriorly to lie medial to the operculum. The expanded medial end of the pars media and the entire pars interna constitute the stapelial footplate, which fills the rostral portion of the oval window of the otic capsule^{1,26}. The footplate is connected to the suprascapula via the columellar muscle in some species¹, although the individuality of this muscle has been questioned in at least some cases^{27,28}. The otic operculum (operculum fenestrae ovalis), found only in caudates and anurans⁵, is an ovoid element that is usually cartilaginous or sometimes partially calcified^{4,25}, that contacts the stapelial footplate and covers the caudal portion of the oval window. The operculum is present in all anurans. The opercularis muscle inserts on the operculum and originates on the suprascapular cartilage of the pectoral girdle^{1,26}.

In all observed anuran species (also see²⁹), the presence/absence of TME structures follows a consistent pattern. Absence of a medial structure is accompanied by the absence of the more lateral structures, such that absence of the columella entails absence of the tympanic annulus and tympanic membrane and absence of the tympanic annulus entails absence of the tympanic membrane but not the columella. Similarly, presence of a lateral structure is accompanied by the presence of the more medial structures, such that presence of the tympanic membrane entails presence of the tympanic annulus and columella and presence of the tympanic annulus entails presence of the columella but not the tympanic membrane. Consequently, we made the following assumptions when scoring the presence/absence of the tympanic structures (Fig. 1): (1) absence of the tympanic membrane and tympanic annulus when the columella is absent; (2) absence of the tympanic membrane when the tympanic annulus is absent; (3) presence of the tympanic annulus and columella when the tympanic membrane is present; and (4) presence of the columella when the tympanic annulus is present.

In total, we scored the condition of the TME for 556 species and 51 genera of Bufonidae, representing >94% of all described species in the family. Among the sampled species, 239 were included in Pyron’s³⁰ phylogenetic analysis. We also scored the conditions of these structures for 1860 of the 2538 non-bufonid anuran species (representing 53 families; see¹⁵) included by Pyron³⁰, as well as 147 non-bufonid anuran species not included in this analysis. The only frog family not sampled by Pyron³⁰ is the recently described Odontobatrachidae³¹. Although our outgroup sampling is not exhaustive, we included data for the vast majority of genera of all the families sampled by Pyron³⁰. Details on material examined, considerations about character coding and character states scored for each transformation series, and literature sources are listed as Supplementary Material (section S1 of the Supplementary Information).

Ancestral state reconstructions. We employed the most recent and densely sampled phylogenetic hypothesis available for Anura, that of Pyron³⁰, for ancestral state reconstruction, and we discuss bufonid species not included in Pyron’s³⁰ study but present in other analyses (e.g., well-supported results of^{32–34}). We focused original data collection primarily on Bufonidae and relied more extensively on literature accounts for non-bufonids, some of which were unclear or ambiguous about the occurrence of specific structures. In particular, taxonomic accounts often use imprecise terminology to describe the external morphology of the otic region³⁵. Consequently, we analysed the phylogenetic distribution of each of the TME structures in Bufonidae but only the columella in analyses of Anura.

To test the homology of the middle ear structures individually and explain their variation among anurans, most parsimonious ancestral state reconstructions³⁶ on Pyron’s³⁰ phylogenetic hypothesis were performed using Mesquite v3.03³⁷. We further explored alternative evolutionary scenarios for the complete loss of all TME structures within Bufonidae with maximum likelihood ancestral reconstructions using the package APE³⁸ and stochastic character mapping³⁹ using phyttools⁴⁰ in R⁴¹. We compared Akaike Information Criterion (AIC) values for a maximum likelihood model in which transition rates were allowed to vary (ARD) throughout the tree and a maximum likelihood model in which transition rates between states were equal (ER). We used the most supported transition rate from our maximum likelihood analyses (ER) to estimate the number of gains and losses across Bufonidae using stochastic character mapping. Stochastic character mapping allowed us to explore the probability of ear transitions under various evolutionary scenarios, giving us a better understanding of the likelihood of

Figure 1. Schematic representation of tympanic middle ear structures in anurans, showing the assumptions followed here for coding absence and presence of the different elements. TM, tympanic membrane; TA, tympanic annulus; CO, columella

regains throughout this family. We considered three scenarios: (1) equal transition rates, no restrictions; (2) equal transition rates and restricting the ancestor to being eared; and (3) a Dollo's model (no regains possible). We ran 10,000 simulations per scenario and counted a state change whenever nodes switched from greater than 50% support for one character state to greater than 50% support for the other character state.

Results

Tympanic middle ear evolution in Anura. The occurrence of a columella is plesiomorphic in Anura, although the sister clade of all other anurans (Ascaphidae + Leiopelmatidae) lacks this structure (see Discussion). Given that the tympanic membrane and tympanic annulus are not fossilizable structures, their occurrence in fossil material cannot be assessed, making it impossible to determine if the presence of those structures is also plesiomorphic in Anura.

The TME is completely absent in at least some species of no fewer than 20 anuran families: Ascaphidae, Alsodidae, Batrachylidae, Bombinatoridae, Brachycephalidae, Brevicipitidae, Bufonidae, Calyptocephalellidae, Craugastoridae, Dicroglossidae, Hemisotidae, Leiopelmatidae, Leptodactylidae, Megophryidae, Microhylidae, Myobatrachidae, Nasikabatrachidae, Rhinophrynidae, Sooglossidae, and Telmatobiidae (section S1 of the Supplementary Information).

Ancestral character state reconstructions and detailed description of the occurrence of the columella in anuran families other than Bufonidae are provided in section S2 of the Supplementary Information. Ancestral state reconstruction using Pyron's³⁰ phylogenetic hypothesis shows that the complete loss of the TME, as evidenced by lack of the columella, occurred independently at least 25 times outside Bufonidae, plus two additional losses when taxa not included in Pyron's³⁰ study but present in other phylogenetic analyses are considered.

Tympanic middle ear evolution in Bufonidae. Based on the phylogenetic hypothesis of Pyron³⁰ and the results of both the parsimony and probabilistic ancestral state reconstructions (Fig. 2 and section S3 of the Supplementary Information), the tympanic membrane, tympanic annulus, and columella were lost in the most recent common ancestor of bufonids, regained subsequently, and then repeatedly re-lost again. Below we summarize the results of the ancestral reconstructions of the tympanic membrane, tympanic annulus, and columella (the numbers of regains and re-losses of TME structures in Bufonidae differs somewhat when taxa not sampled by Pyron³⁰ are considered; see sections S1 and S4 of the Supplementary Information and Discussion, below).

Parsimony ancestral state reconstruction. The tympanic membrane was lost in the most recent common ancestor of Bufonidae and reappeared in the sister clade of *Nannophryne*. Subsequent independent losses occurred at least 25 times (see sections S3 and S4 of the Supplementary Information). The absence of the tympanic annulus is the

Figure 2. Partial phylogenetic tree of Pyron³⁰ showing parsimony ancestral state reconstructions for the columella in Bufonidae. The absence of the columella is a synapomorphy of Bufonidae (see section S2 of the Supplementary Information), with independent regains in a subclade of *Atelopus*, *Frostius* (not included in this analysis, but see text), and the sister clade of *Nannophryne*, followed by 10 independent losses.

inferred ancestral condition in bufonids, with two independent regains: (a) within *Atelopus* in a clade composed of *A. flavescens*, *A. franciscus*, *A. pulcher*, and *A. spumarius*, and (b) in the sister clade of *Nannophryne*. The gain of the tympanic annulus in the latter clade was followed by 10 independent losses (see sections S3 and S4 of the Supplementary Information). Finally, the phylogenetic distribution of the gains and losses of the columella is identical to that of the tympanic annulus for the taxa included in the hypothesis of Pyron³⁰ (see Fig. 2 and section S4 of the Supplementary Information). However, if bufonid species not included by Pyron³⁰ are also considered, the columella sometimes occurs without a tympanic annulus (see Discussion and section S1 of the Supplementary Information).

Stochastic mapping of complete tympanic middle ear loss under various constraints. Our stochastic character mapping estimated similar patterns of TME loss and regain within Bufonidae under various evolutionary scenarios (see section S3 of the Supplementary Information). When we ran an equal rates model of evolution we found results similar to the parsimony reconstructions of the tympanic annulus and columella with strong support for an ancestor lacking these structures, two regains, and 10 losses within the tree. When we assumed the ancestor had these structures and a model of equal transition rates, we found support for 12 losses and still two regains. When restricting regains from occurring (Dollo's model), we found a total of 17 losses across bufonids.

Discussion

The lack of the columella in Ascaphidae and Leiopelmatidae, which together form the sister clade of all other extant anurans (=Lalagobatrachia), has generated much discussion about the plesiomorphic condition in Anura^{29,42–44}. However, since most proanurans and stem anuran fossils have a columella (i.e. *Mesophryne beipiaoensis*, *Notobatrachus* spp., *Prosalirus bitis*, *Triadobatrachus massinoti*, *Yizhoubatrachus macilentus*^{45–49}), the lack of this structure in Ascaphidae + Leiopelmatidae appears to be synapomorphic. A columella could not be identified in the stem anuran *Vieraella herbstii*; however, the state of preservation of the specimen is poor⁵⁰, leading some authors to consider the occurrence of a columella to be unknown (e.g.,⁴⁷) and others to consider it to be absent (e.g.,⁴³). Regardless, given the phylogenetic position of *Vieraella herbstii*^{46,47,50}, this controversy has no bearing on our inferences of the evolutionary history of this structure in anurans.

Although available evidence clearly indicates the plesiomorphic presence of the columella in Anura, it is unknown if the ancestral anuran also possessed a tympanic membrane and annulus. The tympanic membrane and tympanic annulus are not fossilizable structures, so their precise phylogenetic origin is unknown. As such, two scenarios are compatible with current evidence: the tympanic annulus and membrane might have been present in the most recent common ancestor of Anura and lost with the columella in Ascaphidae + Leiopelmatidae, or they might have arisen in Lalagobatrachia.

Among non-bufonid anurans, the TME was completely lost at least 27 times (see above). All these losses involve small clades scattered across the major lineages of Anura, implying several putative synapomorphies (e.g., *Atelognathus* + *Chaltenobatrachus*, *Brachycephalus*, Nasikabatrachidae + Sooglossidae, *Pseudophryne*, *Telmatobufo*) or autapomorphies (e.g., *Balebreviceps hilmani*, *Melanobatrachus indicus*, *Rhinophrynus dorsalis*). For other anuran clades (e.g., *Alsodes*, *Microhyla*, *Nanorana*, *Scutiger*, *Telmatobius*), denser taxon sampling is necessary to obtain adequate evidence to understand the evolution of the TME (see section S2 of the Supplementary Information for a more exhaustive discussion about the evolution of this structure in these and other non-bufonid species).

Although TME structures were lost repeatedly in Anura, TME evolution in Bufonidae is especially complex. All of the families that are closely related to Bufonidae³⁰ (also see^{51,52}) have a complete TME, making the absence of these structures a synapomorphy of Bufonidae. As such, the lack of a TME is plesiomorphic in the earliest diverging lineages (i.e., *Amazophrynella*, most species of *Atelopus*, *Dendrophryniscus*, *Oreophrynella*, *Osornophryne*, *Melanophryniscus*, and *Nannophryne*).

Independent of the methodological approach, available evidence indicates that the complete TME was regained within Bufonidae in *Frostius* and the sister clade of *Nannophryne*, whereas the tympanic annulus and columella were regained in a subclade of *Atelopus*. Thus, these structures are not homologous with the equivalent structures found in the TME of other anurans, although it seems likely that the underlying genetic basis for their development is homologous (i.e., deep homology⁵³, see below). Subsequent losses occurred several times in different clades, indicating a complex evolutionary history of the TME in Bufonidae (see Fig. 2 and section S3 of the Supplementary Information).

The complex evolutionary history of the TME in Bufonidae is even more unusual when compared to other tetrapods. As noted above, although extant caecilians and salamanders do not possess a tympanic membrane or middle ear cavity, it has been hypothesized that a TME might have been present plesiomorphically and that the lateral elements might have been lost independently^{5,54}. Regardless, although the remaining middle ear structures underwent extensive modification, the columella was lost only once in each group, having been greatly reduced or lost in salamandrid salamanders^{1,55} and lost in adult scolecomorphid caecilians (present as a cartilaginous element in fetal and juvenile *Scolecomorphus kirkii*⁵⁶).

Among amniotes, TME loss is extremely rare. There are no documented losses among turtles (e.g.,⁵⁷), archosaurs (e.g.,^{58–60}), or mammals (e.g.,⁶¹), despite the remarkable middle ear transformations in fossorial and marine mammals^{62,63}. Among lepidosaurs, numerous lineages have lost the lateral-most components of the TME, including Serpentes, Amphisbaenia, Agamidae, Diploglossidae, Gymnophthalmidae, Lanthonotidae, Phrynosomatidae, and Scincidae^{11,60,64–66}, but the columella appears to be present in all but the pygopod lizard *Aprasia repens*¹² and, possibly, the lampprophid snakes *Atractaspis* and *Xenocalamus*¹³.

With few exceptions, the development of TME structures in anurans follows a consistent sequence that might explain the consistent pattern of co-occurrence of middle ear structures and provides clues about the mechanisms involved in their loss and gain. First, the medial end of the pars media plectri develops as a chondrification within the connective tissue membrane spanning the fenestra ovalis adjacent to the already formed operculum^{27,67–70}. Next, the pars interna plectri begins to chondrify and, with the incipient pars media, form the future stapedial footplate. Subsequently, a socket-like structure begins to be defined, articulating with the anterior edge of the operculum. The lateral-most portion of the stapedial footplate elongates to complete the formation of the shaft of the pars media plectri, which extends laterally towards the outside of the head. Meanwhile, the tympanic annulus and pars externa plectri develop as cartilaginous condensations associated with the posterior margin of the palatoquadrate. As the palatoquadrate swings posteriorly during metamorphosis, so too do the tympanic annulus and pars externa plectri. As the ontogenetic sequence of development of these structures progresses, they are positioned in the same medial-lateral plane. At this point, the partes media and externa plectri connect synchronotically to each other and the tympanic annulus induces the differentiation of the tympanic membrane^{27,70}.

The sequences of losses and gains appear to be related to the relative timing of the development of structures (heterochronies) and tissue differentiation phenomena. For example, Helff⁶⁹ demonstrated the inductive effects of the tympanic annulus on the tegument to produce the differentiation of the tympanic membrane, which explains why the tympanic membrane never occurs in the absence of a tympanic annulus. Similarly, Hetherington²⁷, Smirnov⁷¹, and Fabrezi and Goldberg⁶⁸ emphasized the relatively late development of TME structures. Hetherington²⁷ and Smirnov⁷¹ also observed that several species undergo post-metamorphic development of previously absent or undeveloped structures (e.g., *Sclerophrys regularis*, *Pseudacris crucifer*, *Bombina*

orientalis). Meanwhile, Smirnov⁷² pointed out that developmental heterochronies (progenesis, neoteny, and post-displacement) seem to play a major role in the post-metamorphic development of these structures. All these events occur in specific sequences and their disruption in particular stages could produce the observed patterns of losses in the subsequent stages of development of the TME. Therefore, research into the genetic basis for the absence of induction of lateral elements promises to be a fruitful line of investigation.

Additionally, genetic mechanisms that directly regulate the expression of these ear structures might be involved. Knowledge of the origin of the components of the vertebrate auditory system is incipient generally and for anurans particularly. However, recent studies of *Xenopus laevis* support a model in which the cartilaginous elements of the TME are derived from three neural crest cell streams (see⁷³): (1) the mandibular stream forms the tympanic annulus, (2) the hyoid stream gives rise to the partes media and externa plectri, and (3) the branchial stream forms the pars interna plectri. The consistent patterns of co-occurrences observed in anurans suggest that a direct role of regulatory genes and/or transcription factors might be involved in the tissular differentiation of the tympanic membrane due to inductive phenomena from the tympanic annulus. Also, it is likely that the development of the tympanic annulus and pars externa plectri (in the margins of the palatoquadrate) and the partes interna and media plectri (in the otic capsule) results from the initiation of a common developmental module, as in the morphogenesis of many other structures^{74,75}. Unfortunately, information on the developmental control genes that lead to the formation of elements in the amphibian middle ear is unavailable. However, some genetic pathways involved in this differentiation process have been identified in other vertebrates and could be examined in frogs⁷⁶.

The lateral–medial dependency between the presence and absence of tympanic middle ear structures appears also to be related to functional constraints: a tympanic membrane without a tympanic annulus or columella would have no acoustic function, as would a tympanic annulus without a columella. In contrast, the tympanic annulus retains its acoustic function in the absence of a tympanic membrane, and the columella remains acoustically functional even in the absence of both structures, as evidenced by the middle ears of salamanders¹. This asymmetric functional dependency appears to have allowed these three structures to evolve sequentially rather than as a single transformation series (i.e., presence or absence of all the three structures), with losses and gains of each element occurring sequentially in a lateral–medial dependency, across the bufonid tree.

The losses, regains, and re-losses of TME structures in Bufonidae make true toads an excellent model to study the behavioural correlates of TME morphology. Previous studies have hypothesized a relationship between earlessness and aquatic or fossorial habitats and lack of acoustic communication or production of low-frequency calls⁷⁷. Additionally, based on the limited evidence presently available (see section S5 of the Supplementary Information), the loss of TME structures in Bufonidae appears to be coincident with the origin of a scramble competition mating system in which males in dense aggregations attempt amplexus indiscriminately and struggle for possession of females⁷⁸. In this mating system, acoustic territorial defence is absent and reliance on hearing for mate choice is greatly reduced or eliminated, as is the effectiveness of prezygotic isolating barriers like advertisement calls⁷⁸, which presumably results in the natural interspecific hybridization observed in many bufonid species (see⁷⁹ and references therein).

Nevertheless, although most of the species of early diverging clades of Bufonidae for which the mating system is known exhibit scramble competition (see section S5 of the Supplementary Information), the reproductive behaviour of most species is unknown, making the character state reconstruction of this behaviour at the root node of Bufonidae ambiguous. Similarly, both within Bufonidae and across Anura, many of the groups that lack a TME employ high frequency (>1 kHz) advertisement calls during reproductive communication (e.g., *Atelopus*⁸⁰, *Bombina*⁸¹, *Brachycephalus*⁸², *Melanophryniscus*⁸³, *Osornophryne*⁸⁴, *Rhinophrynus*⁸⁵, and *Sechellophryne* + *Sooglossus*⁸⁶). Indeed, despite the absence of a TME and the occurrence of a scramble competition mating strategy, interspecific acoustic diversity is maintained in most genera (e.g.,^{80,87–89}) suggesting that acoustic signals play a still unclear role in communication and/or mate choice. The maintenance of call diversity and widespread production of advertisement calls may be explained by extratympanic hearing pathways in earless frogs. Multiple extratympanic pathways, including a lung pathway (e.g., *Atelopus*^{22,23}, *Bombina*⁸¹, *Nectophrynoides asperginis*²¹), an opercularis pathway (reviewed by^{3,25,90,91}), and bone conduction enhanced by resonance of the oral cavity (*Sechellophryne*⁹²) have been shown effective or hypothesized so far in a few earless species. Given that in at least some anurans airborne sounds are transferred via both tympanic and extratympanic pathways (reviewed by^{3,25}), anurans may experience relaxed selective pressures on the TME if TME plasticity does not greatly affect acoustic acuity. If the generality of alternative sound transfer pathways for aerial sounds is corroborated across anuran diversity, then the pre-existence of alternative pathways for airborne sound transmission might explain the high rate of TME loss in anurans. Nevertheless, currently proposed sound localization pathways in anurans all require middle ear coupling⁹³, leaving an alternative mechanism by which earless species localize audible sounds unknown.

Concluding remarks

Since the tympanic annulus and membrane first arose in combination with the plesiomorphically present columella, either prior to the origin of Anura or in Lagobatrachia (the clade formed by all anurans except Ascaphidae and Leiopelmatidae), our analysis indicates that the TME was completely lost at least 38 times in anurans, usually in small clades within diverse families. Bufonidae is exceptional within both Anura and among all tetrapods in that the loss of all TME structures preceded a radiation of more than 150 earless species followed by independent regains and many additional losses in most derived clades. In contrast, among the approximately 26500 species of amniotes the TME was completely lost only three times and was never regained. Available evidence suggests that losses/gains of each TME structure constitute independent transformation series that occur in a lateral–medial dependency, where heterochronic events and regulation via specific genetic mechanisms are implied during development. The complex pattern of TME evolution, extensive morphological and reproductive

diversity, and maintenance of bioacoustic diversity despite the loss of TME structures make Bufonidae a promising model to study extratympanic pathways of sound transmission, the physiological and behavioural consequences of middle ear loss, and the underlying genetic and developmental mechanisms that shaped its remarkable TME diversity.

References

- Wever, E. G. *The amphibian ear*. (Princeton University Press, 1985).
- Mason, M. J., Segenhout, J. M., Cobo-Cuan, A., Quiñones, P. M. & van Dijk, P. The frog inner ear: picture perfect? *Journal of the Association for Research in Otolaryngology* **16**, 171–188 (2015).
- Smotherman, M. & Narins, P. Evolution of the amphibian ear In *Evolution of the vertebrate auditory system* (eds Manley, G. A., Popper, A. N. & Fay, R. R.) Ch. 6, 164–199 (Springer Science & Business Media, 2004).
- Wever, E. G. The ear and hearing in the frog, *Rana pipiens*. *Journal of Morphology* **141**, 461–477 (1973).
- Maddin, H. C. & Anderson, J. S. Evolution of the amphibian ear with implications for lissamphibian phylogeny: insight gained from the caecilian inner ear. *Fieldiana Life and Earth Sciences* **2**, 59–76 (2012).
- Christensen-Dalsgaard, J. & Carr, C. E. Evolution of a sensory novelty: tympanic ears and the associated neural processing. *Brain Research Bulletin* **75**, 365–370 (2008).
- Clack, J. A. The evolution of tetrapod ears and the fossil record. *Brain, Behavior and Evolution* **50**, 198–212 (1997).
- Manley, G. A. An evolutionary perspective on middle ears. *Hearing Research* **263**, 3–8 (2010).
- Manley, G. A. & Sienknecht, U. J. The evolution and development of middle ears in land vertebrates In *The middle ear: science, otosurgery and technology* (eds Puria, S., Popper, A. N. & Fay, R. R.) Ch. 3, 7–30 (Springer, 2013).
- Müller, J. & Tsuji, L. A. Impedance-matching hearing in Paleozoic reptiles: evidence of advanced sensory perception at an early stage of amniote evolution. *PLoS One* **2**, e889 (2007).
- Manley, G. A. *Peripheral hearing mechanisms in reptiles and birds*. (Springer, 1990).
- Daza, J. D. & Bauer, A. M. Cranial anatomy of the pygopodid lizard *Aprasia repens*, a gekkotan masquerading as a scolecophidian In *All animals are interesting: a festschrift in honour of Anthony P. Russell* BIS Verlag, Oldenburg, Germany (eds Bininda-Emonds, O. R. P. et al.) 303–350 (BIS Verlag, 2015).
- Cundall, D. & Irish, F. The snake skull In *Biology of the reptilia* Vol. 20 (eds Gans, C., Gaunt, A. S. & Adler, C.) Ch. 2, 349–692 (Society for the Study of Amphibians and Reptiles, 2008).
- Jaslow, A. P., Hetherington, T. E. & Lombard, R. E. Structure and function of the amphibian middle ear In *The evolution of the amphibian auditory system* (eds Fritsch, B. et al.) Ch. 2, 69–91 (Wiley, 1988).
- Frost, D. R. Amphibian species of the world: an online reference. Version 6.0 <http://research.amnh.org/herpetology/amphibia/index.php> (11 April 2016). (2016).
- Grandison, A. G. C. Morphology and phylogenetic position of the west african *Didynamipus sjoestedti* Andersson, 1903 (Anura Bufonidae). *Monitore Zoologico Italiano* **15**, 187–215 (1981).
- Grant, T. & Bolívar-G, W. A new species of semiarboreal toad with a salamander-like ear (Anura: Bufonidae: *Rhinella*). *Herpetologica* **70**, 198–210 (2014).
- Lötters, S. et al. Assessing the molecular phylogeny of a near extinct group of vertebrates: the neotropical harlequin frogs (Bufonidae; *Atelopus*). *Systematic and Biodiversity* **1**, 45–57 (2011).
- McDiarmid, R. W. Comparative morphology and evolution of frogs of the genera *Atelopus*, *Dendrophryniscus*, *Melanophryniscus*, and *Oreophrynella*. *Bulletin of Los Angeles County Museum of Natural History* **12**, 1–66 (1971).
- Tihen, J. A. Two new genera of african bufonids, with remarks on the phylogeny of related genera. *Copeia* **1960**, 225–233 (1960).
- Arch, V. S., Richards-Zawacki, C. L. & Feng, A. S. Acoustic communication in the Kihansi spray toad (*Nectophrynoides asperginis*): insights from a captive population. *Journal of Herpetology* **45**, 45–49 (2011).
- Boistel, R. et al. Whispering to the deaf: communication by a frog without external vocal sac or tympanum in noisy environments. *PLoS One* **6**, e22080 (2011).
- Lindquist, E. D., Hetherington, T. E. & Volman, S. F. Biomechanical and neurophysiological studies on audition in eared and earless harlequin frogs (*Atelopus*). *Journal of Comparative Physiology A* **183**, 265–271 (1998).
- Gaupp, E. A. *Ecker's und R. Wiedersheim's Anatomie des Frosches, part 1*. (Braunschweig: Friedrich Vieweg und Sohn, 1896).
- Mason, M. J. Pathways for sound transmission to the inner ear in amphibians In *Hearing and sound communication in amphibians* (eds Narins, P. M. et al.) Ch. 6, 147–183 (Springer Science+Business Media, LLC, 2007).
- Vorobyeva, E. & Smirnov, S. Characteristic features in the formation of anuran sound-conducting systems. *Journal of Morphology* **192**, 1–11 (1987).
- Hetherington, T. E. Timing of development of the middle ear of Anura (Amphibia). *Zoomorphology* **106**, 289–300 (1987).
- Mason, M. J. & Narins, P. M. Vibrometric studies of the middle ear of the bullfrog *Rana catesbeiana* II. The operculum. *Journal of Experimental Biology* **205**, 3167–3176 (2002).
- Trueb, L. Patterns of cranial diversity among the Lissamphibia In *Patterns of structural and systematic diversity Vol. 2 The Skull* (eds Hanken, J. & Hall, B. K.) Ch. 6, 255–343 (University of Chicago Press, 1993).
- Pyron, R. A. Biogeographic analysis reveals ancient continental vicariance and recent oceanic dispersal in amphibians. *Systematic Biology* **63**, 779–797 (2014).
- Barej, M. F. et al. The first endemic west african vertebrate family—a new anuran family highlighting the uniqueness of the Upper Guinean biodiversity hotspot. *Frontiers in Zoology* **11**, [1–10] (2014).
- Chandramouli, S. R. et al. A new genus and species of arboreal toad with phytotelmonous larvae, from the Andaman Islands, India (Lissamphibia, Anura, Bufonidae). *ZooKeys* **555**, 57–90 (2016).
- Peloso, P. L. V., Faivovich, J., Grant, T., Gasparini, J. L. & Haddad, C. F. B. An extraordinary new species of *Melanophryniscus* (Anura, Bufonidae) from southeastern Brazil. *American Museum Novitates* **3762**, 1–31 (2012).
- van Bocxlaer, I. et al. Gradual adaptation toward a range-expansion phenotype initiated the global radiation of toads. *Science* **327**, 679–682 (2010).
- Lynch, J. D. & Duellman, W. E. Frogs of the genus *Eleutherodactylus* in western Ecuador. *University of Kansas Natural History Museum, Special Publication* **23**, 1–236 (1997).
- Fitch, W. M. Toward defining the course of evolution: minimum change for a specific tree topology. *Systematic Zoology* **20**, 406–416 (1971).
- Maddison, W. P. & Maddison, D. R. Mesquite: a modular system for evolutionary analysis. Version 3.03. URL <http://mesquiteproject.org/> (2015).
- Paradis, E., Claude, J. & Strimmer, K. APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* **20**, 289–290 (2004).
- Huelsenbeck, J. P., Nielsen, R. & Bollback, J. P. Stochastic mapping of morphological characters. *Systematic Biology* **52**, 131–158 (2003).
- Revell, L. J. Phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* **3**, 217–223 (2012).

41. R Core Team R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/> (2013).
42. Bolt, J. R. & Lombard, R. E. Evolution of the amphibian tympanic ear and the origin of frogs. *Biological Journal of the Linnean Society* **24**, 83–99 (1985).
43. Green, D. M. & Cannatella, D. C. Phylogenetic significance of the amphicoelous frogs, Ascaphidae and Leiopelmatidae. *Ethology Ecology & Evolution* **5**, 233–245 (1993).
44. Stephenson, E. M. T. The anatomy of the head of the New Zealand frog, *Leiopelma*. *Transactions of the Zoological Society of London* **27**, 255–305 (1951).
45. Báez, A. M. & Nicoli, L. A new species of *Notobatrachus* (Amphibia, Salientia) from the Middle Jurassic of northwestern Patagonia. *Journal of Paleontology* **8**, 403–407 (2008).
46. Gao, K.-Q. & Chen, S. A new frog (Amphibia: Anura) from the Lower Cretaceous of western Liaoning, China. *Cretaceous Research* **25**, 761–769 (2004).
47. Gao, K.-Q. & Wang, Y. Mesozoic anurans from Liaoning Province, China, and phylogenetic relationships of archaebatrachian anuran clades. *Journal of Vertebrate Paleontology* **21**, 460–476 (2001).
48. Rage, J.-C. & Rocek, Z. Redescription of *Triadobatrachus massinoti* (Piveteau, 1936) an anuran amphibian from the early Triassic. *Palaeontographica A* **206**, 1–16 (1989).
49. Shubin, N. H. & Jenkins, F. A. An early Jurassic jumping frog. *Nature* **377**, 49–52 (1995).
50. Báez, A. M. & Basso, N. G. The earliest known frogs of the Jurassic of South America: review and cladistic appraisal of their relationships. *Müncher Geowissenschaftliche Abhandlungen (A)* **30**, 131–158 (1996).
51. Frost, D. R. *et al.* The amphibian tree of life. *Bulletin of the American Museum of Natural History* **297**, 1–370 (2006).
52. Grant, T. *et al.* Phylogenetic systematics of dart-poison frogs and their relatives (Amphibia: Athesphatanura: Dendrobatidae). *Bulletin of the American Museum of Natural History* **299**, 1–262 (2006).
53. Shubin, N., Tabin, C. & Carroll, S. Fossils, genes and the evolution of animal limbs. *Nature* **388**, 639–648 (1997).
54. Lombard, R. E. & Bolt, J. R. Evolution of the tetrapod ear: an analysis and reinterpretation. *Biological Journal of the Linnean Society* **11**, 19–76 (1979).
55. Rose, C. S. The developmental morphology of salamander skulls In *Amphibian biology Vol. 5, Osteology* (eds Heatwole, H. & Davies, M.) Ch. 4, 1684–1781 (Surrey Beatty & Sons, 2003).
56. Müller, H., Wilkinson, M., Loader, S. P., Wirkner, C. S. & Gower, D. J. Morphology and function of the head in foetal and juvenile *Scolecophorus kirkii* (Amphibia: Gymnophiona: Scolecophoridae). *Biological Journal of the Linnean Society* **96**, 491–504 (2009).
57. Gaffney, E. S. Comparative cranial morphology of recent and fossil turtles. *Bulletin of the American Museum of Natural History* **164**, 65–376 (1979).
58. Feduccia, A. Morphology of the bony stapes (columella) in the Passeriformes and related groups: evolutionary implications. *University of Kansas Museum of Natural History, Miscellaneous Publications* **63**, 1–34 (1975).
59. Montefeltro, F. C., Andrade, D. V. & Larsson, H. C. E. The evolution of the meatal chamber in crocodyliforms. *Journal of Anatomy* **228**, 838–863 (2016).
60. Saunders, J. C., Duncan, R. K., Doan, D. E. & Werner, Y. L. The middle ear of reptiles and birds In *Comparative hearing: birds and reptiles* (ed Fay, R. R., Popper, A. N. & Dooling, R. J.) Ch. 2, 13–69 (Springer-Verlag, 2000).
61. O’Leary, M. A. *et al.* The placental mammal ancestor and the post-K-Pg radiation of placentals. *Science* **339**, 662–667 (2013).
62. Mason, M. J. Middle ear structures in fossorial mammals: a comparison with non-fossorial species. *Journal of Zoology* **255**, 467–486 (2001).
63. Solntseva, G. Adaptive features of the middle ear of mammal in ontogeny. *Acta Zoologica Bulgarica* **65**, 101–116 (2013).
64. Greer, A. E. The loss of the external ear opening in scincid lizards. *Journal of Herpetology* **36**, 544–555 (2002).
65. Pellegrino, K., Rodrigues, M. T., Yonenaga-Yassuda, Y. & Sites, J. W. A molecular perspective on the evolution of microteiid lizards (Squamata, Gymnophthalmidae), and a new classification for the family. *Biological Journal of the Linnean Society* **74**, 315–338 (2001).
66. Smith, M. A. Evolutionary changes in the middle ear of certain agamid and iguanid lizards. *Proceedings of the Zoological Society of London* **108**, 543–549 (1938).
67. Barry, T. H. The ontogenesis of the sound-conducting apparatus of *Bufo angusticeps* Smith. *Morphologisches Jahrbuch* **97**, 477–544 (1956).
68. Fabrezi, M. & Goldberg, J. Heterochrony during skeletal development of *Pseudis platensis* (Anura, Hylidae) and the early offset of skeleton development and growth. *Journal of Morphology* **270**, 205–220 (2009).
69. Helff, O. M. Studies on amphibian metamorphosis. III. The influence of the annular tympanic cartilage on the formation of the tympanic membrane. *Physiological Zoology* **1**, 463–495 (1928).
70. Sedra, S. N. & Michael, M. I. The ontogenesis of the sound conducting apparatus of the Egyptian toad, *Bufo regularis* Reuss, with a review of this apparatus in Salientia. *Journal of Morphology* **103**, 359–375 (1959).
71. Smirnov, S. V. Postmetamorphic skull development in *Bombina orientalis* (Amphibia Discoglossidae), with comments on neoteny. *Zoologischer Anzeiger* **223**, 91–99 (1989).
72. Smirnov, S. V. The anuran middle ear: developmental heterochronies and adult morphology diversification. *Belgian Journal of Zoology* **121**, 99–110 (1991).
73. Gross, J. B. & Hanken, J. Segmentation of the vertebrate skull: neural-crest derivation of adult cartilages in the clawed frog, *Xenopus laevis*. *Integrative and Comparative Biology* **48**, 681–696 (2008).
74. Collin, R. & Miglietta, M. P. Reversing opinions on Dollo’s law. *Trends in Ecology & Evolution* **23**, 602–609 (2008).
75. Gilbert, S. F., Opitz, J. M. & Raff, R. A. Resynthesizing evolutionary and developmental biology. *Developmental Biology* **173**, 357–372 (1996).
76. Chapman, S. C. Can you hear me now? Understanding vertebrate middle ear development. *Frontiers in Bioscience* **16**, 1675–1692 (2011).
77. Hetherington, T. E. The effects of body size on the evolution of the amphibian middle ear In *The evolutionary biology of hearing* (eds Webster, D. B., Fay, R. R. & Popper, A. N.) Ch. 21, 421–437 (Springer, 1992).
78. Wells, K. D. *The ecology and behaviour of amphibians*. (The University of Chicago Press, 2007).
79. Pereyra, M. O. *et al.* Phylogenetic relationships of toads of the *Rhinella granulosa* group (Anura: Bufonidae): a molecular perspective with comments on hybridization and introgression. *Cladistics* **32**, 36–53 (2016).
80. Crocroft, R. B., McDiarmid, R. W., Jaslow, A. P. & Ruiz-Carranza, P. M. Vocalization of eight species of *Atelopus* (Anura: Bufonidae) with comments on communication in the genus. *Copeia* **1990**, 631–643 (1990).
81. Hetherington, T. E. & Lindquist, E. D. Lung-based hearing in an “earless” anuran amphibian. *Journal of Comparative Physiology A* **184**, 395–401 (1999).
82. Condez, T. H., Clemente-Carvalho, R. B. G., Haddad, C. F. B. & dos Reis, S. F. A new species of *Brachycephalus* (Anura: Brachycephalidae) from the highlands of the Atlantic Forest, southeastern Brazil. *Herpetologica* **70**, 89–99 (2014).
83. Duré, M. I., Schaefer, E. F. & Kehr, A. I. Acoustic repertoire of *Melanophryniscus cupreuscapularis* (Céspedes and Álvarez 2000) (Anura: Bufonidae): advertisement, encounter, and release calls. *Journal of Herpetology* **49**, 53–59 (2015).
84. Gluesenkamp, A. G. & Acosta, N. Sexual dimorphism in *Osornophryne guacamayo* with notes on natural history and reproduction in the species. *Journal of Herpetology* **35**, 148–151 (2001).

85. Sandoval, L., Barrantes, G., Ocampo, D. & Sánchez-Quirós, C. Sexual size dimorphism and acoustical features of the pre-advertisement and advertisement calls of *Rhinophrynus dorsalis* Duméril & Bibron, 1841 (Anura: Rhinophrynidae). *Mesoamerican Herpetology* **2**, 154–166 (2015).
86. Gerlach, J. & Willi, J. A new species of frog, genus *Sooglossus* (Anura, Sooglossidae) from Silhouette Island, Seychelles. *Amphibia-Reptilia* **23**, 445–458 (2002).
87. Blair, W. F. Evolution in the genus *Bufo* (University of Texas Press, Austin and London, 1972).
88. Caldart, V. M., dos Santos, T. G. & Maneyro, R. The advertisement and release calls of *Melanophryniscus pachyrhynchus* (Miranda-Ribeiro, 1920) from the central region of Rio Grande do Sul, southern Brazil. *Acta Herpetologica* **8**, 115–122 (2013).
89. McDiarmid, R. W. & Gorzula, S. Aspects of the reproductive ecology and behavior of the Tepui toads, genus *Oreophrynella* (Anura, Bufonidae). *Copeia* **1989**, 445–451 (1989).
90. Lombard, R. E. & Straughan, I. R. Functional aspects of anuran middle ear structures. *Journal of Experimental Biology* **61**, 71–93 (1974).
91. Hetherington, T. E. & Lombard, R. E. Electromyography of the opercularis muscle of *Rana catesbeiana*: an amphibian tonic muscle. *Journal of Morphology* **175**, 17–26 (1983).
92. Boistel, R. *et al.* How minute sooglossid frogs hear without a middle ear. *Proceedings of the National Academy of Sciences* **110**, 15360–15364 (2013).
93. Willis, K. L., Christensen-Dalsgaard, J. & Carr, C. E. Auditory brain stem processing in reptiles and amphibians: roles of coupled ears In *Insights from comparative hearing research* (eds Köppl, C. *et al.*) Ch. 8, 193–225 (Springer, 2013).

Acknowledgements

For access to collections and specimen loans we thank Darrel R. Frost (AMNH), James Aparicio (CBF), Célio F.B. Haddad (CFBH), John D. Lynch (ICN), Linda Trueb, William E. Duellman, and Rafe Brown (KU), Glauçia Maria Funk Pontes (MCP), Ignacio de la Riva and José González (MNCN), Miguel T. Rodriguez and Paulo E. Vanzolini (MZUSP), Ingrid Fernandez and Lucindo González (MNK), David C. Cannatella (TNHC), Ronald Nussbaum (UMMZ), W. Ron Heyer and Ronald Crombie (USNM), and Ariovaldo Giarretta (ZUEC). We also thank MicroCT Core Facility (San Antonio, Texas), and Katyuscia Araujo-Vieira, Aaron Bauer, Sara Bertelli, Laura Nicoli, Diego Pol, Marco Rada, Guillermo Rougier, Juliana Sterli, Miguel T. Rodrigues, and Marcelo Weksler for sharing information about TME structures in different tetrapod groups. Agustín Elias-Costa contributed in the production of Figure 1. We are grateful to the anonymous reviewers for their insightful comments and suggestions that improved the manuscript. We thank ANPCyT, CONICET, FAPESP, CNPq, Fulbright Commission Argentina, and NSF funding for financial support: NSF IOS-1350346; PIP 1112008010-2422, 112201101-00875, and 112201101-00889; PICT 2011-1524, 2010-1740, 2012-2687, and 2013-404; CNPq proc. 305234/2014-5, and FAPESP procs. 2012/09401-5, 2012/10000-5, 2013/20423-3, 2013/50741-7, and 2014/03585-2.

Author Contributions

All authors conceived of the study. M.O.P., M.C.W., K.L.H., T.G. and J.F. coordinated data collection and analysis. M.O.P., M.C.W., J.S.B., B.L.B., D.B., M.T., J.J.O.-S., L.A.C., T.G. and J.F. collected the data. M.O.P. and M.C.W. performed the analyses. M.O.P., M.C.W., J.S.B., D.B., T.G. and J.F. wrote the manuscript. B.L.B., M.T., J.J.O.-S., J.M.G., L.A.C. and K.L.H. provided critical comments. All authors approved the final version of the manuscript for submission.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Pereyra, M. O. *et al.* The complex evolutionary history of the tympanic middle ear in frogs and toads (Anura). *Sci. Rep.* **6**, 34130; doi: 10.1038/srep34130 (2016).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016