SCIENTIFIC REPORTS

OPEN

Monocytic MKP-1 is a Sensor of the Metabolic Environment and Regulates Function and Phenotypic Fate of Monocyte-Derived Macrophages in Atherosclerosis

Hong Seok Kim^{1,2}, Sina Tavakoli³, Leigh Ann Piefer⁴, Huynh Nga Nguyen⁵ & Reto Asmis^{3,4,5}

Diabetes promotes the S-glutathionylation, inactivation and subsequent degradation of mitogen-activated protein kinase phosphatase 1 (MKP-1) in blood monocytes, and hematopoietic MKP-1-deficiency in atherosclerosis-prone mice accelerates atherosclerotic lesion formation, but the underlying mechanisms were not known. Our aim was to determine the mechanisms through which MKP-1 deficiency in monocytes and macrophages promotes atherogenesis. Transplantation of MKP-1-deficient bone marrow into LDL-R^{-/-} (MKP-1^{LeuKO}) mice accelerated high-fat diet (HFD)-induced atherosclerotic lesion formation. After 12 weeks of HFD feeding, MKP-1^{LeuKO} mice showed increased lesion size in both the aortic root (1.2-fold) and the aorta (1.6-fold), despite reduced plasma cholesterol levels. Macrophage content was increased in lesions of MKP-1^{LeuKO} mice compared to mice that received wildtype bone marrow. After only 6 weeks on a HFD, *in vivo* chemotactic activity of monocytes was already significantly increased in MKP-1^{LeuKO} mice. MKP-1 deficiency in monocytes and macrophages promotes and accelerates atherosclerotic lesion formation by hyper-sensitizing monocytes to chemokine-induced recruitment, predisposing macrophages to M1 polarization, decreased autophagy and oxysterol-induced cell death whereas overexpression of MKP-1 protects macrophages against metabolic stress-induced dysfunction. MKP-1 serves as a master-regulator of macrophage phenotype and function and its dysregulation by metabolic stress may be a major contributor to atherogenesis and the progression of atherosclerotic plaques.

Atherosclerosis is a chronic inflammatory disease characterized by monocyte infiltration and macrophage accumulation in the vessel wall¹. Inflammatory processes are tightly regulated, and macrophages play a critical role in the initiation, maintenance, and resolution of inflammation by converting into distinct activation states, ranging from a classically activated or inflammatory state, to an alternatively activated or inflammation-resolving state, and to a deactivated state². The dysregulation of macrophage activation and functions appears to be a major contributor to the conversion of acute inflammation into a chronic process³ and to the development and progression of atherosclerotic lesions³.

Macrophage numbers and functionalities within atherosclerotic lesions and their removal from plaques are controlled by four key processes: recruitment, autophagy, apoptosis, and macrophage polarization. A better understanding of the molecular mechanisms that result in the dysregulation of these processes would therefore be critical for the development of targeted therapies for the prevention and treatment of atherosclerosis and possibly other chronic inflammatory diseases associated with metabolic disorders. Metabolic stress promotes monocyte “priming”, i.e. the dysregulation and hypersensitization of blood monocytes to chemokines, resulting

¹Department of Molecular Medicine, College of Medicine, Inha University, Incheon 22212, Republic of Korea. ²Hypoxia-related Disease Research Center, College of Medicine, Inha University, Incheon 22212, Republic of Korea. ³Department of Radiology, University of Texas Health Science Center at San Antonio, San Antonio, TX 78229, USA. ⁴Department of Clinical Laboratory Sciences, University of Texas Health Science Center at San Antonio, San Antonio, TX 78229, USA. ⁵Department of Biochemistry, University of Texas Health Science Center at San Antonio, San Antonio, TX 78229, USA. Correspondence and requests for materials should be addressed to R.A. (email: asmis@uthscsa.edu)

in the over-recruitment of monocyte-derived macrophages to sites of vascular injury and the onset of atherogenesis^{4,5}. Impairment of macrophage autophagy, a process that protects macrophages from cell damage by degrading damaged and potentially noxious and proinflammatory material, has also been implicated in atherosclerotic lesion progression⁶. In addition, dysregulated macrophage apoptosis and impaired phagocytosis of apoptotic macrophages or “efferocytosis” lead to secondary necrosis and increased inflammation, further promoting plaque progression and destabilization^{7,8}. Finally, dysregulated macrophage polarization may further promote and accelerate atherogenesis as the persistence of classically activated, pro-inflammatory “M1” macrophages and the delayed or impaired appearance of alternatively activated “M2” macrophages, impairs inflammation resolution, further fueling the atherogenic process^{9,10}.

Macrophage chemotaxis⁴, autophagy¹¹, apoptosis^{12,13}, and polarization¹⁴ are mediated by mitogen-activated protein kinase (MAPK) pathways, which in turn are counter-regulated by MAPK phosphatases (MKPs)¹⁵. MKPs are a family of dual-specificity protein phosphatases that are responsible for the dephosphorylation of both phosphothreonine and phosphotyrosine residues in MAPKs, including JNK, p38 MAPK, and ERK¹⁵. Dephosphorylation of MAPKs by MKPs inhibits MAPK activity, thereby negatively regulating MAPK signaling.

We recently identified MKP-1 as a critical regulator of monocyte adhesion and migration which, when inactivated and degraded in response to metabolic stress, promotes macrophage recruitment and accumulation in early atherosclerotic lesions⁴. MKP-1 was initially characterized as an ERK1 and ERK2 phosphatase¹⁶. Subsequent studies indicated that p38 and JNK MAPKs are also MKP-1 substrates and that MKP-1 is capable of inactivating all three classes of MAPK *in vivo*¹⁷. MKP-1 has been reported to inhibit apoptotic events in leukemia cells¹⁸, and a recent study suggests that MKP-1 may control macrophage phenotypic transitions during tissue repair¹⁹. However, little was known about the roles of MKP-1 in macrophage apoptosis, autophagy and polarization. Our study shows that, in addition to promoting macrophage recruitment and accumulation, MKP-1 deficiency in monocytes and macrophages accelerates atherogenesis and lesion progression by impairing macrophage autophagy, enhancing apoptosis, and by dysregulating macrophage polarization whereas overexpression of MKP-1 protects macrophages against metabolic stress-induced dysfunction. Our findings identified MKP-1 as a master regulator of macrophage function and fate.

Results

MKP-1 deficiency in macrophages promotes atherosclerosis. We showed that hematopoietic MKP-1-deficiency in atherosclerosis-prone mice accelerates atherosclerotic lesion formation⁴, but the underlying mechanisms were not known. To examine the effects of MKP-1 deficiency on macrophage function *in vivo*, we performed bone marrow transplantation experiments in low density lipoprotein receptor-deficient (LDL-R^{-/-}) mice using MKP-1^{-/-} and wildtype control mice as bone marrow donors. We induced moderate metabolic stress by feeding the transplant recipients a high-fat diet for six and 12 weeks, and assessed monocyte chemotactic responses *in vivo* and atherosclerotic lesions formation. After only six weeks of HFD feeding, MKP-1 deficient monocytes showed a 4.2-fold higher chemotactic activity in response to monocyte chemoattractant protein-1 (MCP-1) than monocytes in mice that received wildtype bone marrow, indicating that these monocytes were already “primed” and dysfunctional (Suppl. Fig. 1)^{4,5}. At this early stage, both groups of mice showed small atherosclerotic lesions in both the aortic arch and the descending aorta (Fig. 1A+B), but we did not detect a significant difference in lesion size between the two groups. These findings suggest that monocyte priming and dysfunction induced by MKP-1 deficiency precedes the formation of atherosclerotic lesions.

After 12 weeks on a HFD, the chemotactic response to MCP-1 of MKP-1-deficient monocytes showed no further increase but remained significantly higher (3.3-fold) than that of monocytes in LDL-R^{-/-} recipients of wildtype bone marrow (Suppl. Fig. 1). Atherosclerotic lesion size was increased in both groups, compared to the 6-week time point. However, compared to recipients of wildtype bone marrow, LDL-R^{-/-} mice with MKP-1-deficient bone marrow showed a 1.6-fold increase in atherosclerotic lesion formation as assessed by *en face* analysis (Fig. 1A+B). The acceleration of atherogenesis by hematopoietic MKP-1 deficiency was not due to changes in blood glucose levels and plasma lipids as these mice actually showed slightly lower total plasma cholesterol than mice that received wildtype bone marrow, and plasma triglyceride levels and lipoprotein profiles were not significantly affected by hematopoietic MKP-1 deficiency (Suppl. Fig. 2+3). Monocyte counts were also not different between the two groups (WT: $(0.16 \pm 0.05) \times 10^9/l$ versus MKP-1^{-/-}: $(0.19 \pm 0.03) \times 10^9/l$; $P = 0.68$).

Histological assessment of atherosclerotic lesions at the aortic sinus after 12 weeks of HFD feeding confirmed the results of the *en face* analysis, revealing a 24% larger foam cell-rich lesion area in mice with hematopoietic MKP-1 deficiency compared to mice that received wildtype bone marrow (Fig. 1C+D). In addition, lesions in mice with hematopoietic MKP-1 deficiency showed a greater CD68-positive area per lesion area (Fig. 1E+F), indicating a higher macrophage content than lesions in control mice. Together these findings suggest that hematopoietic MKP-1-deficiency in atherosclerosis-prone mice accelerates atherogenesis at least in part by sensitizing blood monocytes to chemokines, increasing their chemotactic activity and by enhancing macrophage infiltration and accumulation in atherosclerotic lesions.

MKP-1-deficient macrophages are autophagy-defective. Autophagy is one of the cell’s survival responses to the stress, and protects macrophages in the plaque by degrading damaged, potentially noxious and proinflammatory material formed within the cells^{20,21}. Autophagy is regulated by MAPK-dependent pathways^{11,22–24}. MAPKs are counter-regulated by MKPs, including MKP-1¹⁵. We therefore examined whether MKP-1 deficiency impairs macrophage autophagy. To this end, we isolated peritoneal macrophages from wildtype and MKP-1^{-/-} mice, and assessed the ratio of light chain 3 (LC3)-II to LC3-I levels as measures of autophagic flux²⁵ and p62/sequestosome 1 (SQSTM1) as an indicator of flux impairment²⁶, respectively. As a positive control of the LC3-II to LC3-I ratio shift, peritoneal macrophages were treated with mechanistic target of rapamycin (mTOR) inhibitor (Suppl. Fig. 4). The

Figure 1. MKP-1 deficiency in macrophages promotes atherosclerosis. Bone marrow transplantation was performed in LDL-R^{-/-} mice using wildtype and MKP-1^{-/-} mice as bone marrow donors. Mice were fed a high-fat diet for 6 and 12 wk. Hearts and aortas of bone marrow recipients were removed, and the size of atherosclerotic lesions was determined by en face analysis (A+B) and Oil Red O-stained sections from the aortic roots (C+D). Experiments were performed using 5–10 mice per experimental group. Macrophage contents in the atherosclerotic aortic roots was assessed by analysis of CD68-stained sections from the aortic root (E+F). Experiments were performed using 6 mice per experimental group. Results shown are mean ± SE.

ratio of LC3-II to LC3-I was decreased by 29% and p62/SQSTM1 levels were increased by 86% in MKP-1-deficient macrophages (Fig. 2A+B), confirming that MKP-1 activity is critical for the maintenance of macrophage autophagy.

Figure 2. Both MKP-1-deficient and metabolically primed macrophages exhibit defective autophagy. (A+B) Autophagic activity was assessed by Western blot analysis as the ratio of LC3-II to LC3-I and p62/SQSTM1 levels in peritoneal macrophages isolated wildtype (WT) and from MKP-1^{-/-} (KO) mice and (C+D) in unprimed (Control) and metabolically primed (LDL + HG) peritoneal macrophages. Results shown are mean $\pm$ SE of 3 independent experiments.

Since metabolic stress promotes the S-glutathionylation, inactivation and subsequent degradation of MKP-1 in blood monocytes⁴, we next examined whether metabolic stress is sufficient to impair macrophage autophagy. To this end, we metabolically primed peritoneal macrophages from C57BL/6 mice by preincubating the cells for 24 h with human LDL (100 μ g/ml) plus high glucose (25 mmol/L)^{4,5}. We then examined autophagic flux by measuring the ratio of LC3-II to LC3-I and p62/SQSTM1 levels. Metabolic stress reduced the ratio of LC3-II to LC3-I in mouse peritoneal macrophages by 51% (Fig. 2C) and increased p62/SQSTM1 levels 1.5-fold (Fig. 2D), indicating that metabolic stress impairs autophagy in macrophages. As expected, MKP-1 levels were reduced by 39% in these macrophages (Suppl. Fig. 5). Together, these results suggest that monocyte priming induced by metabolic stress, and the associated partial loss of MKP-1 activity, may predispose monocyte-derived macrophages recruited to atherosclerotic lesions to impaired autophagy.

This mechanistic link between metabolic priming of blood monocytes, loss of MKP-1 activity and impaired macrophage autophagy is further supported by immunofluorescence data we obtained in atherosclerotic lesions from chimeric LDL-R^{-/-} mice that received either wildtype or MKP-1-deficient bone marrow. We found a 49% increase in p62 levels in macrophages in atherosclerotic lesions of LDL-R^{-/-} mice with hematopoietic MKP-1 deficiency compared to mice with MKP-1-expressing macrophages (Fig. 3A+B). This finding confirms our *in vitro* data and strongly suggests that loss of MKP-1 activity impairs macrophage autophagy in atherosclerotic lesions.

Figure 3. Loss of MKP-1 activity impairs macrophage autophagy in atherosclerotic lesions. (A+B) Autophagy in the atherosclerotic aortic roots of mice that received either wildtype or MKP-1^{-/-} bone marrow and were fed a high-fat diet for 12 weeks was assessed by immunofluorescence. The autophagy marker p62/SQSTM1 (red) was concurrently imaged with CD68 (green). Experiments were performed using 5 mice per experimental group. Results shown are mean $\pm$ SE.

Loss of MKP-1 activity sensitizes oxysterol-induced cell death in macrophages. Both p38 and JNK MAP kinases play key roles in macrophage apoptosis^{8,27,28}. It is well established that the sustained induction of apoptosis in macrophages within advanced plaques results in a significant increase in lesion size²⁹. We therefore examined whether MKP-1 deficiency sensitizes macrophages to apoptosis induced by oxysterols. Since 7-ketocholesterol (7-KC), a major oxidation product of cholesterol found in human atherosclerotic plaque³⁰, is cytotoxic to macrophages³¹, we treated peritoneal macrophages isolated from wildtype and MKP-1^{-/-} mice for 24 h with either vehicle or 7-KC to induce apoptosis. Cell death was determined by trypan blue dye exclusion (Suppl. Fig. 6) and by measuring caspase 3/7 activation (Suppl. Fig. 7). Cell death and caspase 3/7 activities in response to 7-KC were markedly increased in MKP-1-deficient macrophages compared to macrophages from wildtype mice (Fig. 4A+B), indicating that MKP-1 protects macrophages against oxysterol-induced cell death. Since metabolic stress results in the loss of MKP-1 activity⁴, we explored whether metabolic stress is sufficient to predispose peritoneal macrophages to oxysterol-induced apoptosis. As expected, metabolic stress had a very similar sensitizing effect on 7-KC-induced cell death as genetic MKP-1 deficiency (Fig. 4C+D).

To investigate whether MKP-1 deficiency also enhances macrophages apoptosis in atherosclerotic lesions, we analyzed terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL)-stained aortic root sections from $LDL-R^{-/-}$ mice that received either wildtype or MKP-1-deficient bone marrow. Lesions from $LDL-R^{-/-}$ mice with hematopoietic MKP-1 deficiency contained more than twice as many apoptotic cells per lesion area than lesions from $LDL-R^{-/-}$ mice with MKP-1 expressing macrophages (Fig. 4E+F). These findings suggest that MKP-1 deficiency in macrophages promotes that accumulation of apoptotic cells in atherosclerotic lesions.

Figure 4. Both MKP-1-deficient and metabolically primed macrophages are sensitized to oxysterol-induced apoptosis. Apoptosis was assessed in peritoneal macrophages treated with vehicle or 7-KC for 24 h. Cell death was measured by trypan blue dye exclusion and caspase 3/7 activation in peritoneal macrophages isolated from wildtype (WT) and MKP-1^{-/-} (KO) mice (A+B), and in unprimed (Control) and metabolically primed (LDL + HG) peritoneal macrophages from C57/BL6 mice (C+D). Results are shown as mean $\pm$ SE (n = 3–4). Cell death in the atherosclerotic aortic roots of mice that received either wildtype or MKP-1^{-/-} bone marrow and were fed a high-fat diet for 12 wk was assessed by TUNEL-positive cells relative to the atherosclerotic lesion area (E+F). Experiments were performed using 5–8 mice per experimental group (MKP-1^{+/+} $\rightarrow$ LDL-R^{-/-}: n = 5; MKP-1^{-/-} $\rightarrow$ LDL-R^{-/-}: n = 8). Results are shown as mean $\pm$ SE.

Figure 5. Metabolic priming and MKP-1 deficiency enhance macrophage M1 polarization and suppress the conversion of macrophages into an M2, inflammation-resolving phenotype. Peritoneal macrophages from wildtype (WT) or MKP-1^{-/-} (KO) mice, and unprimed (Control) and metabolically primed (LDL + HG) peritoneal macrophages were polarized as follows: TNF- α plus IFN- γ stimulation to generate M1 macrophages, and IL-4 stimulation as a model of M2 polarization. Untreated peritoneal cells were considered unpolarized. Macrophage polarization was assessed by quantifying and comparing levels of the indicated mRNAs between wildtype and MKP-1^{-/-}, and unprimed and primed macrophages by RT-qPCR. Expression levels were

determined using TaqMan® probes in conjunction with the BioMark™ HD Fluidigm System and Fluidigm Real-Time PCR Analysis software. mRNA levels were normalized to a housekeeping gene (*Hprt*) as well as unprimed macrophages (Control), and the $2^{-\Delta\Delta Ct}$ for each mRNA is reported (A+C). Results are shown as mean of 4 independent samples. STAT1 phosphorylation were assessed with peritoneal macrophages from wildtype (WT) or MKP-1^{-/-} (KO) mice, and unprimed (Control) and metabolically primed (LDL + HG) peritoneal macrophages (B+D). Results are shown as mean ± SE (n = 3–4).

MKP-1 deficiency enhances M1 polarization of macrophages and dampens the conversion of macrophages into an inflammation-resolving M2 phenotype. In atherosclerotic lesions, as in other inflamed tissues, the microenvironment plays a key role in determining the activation state of macrophages. Cytokines are an important constituent of the inflammatory milieu and drive macrophage activation either toward a classical (M1, proinflammatory) or alternative (M2, anti-inflammatory) polarization state³. The M1 macrophage phenotype is controlled by NF-κB, signal transducer and activator of transcription 1 (STAT1) and interferon-regulatory factor 5 (IRF5)^{32,33}, but STAT1 activity is essential for macrophages to polarize into an M1 phenotype^{33,34}. Phosphorylation of STAT1 at Ser727 is required for its activation^{35,36}. P38 MAPK is required for STAT1 phosphorylation and transcriptional activation induced by interferons^{35,37}. Interestingly, STAT1 has been suggested as a physiological non-MAPK substrate for MKP-1^{38,39}. We used real-time PCR in custom-designed microfluidic dynamic arrays^{40,41} to explore how MKP-1 deficiency affects macrophage polarization into M1 or M2 activation states. We found that MKP-1 deficiency enhanced interferon gamma+ tumor necrosis factor alpha (INFγ + TNFα)-induced polarization of macrophages toward an M1 phenotype and suppressed the interleukin-4 (IL-4)-induced conversion of macrophages into an M2 phenotype (Fig. 5A). STAT1 phosphorylation was increased by 53% in MKP-1-deficient macrophages as compared to wildtype cells (Fig. 5B), confirming that STAT1, at least in macrophages, is indeed a physiological non-MAPK substrate for MKP-1.

To determine whether metabolic stress again had a similar effect on macrophage polarization as MKP-1 deficiency, we repeated these experiments with peritoneal macrophages isolated from C57BL/6 mice that were metabolically primed, i.e. pre-incubated for 24 h with human LDL (100 μg/ml) plus high glucose (25 mmol/L) prior to being activated with either INFγ + TNFα (M1) or IL-4 (M2). Like MKP-1-deficient macrophages, metabolically primed macrophages displayed enhanced expression of the M1 markers (red font) in response to INFγ + TNFα, but suppressed expression of M2 markers (green font) after IL-4 stimulation (Fig. 5C). Like MKP-1-deficient macrophages, metabolically primed macrophages showed a significantly higher level of STAT1 phosphorylation (202%; Fig. 5D). Together, these findings confirm that metabolically primed macrophages, like MKP-1-deficient macrophages, are pre-programmed for exaggerated M1 polarization but suppressed M2 activation. In both cases, this skewed polarization profile appears to be due to the hyperactivation of STAT1 as a results of reduced MKP-1 activity.

Overexpression of MKP-1 protects macrophages against impaired autophagy, accelerated cell death, and skewed polarization induced by metabolic stress. Metabolic stress promotes MKP-1 inactivation and degradation, and metabolically primed macrophages show all the phenotypic and functional hallmarks of macrophages with genetic MKP-1 deficiency. We therefore determined whether overexpression of MKP-1 protects metabolically stressed macrophages against defective autophagy, sensitization to oxysterol-induced cell death, and skewed M1/M2 polarization. To this end, we used a lentivirus-based transduction system to overexpress Flag-tagged MKP-1 (or green fluorescent protein, GFP) in bone marrow-derived macrophages. In contrast to GFP-expressing control macrophages, macrophages expressing Flag-MKP-1 were resistant to metabolic stress and showed no defects in autophagy (Fig. 6A) or any increased sensitivity to oxysterol-induced activation of caspase 3/7 (Fig. 6B). Furthermore, STAT1 phosphorylation induced by metabolic priming was suppressed in macrophages overexpressing MKP-1 (Fig. 7A) and the skewing of the polarization profile was prevented in MKP-1 overexpressing cells (Fig. 7B). These data provide further evidence that MKP-1 is essential for macrophages to maintain autophagy, resist cytotoxic stress induced by oxysterols and to properly respond to both pro-inflammatory as well as inflammation-resolving signals.

Discussion

Macrophages are critical for tissue repair as they promote both inflammation, during the early phases, as well as inflammation resolution critical for wound healing⁴². Coordinating these complex functions requires both a high degree of plasticity as well as a tight regulation of cell signaling and cellular functions.

We identified MKP-1 as a regulator of monocyte signaling, activation and function, sensitive to metabolic stress-induced inactivation and S-glutathionylation dependent degradation⁴. We therefore hypothesized that dysregulation and loss of MKP-1 activity in blood monocytes induced by metabolic disorders may have profound effects on macrophages derived from these monocytes, which are recruited to sites of tissue injury and may affect macrophage functions associated with both inflammatory as well as inflammation resolving processes. Thus, loss of MKP-1 in macrophages may play a central role in chronic inflammatory diseases such as atherosclerosis. We found that genetic MKP-1 deficiency impairs macrophage autophagy, sensitizes macrophages to oxysterol-induced cell death, and enhances M1 polarization while suppressing M2 polarization. We also provide direct evidence that metabolic stress-induced MKP-1 deficiency in blood monocytes directly promotes atherogenesis by enhancing monocyte recruitment into the vessel wall, dysregulating macrophage plasticity and by predisposing monocyte-derived macrophages for impaired autophagy and increased cell death within atherosclerotic lesions.

Figure 6. Overexpression of MKP-1 protects macrophages against impaired autophagy, and accelerated cell death. Bone marrow-derived macrophages were infected with lentiviral vectors carrying either Flag-tagged MKP-1 or GFP. Cells were then treated for 24 hours with vehicle or primed with LDL + HG, and p62 protein levels (A), Caspase-3/7 activities (B) Results are shown as mean $\pm$ SE ($n = 3-4$).

MKP-1 preferentially dephosphorylates activated p38 MAPK and JNK relative to ERK⁴³. Studies with MKP-1 knockout mice provided compelling evidence that MKP-1 is an important regulator of innate and adaptive immune responses, inflammation and infection control by regulating p38 MAPK and JNK activation in response to stimuli and stress^{44,45}. Additionally, MKP-1-deficient fibroblasts exhibit enhanced sensitivity to apoptosis mediated by p38 MAPK hyperactivation, suggesting a role for MKP-1 in cell survival signaling⁴⁶. Our finding that MKP-1 deficient macrophages in culture are significantly more sensitive than wildtype macrophages to inducers of apoptosis, such as oxysterols, supports that hypothesis. The fact that MKP-1-deficiency sensitized macrophages to 7-KC-induced apoptosis and that atherosclerotic lesions of recipients of MKP-1-deficient bone marrow showed a dramatic increase in apoptotic macrophages strongly suggests that MKP-1 plays a critical role in protecting macrophages in atherosclerotic plaques against oxysterol-induced cell injury and death.

To date, MKP-1 had not been linked to autophagy, however, p38 MAPK activation was shown to inhibit autophagy in macrophages¹¹ and in human CD8⁺ T cells⁴⁷, which suggested that MKP-1 may also play an important role in macrophage autophagy. Indeed, we found that MKP-1 deficiency significantly impairs autophagy and autophagic flux. Considering the pro-survival role of autophagy^{48,49}, impaired autophagy may have at least partly contributed to the increased vulnerability of MKP-1-deficient macrophages to oxysterol-induced apoptosis we observed both *in vitro* and *in vivo* (Fig. 4).

Polarization of macrophages into either classically activated M1 or alternatively activated M2 macrophages is regulated by STAT and MAPK pathways. Here we provide the first evidence that MKP-1 plays a critical role in regulating macrophage polarization, and that loss of MKP-1 activity results in dysregulated macrophage activation and polarization. STAT1 is a key activator of M1 macrophage polarization, and Ser727 phosphorylation is essential for M1 macrophage polarization^{35,36,50,51}. Importantly, p38 MAPK is required for STAT1 activation^{35,37}. Moreover, STAT1 has been suggested to be a physiological non-MAPK substrate for MKP-1^{38,39} and we observed that MKP-1 inhibition, either by MKP-1 deficiency or metabolic stress, increased STAT1 phosphorylation at Ser727. This finding suggests that reduced MKP-1 activity results in the hyperactivation of STAT1 (Fig. 5B,D), and this mechanism would account for the hyper-inflammatory profile of M1 (IFN γ + TNF α) polarized macrophages that were exposed to metabolic stress (Fig. 5C). We therefore provide evidence that STAT1 in macrophages is in fact a physiological non-MAPK substrate for MKP-1.

Previously, we showed that metabolic stress decreases MKP-1 levels and activity in monocytes and macrophages, and both the inactivation and degradation of MKP-1 are mediated by the S-glutathionylation of the catalytic cysteine residue 258⁴. Here, we show that metabolic stress mimics the effects of genetic MKP-1 deficiency, including defective autophagy, sensitization to oxysterol-induced cell death, and skewed M1/M2 polarization. We demonstrated that MKP-1 deficiency occurs in blood monocytes of diabetic mice⁴. Oxidative stress induced by chronic metabolic stress is likely to cause macrophage MKP-1 deficiency, and macrophages deficient in MKP-1 are likely to be found in microenvironments with high metabolic stress, such as those found in atherosclerotic lesions (Fig. 1E) and lipid-enriched diabetic environments. If these microenvironments persist and

Figure 7. Overexpression of MKP-1 protects macrophages against skewed polarization induced by metabolic stress. Bone marrow-derived macrophages were infected with lentiviral vectors carrying either Flag-tagged MKP-1 or GFP. Cells were then treated for 24 hours with vehicle or primed with LDL + HG, and STAT1 phosphorylation (A), and macrophage polarization were assessed (B). Results are shown as mean $\pm$ SE (n = 3–4).

become chronic, they are likely to promote local macrophage dysfunction, which, in turn, is likely to increase inflammation and impair inflammation resolution at those sites.

Overexpression of MKP-1 not only prevents monocyte priming and hypersensitization to chemokines⁴, but also protected metabolically stressed macrophages against defective autophagy, sensitization to oxysterol-induced cell death, and skewed M1/M2 polarization. These results provide further evidence that MKP-1 is a key regulator of monocyte and macrophage function, activation, and plasticity, and that MKP-1 deficiency induced by metabolic stress, promotes macrophage dysfunction, a process that may play a fundamental role in the transition of acute inflammation to states of chronic inflammation typically associated with metabolic disorders. These findings also suggest that therapeutic or pharmacological strategies aimed at preserving monocyte and macrophage MKP-1 activity are likely to protect against chronic inflammatory disease associated with metabolic disorders.

Another intriguing possibility is that MKP-1 deficiency occurs early on during macrophage development, i.e. in circulating monocytes exposed to hyperglycemia and dyslipidemia, or even in as early as during myelopoiesis, and leads to the epigenetic reprogramming of blood monocytes. MKP-1 is the only known mammalian histone H3 serine 10 phosphatase⁵². It is well-known that during mitosis serine 10 on histone H3 is highly phosphorylated on condensed chromosomes^{53,54}, and phosphorylated serine 10 has therefore been used as mitotic marker. However, recent studies have demonstrated that a diverse array of stimuli are able to rapidly induce phosphorylation of serine 10 on histone H3, a process that is associated with transcriptional activation of immediate early genes, such as c-fos and c-jun^{55,56}. Phosphorylation of serine 10 on histone H3 is also known to mediate expression of genes related to inflammation such as cyclooxygenase (COX)-2⁵⁷, interleukin (IL)-6, and IL-8^{58,59}. Therefore, it is conceivable that MKP-1 deficiency induced by metabolic stress leads to epigenetic changes in blood monocytes, at least in part mediated through serine 10 phosphorylation on histone H3 (Suppl. Fig. 8). Epigenetic reprogramming of blood monocytes in response to chronic metabolic stress could account for the dysregulation of monocyte-derived macrophages associated with chronic inflammation and impaired wound healing, hallmarks of metabolic disorders such as hyperlipidemia and diabetes.

In summary, we discovered a new mechanistic link between metabolic disorders, macrophage dysfunction, and atherogenesis. With the identification of macrophage MKP-1 as an early sensor of metabolic stress and a master regulator of macrophage autophagy, apoptosis, and polarization, and loss of MKP-1 activity as a key mechanism for macrophage dysfunction induced by metabolic stress, both *in vitro* and *in vivo*, we may also have identified a novel biomarker and potential therapeutic target for the prevention and treatment of atherosclerosis.

Methods

Detailed methods are described in the Supplemental Materials.

Animals and diets. Female LDL-R^{-/-} recipient mice and female C57BL/6 donor mice were obtained from Jackson Labs (Bar Harbor, ME). Female MKP-1^{-/-} donor mice were kindly provided by the laboratory of Dr. Robert Kramer. All mice had been backcrossed to C57BL/6 background for more than 10 generations. To induce hypercholesterolemia, bone marrow recipient mice were switched to a high fat diet (HFD; 21% milk fat and 0.2% cholesterol, diet no. F5540, Bio-Serv, Frenchtown, NJ). Mice were maintained on HFD for 6 or 12 weeks. All studies were performed in accordance with the guidelines and regulations of and with the approval of the UTHSCSA Institutional Animal Care and Use Committee.

Bone Marrow Cell Collection. On the day of bone marrow transplantation, bone marrow cell suspensions were collected from C57BL/6 and MKP-1^{-/-} mice.

Irradiation and Bone Marrow Transplantation. LDL-R^{-/-} recipient mice were maintained on antibiotics for the duration of the study. Before transplantation, recipient LDL-R^{-/-} mice received 2 equal doses of 4.7 Gy, with 3 h between each dose (9.4 Gy total, Cobalt-60 Irradiator). Animals were given a 4 h recovery period prior to bone marrow transplantation. LDL-R^{-/-} mice were divided into two groups based on the strain of their donors (n = 30/group). Bone marrow cells (10–15 × 10⁶ cells in 150–300 µl) were then injected via the retro-orbital sinus. Animals were given a 4 weeks of recovery prior to initiation of HFD feeding.

Plasma Cholesterol, Triglycerides, and Lipoprotein Profile. Mice were fasted overnight prior to euthanasia and blood was collected by cardiac puncture. Plasma cholesterol and triglycerides were quantified using enzymatic assay kits per manufacturer's protocol (Wako Chemicals USA, Inc., Richmond, VA). Plasma was pooled from each treatment group and centrifuged to clarify the sample and 100 µl of pooled plasma was used for size exclusion chromatography.

In Vivo Macrophage Recruitment Assay. Each mouse received two Matrigel plugs three days prior to euthanasia as described previously⁵.

Histological and Immunohistochemical Analyses of Heart, Aorta, and Aortic Root. After peritoneal lavage, the entire aorta with the intact heart was dissected free of fat and removed. The aortas (proximal ascending aorta to the bifurcation) were fixed in 4% PFA for 48 h prior to staining for *en face* analysis. To determine the extent of the atherosclerosis, aortas were stained with Oil Red O (ORO; Sigma-Aldrich, St. Louis, MO). To further characterize lesions, neutral and intracellular lipid content were quantified by staining sections of heart tissue with ORO. Sections of aortic root were blocked with 10% normal Goat serum and incubated overnight at 4 °C with rat anti-mouse CD68:biotin (1:200; Bio-Rad Laboratories, Inc., Raleigh, NC) primary antibody. Tissues were also stained with antibodies to detect Signaling adaptor p62 (1:200; SQSTM1, Pierce, Rockford, IL), LC3B (1:200; Pierce, Rockford, IL), and the phosphorylation status of signal transducer and activator of transcription 1

(STAT1) at Ser 727 [Phospho-Stat1 (Ser727); 1:100; Cells Signaling Technology, Boston, MA)]. Selected sections of the aortic root were used for the detection of apoptotic and necrotic cells in atherosclerotic lesions. Tissues were analyzed by terminal dUTP terminal nick end-labeling (TUNEL assay; Roche Diagnostics, Indianapolis, IN) according to the manufacturer's instruction.

Cell Culture. Resident peritoneal macrophages were collected from C57BL/6 and MKP-1^{-/-} mice by lavage. Adherent macrophages were cultured in RPMI complete supplemented with 10% fetal bovine serum. Bone marrow-derived macrophages (BMDMs) were generated from C57BL/6 mice. Fresh bone marrow cells were cultured for 7 days in RPMI complete medium supplemented with 10% FBS and 50 ng/ml M-CSF. Metabolic stress was induced by incubating macrophages for 24 h with 100 µg/ml LDL and 20 mmol/L D-glucose (LDL + HG). M1 polarization was induced with IFN- γ plus TNF- α (PeproTech; 50 and 10 ng/ml, respectively)⁶⁰. Alternative (M2) polarization was induced by stimulating cells for 24 h with IL-4 (PeproTech; 10 ng/ml)⁶⁰. Macrophages that were grown in the absence of the above cytokines were considered nonpolarized or M0.

Lentiviral Transduction of Macrophages. All lentiviral supernatants were prepared by cotransfection of HEK-293T cells with one of the vector transfer constructs, murine ecotropic envelope vector (pCAG-Eco, Addgene Plasmid # 35617), and the packaging vectors pMDLg/pRRE and RSV-Rev. For gene transduction, bone marrow-derived macrophages were used on day 5 of differentiation. Two million cells were plated in 12-well plates, and viruses (MOI = 20) were added to each well in the presence of 6 µg/ml of DEAE-dextran sulfate (Sigma-Aldrich, St. Louis, MO).

Gene Expression. Total RNA was isolated from cells using PureLink RNA Mini Kit (Ambion, Grand Island, NY). Reverse transcription was performed using a QuantiTect Reverse Transcription Kit (Qiagen, Valencia, CA). Each cDNA sample was then separated into 48 separate reactions for qPCR analysis using the BioMark 48 × 48 dynamic array nanofluidic chip (Fluidigm Inc., USA) according to manufacturer's instruction. The 40 individual Taqman primer-probe mixtures (Applied Biosystems) specific for individual transcripts of interest are listed in Table S1. Amplification data were analyzed using SDS2.4 software (Applied Biosystems, Grand Island, NY), and gene expression levels were normalized to *Hprt* as the housekeeping gene.

Statistics. Data were analyzed using ANOVA (Sigma Stat 12.0). Data were tested for use of parametric or nonparametric post hoc analysis, and multiple comparisons were performed by using the Least Significant Difference method. All data are presented as mean ± SE of at least 3 independent experiments. Results were considered statistically significant at the $P < 0.05$ level.

References

- Libby, P. Inflammation in atherosclerosis. *Nature* **420**, 868–874 (2002).
- Back, M. & Hansson, G. K. Anti-inflammatory therapies for atherosclerosis. *Nat Rev Cardiol* **12**, 199–211 (2015).
- Moore, K. J. & Tabas, I. Macrophages in the pathogenesis of atherosclerosis. *Cell* **145**, 341–355 (2011).
- Kim, H. S., Ullevig, S. L., Zamora, D., Lee, C. F. & Asmis, R. Redox regulation of MAPK phosphatase 1 controls monocyte migration and macrophage recruitment. *Proc Natl Acad Sci USA* **109**, E2803–2812 (2012).
- Ullevig, S. *et al.* NADPH oxidase 4 mediates monocyte priming and accelerated chemotaxis induced by metabolic stress. *Arterioscler Thromb Vasc Biol* **32**, 415–426 (2012).
- Martinet, W. & De Meyer, G. R. Autophagy in atherosclerosis: a cell survival and death phenomenon with therapeutic potential. *Circ Res* **104**, 304–317 (2009).
- Kockx, M. M. & Herman, A. G. Apoptosis in atherosclerosis: beneficial or detrimental? *Cardiovasc Res* **45**, 736–746 (2000).
- Tabas, I. Consequences and therapeutic implications of macrophage apoptosis in atherosclerosis: the importance of lesion stage and phagocytic efficiency. *Arterioscler Thromb Vasc Biol* **25**, 2255–2264 (2005).
- Wolfs, I. M., Donners, M. M. & de Winther, M. P. Differentiation factors and cytokines in the atherosclerotic plaque micro-environment as a trigger for macrophage polarisation. *Thromb Haemost* **106**, 763–771 (2011).
- Mantovani, A., Garlanda, C. & Locati, M. Macrophage diversity and polarization in atherosclerosis: a question of balance. *Arterioscler Thromb Vasc Biol* **29**, 1419–1423 (2009).
- Mei, S. *et al.* p38 mitogen-activated protein kinase (MAPK) promotes cholesterol ester accumulation in macrophages through inhibition of macroautophagy. *J Biol Chem* **287**, 11761–11768 (2012).
- Gumireddy, K., Reddy, C. D. & Swamy, N. Mitogen-activated protein kinase pathway mediates DBP-maf-induced apoptosis in RAW 264.7 macrophages. *J Cell Biochem* **90**, 87–96 (2003).
- Singhal, P. C. *et al.* Role of p38 mitogen-activated protein kinase phosphorylation and Fas-Fas ligand interaction in morphine-induced macrophage apoptosis. *J Immunol* **168**, 4025–4033 (2002).
- Zhang, W., Xu, W. & Xiong, S. Macrophage differentiation and polarization via phosphatidylinositol 3-kinase/Akt-ERK signaling pathway conferred by serum amyloid P component. *J Immunol* **187**, 1764–1777 (2011).
- Patterson, K. I., Brummer, T., O'Brien, P. M. & Daly, R. J. Dual-specificity phosphatases: critical regulators with diverse cellular targets. *Biochem J* **418**, 475–489 (2009).
- Chu, Y., Solski, P. A., Khosravi-Far, R., Der, C. J. & Kelly, K. The mitogen-activated protein kinase phosphatases PAC1, MKP-1, and MKP-2 have unique substrate specificities and reduced activity *in vivo* toward the ERK2 sevenmaker mutation. *J Biol Chem* **271**, 6497–6501 (1996).
- Groom, L. A., Sneddon, A. A., Alessi, D. R., Dowd, S. & Keyse, S. M. Differential regulation of the MAP, SAP and RK/p38 kinases by Pyst1, a novel cytosolic dual-specificity phosphatase. *EMBO J* **15**, 3621–3632 (1996).
- Franklin, C. C., Srikanth, S. & Kraft, A. S. Conditional expression of mitogen-activated protein kinase phosphatase-1, MKP-1, is cytoprotective against UV-induced apoptosis. *Proc Natl Acad Sci USA* **95**, 3014–3019 (1998).
- Perdiguerro, E. *et al.* p38/MKP-1-regulated AKT coordinates macrophage transitions and resolution of inflammation during tissue repair. *J Cell Biol* **195**, 307–322 (2011).
- Kiffin, R., Bandyopadhyay, U. & Cuervo, A. M. Oxidative stress and autophagy. *Antioxid Redox Signal* **8**, 152–162 (2006).
- Liao, X. *et al.* Macrophage autophagy plays a protective role in advanced atherosclerosis. *Cell Metab* **15**, 545–553 (2012).
- Cagnol, S. & Chambard, J. C. ERK and cell death: mechanisms of ERK-induced cell death—apoptosis, autophagy and senescence. *FEBS J* **277**, 2–21 (2010).

23. Webber, J. L. & Tooze, S. A. Coordinated regulation of autophagy by p38alpha MAPK through mAtg9 and p38IP. *EMBO J* **29**, 27–40 (2010).
24. Wei, Y., Pattingre, S., Sinha, S., Bassik, M. & Levine, B. JNK1-mediated phosphorylation of Bcl-2 regulates starvation-induced autophagy. *Mol Cell* **30**, 678–688 (2008).
25. Mizushima, N. & Yoshimori, T. How to interpret LC3 immunoblotting. *Autophagy* **3**, 542–545 (2007).
26. Mathew, R. *et al.* Autophagy suppresses tumorigenesis through elimination of p62. *Cell* **137**, 1062–1075 (2009).
27. Devries-Seimon, T. *et al.* Cholesterol-induced macrophage apoptosis requires ER stress pathways and engagement of the type A scavenger receptor. *J Cell Biol* **171**, 61–73 (2005).
28. Liang, S. L., Liu, H. & Zhou, A. Lovastatin-induced apoptosis in macrophages through the Rac1/Cdc42/JNK pathway. *J Immunol* **177**, 651–656 (2006).
29. Gautier, E. L. *et al.* Macrophage apoptosis exerts divergent effects on atherogenesis as a function of lesion stage. *Circulation* **119**, 1795–1804 (2009).
30. Lyons, M. A. & Brown, A. J. 7-Ketocholesterol. *Int J Biochem Cell Biol* **31**, 369–375 (1999).
31. Larsson, D. A., Baird, S., Nyhalah, J. D., Yuan, X. M. & Li, W. Oxysterol mixtures, in atheroma-relevant proportions, display synergistic and proapoptotic effects. *Free Radic Biol Med* **41**, 902–910 (2006).
32. Sica, A. & Mantovani, A. Macrophage plasticity and polarization: *in vivo* veritas. *J Clin Invest* **122**, 787–795 (2012).
33. Lawrence, T. & Natoli, G. Transcriptional regulation of macrophage polarization: enabling diversity with identity. *Nat Rev Immunol* **11**, 750–761 (2011).
34. Tugal, D., Liao, X. & Jain, M. K. Transcriptional control of macrophage polarization. *Arterioscler Thromb Vasc Biol* **33**, 1135–1144 (2013).
35. Goh, K. C., Haque, S. J. & Williams, B. R. p38 MAP kinase is required for STAT1 serine phosphorylation and transcriptional activation induced by interferons. *EMBO J* **18**, 5601–5608 (1999).
36. Varinou, L. *et al.* Phosphorylation of the Stat1 transactivation domain is required for full-fledged IFN-gamma-dependent innate immunity. *Immunity* **19**, 793–802 (2003).
37. Kovarik, P. *et al.* Stress-induced phosphorylation of STAT1 at Ser727 requires p38 mitogen-activated protein kinase whereas IFN-gamma uses a different signaling pathway. *Proc Natl Acad Sci USA* **96**, 13956–13961 (1999).
38. Liu, D., Scafidi, J., Prada, A. E., Zahedi, K. & Davis, A. E. 3rd. Nuclear phosphatases and the proteasome in suppression of STAT1 activity in hepatocytes. *Biochem Biophys Res Commun* **299**, 574–580 (2002).
39. Ichikawa, T. *et al.* Nitroalkenes suppress lipopolysaccharide-induced signal transducer and activator of transcription signaling in macrophages: a critical role of mitogen-activated protein kinase phosphatase 1. *Endocrinology* **149**, 4086–4094 (2008).
40. Spurgeon, S. L., Jones, R. C. & Ramakrishnan, R. High throughput gene expression measurement with real time PCR in a microfluidic dynamic array. *PLoS One* **3**, e1662 (2008).
41. Laurell, H. *et al.* Correction of RT-qPCR data for genomic DNA-derived signals with ValidPrime. *Nucleic Acids Res* **40**, e51 (2012).
42. Tabas, I. Macrophage death and defective inflammation resolution in atherosclerosis. *Nat Rev Immunol* **10**, 36–46 (2010).
43. Franklin, C. C. & Kraft, A. S. Conditional expression of the mitogen-activated protein kinase (MAPK) phosphatase MKP-1 preferentially inhibits p38 MAPK and stress-activated protein kinase in U937 cells. *J Biol Chem* **272**, 16917–16923 (1997).
44. Chi, H. *et al.* Dynamic regulation of pro- and anti-inflammatory cytokines by MAPK phosphatase 1 (MKP-1) in innate immune responses. *Proc Natl Acad Sci USA* **103**, 2274–2279 (2006).
45. Korhonen, R. *et al.* Attenuation of the acute inflammatory response by dual specificity phosphatase 1 by inhibition of p38 MAP kinase. *Mol Immunol* **48**, 2059–2068 (2011).
46. Wu, J. J. & Bennett, A. M. Essential role for mitogen-activated protein (MAP) kinase phosphatase-1 in stress-responsive MAP kinase and cell survival signaling. *J Biol Chem* **280**, 16461–16466 (2005).
47. Henson, S. M. *et al.* p38 signaling inhibits mTORC1-independent autophagy in senescent human CD8(+) T cells. *J Clin Invest* **124**, 4004–4016 (2014).
48. Lum, J. J. *et al.* Growth factor regulation of autophagy and cell survival in the absence of apoptosis. *Cell* **120**, 237–248 (2005).
49. Lum, J. J., DeBerardinis, R. J. & Thompson, C. B. Autophagy in metazoans: cell survival in the land of plenty. *Nat Rev Mol Cell Biol* **6**, 439–448 (2005).
50. Ramsauer, K. *et al.* p38 MAPK enhances STAT1-dependent transcription independently of Ser-727 phosphorylation. *Proc Natl Acad Sci USA* **99**, 12859–12864 (2002).
51. Zykova, T. A., Zhang, Y., Zhu, F., Bode, A. M. & Dong, Z. The signal transduction networks required for phosphorylation of STAT1 at Ser727 in mouse epidermal JB6 cells in the UVB response and inhibitory mechanisms of tea polyphenols. *Carcinogenesis* **26**, 331–342 (2005).
52. Kinney, C. M. *et al.* Histone H3 as a novel substrate for MAP kinase phosphatase-1. *Am J Physiol Cell Physiol* **296**, C242–249 (2009).
53. Hendzel, M. J. *et al.* Mitosis-specific phosphorylation of histone H3 initiates primarily within pericentromeric heterochromatin during G2 and spreads in an ordered fashion coincident with mitotic chromosome condensation. *Chromosoma* **106**, 348–360 (1997).
54. Cheung, P., Allis, C. D. & Sassone-Corsi, P. Signaling to chromatin through histone modifications. *Cell* **103**, 263–271 (2000).
55. Clayton, A. L., Rose, S., Barratt, M. J. & Mahadevan, L. C. Phosphoacetylation of histone H3 on c-fos- and c-jun-associated nucleosomes upon gene activation. *EMBO J* **19**, 3714–3726 (2000).
56. Thomson, S., Clayton, A. L. & Mahadevan, L. C. Independent dynamic regulation of histone phosphorylation and acetylation during immediate-early gene induction. *Mol Cell* **8**, 1231–1241 (2001).
57. Park, G. Y., Joo, M., Pedchenko, T., Blackwell, T. S. & Christman, J. W. Regulation of macrophage cyclooxygenase-2 gene expression by modifications of histone H3. *Am J Physiol Lung Cell Mol Physiol* **286**, L956–962 (2004).
58. Yamamoto, Y., Verma, U. N., Prajapati, S., Kwak, Y. T. & Gaynor, R. B. Histone H3 phosphorylation by IKK-alpha is critical for cytokine-induced gene expression. *Nature* **423**, 655–659 (2003).
59. Anest, V. *et al.* A nucleosomal function for IkappaB kinase-alpha in NF-kappaB-dependent gene expression. *Nature* **423**, 659–663 (2003).
60. Tavakoli, S., Zamora, D., Ullevig, S. & Asmis, R. Bioenergetic profiles diverge during macrophage polarization: implications for the interpretation of 18F-FDG PET imaging of atherosclerosis. *J Nucl Med* **54**, 1661–1667 (2013).

Acknowledgements

The authors would like to thank D. Vanegas and J. Han for technical assistance. This work was supported by grants to R.A. from the NIH (HL70963 and AT006885) and the AHA (085501F). H.S.K. was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIP) (NRF-2014R1A5A2009392 and NRF-2015R1D1A1A01058340).

Author Contributions

H.S.K. and R.A. designed the study. H.S.K., S.T., L.A.P. and H.N.N. performed experiments. H.S.K. and R.A. analyzed data. H.S.K. and R.A. wrote the paper.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Kim, H. S. *et al.* Monocytic MKP-1 is a Sensor of the Metabolic Environment and Regulates Function and Phenotypic Fate of Monocyte-Derived Macrophages in Atherosclerosis. *Sci. Rep.* **6**, 34223; doi: 10.1038/srep34223 (2016).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016