

SCIENTIFIC REPORTS



OPEN

Transparency and tunable slow and fast light in a nonlinear optomechanical cavity

Ling Li¹, Wenjie Nie¹ & Aixi Chen^{1,2}

Received: 01 August 2016
Accepted: 23 September 2016
Published: 11 October 2016

We investigate theoretically the optical response of the output field and the tunable slow and fast light in a nonlinear optomechanical cavity with a degenerate optical parametric amplifier (OPA) and a higher order excited atomic ensemble. Studies show that the higher-order-excitation atom which is similar to the degenerate OPA that acts as a nonlinear medium, induces an additional dip in absorption spectrum of the probe field. The coherence of the mechanical oscillator leads to split the peak in absorption in the probe field spectrum so that the phenomenon of optomechanically induced transparency (OMIT) is generated from the output probe field. In particular, the presence of nonlinearities with the degenerate OPA and the higher order excited atoms can affect significantly the width of the transparency windows, providing an additional flexibility for controlling optical properties. Furthermore, in the presence of the degenerate OPA, the optical-response properties for the probe field become phase-sensitive so that a tunable switch from slow to fast light can be realized.

It is well-known that a three-level atomic medium driven by a strong controlling field can become transparent for a weak probe field, which results from the quantum interference between the two different pathways of excitation in atomic system. This phenomenon is a typical quantum coherent effect called the electromagnetically induced transparency (EIT)^{1–3}, which has been widely investigated both in theory^{4–6} and in experiment^{2,7}. It is shown that EIT is important for various applications^{8,9} such as slow light, light storage and the production of a giant nonlinear effect. The phenomenon of EIT has been recently observed in the other solid state systems, i.e., quantum wells¹⁰, metamaterial¹¹ and nitrogen-vacancy centers¹². In addition, the double EIT windows in multi-level atomic systems have been studied in detail¹³.

On the other hand, the propagation of a weak probe field in optomechanical system¹⁴ can be coherently manipulated through the driving of a strong coupling field which leads to an optomechanical coupling between the cavity mode and the mechanical oscillator. Several important quantum optomechanical characteristics, i.e., optomechanical entanglement^{15,16} and optically cooling mechanical mode^{17–20} as well as transitions between classical and quantum behaviors of a mechanical system^{21,22} have been extensively investigated by pumping the cavity with external laser fields. In particular, a phenomenon of EIT-like, called generally the optomechanically induced transparency (OMIT), has been shown theoretically^{23–29} and observed experimentally in optical cavities^{30–33} and microwave cavities³⁴. OMIT can be designed to slow and switch a probe signal³⁵ and further used to store light³⁶. The transparency behavior in an optomechanical system also advance the ground-state cooling of mechanical motions and optomechanical entanglement between the optical and mechanical modes^{37–39}.

Upon combining the optomechanical system with cavity quantum electrodynamics (QED)⁴⁰, the influence of an additional atomic medium on the OMIT in a hybrid optomechanical system^{41,42} has been investigated in detail, where the low-atomic excitation limit of atoms requires the single-atom excitation probability to be much less than 1^{43,44}. However, this may restrict the amplitude of the driving field in this kind of hybrid optomechanical system so that the optomechanical coupling between the optical and mechanical modes can not be advanced further by increasing the pump power. In order to relax the constraints on the driving of the system or the atom-field coupling strength, we need to go beyond the low-atomic excitation limit for the atomic medium embedded in an optomechanical system. In previous paper, we have shown that when the low excitation condition of atoms breaks slightly, a large driving but a relatively small atom-field detuning can be applied to help observe OMIT behavior in a levitated optomechanical system⁴⁵.

¹Department of Applied Physics, East China Jiaotong University, Nanchang, 330013, China. ²Institute for Quantum Computing, University of Waterloo, Ontario N2L 3G1, Canada. Correspondence and requests for materials should be addressed to W.N. (email: niewenjiezh@sinacn) or A.C. (email: aixichen@ecjtu.edu.cn)




Figure 1. Schematic of the setup studied in the paper. The optomechanical cavity contains N identical two-level atoms and a degenerate OPA. The cavity mode is driven by a strong input laser field and a weak probe field through the left cavity mirror. The nonlinear crystal is pumped by an additional laser beam to produce parametric amplification.

In addition, we know the nonlinear optical effect of the optomechanical system can be obviously enhanced by adding a degenerate optical parametric amplifier (OPA). The degenerate OPA placed in the optomechanical system can increase the effective optomechanical coupling between the movable mirror and the cavity field, which results from increase of the photon numbers in the cavity. Some researches about the influences of the degenerate OPA on the propagation of the probe field are reported in refs 28,46,47, where the optical properties of the output field, i.e., the width of OMIT, can be controlled easily by adjusting the pump amplitude of the degenerate OPA. On the other hand, when the degenerate OPA is included, the quantum interference effect between the probe field and the generated anti-Stokes field depends strongly on the phase of the degenerate OPA so that the optical-response properties for the probe field will become phase-sensitive. As so far, the response of nonlinear optomechanical cavity with an excited atomic ensemble to a weak probe light is not reported under the condition of the existence of the degenerate OPA. In this paper, we investigate the properties of absorption and dispersion of the probe field propagating the optomechanical system including low- or high-excitation atomic medium and a degenerate OPA. Also, we contrast the role of the nonlinearities of the higher-order excitation of atoms with one of the degenerate OPA in the optical properties of the output field. Further, OMIT behavior in absorption of the probe field generated through the optomechanical coupling is discussed in detail, which is influenced by the higher order excitation of the atoms and the degenerate OPA. We also discuss in detail how to control the switch from slow to fast light of the output probe field by adjusting the phase of the field driving the degenerate OPA. The result suggests that the phase-sensitive interference effects have potential applications to provide new tools for controlling and engineering light propagation.

Optomechanical Model and Hamiltonian

As shown schematically in Fig. 1, the model that we consider is a nonlinear optomechanical system, where a degenerate OPA and N identical two-level atoms (with transition frequency ω_b and decay rate γ_b) are placed in a Fabry-Férot cavity with length L consisting of one fixed mirror and one movable mirror. Specifically, we consider a quadrupole transition between the $6s^2\ ^1S_0$ ground state $|g\rangle$ and the $6s6p\ ^3P_1$ excited state $|e\rangle$ of atomic barium for the two-level atoms^{48,49}. The transition wavelength between the two states in atomic barium and the decay rate of the excited state to the ground state are $\lambda = 791\text{ nm}$ and $\gamma_b = 47\text{ kHz}$, respectively⁴⁸. Further, we consider a realistic optical cavity, where the intracavity photon leakage can be occurred through the input and the movable mirror^{33,50,51}. We assume the decay rates of two cavity mirrors are equal without loss of generality, i.e., $\kappa_0 = \kappa_1$, where κ_0 and κ_1 are the decay rates of the cavity field through the input and the movable mirrors, respectively.

The cavity mode with frequency ω_c is driven by a classical control field with frequency ω_f and amplitude ε_c as well as a weak probe field with frequency ω_p and amplitude ε_p , which then exerts an optical radiation pressure on the movable cavity mirror. Moreover, the system is pumped by an additional laser beam with coupling coefficient β to produce parametric amplification^{28,46,47}. In general, the movable mirror is treated as a quantum-mechanical harmonic oscillator with resonance frequency ω_m , effective mass m and damping rate γ_m . Then, the total Hamiltonian of the system can be written as^{15,28,43}

$$\begin{aligned}
 H = & \hbar\omega_c a^\dagger a + \frac{\hbar\omega_b}{2} \sum_{i=1}^N \sigma_z^{(i)} + \frac{p^2}{2m} + \frac{m\omega_m^2 q^2}{2} - \hbar G_0 a^\dagger a q + \left(\hbar g a \sum_{i=1}^N \sigma_+^{(i)} + H.c. \right) \\
 & + i\hbar G_A (e^{i\theta} a^{\dagger 2} - H.c.) + i\hbar (\varepsilon_c a^\dagger e^{-i\omega_f t} + \varepsilon_p a^\dagger e^{-i\omega_p t} - H.c.), \tag{1}
 \end{aligned}$$

where the first term describes the free Hamiltonian of cavity field and a (a^\dagger) is the annihilation (creation) operator of the cavity mode satisfying the commutation relation $[a, a^\dagger] = 1$. The second term is the free Hamiltonian of the atomic ensemble, where the ground state and the excited state of the i th two-level atom are described by $|g\rangle^{(i)}$ and $|e\rangle^{(i)}$ and therefore $\sigma_z^{(i)} = |e\rangle^{(i)}\langle e| - |g\rangle^{(i)}\langle g|$. In addition, the pseudospin-1/2 operators $\sigma_\pm^{(i)} = |e\rangle^{(i)}\langle g|$ and $\sigma_-^{(i)} = |g\rangle^{(i)}\langle e|$ for the i th atom satisfy the commutation relations $[\sigma_+^{(i)}, \sigma_-^{(i)}] = \sigma_z^{(i)}$ and $[\sigma_z^{(i)}, \sigma_\pm^{(i)}] = \pm 2\sigma_\pm^{(i)}$. The third and fourth terms are the kinetic and potential energies of the movable mirror, respectively; q and p are the position and momentum operators for the movable mirror with the commutation relation $[q, p] = i\hbar$. The fifth term describes the optomechanical coupling of the cavity mode with the movable mirror; $G_0 = \omega_c L$ is the optomechanical coupling strength between the mechanical mode and cavity mode⁵³. L is the length of the cavity. The last term in the first line denotes the interaction of the atomic ensemble with the driven cavity field, where g represents the averaged atom-field coupling strength^{43,44,52}. The first term in the second line describes the coupling of the cavity mode with the degenerate OPA; G_A is the nonlinear gain of the degenerate OPA, which is proportional to the pump amplitude, E_{OPA} , i.e., $G_A = \beta|E_{OPA}|$; θ is the phase of the field driving the OPA^{28,46,47}. The last terms describe the interaction of the cavity field with the coupling field and that of the cavity field with the probe field, with the amplitude $\varepsilon_c = \sqrt{\frac{2\kappa_0 P_c}{\hbar\omega_f}}$ and $\varepsilon_p = \sqrt{\frac{2\kappa_0 P_p}{\hbar\omega_p}}$, respectively. P_c and P_p are the laser powers.

For simplification, we use the Holstein-Primakoff transformation⁵⁴, which is defined as $\sum_{i=1}^N \sigma_+^{(i)} = B^\dagger \sqrt{N - B^\dagger B}$, $\sum_{i=1}^N \sigma_-^{(i)} = \sqrt{N - B^\dagger B} B$ and $\sum_{i=1}^N \sigma_z^{(i)} = 2B^\dagger B - N$, to rewrite Hamiltonian (0), where B and B^\dagger satisfy the fundamental commutation relation $[B, B^\dagger] = 1$ ⁵⁴. In general, in the low-excitation limit $\langle B^\dagger B \rangle / N \ll 1$ for the atoms, the Holstein-Primakoff transformation becomes $\sum_{i=1}^N \sigma_+^{(i)} = \sqrt{N} B^\dagger$ and $\sum_{i=1}^N \sigma_-^{(i)} = \sqrt{N} B$. In the present system, we assume that the low-excitation condition is broken slightly for the atomic ensemble so that the higher orders of the Holstein-Primakoff transformation should be included^{45,55}, i.e., $\sum_{i=1}^N \sigma_+^{(i)} = \sqrt{N} B^\dagger \left(1 - \frac{B^\dagger B}{2N}\right)$, $\sum_{i=1}^N \sigma_-^{(i)} = \sqrt{N} \left(1 - \frac{B^\dagger B}{2N}\right) B$ and $\sum_{i=1}^N \sigma_z^{(i)} = 2B^\dagger B - N$. Then, in the interaction picture with respect to $H_0 = \hbar\omega_f (a^\dagger a + B^\dagger B)$, the Hamiltonian of the total system, Eq. (0), is rewritten as

$$\begin{aligned}
 H = & \hbar\Delta_a a^\dagger a + \hbar\Delta_b B^\dagger B + \frac{p^2}{2m} + \frac{m\omega_m^2 q^2}{2} - \hbar G_0 a^\dagger a q \\
 & + \left(\hbar G a B^\dagger - \frac{\hbar G a B^{\dagger 2} B}{2N} + H.c. \right) \\
 & + i\hbar G_A (e^{i\theta} a^{\dagger 2} - H.c.) + i\hbar \varepsilon_c (a^\dagger - a) + i\hbar (\varepsilon_p e^{-i\delta t} a^\dagger - \varepsilon_p^* e^{i\delta t} a), \tag{2}
 \end{aligned}$$

where $G = g\sqrt{N}$ is the collective coupling strength of the atomic ensemble with the cavity field. $\Delta_a = \omega_c - \omega_f$, $\Delta_b = \omega_b - \omega_f$ and $\delta = \omega_p - \omega_f$ are the detunings. In the derivation of Eq. (2), a constant term $N\omega_a/2$ has been neglected.

System Dynamics and Equation of Motion

For a detailed analysis of the system, we consider photon losses in the cavity through the input mirror with decay rate κ_0 and the movable mirror with decay rate κ_b ; and Brownian noise acting on the mirror as well as decays associated with the atoms. Based on the Hamiltonian in Eq. (2), the quantum dynamics of the system can be described by the following Heisenberg-Langevin equation³³:

$$\begin{aligned}
 \dot{q} &= p/m, \\
 \dot{p} &= -m\omega_m^2 q + \hbar G_0 a^\dagger a - \gamma_m p + \xi(t), \\
 \dot{a} &= -(\kappa_0 + \kappa_1 + i\Delta_a) a - iGB + \frac{iG}{2N} B^\dagger B^2 + iG_0 a q + 2\hbar G_A e^{i\theta} a^\dagger \\
 &\quad + \varepsilon_c + \varepsilon_p e^{-i\delta t} + \sqrt{2\kappa_0} a_0^{in} + \sqrt{2\kappa_1} a_1^{in}, \\
 \dot{B} &= -(\gamma_b + i\Delta_b) B + \frac{iGaB^\dagger B}{N} - iGa + \frac{iGa^\dagger B^2}{2N} + \sqrt{2\gamma_b} B_{in}, \tag{3}
 \end{aligned}$$

where a_0^{in} , a_1^{in} and B_{in} are the input vacuum noise operators with zero mean value corresponding to the input mirror, the movable mirror and the atomic ensemble, which are fully characterized by the nonzero correlation functions $\langle a_j^{in}(t) a_j^{in\dagger}(t') \rangle = \delta(t - t')(j = 0, 1)$ and $\langle B_{in}(t) B_{in}^\dagger(t') \rangle = \delta(t - t')$, respectively⁵⁶. $\xi(t)$ is the Brownian stochastic force with zero mean value and its correlation function is characterized by $\langle \xi(t)\xi(t') \rangle = m\hbar\gamma_m \int_{2\pi}^{d\omega} e^{-i\omega(t-t')\omega} \left[1 + \coth\left(\frac{\hbar\omega}{2k_B T}\right) \right]$ ⁵⁷, where k_B is the Boltzmann constant and T is the temperature of the reservoir related to the movable mirror.

In order to studying the effect of the higher order excited atomic ensemble and the degenerate OPA on the optical properties of the output field in the optomechanical system, we need to investigate the motional equations

of the quantum fluctuations around the mean values. Further, the mean values at steady state for the movable mirror, atomic ensemble and cavity field can be obtained from Eq. (3) by setting all time derivatives to 0. These are found to be

$$\begin{aligned} p_s &= 0, \quad q_s = \frac{\hbar G_0 |a_s|^2}{m\omega_m^2}, \\ a_s &= \frac{(\kappa - i\Delta)E_c + 2G_A e^{i\theta} E_c^*}{\kappa^2 + \Delta^2 - 4G_A^2}, \\ 0 &= -(\gamma_b + i\Delta_b)b_s + ig a_s (|b_s|^2 - 1) + \frac{ig a_s^* b_s^2}{2}, \end{aligned} \quad (4)$$

where $\Delta = \Delta_a - G_0 q_s$, and $E_c = \varepsilon_c - iG\sqrt{N}(1 - |b_s|^2/2)b_s$ with $b_s = B_s/\sqrt{N}$. $\kappa = \kappa_0 + \kappa_1$ is the total cavity decay rate. It is seen from Eq. (4) that the steady-state values of the system depend strongly on the nonlinear gain G_A and the phase θ . Thus, the combine of the nonlinear optics and optomechanics may be used to control the optical properties of the output field.

To this end, one can split each operator in Eq. (3) into the steady-state mean value at the fixed point and a small quantum fluctuation, i.e., $O = O_s + \delta O$ ($O = a, B, p, q$). Further, we consider that there are a large number of photons in the cavity, i.e., $|a_s| \gg 1$, so that all the higher terms ($\delta o \delta o$) in Eq. (3) can be neglected. Then, the quantum Langevin equations for the fluctuations can be written as

$$\begin{aligned} \delta \dot{q} &= \delta p/m, \\ \delta \dot{p} &= -m\omega_m^2 \delta q + \hbar G_0 (a_s^* \delta a + a_s \delta a^\dagger) - \gamma_m \delta p + \xi(t), \\ \delta \dot{a} &= -(\kappa + i\Delta) \delta a + G_4 \delta a^\dagger + iG_0 a_s \delta q - iG_1 \delta B + iG_3 \delta B^\dagger + \varepsilon_p e^{-i\delta t} \\ &\quad + \sqrt{2\kappa_0} a_0^{in} + \sqrt{2\kappa_1} a_1^{in}, \\ \delta \dot{B} &= -(\gamma_b + i\Delta_b^0) \delta B - iG_1 \delta a + iG_2 \delta B^\dagger + iG_3 \delta a^\dagger + \sqrt{2\gamma_b} B_{in}, \end{aligned} \quad (5)$$

where $G_1 = G(1 - |b_s|^2)$, $G_2 = \frac{G a_s b_s}{\sqrt{N}}$, $G_3 = \frac{G b_s^2}{2}$, $G_4 = 2G_A e^{i\theta}$ and $\Delta_b^0 = \Delta_b - G(a_s b_s^* + a_s^* b_s)/\sqrt{N}$. In order to solve Eq. (5), we use the ansatz $\delta O = O_+ e^{-i\delta t} + O_- e^{i\delta t}$. Substituting this ansatz into Eq. (5), we can obtain the following solution of interest for the response of the cavity:

$$a_+ = \varepsilon_p \left[\beta_1 + iG_1 A_2 + iG_3 A_4 + (i\beta_5 + iG_1 A_1 - iG_3 A_3) \frac{i\beta_6 - iG_3^* A_2 - iG_1 A_4}{\beta_2 + iG_3^* A_1 - iG_1 A_3} \right]^{-1}, \quad (6)$$

where

$$A_1 = \frac{iG_3}{\beta_3} + \frac{iG_2 (iG_1 + \frac{G_2^* G_3}{\beta_3})}{\beta_3 \beta_4 - |G_2|^2}, \quad A_2 = \left[\frac{-iG_2 (\frac{G_1 G_2^*}{\beta_3} + iG_3^*)}{\beta_3 \beta_4 - |G_2|^2} - \frac{iG_1}{\beta_3} \right], \quad A_3 = \frac{G_2^* G_3}{\beta_4 - \frac{|G_2|^2}{\beta_3}} + iG_1, \quad A_4 = \frac{\frac{G_2^* G_1}{\beta_3} + iG_3^*}{\beta_4 - \frac{|G_2|^2}{\beta_3}},$$

$$\beta_0 = \frac{\hbar G_0^2}{m(\omega_m^2 - \delta^2 - i\delta\gamma_m)}, \quad \beta_1 = \kappa + i(\Delta - \delta) - \beta_0 |\alpha_s|^2, \quad \beta_2 = \kappa - i(\Delta + \delta) + \beta_0 |\alpha_s|^2, \quad \beta_3 = \gamma_b + i(\Delta_b^0 - \delta),$$

$$\beta_4 = \gamma_b - i(\Delta_b^0 + \delta), \quad \beta_5 = iG_4 - \beta_0 a_s^2 \text{ and } \beta_6 = iG_4 - \beta_0 a_s^{*2}.$$

The expression (6) and the steady-state values in Eq. (4) help us investigate the component of the output field oscillating at the probe frequency ω_p , which relates to the higher order excitation of the atomic ensemble $|b_s|^2$ as well as the degenerate OPA. In addition, the expression of a_- is not necessary since this describes the four-wave mixing with frequency $\omega_p - 2\omega_f$ for the driving field and the weak probe field. We can calculate the response of the system to all frequencies detected by the output field via the standard input-output theory⁵⁸ ($a_{out}(t) + \frac{\varepsilon_c + \varepsilon_p e^{-i\delta t}}{\sqrt{2\kappa_0}} = \sqrt{2\kappa_0} \langle a \rangle$). Using the above input-output relation and the ansatz $\delta a = a_+ e^{-i\delta t} + a_- e^{i\delta t}$, we can express the mean value of the output field as

$$\sqrt{2\kappa_0} \langle a_{out}(t) \rangle = (2\kappa_0 \alpha_s - \varepsilon_c) + (2\kappa_0 a_+ - \varepsilon_p) e^{-i\delta t} + 2\kappa_0 a_- e^{i\delta t}. \quad (7)$$

It is noted that the second term on the right-hand side of Eq. (7) corresponds to the response of the whole system to the weak probe field at frequency ω_p . Thus, we can examine the total output field at the frequency ω_p by defining an amplitude of the rescaled output field corresponding to the weak probe field as

$$E_{out} = 2\kappa_0 a_+ / \varepsilon_p, \quad (8)$$

where we have removed the constant term. The real and imaginary parts of the output probe field account for in-phase and out-phase quadratures of the output field spectra and can be written as $\text{Re}(E_{out}) = \kappa_0 (a_+ + a_+^*) / \varepsilon_p$ and $\text{Im}(E_{out}) = \kappa_0 (a_+ - a_+^*) / \varepsilon_p$, which describe the absorption and dispersion of the whole system to the weak probe field, respectively. In general, the modification of the probe response and the phenomenon of the transparency can be generated by the coupling the atoms or the mechanical motion with cavity field^{41,42}. In the present optomechanical system, we investigate in detail the generation of dips in absorption induced by the atom-field and optomechanical couplings as well as the degenerate OPA.

Moreover, in the region of the narrow transparency window the rapid phase dispersion $\varphi(\omega_p) = \arg(E_{out})$ can cause the group delay expressed as $\tau_g = \frac{\partial \varphi(\omega_p)}{\partial \omega_p}$. A positive group delay with $\tau_g > 0$ corresponds to slow light




Figure 2. The absorption $\text{Re}(E_{out})$ (a) and dispersion $\text{Im}(E_{out})$ (b) are plotted as a function of δ/ω_m in the absence and presence of the degenerate OPA. Here the optomechanical coupling is turned off and the low-excitation limit for the atoms is satisfied with $b_s = -0.01 + 0.04i$. When the degenerate OPA is included, we select the parameters $G_A = 0.7\kappa$ and $\theta = 3\pi/2$. Other parameters are chosen to be $\lambda_f = 791$ nm, $L = 0.001$ m, $\kappa = 2\pi \times 215 \times 10^3$ Hz, $\Delta = \Delta_a = \omega_m = 2\pi \times 947 \times 10^3$ Hz, $N = 10^6$, $g = 2\pi \times 2.3 \times 10^2$ Hz and $\gamma_b = 2\pi \times 7.5 \times 10^3$ Hz.

propagation and a negative group delay with $\tau_g < 0$ corresponds to fast light propagation^{30,32,59}. In the following section, we also investigate theoretically a tunable switch from slow to fast light in the nonlinear optomechanical cavity by adjusting the nonlinear gain of the OPA and the phase of the field driving the OPA.

Results

In this section, we numerically evaluate the values of phase quadratures $\text{Re}(E_{out})$ and $\text{Im}(E_{out})$ and quantify the slow and fast light effects through the corresponding output field a_+ . Further, in order to demonstrate the phenomenon of transparency and the group delay of the probe field in the system, we select the accessible parameters in optomechanical systems^{43,60}, i.e., the wavelength of the driving field $\lambda_f \approx 791$ nm, the total cavity length $L = 0.001$ m, the total cavity decay rate $\kappa = 2\pi \times 215 \times 10^3$ Hz and $\kappa_0/\kappa = 0.5$, the frequency of the moving mirror $\omega_m = 2\pi \times 947 \times 10^3$ Hz, the mechanical factor $Q = \omega_m/\gamma_m = 6700$, and the mass of the oscillating mirror $m = 25$ ng. In addition, we choose the parameters of atoms, i.e., the number of the atoms $N = 10^6$, the atom-field coupling strength $g = 2\pi \times 2.3 \times 10^2$ Hz and the decay rate of atom $\gamma_b = 2\pi \times 7.5 \times 10^3 \approx 47000$ Hz^{43,48}. We also consider that the cavity is driven on its red sideband, i.e., $\Delta = \omega_m$. The atom-field detuning Δ_b and the driving strength of the optical cavity ε_c can be calculated in terms of the excitation number of the atoms $|b_s|^2$ and the steady-state value b_s .

Transparency effect. We first consider that the low-excitation condition of atoms is satisfied, i.e., $|b_s|^2 \ll 1$. Moreover, the degenerate OPA and the optomechanical coupling between the cavity field and the movable mirror are also removed or turned off, i.e., $G_A = 0$ and $G_0 = 0$. In this case, the expression of a_+ in Eq. (6) can be simplified as

$$a_+ = \frac{\varepsilon_p}{\kappa + i(\Delta_a - \delta) + \frac{G^2}{\gamma_b + i(\Delta_b - \delta)}}. \tag{9}$$

It is found that the denominator of the response function is quadratic in δ so that the coherent coupling between the atomic collective mode and the optical mode can lead to a dip in absorption obtained as $\delta \simeq 0.13\omega_m$ and therefore the generation of the transparency behavior which is shown in Fig. 2(a), where we show the absorption $\text{Re}(E_{out})$ of the output field as a function of δ/ω_m with $b_s = -0.01 + 0.04i$; the corresponding driving strength $\varepsilon_c = 35.12\kappa$, atom-field detuning $\Delta_b = 0.43\kappa$ and excitation number of atoms $|b_s|^2 \approx 0.002$. We can see from Fig. 2(a) that the positions of two peaks in absorption appear at $\delta \simeq 0$ and $\delta \simeq \omega_m$, where the left peak in absorption results from the atom-field coupling⁴².

In order to compare with the above situation, we now explore the effect of the degenerate OPA on the atom-field coupled system. Correspondingly, the expression of the response of a_+ in Eq. (6) is simplified as




Figure 3. The absorption $\text{Re}(E_{\text{out}})$ and dispersion $\text{Im}(E_{\text{out}})$ are plotted as a function of δ/ω_m with $b_s = -0.25 + 0.40i$. The frequency of the moving mirror $\omega_m = 2\pi \times 947 \times 10^3$ Hz, the mechanical factor $Q = \omega_m/\gamma_m = 6700$ and the mass of the oscillating mirror $m = 25$ ng. Other parameter values are the same as in Fig. 2. (a) In the absence of the degenerate OPA and the optomechanical coupling. (b) In the presence of the degenerate OPA and the optomechanical coupling.

$$a_+ = \frac{\varepsilon_p}{\kappa + i(\Delta_a - \delta) + \frac{G^2}{\gamma_b + i(\Delta_b - \delta)} - \frac{G_4^2}{\kappa - i(\Delta_a + \delta) + \frac{G^2}{\gamma_b - i(\Delta_b + \delta)}}} \quad (10)$$

The denominator of the response function is quadruplicate in δ . Then, under the condition of certain parameters, we can obtain more dips in absorption. Indeed, when the degenerate OPA is included in the system, i.e., $G_A = 0.7\kappa$ and $\theta = 3\pi/2$ in Fig. 2(a), an additional dip in absorption appears near the left peak in absorption, i.e., $\delta \simeq -0.04\omega_m$, which results from the coupling between the degenerate OPA and the cavity field. Thus, we can obtain two transparent windows. Figure 2(b) depicts the dispersion shapes of the output field in the absence and presence of the degenerate OPA.

The additional dip in absorption that is similar to the one induced by the degenerate OPA can be generated by the higher order excitation of the atomic ensemble. In this case of presence of the higher order excited atomic ensemble and absence of the degenerate OPA and the optomechanical coupling, the amplitude of the output field [Eq. (6)] can be simplified as

$$a_+ = \frac{\varepsilon_p}{\kappa + i(\Delta_a - \delta) + \frac{G_1^2}{\beta_3} + \alpha_1 - \frac{\alpha_3 \alpha_4}{\kappa - i(\Delta_a + \delta) + \alpha_2 + \frac{G_1^2}{\beta_4 - G_1^2/\beta_3}}}, \quad (11)$$

where $\alpha_1 = iG_3A_4 + \frac{G_1G - 2(G_1G_2^*/\beta_3 + iG_3^*)}{\beta_3\beta_4 - |G_2|^2}$, $\alpha_2 = iG_3^*A_1 - \frac{iG_1G_2^*G_3}{\beta_4 - |G_2|^2/\beta_3}$, $\alpha_3 = G_1A_1 - G_3A_3$ and $\alpha_4 = -G_3^*A_2 - G_1A_4$ are the terms related to the higher order of the Holstein-Primakoff transformation. One can clearly observe from Eq. (10) and Eq. (11) that the higher order coupling terms of the atoms play a role of nonlinear medium similar to the degenerate OPA for the generation of the transparency behavior.

In Fig. 3(a), we show the absorption $\text{Re}(E_{\text{out}})$ and dispersion $\text{Im}(E_{\text{out}})$ of the output field in the absence of the degenerate OPA as a function of δ/ω_m , where we select $b_s = -0.25 + 0.40i$ so that the excitation number of the atomic ensemble $|b_s|^2 = 0.22$, the driving strength of the cavity $\varepsilon_c = 267.5\kappa$ and the atom-field detuning $\Delta_b = 0.20\kappa$. We see from Fig. 3(a) that for the selected system parameters there exists indeed an additional dip in absorption at $\delta \simeq -0.04\omega_m$ which results from the higher order excitation of the atomic ensemble, corresponding to the terms of $\alpha_{1,2,3,4}$. Further, the higher order terms related to the $|b_s|^2$ influence slightly the position of the dip in absorption at $\delta \simeq 0.13\omega_m$ as well as the peaks in absorption; such as, the dip in absorption is attained at $\delta \simeq 0.10\omega_m$. In particular, when the low-excitation condition of atoms breaks slightly, a large driving strength i.e., $\varepsilon_c = 267.5\kappa$ but a relatively small atom-field detuning i.e., $\Delta_b = 0.20\kappa$ can be applied to help observe the transparency behavior in the atom-field coupled system. In contrast, the low-excitation limit of atoms requires a small driving of the cavity but a relatively large atom-field detuning. For example, in the low-excitation limit, i.e., $|b_s|^2 = 0.002$, the driving strength $\varepsilon_c = 35.12\kappa$ and the atom-field detuning $\Delta_b = 0.43\kappa$ could be used [see Fig. 2].

Figure 3(b) demonstrates the transparency behavior in absorption of output probe field in the presence of the optomechanical coupling between the cavity field and the movable mirror when the degenerate OPA and the higher order excitation of the atomic ensemble are both included, i.e., $G_A = 0.7\kappa$ Hz and $b_s = -0.25 + 0.40i$. In this




Figure 4. The absorption $\text{Re}(E_{\text{out}})$ is plotted as a function of δ/ω_m with (a) different G_A 's and (b) different b_s 's. Other parameter values are the same as in Figs 2 and 3.

case, $|b_s|^2 = 0.22$, $\varepsilon_c = 140.78\kappa$ and $\Delta_b = 0.21\kappa$. We can easily find from Fig. 3(b) that there exist three dips in absorption, where the peak in absorption at $\delta \simeq \omega_m$ splits and therefore the optomechanically induced transparency (OMIT) appears due to the coupling between the cavity field and the movable mirror. This is because in general the strongest optomechanical coupling obtains at $\delta \simeq \omega_m$ so that the probe beam interferes destructively with the anti-Stokes field generated by the mechanical oscillator and therefore a transparency window appears in the probe transmission spectrum²⁴. Moreover, the positions of the left side of the two dips in absorption are $\delta \simeq -0.04\omega_m$ and $\delta \simeq 0.11\omega_m$, respectively, which depends on the atom-field coupling and the degenerate OPA.

We can study the effect of the degenerate OPA on the transparency behaviors in the nonlinear optomechanical system with a higher order excited atomic medium. In Fig. 4(a), we show the absorption $\text{Re}(E_{\text{out}})$ in the presence of the OPA as a function of δ/ω_m with different G_A 's. Other parameter values are the same as in Figs 2 and 3. From Fig. 4(a), we see clearly that the larger the nonlinear gain of the degenerate OPA, the wider the OMIT windows. This is because the photon number in the cavity increases with increasing G_A [see Eq. (4)]. Therefore, the effective optomechanical coupling $G_0 a_c$ can be increased by adding a degenerate OPA inside the optical cavity, which leads to widen the OMIT window. This suggests that the nonlinear gain of the degenerate OPA can be used to control the optical-response properties in the nonlinear optomechanical system and the width of OMIT window. Further, in the presence of both the higher order excitation of the atoms and the degenerate OPA, the left dip in absorption shallows, i.e., $G_A = 0.5\kappa$ corresponds to the blue dashed line in Fig. 4(a). This means that the roles of the higher order nonlinearity of atoms and the degenerate OPA cancel each other for generating the dip in absorption in the output field. When the nonlinear gain of the OPA becomes larger, i.e., $G_A = 1.0\kappa$ in Fig. 4(a), the degenerate OPA plays a determined role for the transparency of the probe field and therefore the left transparency window reappears.

We now discuss the important role of excitation of the atomic ensemble in the properties of the output field and OMIT. In Fig. 4(b), we show the absorption $\text{Re}(E_{\text{out}})$ as a function of δ/ω_m with different b_s 's in the absence of the degenerate OPA. We consider the three cases with $b_s = -0.25 + 0.40i$, $b_s = -0.25 + 0.30i$ and $b_s = -0.25 + 0.20i$. The corresponding excitation numbers are calculated as $|b_s|^2 = 0.22$, $|b_s|^2 = 0.15$ and $|b_s|^2 = 0.10$, respectively. In addition, the corresponding atom-field detunings are $\Delta_b = 0.20\kappa$, $\Delta_b = 0.22\kappa$ and $\Delta_b = 0.23\kappa$, respectively, and driving strengths of the optical cavity are $\varepsilon_c = 267.50\kappa$, $\varepsilon_c = 191.99\kappa$ and $\varepsilon_c = 137.23\kappa$, respectively. From Fig. 4(b), we see that the increase of the excitation number of the atoms widens the OMIT window. This is because when the low-excitation condition of atoms is broken slightly, the effective optomechanical coupling strength is enhanced by the increase of the photon number, which depends strongly on the higher order excitation number of atoms. In addition, the additional dip in absorption shallows so that the left transparency behavior disappears when the excitation number decreases. These results can be applied to determinate the excitation number of atomic ensemble and its important role in the properties of the output field.

Tunable slow and fast light. As mentioned in previous section, the optical response of the system to the weak probe field can be described by the group delay. In Fig. 5, the group delay of the output field at the frequency of the probe field is plotted as a function of δ/ω_m with different G_A 's and θ 's (a) and different b_s 's (b). Here we consider the driving of the cavity field is so large that the low-excitation condition of atoms is broken slightly. For example, in Fig. 5(a) we still select $b_s = -0.25 + 0.4i$ and therefore $|b_s|^2 = 0.22$. This leads to generate an additional dip in absorption at $\delta \approx 0$ even in the absence of the degenerate OPA [see the red line in Fig. 4(b)]. In Fig. 5(b), we always remove the degenerate OPA, i.e., $G_A = 0$ and focus justly on the effect of the higher-order excitation of atoms on the optical properties of the system. Other parameter values are the same as in Fig. 4.

We see clearly from Fig. 5(a) that in the absence of the degenerate OPA, a positive group delay of the output field is obtained at $\delta \approx 0$ or $\delta \approx \omega_m$, which corresponds to the slow light effect of the output probe field and results




Figure 5. Group delay τ_g is plotted as a function of δ/ω_m with different G_A 's and θ 's ($b_s = -0.25 + 0.4i$) (a) and different b_s 's ($G_A = 1.5\kappa$) (b). Other parameter values are the same as in Fig. 4.




Figure 6. Group delays τ_g at (a) $\delta \approx 0$ and (b) $\delta \approx \omega_m$ are plotted as a function of the phase θ/π with different G_A 's. Other parameter values are the same as in Fig. 4.

from the enhancement of the optical transparency induced by the higher order excitation of the atoms and the optomechanical coupling, respectively [see the red line in Fig. 5(a)]. When the degenerate OPA is included in the system, i.e., $G_A = 1.5\kappa$ and $\theta = \pi/2$, the group delay at $\delta \approx 0$ can be turned to negative value and therefore the output probe field contains fast light. In contrast, the positive group delay at $\delta \approx \omega_m$ is decreased due to the effect of the degenerate OPA. Moreover, when the degenerate OPA with the phase $\theta = 3\pi/2$ is included, the group delays at $\delta \approx 0$ and $\delta \approx \omega_m$ can be decreased due to the influence the degenerate OPA. In Fig. 5(b), we see that in the absence of the degenerate OPA, the group delay of the output field is always positive with different steady-state excitation number. Thus, only the phenomenon of slow light can be appeared.

In Fig. 6, we show the group delays τ_g at $\delta \approx 0$ and $\delta \approx \omega_m$, respectively, as a function of the phase θ/π with different G_A 's. In Fig. 6(a), we can see that in the absence of the degenerate OPA, i.e., $G_A = 0$ Hz, the group delay with $\delta \approx 0$ induced by the higher order excitation of the atoms is positive with $\tau_g = 2.2\mu\text{s}$ and the transmitted probe field contains slow light. In the presence of the degenerate OPA, i.e., $G_A = 0.3\kappa$, the positive group delay is not a monotonous function of the phase and a peak appears in the intermediate value of the phase. When the nonlinear gain of the OPA becomes larger, i.e., $G_A = 0.6\kappa$, the group delay can be negative corresponding to the fast light within a finite interval of the phase around $\theta = 0.88\pi$, whose position moves in the positive direction

with increasing G_A for the selected range of the parameters [see the dash-dot and dot lines in Fig. 6(a)]. Physically, when the degenerate OPA is added inside the atom-field coupled system, the quantum interference effect between the probe field and the anti-Stokes field generated by the atoms is related directly to the phase of the degenerate OPA by the coupling coefficient G_A [see Eq. (6) or Eq. (10)]. Therefore, the optical-response properties for the probe field become phase-sensitive so that a tunable switch from slow to fast light can be realized by adjusting the phase of the degenerate OPA.

In Fig. 6(b), we can see that in the absence of the degenerate OPA, i.e., $G_A = 0$, the group delay with $\delta \approx \omega_m$ induced by the OMIT is $\tau_g = 54.5 \mu\text{s}$, which is much greater than that induced by the higher order atomic excitation. When the degenerate OPA is included, i.e., $G_A = 0.5\kappa$, the effect of the degenerate OPA on the optomechanical system leads to the increase in the group delay for the intermediate value of the phase. In particular, when the nonlinear gain of the OPA becomes larger, i.e., $G_A = 2.5\kappa$, a negative group delay and therefore the fast light appears in a small interval of the phase around $\theta = 0.36\pi$. Similarly, the tunability of the group delay induced by the OMIT results from the quantum interference effect between the probe field and the anti-Stokes field generated by the mechanical oscillator, which depends strongly on the phase of the degenerate OPA. These results suggest that the phase-sensitive interference effect can be used to control the light propagation from slow to fast light of the transmitted probe field.

Discussion

In conclusion, we have studied a nonlinear optomechanical cavity with a degenerate OPA and a higher order excited atomic ensemble, which is driven by pump and probe laser fields. We derive the expression of the response of probe field in the system and demonstrate numerically the optical properties of the output field with experimentally accessible parameters. It is shown that there exist three dips in absorption in the output field in the presence of a higher order excited atomic medium, which induced by the optomechanical and atom-field couplings as well as the higher order excited terms related to the atom-field coupling, respectively. Further, the OMIT behavior in absorption of probe field can be generated through the optomechanical coupling between the cavity field and the movable mirror. We show that the higher order excitation of the atoms and the degenerate OPA can significantly affect the width of these transparency windows, which can be applied to determinate the excitation number of atoms and the important roles of nonlinear media in the optical properties of the output field. In particular, we find that the higher order excitation of the atoms and the degenerate OPA play a similar role of nonlinear medium in the generation of the additional transparency behavior in the probe field spectrum. We also discuss in detail the change in the magnitude of the group delay as well as a tunable switch from slow to fast light of the output probe field, where the group delay tunability is mainly due to the phase of the degenerate OPA. The phase-sensitive interference effects in the optical-response properties have potential applications to provide new tools for controlling and engineering light propagation.

References

- Harris, S. E., Field, J. E. & Imamoglu, A. Nonlinear optical processes using electromagnetically induced transparency. *Phys. Rev. Lett.* **64**, 1107 (1990).
- Boller, K. J., Imamoglu, A. & Harris, S. E. Observation of electromagnetically induced transparency. *Phys. Rev. Lett.* **66**, 2593 (1991).
- Fleischhauer, M., Imamoglu, A. & Marangos, J. P. Electromagnetically induced transparency: Optics in coherent media. *Rev. Mod. Phys.* **77**, 633 (2005).
- Zhu, S. Y. & Scully, M. O. Spectral Line Elimination and Spontaneous Emission Cancellation via Quantum Interference. *Phys. Rev. Lett.* **76**, 388 (1996).
- Wu, Y. & Yang, X. Electromagnetically induced transparency in \vee -, Λ -, and cascade-type schemes beyond steady-state analysis. *Phys. Rev. A* **71**, 053806 (2005).
- Roghani, M., Breuer, H. P. & Helm, H. Dissipative light scattering by a trapped atom approaching electromagnetically-induced-transparency conditions. *Phys. Rev. A* **81**, 033418 (2010).
- Abdumalikov, A. A. *et al.* Electromagnetically Induced Transparency on a Single Artificial Atom. *Phys. Rev. Lett.* **104**, 193601 (2010).
- Lukin, M. D. Colloquium: Trapping and manipulating photon states in atomic ensembles. *Rev. Mod. Phys.* **75**, 457 (2003).
- Kash, M. M. *et al.* Ultraslow Group Velocity and Enhanced Nonlinear Optical Effects in a Coherently Driven Hot Atomic Gas. *Phys. Rev. Lett.* **82**, 5229 (1999).
- Phillips, M. C. *et al.* Electromagnetically Induced Transparency in Semiconductors via Biexciton Coherence. *Phys. Rev. Lett.* **91**, 183602 (2003).
- Liu, N. *et al.* Plasmonic analogue of electromagnetically induced transparency at the Drude damping limit. *Nature materials* **8**, 758 (2009).
- Santori, C. *et al.* Coherent Population Trapping of Single Spins in Diamond under Optical Excitation. *Phys. Rev. Lett.* **97**, 247401 (2006).
- Paspalakis, E. & Knight, P. L. Transparency and parametric generation in a four-level system †Reviewing of this paper was handled by a member of the Editorial Board. *J. Mod. Opt.* **49**, 87 (2002).
- Aspelmeyer, M., Kippenberg, T. J. & Marquardt, F. Cavity optomechanics. *Rev. Mod. Phys.* **86**, 1391 (2014).
- Vitali, D. *et al.* Optomechanical Entanglement between a Movable Mirror and a Cavity Field. *Phys. Rev. Lett.* **98**, 030405 (2007).
- Nie, W. J. *et al.* Enhancing steady-state entanglement via vacuum-induced emitter-mirror coupling in a hybrid optomechanical system. *J. Phys. B: At. Mol. Opt. Phys.* **49**, 025501 (2016).
- Liu, Y.-C. *et al.* Dynamic Dissipative Cooling of a Mechanical Resonator in Strong Coupling Optomechanics. *Phys. Rev. Lett.* **110**, 153606 (2013).
- Gigan, S. *et al.* Self-cooling of a micromirror by radiation pressure. *Nature* **444**, 67 (2006).
- Nie, W. J. *et al.* Dynamics of a levitated nanosphere by optomechanical coupling and Casimir interaction. *Phys. Rev. A* **88**, 063849 (2013).
- Nie, W. J., Chen, A. X. & Lan, Y. H. Cooling mechanical motion via vacuum effect of an ensemble of quantum emitters. *Opt. Exp.* **23**, 30970 (2015).
- Schwab, K. C. & Roukes, M. L. Putting mechanics into quantum mechanics. *Phys. Today* **58**, 36 (2005).
- Ma, J. *et al.* Formation and manipulation of optomechanical chaos via a bichromatic driving. *Phys. Rev. A* **90**, 043839 (2014).
- Jing, H. *et al.* Optomechanically-induced transparency in parity-time-symmetric microresonators. *Scientific Reports* **5**, 9663 (2015).
- Agarwal, G. S. & Huang, S. Electromagnetically induced transparency in mechanical effects of light. *Phys. Rev. A* **81**, 041803 (2010).

25. Huang, S. & Agarwal, G. S. Electromagnetically induced transparency from two-phonon processes in quadratically coupled membranes. *Phys. Rev. A* **83**, 023823 (2011).
26. Huang, S. & Agarwal, G. S. Electromagnetically induced transparency with quantized fields in optocavity mechanics. *Phys. Rev. A* **83**, 043826 (2011).
27. Qu, K. & Agarwal, G. S. Fano resonances and their control in optomechanics. *Phys. Rev. A* **87**, 063813 (2013).
28. Shahidani, S., Naderi, M. H. & Soltanolkotabi, M. Control and manipulation of electromagnetically induced transparency in a nonlinear optomechanical system with two movable mirrors. *Phys. Rev. A* **88**, 053813 (2013).
29. Davuluri, S. & Zhu, S. Y. Controlling optomechanically induced transparency through rotation. *Europhysics Lett.* **112**, 64002 (2015).
30. Weis, S. *et al.* Optomechanically Induced Transparency. *Science* **330**, 1520 (2010).
31. Lin, Q. *et al.* Coherent mixing of mechanical excitations in nano-optomechanical structures. *Nat. Photon.* **4**, 236 (2010).
32. Safavi-Naeini, A. H. *et al.* Electromagnetically induced transparency and slow light with optomechanics. *Nature* **472**, 69 (2011).
33. Karuza, M. *et al.* Optomechanically induced transparency in a membrane-in-the-middle setup at room temperature. *Phys. Rev. A* **88**, 013804 (2013).
34. Massel, F. *et al.* Multimode circuit optomechanics near the quantum limit. *Nat. Commun.* **3**, 987 (2012).
35. Zhou, X. *et al.* Slowing, advancing and switching of microwave signals using circuit nanoelectromechanics. *Nat. Phys.* **9**, 179 (2013).
36. Chang, D. E. *et al.* Slowing and stopping light using an optomechanical crystal array. *New J. Phys.* **13**, 023003 (2011).
37. Guo, Y. J. *et al.* Electromagnetically-induced-transparency-like ground-state cooling in a double-cavity optomechanical system. *Phys. Rev. A* **90**, 053841 (2014).
38. Genes, C. *et al.* Atom-membrane cooling and entanglement using cavity electromagnetically induced transparency. *Phys. Rev. A* **84**, 051801(R) (2011).
39. He, Y. Optomechanically induced transparency associated with steady-state entanglement. *Phys. Rev. A* **91**, 013827 (2015).
40. Haroche, S. & Raimond, J. M. *Exploring the Quantum: Atoms, Cavities and Photons* (Oxford University Press, New York, 2006).
41. Wang, H. *et al.* Optomechanical analog of two-color electromagnetically induced transparency: Photon transmission through an optomechanical device with a two-level system. *Phys. Rev. A* **90**, 023817 (2014).
42. Akram, M. J., Ghafoor, F. & Saif, F. Electromagnetically induced transparency and tunable fano resonances in hybrid optomechanics. *J. Phys. B: At. Mol. Opt. Phys.* **48**, 065502 (2015).
43. Genes, C., Vitali, D. & Tombesi, P. Emergence of atom-light-mirror entanglement inside an optical cavity. *Phys. Rev. A* **77**, 050307 (2008).
44. Genes, C., Ritsch, H. & Vitali, D. Micromechanical oscillator ground-state cooling via resonant intracavity optical gain or absorption. *Phys. Rev. A* **80**, 061803 (2009).
45. Nie, W. J., Chen, A. X. & Lan, Y. H. Optical-response properties in levitated optomechanical systems beyond the low-excitation limit. *Phys. Rev. A* **93**, 023841 (2016).
46. Huang, S. & Agarwal, G. S. Enhancement of cavity cooling of a micromechanical mirror using parametric interactions. *Phys. Rev. A* **79**, 013821 (2009).
47. Farman, F. & Bahrapour, A. R. Effects of optical parametric amplifier pump phase noise on the cooling of optomechanical resonators. *J. Opt. Soc. Am. B* **30**, 1898 (2013).
48. Wilson, G. A. *et al.* Inversionless gain in driven three-level ssV-type atoms: A comparison of broadband and monochromatic excitation. *Phys. Rev. A* **50**, 3394 (1994).
49. Yi, Z., Gu, W.-j. & Li, G.-x. Sideband cooling of atoms with the help of an auxiliary transition. *Phys. Rev. A* **86**, 055401 (2012).
50. Karuza, M. *et al.* Optomechanical sideband cooling of a thin membrane within a cavity. *New J. Phys.* **14**, 095015 (2012).
51. Biancofiore, C. *et al.* Quantum dynamics of an optical cavity coupled to a thin semitransparent membrane: Effect of membrane absorption. *Phys. Rev. A* **84**, 033814 (2011).
52. Chen, X. *et al.* Cooling of macroscopic mechanical resonators in hybrid atom-optomechanical systems. *Phys. Rev. A* **92**, 033841 (2015).
53. Law, C. K. Interaction between a moving mirror and radiation pressure: A Hamiltonian formulation. *Phys. Rev. A* **51**, 2537 (1995).
54. Holstein, T. & Primakoff, H. Field Dependence of the Intrinsic Domain Magnetization of a Ferromagnet. *Phys. Rev.* **58**, 1098 (1940).
55. Turek, Y. *et al.* Indirect driving of a cavity-QED system and its induced nonlinearity. *Phys. Rev. A* **90**, 013836 (2014).
56. Gardiner, C. & Zoller, P. *Quantum noise: a handbook of Markovian and non-Markovian quantum stochastic methods with applications to quantum optics* (Springer Science Business Media, 2004).
57. Giovannetti, V. & Vitali, D. Phase-noise measurement in a cavity with a movable mirror undergoing quantum Brownian motion. *Phys. Rev. A* **63**, 023812 (2001).
58. Walls, D. F. & Milburn, G. J. *Quantum Optics* (Springer-Verlag, Berlin, 1994).
59. Milonni, P. W. *Fast light, slow light and lefthanded light* (Institute of Physics Publishing, Bristol, 2005).
60. Gröblacher, S. *et al.* Observation of strong coupling between a micromechanical resonator and an optical cavity field. *Nature* **460**, 724 (2009).

Acknowledgements

W. N. is supported by the National Natural Science Foundation of China (NSFC) under Grants No. 11304010 and No. 11565014 and the Natural Science Foundation of Jiangxi Province (Grant No. 20161BAB211023). A. C. is supported by the NSFC under Grant No. 11365009.

Author Contributions

L.L. and W.N. wrote the main manuscript text and prepared Figures 1–6. All the authors reviewed the manuscript and discussed the results and edited the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Li, L. *et al.* Transparency and tunable slow and fast light in a nonlinear optomechanical cavity. *Sci. Rep.* **6**, 35090; doi: 10.1038/srep35090 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016