

SCIENTIFIC REPORTS



OPEN

Functional characterization of a *csoR-cueA* divergon in *Bradyrhizobium liaoningense* CCNWSX0360, involved in copper, zinc and cadmium cotolerance

Jianqiang Liang, Mingzhe Zhang, Mingmei Lu, Zhefei Li, Xihui Shen, Minxia Chou[†] & Gehong Wei[†]

Received: 20 June 2016
Accepted: 26 September 2016
Published: 11 October 2016

Random mutagenesis in a symbiotic nitrogen-fixing *Bradyrhizobium liaoningense* CCNWSX0360 (Bln0360) using Tn5 identified five copper (Cu) resistance-related genes. They were functionally sorted into three groups: transmembrane transport (*cueA* and *tolC*); oxidation (*copA*); and protection of the membrane barrier (*lptE* and *ctpA*). The gene *cueA*, together with the upstream *csoR* (Cu-sensitive operon repressor), constituted a *csoR-cueA* divergon which plays a crucial role in Cu homeostasis. Deletion of *cueA* decreased the Cu tolerance of cells, and complementation of this mutant restored comparable Cu resistance to that of the wild-type. Transcriptional and fusion expression analysis demonstrated that *csoR-cueA* divergon was up-regulated by both the monovalent Cu⁺ and divalent Zn²⁺/Cd²⁺, and negatively regulated by transcriptional repressor CsoR, via a bidirectional promoter. Deletion of *csoR* renders the cell hyper-resistant to Cu, Zn and Cd. Although predicted to encode a Cu transporting P-type ATPase (CueA), *cueA* also conferred resistance to zinc and cadmium; two putative N-MBDs (N-terminal metal binding domains) of CueA were required for the Cu/Zn/Cd tolerance. Moreover, *cueA* is needed for nodulation competitiveness of *B. liaoningense* in Cu rich conditions. Together, the results demonstrated a crucial role for the *csoR-cueA* divergon as a component of the multiple-metal resistance machinery in *B. liaoningense*.

Some transition metals, such as copper (Cu) and zinc (Zn), are essential for many cellular processes, however, they are also toxic in excess by generating free radical species or displacing other metals from their native binding sites in metalloenzymes¹. Cadmium (Cd), a ubiquitous metal with unknown biological function, can be extremely toxic, even at low levels. Thus, bacteria have evolved various mechanisms to control intracellular metal ion concentrations, ensuring that they do not reach toxic levels.

Mechanisms of Cu resistance and their regulation have been studied extensively in model organisms *Escherichia coli*, *Enterococcus hirae* and *Mycobacterium tuberculosis*. In *E. coli*, detoxification of intracellular Cu is primarily accomplished by Cu transporting P-type ATPase (CopA), multicopper oxidase (CueO) and resistance-nodulation-cell division (RND)-type Cus system (CusCFBA)². However, in Gram-positive *E. hirae*, the Cus system and CueO are absent, and Cu homeostasis mainly depends on the Cop system comprising a transcriptional repressor (CopY), a Cu chaperone (CopZ), and two Cu transporting P-type ATPases (CopA and CopB)³. In *M. tuberculosis*, CtpV, a Cu transporting P-type ATPase and MctB, a Cu transport outer membrane protein, together with metallothionein MymT, constitute the defense system against excess Cu⁴.

Although there are a variety of Cu resistance mechanisms, active efflux mediated by Cu transporting P_{1B}-type ATPase is the most central⁴. The P_{1B}-type ATPases confer heavy metal resistance through pumping out cytoplasmic metal ions including Ag⁺, Cu⁺, Cu²⁺, Zn²⁺, Cd²⁺, Pb²⁺, Co²⁺, Fe²⁺, and Ni²⁺. They are divided into five

State Key Laboratory of Crop Stress Biology in Arid Areas, College of Life Sciences, Northwest A&F University, Yangling, Shaanxi, China. [†]Present address: College of Life Sciences, Northwest A&F University, 22 Xinong Road, Yangling, Shaanxi 712100 China. Correspondence and requests for materials should be addressed to M.C. (email: minxia95@126.com) or G.W. (email: weigehong@nwsuaf.edu.cn)

Strains/mutants	Gene ^c	Protein/putative function ^{a,c}	Maximum tolerable metal concentrations (MTC; mM) ^b						
			Cu ²⁺	Zn ²⁺	Cd ²⁺	Pb ²⁺	Ni ²⁺	Co ²⁺	Ag ⁺
Bln0360	NA	NA	2.0	3.2	0.15	2.8	0.6	1.4	0.1
Bln-d	<i>cueA</i>	Heavy metal transporting P-type ATPase	0.8	3.2	0.15	2.8	0.6	1.4	0.1
Bln-163	<i>cueA</i>	Heavy metal transporting P-type ATPase	0.8	3.2	0.15	2.8	0.6	1.4	0.1
Bln-32	<i>tolC</i>	Type I secretion outer membrane protein	1.0	1.2	0.1	2.4	0.4	0.6	0.1
Bln-c	<i>copA</i>	Multicopper oxidase	0.6	3.2	0.15	2.8	0.6	1.4	0.1
Bln-29	<i>ctpA</i>	Carboxy-terminal protease, membrane integrity associated	0.6	2.8	0.1	1.6	0.6	1.4	0.1
Bln-54	<i>lptE</i>	Lipopolysaccharide-assembly lipoprotein	0.6	1.2	0.05	1.6	0.4	1.0	0.1

Table 1. The location of transposon insertions in Bln0360 and the levels of metal tolerance in insertion mutants. ^aInterrupted gene encoded proteins/functions are based on the annotated genome sequence of *B. diazoefficiens* USDA 110¹⁷. ^bThe MTCs were determined on TY plate containing elevated concentrations of metal ions (Cu²⁺/Co²⁺, 0–2.4 mM at 0.2 mM intervals; Zn²⁺, 0–4.0 mM at 0.4 mM intervals; Ag⁺, 0.025–0.05, 0.075, 0.1 and 0.125 mM; Ni²⁺, 0.2, 0.4, 0.6, and 0.8 mM). ^cNot available.

groups according to their substrate specificity (P_{1B1}–P_{1B5}), among which P_{1B1}- and P_{1B2}-type ATPases are responsible for Cu⁺/Ag⁺ and Zn²⁺/Cd²⁺/Pb²⁺ translocation, respectively⁵. It is generally accepted that P_{1B}-type ATPases have high specificity for the heavy-metal ions they transport. The substrate specificity is presumably relies on the conserved residues in transmembrane segments H6, H7, and H8, but remains to be established⁶. Another important structure of P_{1B}-ATPases is the presence of cytoplasmic N-MBDs (N-terminal metal binding domains). Current studies mainly focus on the N-MBDs featuring prototypical GXXCXXC motif(s). The N-MBDs appear to be responsible for sensing cellular metal ions and regulating the ATPase enzyme activity as well as conferring ionic specificity^{7,8}. In contrast, much less is known about the role of N-MBDs containing His-rich motif which is mainly present in the P_{1B2}- and P_{1B3}-type ATPase⁶.

In most Gram-negative bacteria, the expression of Cu ATPase is mainly regulated by transcriptional activator CueR; whereas Gram-positive bacterial Cu ATPase is repressed by CsoR or CopY⁶. CsoR was initially identified in *M. tuberculosis*, and subsequently in other Gram-positive bacteria such as *Bacillus subtilis*, *Corynebacterium glutamicum*, *Listeria monocytogenes*, and *Streptomyces lividans*⁶. Experimental evidence for Cu homeostasis gene regulation by CsoR in Gram-negative bacteria is still lacking to date. *Thermus thermophilus* CsoR is the only instance identified in Gram-negative bacteria, however, metal-binding motif of the CsoR (H-C-H-H) is the same as that of *E. coli* RcnR, which is distinctly different from that of CsoRs (C-H-C) from Gram-positive bacteria above⁹.

Rhizobia are Gram-negative soil-dwelling bacteria that form a symbiosis with legumes to fix nitrogen from the atmosphere¹⁰. Recently, the nitrogen fixer has attracted great attention for their role in aiding phytoremediation of metal contaminated soils^{11,12}. Cu, a ubiquitous transition metal, enters soils via agricultural and industrial activities, and exposure at high levels have presented serious threats to the environment and human health¹³. Some rhizobia can tolerate high concentrations of Cu and display the potential phytoremediation by their host plants in Cu contaminated soil^{11,12,14,15}. However, Cu resistance determinants of rhizobia are poorly characterized.

In the present study, the mechanisms of Cu resistance in *Bradyrhizobium liaoningense* CCNWSX0360 were investigated through random transposon mutagenesis. A *csoR-cueA* divergon encoding a CsoR-like repressor and a heavy metal transporting P-type ATPase (CueA) was functionally characterized; *csoR-cueA* divergon plays a crucial role in Cu homeostasis, and also involves in Zn/Cd resistance suggesting a versatile metal resistance component. Furthermore, the role of *cueA* in symbiotic nodulation under Cu stress was investigated, which will contribute to improving the metal bioremediation potential of legume-rhizobium symbiosis.

Results

Isolation and phylogenetic identification of the Cu resistant isolate Bln0360. A total of 108 rhizobia were isolated from the nodules of 13 leguminous plant species in the study. Among them, strain CCNWSX0360 from *Vigna unguiculata* showed the highest resistance to Cu (2.0 mM) and was selected for the study. The 16S rRNA gene sequence of strain CCNWSX0360 (KU507314) showed 100% similarity to *B. liaoningense* SEMIA 5022 (FJ390920) and 99.9% similarity to strain 2281^T (AJ250813). Phylogenetic analysis revealed that strain CCNWSX0360 belonged to *B. liaoningense* (Fig. S1) and it was named *B. liaoningense* CCNWSX0360 (Bln0360). The maximum tolerable metal concentrations (MTCs) of Bln0360 to the test metals were 2.0, 3.2, 0.15, 2.8, 0.6, 1.4, and 0.1 mM for Cu²⁺, Zn²⁺, Cd²⁺, Pb²⁺, Ni²⁺, Co²⁺, and Ag⁺ (Table 1).

Identification of genes involved in Cu resistance by transposon mutagenesis. To identify genes involved in Cu resistance in Bln0360, a transposon mutant library (17,247 Tn5-insertions) was constructed. Upon screening, six Cu sensitive mutants were obtained (Table 1). To further test the sensitivity and specificity, the MTCs of various metal ions (Cu²⁺, Zn²⁺, Pb²⁺, Cd²⁺, Co²⁺, Ni²⁺, and Ag⁺) for each mutant were determined in TY (Tryptone-Yeast) medium. Of the mutants, three mutants (Bln-c, Bln-54, and Bln-29) exhibited a drastic reduction in Cu tolerance, with MTCs much lower (0.6 mM Cu²⁺) than that of Bln0360 (2.0 mM Cu²⁺). Bln-29, Bln-32 and Bln-54 also showed the varying of decreased tolerance toward other metals ions (Zn²⁺, Pb²⁺, Cd²⁺, Co²⁺, and Ni²⁺), but did not exclusively affect the resistance to Cu²⁺. In contrast, no difference in tolerance to metals other than Cu²⁺ was observed between the mutants (Bln-c, Bln-163, and Bln-d) and the wild-type strain.




Figure 1. Organization of the *csoR-cueA* divergon and their promoter regions from Bln0360. (A) Schematic representation of the arrangement of the *csoR-cueA* divergon; the positions of Tn5 insertions are indicated by inverted black triangles. **(B)** Nucleotide sequences of the bidirectional *csoR-cueA* promoter region. Partial coding sequences for *csoR* and *cueA* are indicated in bold. The transcriptional start sites (+1) of *csoR* and *cueA* are indicated by large bold letter, and the ribosome binding site (RBS) of *cueA* is underlined. The predicted -35 and -10 elements of the *csoR* and *cueA* promoter are boxed. The 19-bp putative CsoR-box with a 7-bp inverted repeat is indicated with a grey background. **(C)** RACE experiments were performed using RNA isolated from uninduced (lanes 1 and 3) or 0.8 mM CuSO₄-induced (lanes 2 and 4) cultures of Bln0360.

Among these mutants, Tn5 was inserted into the same gene encoding a putative heavy metal-transporting P-type ATPase, named *cueA* (KU665989), in both strains Bln-d and Bln-163, which was consistent with their identical tolerance to the tested metals in this study (Fig. 1A and Table 1). In strain Bln-32, the interrupted gene *tolC* (KU665993) encoded an outer membrane protein showing 21% identity to *E. coli* TolC, which is an outer membrane component of a multidrug efflux system, AcrAB-TolC¹⁶. In Bln-c, the interrupted gene *copA* (KU665990) encoded a multicopper oxidase showing 98% identity to CopA from *Bradyrhizobium diazoefficiens* USDA 110¹⁷. Moreover, individual genes *lptE* (KU665991) and *ctpA* (KU665992) encoding putative membrane formation associated proteins were respectively interrupted in Bln-29 and Bln-54^{18,19}.

Further analysis of the sequences derived from Bln-163 and Bln-d identified a small open reading frame, named *csoR* (KU665989), which was inversely oriented and located immediately upstream of *cueA* (Fig. 1A). The ORF encodes a putative CsoR-like regulator which has mainly been reported in Gram-positive bacteria⁶. The remainder of the experiment focuses on the genetic arrangement and function of the *cueA-csoR* divergon from the Gram-negative bacteria.

***cueA* is critical for Cu resistance of Bln0360.** *In silico* analysis showed that *cueA* encodes a putative protein of 815 amino acid residues with a theoretical molecular mass of 85.5 kDa. The deduced CueA amino acid sequence showed high identity with several previously characterized Cu transporting P-type ATPases: CopA of *M. amorphae* (EHH02252, 58.5%)¹¹, CopA of *A. tumefaciens* (Atu0937, 42.3%)²⁰ and CopA of *E. coli* (BAE76263, 38.2%)²¹. Alignment of sequences revealed three conserved domains of the P-type ATPase family, including a SGES phosphatase domain (A-domain), a DKTGT aspartyl kinase domain (P-domain), and a GXGXND ATP-binding domain (N-domain), which were also present in CueA (Fig. S2). CueA contained eight predicted transmembrane segments (TMS), with the CPX motif (CPC) located in TMS6 and the signature sequences NY in TMS7 and MXXSS in TMS8. These *in silico* data allowed assignment of CueA to subclass type P_{1B1} according to Palmgren's classification⁵.

qRT-RPCR analysis of the Cu²⁺ responsiveness of *cueA* showed that there was a gradual response with elevated concentration of added CuSO₄ (Fig. 2A), suggesting a significant dose-dependent effect. A notable induction was observed when 0.005 mM CuSO₄ was added to cultures ($P < 0.01$). Induction of *cueA* reached up to ~350-fold relative to untreated cells, at 2.5 mM (Fig. 2A), though this was beyond the maximum Cu²⁺ concentration the bacteria could tolerate in TY plate. Dose-dependent induction of *csoR* by CuSO₄ also began at 0.005 mM, but reached a maximum at 1.25 mM (Fig. 2A). Addition of the Cu⁺-specific chelating agent bathocuproine disulfonate (BCS, 1.0 mM) to the cultures completely eliminated the induction caused by 0.1 mM Cu²⁺, and partially eliminated that caused by 1.0 mM Cu²⁺ (Fig. 2B). These results suggested that expression of *cueA* could be induced by Cu⁺. Although Cu²⁺ was added to the medium, it would be reduced to Cu⁺, intracellularly; therefore, Cu⁺ might be the actual inducer^{21,22}.

To verify the Cu sensitive phenotype of Tn5 insertions Bln-d and Bln-163 was indeed caused by inactivation of *cueA*, an in-frame deletion mutant (Δ *cueA*) of Bln0360 was constructed and subjected to metal tolerance assays. No difference was observed in growth under various metal stresses between the transposon and constructed mutants (data not shown), indicating that no polar mutations were produced by insertion of Tn5.




Figure 2. *cueA* contributes to Cu resistance in Bln0360. **(A)** Transcription of *cueA* and *csoR* in response to Cu²⁺. The cDNA was derived from Bln0360 culture at mid-log phase after induction for 2 h with elevated concentrations of CuSO₄. Normalized expression of *cueA* and *csoR* with respect to 16S rRNA expression is presented as the mean ± SD (standard deviation) of data from three independent samples. Relative mRNA levels were expressed as fold change with respect to the untreated control. ***P* < 0.01. **(B)** β-Galactosidase activity of *P_{csoR}::lacZ* and *P_{cueA}::lacZ* reporters were determined in TY medium at 0.1 mM or 1.0 mM CuSO₄ with and without the addition of 1.0 mM bathocuproine disulfonate (BCS). Values are presented as the mean ± SD of data from three independent experiments. ****P* < 0.001; ***P* < 0.01. **(C)** Tolerance levels of Bln0360, the $\Delta cueA$ mutant and the $\Delta csoR(csoR)$ complemented strain to Cu²⁺. Ten-fold serial dilutions of log phase culture were spotted onto TY agar plates with no addition, 1.0 mM BCS, 0.8 mM CuSO₄ or 0.8 mM CuSO₄ + 1.0 mM BCS. Plates were photographed after 7 d incubation at 28 °C.

Complementation of the $\Delta cueA$ mutant with full-length *cueA* gene restored the Cu resistance to near that of the wild-type (Fig. 2C).

The Cu sensitivity of $\Delta cueA$ on TY agar medium containing 0.8 mM CuSO₄ was completely eliminated by adding 1.0 mM BCS (Fig. 2C). Thus, Cu toxicity to Bln0360 may be dependent on the conversion of Cu²⁺ to Cu⁺, and also implies that CueA plays a role in protection of Bln0360 from Cu⁺, which agreed with the responses of the Cu²⁺-inducible expression of *cueA* to BCS (Fig. 2B). Overall, these data demonstrated that CueA is involved in and plays the major role in intracellular Cu detoxification via Cu⁺ efflux.

Two putative N-terminal MBDs is required for full function of CueA. Unlike other P_{1B1}-type ATPases that possess the typical N-terminal GXXCXXC motif, CueA contains two His-rich stretches (Fig. 3A). The presence of His-rich stretch in the CueA is unusual, in that P_{1B}-type ATPase with His-rich N-MBDs are usually involved in divalent metal ions transportation^{23,24}. To explore the role of the two putative MBDs in the Cu resistance mediated by CueA, three variants CueA- Δ MBD_a (with the first His-rich stretch deleted), CueA- Δ MBD_b (with the second His-rich stretch deleted) and CueA- Δ MBD_{ab} (with both His-rich stretches deleted) were expressed in the Cu sensitive $\Delta cueA$ mutant. Tolerance assay of these transformants indicated that deletion of the MBD_b largely abrogated the Cu resistance mediated by CueA, whereas observable reduction caused by deletion of the MBD_a only observed on 1.2 mM CuSO₄-supplemented plates (Fig. 3B). The result indicated that both the putative His-rich domains were obligatory for the function of CueA and furthermore the second domain showed a dominant role. Note that the $\Delta cueA$ mutant expressing CueA- Δ MBD_{ab} had a distinct growth advantage compared to the mutant with empty plasmid at Cu²⁺ concentrations of 0.8 mM. It indicated that CueA is capable of functioning in a manner independent of the N-terminal MBDs (Fig. 3B). These data support the model in which N-terminal MBDs is responsible for regulation rather than an absolutely essential for the catalytic mechanism of P_{1B}-ATPase²⁵.




Figure 3. The N-terminal His-rich stretches are required for full Cu resistance of CueA. (A) Characteristic metal binding motifs in the N-terminus of CopA from *E. coli* and CueA from Bln0360. His-rich stretch and conserved residues within GXXCXXC motif are highlighted. Superscript numbers represent amino acid positions. (B) Comparative Cu tolerance of $\Delta cueA$ strains expressing CueA and its variants on TY agar plates containing 0.8 or 1.2 mM CuSO₄. Plates were photographed after 7 d incubation at 28 °C. Strain $\Delta cueA$ (vector) was used as the negative control. MBD_a indicates deletion of the first His-rich stretch, MBD_b indicates deletion of the second, and MBD_{ab} indicates deletion of both.

CsoR negatively regulates the *csoR-cueA* divergon. *csoR* encodes a putative protein of 91 amino acid residues showing 44.4% and 27.6% identities with CsoR from *C. glutamicum* (AIK84123) and *M. tuberculosis* (P9WP49), respectively^{26,27}. Alignment of the deduced protein with its orthologs revealed the presence of a conserved C-H-C motif (Cys³³, His⁵⁸ and Cys⁶²), which served as Cu⁺-binding ligands (Fig. S4), furthermore, I-TASSER software predicted three alpha helices similar to the *M. tuberculosis* CsoR, suggesting a similar mechanism to sense and respond to Cu⁺.

Inspection of the intergenic region between *csoR* and *cueA* identified -10 and -35 promoter sequences separately in the upstream region of each gene; the -10 elements partially overlapped (Fig. 1B). The transcription start site (TSS; position +1) of *cueA* was mapped to the G nucleotide located 42 nucleotides upstream of the putative start codon (ATG) by 5' rapid amplification of cDNA ends (RACE) (Fig. 1B,C). Interestingly, the putative TSS of *csoR* was mapped to the first nucleotide (A) of start codon of this gene, suggesting a leaderless *csoR* mRNA, in which the TSS is starts directly with a 5'-terminal AUG²⁸. Further examination of the sequence upstream of the TSS revealed a 7-bp inverted repeat separated by 5 bp (5'-TATACCCCTACCGGGTATA-3'), which highly similar to the recognition motif (CsoR-box) of the CsoR from *C. glutamicum*²⁷. The CsoR-box overlapped with the overlapping -10 element of *cueA* and *csoR*, indicating a bidirectional promoter structure. The conserved sequence motif of CsoR and the location of the CsoR-box suggest that it regulates expression of both transcripts (i.e., *csoR* and *cueA*) simultaneously, in opposite directions.

To elucidate whether or not the *csoR-cueA* divergon is autoregulated by CsoR, the transcription of *csoR* and *cueA* in response to Cu²⁺ were examined by Quantitative real-time PCR (qRT-PCR) in the $\Delta csoR$ mutant and the $\Delta csoR$ (*csoR*) complemented strain (Fig. 4A,B). In the $\Delta csoR$ mutant, a high level of *csoR* and *cueA* mRNA was detected even without the addition of CuSO₄, indicating uncontrolled transcription. Complementation of the mutant with *csoR* gene resulted in high level transcription of *csoR* and restored the Cu²⁺-dependent induction of *cueA*, suggesting that *csoR* acts as a negative regulatory factor.

The β -galactosidase activity of P_{*csoR*}::*lacZ* and P_{*cueA*}::*lacZ* fusions was determined to further confirm the data of the two genes expression. As shown in Fig. 4B,C, expression of P_{*csoR*}::*lacZ* and P_{*cueA*}::*lacZ* was completely




Figure 4. Effects of *csoR* deletion on the expression of the *csoR-cueA* divergon and the tolerance of Bln0360 to Cu exposure. **(A)** The relative mRNAs levels of *cueA* (left) and *csoR* (right) in Bln0360, the Δ*csoR* mutant and the Δ*csoR*(*csoR*) complemented strain exposed to media containing or lacking 0.8 mM CuSO₄ for 2 h. mRNA levels are presented relative to the untreated wild-type strain. **(B)** β-Galactosidase activity of P_{csoR}::lacZ (left) and P_{cueA}::lacZ reporter (right) in wild-type Bln0360, the Δ*csoR* mutant and the Δ*csoR*(*csoR*) complemented strain grown in TY medium with and without the addition of 0.8 mM CuSO₄ for 2 h. **(C)** Expression of the *csoR-cueA* divergon in wild-type Bln0360 and Δ*csoR* mutant in response to elevated concentrations of CuSO₄. The SD from three independent experiments is indicated on each bar. **P* < 0.05, ***P* < 0.01, ****P* < 0.001.

derepressed in the *csoR* mutant, which was consistent with the results of qRT-PCR. Complementation of the mutant restored the repression and Cu²⁺-responsiveness of P_{csoR}::lacZ and P_{cueA}::lacZ (Fig. 4B). The results provided further evidence that CsoR negatively regulates the *csoR-cueA* divergon. It is noteworthy that fusion reporter P_{cueA}::lacZ expressed 1.9–4.8 fold higher β-galactosidase activity than the P_{csoR}::lacZ fusion at each Cu²⁺ concentration in the wild-type strain (Fig. 4C). In contrast, in the CsoR deletion mutant, the β-galactosidase activity of each fusion was indistinguishable, indicating parallel promoter strength. These data showed that CsoR controlled the promoters of both *csoR* and *cueA* simultaneously, but influenced their activity to varying extents. Notably, the expression level of *csoR* decreased when the induction concentration of Cu²⁺ was 2.5 mM as seen from its transcript abundance relative to that at lower Cu²⁺ concentrations (Figs 4C and 2A), which is probably a result of competition between RNA polymerases for the respective promoters. Collectively, these data clearly showed that CsoR acts as a repressor to regulate the *csoR-cueA* divergon expression via a bidirectional promoter.




Figure 5. The *csoR-cueA* divergon is up-regulated by Zn^{2+} and Cd^{2+} . (A) mRNA expression levels of *csoR* and *cueA* genes in response to different metal ions. cDNA was derived from mid-log phased Bln0360 culture after 2 h incubation with 1.0 mM $CuSO_4$, 0.1 mM $CdSO_4$, 0.075 mM $AgNO_3$, 2.0 mM $PbNO_3$, 0.4 mM $NiSO_4$ or 1.0 mM $CoCl_2$ (bacterial growth was markedly inhibited by the above specified concentrations of metal ions) and subjected to qRT-PCR assay. Fold change of gene expressions was estimated mRNA levels compared to the untreated cultures. (B) Expression of the *csoR-cueA* divergon in response to Cu^{2+} , Zn^{2+} , and Cd^{2+} . Strain Bln0360 carrying *P_{csoR}::lacZ* or *P_{cueA}::lacZ* fusions were grown in TY medium to log phase and individually supplemented with 1.0 mM $CuSO_4$, 1.0 mM $ZnSO_4$ or 0.1 mM $CdSO_4$. Cells were collected and β-Galactosidase activities were determined. Error bars indicate SD of triplicates. *** $P < 0.001$; ** $P < 0.01$.

Expression of the *csoR-cueA* divergon could be induced by Zn^{2+} and Cd^{2+} . Since CsoR in some bacteria (e.g., *T. thermophilus* and *C. glutamicum*) can sense various metal ions and derepress transcription^{9,27}, we investigated whether metal ions other than Cu^{2+} could induce expression of the *csoR-cueA* divergon in Bln0360. The result showed that the mRNA expression of *cueA* gene could be markedly induced by Zn^{2+} and Cd^{2+} (~90-fold for 1.0 mM Zn^{2+} and ~135-fold for 0.1 mM Cd^{2+} , respectively) as well as Cu^{2+} . Similar up-regulation of *csoR* was also observed, but with lower fold-changes (Fig. 5A).

To further confirm the expression pattern of the divergon by Zn^{2+} and Cd^{2+} , we determined β-galactosidase activity of *P_{csoR}::lacZ* and *P_{cueA}::lacZ* in response to the two metal ions. As Fig. 5B shows, 1.0 mM Zn^{2+} caused a 3.4-fold ($P < 0.01$) and 0.1 mM Cd^{2+} caused a 3.8-fold increase ($P < 0.01$) in the expression of *cueA* compared with the treatment in the absence of metal ions. Significant increases (1.9-fold for Zn^{2+} and 2.7-fold for Cd^{2+}) were also observed in the expression of *csoR* in response to the same concentration of Zn^{2+} and Cd^{2+} . Additionally, expressions of *csoR* and *cueA* were markedly higher in the treatment of 1.0 mM Cu^{2+} than that in the same concentration of Zn^{2+} (Fig. 5B). Combined with the data of responses to BCS above, these results indicated that *csoR-cueA* divergon can be induced by both monovalent Cu^+ and divalent Zn^{2+}/Cd^{2+} and may be involved in detoxification of multiple heavy metals.

***csoR* mutation increase resistance to Cu, Zn and Cd.** The *csoR*-deficient mutant was selected to test the altered tolerance to Cu, Zn, Cd, Ni and Ag, respectively. Since CueA was evidenced to facilitate Cu tolerance, the strain deficient in its repressor CsoR should be more resistant to Cu as compared to wild-type strain. As expected, $\Delta csoR$ mutant displayed a distinct growth advantage in the media containing high concentration (>1.6 mM) of Cu^{2+} compared to Bln0360, as determined by culture optical density and plate assay (Fig. 6A,D); the MTC of Cu^{2+} for $\Delta csoR$ mutant was raised from original 2.0 to 2.4 mM (Fig. 6D).

Intriguingly, deletion of the *csoR* gene also increased tolerance of strain to Zn and Cd, though mutation of its target gene (*cueA*) had no measurable impact on the tolerance to these two metals. As shown in Fig. 6B, $\Delta csoR$ mutant showed markedly higher OD_{600s} than Bln0360 in $ZnSO_4$ -supplemented TY broth. Similarly, Bln0360 showed a sharp reduction in growth after 0.08 mM $CdSO_4$, approaching zero growth at approximately 0.16 mM (Fig. 6C); in contrast, $\Delta csoR$ mutant remains tolerant until approximately 0.2 mM. The hyper-tolerant phenotype was also observed by plate assay and the MTCs of $ZnSO_4$ and $CdSO_4$ for $\Delta csoR$ mutant were 4.0 mM and 0.25 mM, respectively, which were higher than 3.2 mM and 0.15 mM for wild-type (Fig. 6D and Table 1). We speculated that one possible reason could be due to the depression of general stress genes in the transcript level, but such increased tolerance was not observed when $\Delta csoR$ mutant was tested on both solid and liquid medium supplemented with Ni^{2+} and Ag^+ (data not shown).

***cueA* confers resistance to Zn and Cd.** Since $\Delta csoR$ mutant displayed an increased tolerance to Zn/Cd and *cueA* expression was up-regulated by the two metals, we hypothesized that *cueA* might contribute to Zn and Cd resistance. To validate this hypothesis, construct pBRR5-*cueA* was introduced into the Zn/Cd sensitive *E. coli* GG48, and the growth of resultant strain was monitored on Luria-Bertani¹⁹ medium containing different concentrations of $CdSO_4$ or $ZnSO_4$. Spot assays clearly indicated that *E. coli* GG48 expressing CueA had a distinct growth advantage on both $ZnSO_4$ - and $CdSO_4$ -supplemented agar plates when compared with the *E. coli* harboring the empty plasmid (Fig. 7A). For Cd, the *E. coli* bearing empty plasmid barely grew on the plate containing $CdSO_4$, whereas cells with CueA expression plasmids appeared relatively robust growth (Fig. 7A). Consistent results were obtained when growing in the liquid media (Fig. 7B). Thus, the Zn/Cd sensitive *E. coli* strain appeared to use CueA as a resistance enhancer.




Figure 6. Deletion of the *csoR* gene from Bln0360 increased tolerance of strain to Cu, Zn and Cd. Growth curves of wild-type Bln0360, $\Delta csoR$ and $\Delta cueA$ strain in TY liquid media exposed to different levels of $CuSO_4$ (A), $ZnSO_4$ (B), and $CdSO_4$ (C). Samples were taken and the optical densities (600 nm) were determined when the wild type in the absence of metal ions reached the stationary phase (7 d). Error bars are SD of triplicates. (D) Comparative Cu/Zn/Cd tolerance between Bln0360, $\Delta csoR$ and $\Delta cueA$ strain on TY agar medium containing 2.0 mM $CuSO_4$, 2.8 mM $ZnSO_4$ or 0.1 mM $CdSO_4$. The MTC values are depicted at the right.

To test whether the Zn/Cd resistance mediated by CueA is related to the unusual N-terminus, three CueA variants, CueA- ΔMBD_a , CueA- ΔMBD_b and CueA- ΔMBD_{ab} , were expressed in *E. coli* GG48 and the cells were subjected to Zn/Cd tolerance assays. As Fig. 7B shows, no significant growth difference was observed in the LB media containing indicated concentration of $ZnSO_4$ or $CdSO_4$ between *E. coli* GG48 (CueA) and the variants *E. coli* GG48 (CueA- ΔMBD_a) or *E. coli* GG48 (CueA- ΔMBD_b). However, the growth of the *E. coli* GG48 (CueA- ΔMBD_{ab}) was decreased to the same extent as that of *E. coli* GG48 (vector), when both MBDs were excised. These results suggested that the two N-terminal MBDs were essential for the Zn and Cd resistance enhancement associated with CueA mediation.

CueA is required for plant colonization in Cu overloaded conditions. Considering that Cu resistance could protect the strain Bln0360 against excess Cu-induced damage, thus, we speculate that loss of the Cu pump would affect the nodulation performance of the strain in such conditions. Hence, the wild-type strain and one of each single mutant were combined at three different ratios and then applied to plants. As Fig. 8A shows, the $\Delta cueA$ mutant displayed a lower competitiveness than Bln0360 in Cu-supplemented conditions, as the observed proportion of nodule occupancy by the $\Delta cueA$ mutant was significantly ($P < 0.05$) lower than the expected proportion at every inoculum ratio. At a ratio of 1:1, the percentages of the $\Delta cueA$ mutant recovered from nodules were 36.47% and 24.77% in the presence of 200 mg/kg and 500 mg/kg $CuSO_4$, respectively (Fig. 8A); no distinct effects on competitiveness were observed when plants were grown without Cu treatment. In contrast, the $\Delta csoR$ mutant was equally competitive with the wild-type strain in nodule occupancy in all treatments (Fig. 8B). Collectively, these results show that CueA is required for the nodulation performance of Bln0360 in Cu rich conditions.

Discussion

Analysis of mutants generated by Tn5 revealed that at least three strategies were adopted to alleviate Cu toxicity in Bln0360 (Table 1). Mutants Bln-d and Bln-163 with P_{1B1} -type ATPase CueA-insertion displayed a specific sensitivity to Cu showing the critical role of this transporter in Cu resistance. In Bln-32, a TolC family protein encoded by *tolC* may be involved in Cu transportation; TolC in *E. coli* is an outer membrane components of type I secretion system that can extrude noxious agents such as antibiotics and toxic metal ions. In addition to the efflux-mediated




Figure 7. CueA is involved in Zn and Cd resistance. (A) Zn and Cd resistance of *E. coli* GG48 expressing CueA. *E. coli* GG48 with empty plasmid was used as negative controls. (B) Effects of N-terminal His-rich stretch deletion on the function of CueA in Zn/Cd resistance. Growth of *E. coli* GG48 expressing different truncated versions of CueA was compared in LB medium containing various concentrations of ZnSO₄ (left) or CdSO₄ (right). Symbols represent *E. coli* GG48 expressing empty plasmid, full-length CueA, CueA- ΔMBD_a , CueA- ΔMBD_b and CueA- ΔMBD_{ab} (○, ●, △, ×, and □, respectively).




Figure 8. Effects of Cu resistance on competition for plant nodulation by Bln0360. The ΔcsoR mutant (A) or ΔcueA mutant (B) was mixed with Bln0360 in indicated ratios (1:9, 1:1 or 9:1) and applied to *V. unguiculata* seedlings. Significant differences between the expected and observed colonization percentages were evaluated by chi-square test at a confidence level of 0.05. The statistics did not include nodules containing both strains. The number indicated on each bar represents the total number of nodules from nine plants.

mechanism, Bln0360 adopts enzymatic detoxification against superfluous Cu, as mutant Bln-c with multicopper oxidase encoding gene *copA*-interruption was extremely sensitive to Cu; multicopper oxidase was thought to protect the periplasm from Cu-induced damage through oxidization of Cu⁺ to less toxic Cu²⁺ in aerobic conditions²⁹. Moreover, two membrane integrity related genes (*ctpA* and *lptE*) presumably play a house-keeping role in protecting Bln0360 cells from Cu toxicity, via their function in maintaining the protective permeability barrier of the cell^{18,19,30}.

Homology and transmembrane signature sequences assigned CueA into group P_{1B1} type (Fig. S2)²⁵. ATPases in this group carry out the function of intracellular Cu⁺/Ag⁺ detoxification through pumping Cu⁺/Ag⁺ from the cytoplasm into the periplasm^{7,21,31}. Mutation and complementation analysis showed that *cueA* to play a critical role in Cu detoxification in Bln0360 (Fig. 2C). Given that the toxicity caused by adding CuSO₄ could be eliminated by adding the Cu⁺-specific chelator BCS (Fig. 2C), it was concluded that CueA could carry out the function of Cu⁺ efflux. It should be noted that no altered Ag⁺ tolerance by deletion of *cueA* or *csor* had been observed, indicating that Ag⁺ may be not the substrate of the transporter.

It is accepted that there are few or no “free” (bioavailable) Cu ions in the bacterial cell³²; therefore, strict regulation of Cu homeostasis is very critical. Expression of CueA was negatively regulated by CsoR repressor, similar to the regulation of Cu transporting P-type ATPase in Gram-positive bacteria⁶. In phylogenetically related *A. tumefaciens*, however, expression of CopA was positively regulated by a CueR-like activator, similar to that in *E. coli*⁶. A BLASTN search of the sequence of *csor-cueA* divergon against the sequenced *Bradyrhizobium* genomes found that the organization genetically linking CsoR repressor and Cu ATPase is conserved, suggesting a primary regulator architecture in this genus (unpublished data from our lab). Under Cu limited condition, CsoR represses transcription via binding to the operator-promoter region of the target gene; upon binding Cu⁺, CsoR is released to form the CsoR-DNA complex, resulting in transcription occurred³³. In our study, *csor* and *cueA* were highly expressed in the *csor*-deleted mutant (Fig. 4A–C), which logically was due to the exposed operator-promoter region allowing the access of the RNA polymerase. The promoters of *csor* and *cueA* had parallel strength in the CsoR-deficient background; however, the expression levels of *cueA* were far higher than those of *csor* in the wild-type strain (Fig. 4C). That could be due to the presence of bidirectional promoter and the CsoR-box located in the overlapped promoters of *cueA* and *csor* (Fig. 1C). As a result, coordination/competition between the RNA polymerase and the CsoR constitutes a significant part of the regulatory process³⁴. In other cases, the two promoters occupied the same sequence element on opposing DNA strand, therefore, the collision between RNA polymerases likely influenced the expression of each gene. In *csor-cueA* divergon, the expression of CsoR repressor was regulated by itself. Coordination between the expression and derepression of transcriptional repressor allows the regulation process under a more narrow control³⁵.

Studies have shown that Cu⁺, but not other metal ions, bind to the CsoR of *M. tuberculosis* and *L. monocytogenes*, thereby relieving the interaction between this regulator and promoter DNA and allowing transcription to proceed^{26,36}. However, expression of the *csor-cueA* divergon from Bln0360 could be induced by Cu²⁺ as well as by Zn²⁺ and Cd²⁺ (Fig. 5A). The Zn²⁺-inducible expression of *csor* regulons has been reported in *C. glutamicum*^{27,37}. For Cd²⁺-inducible expression, it is easy to understand in that Zn²⁺ and Cd²⁺ share similar coordination geometries³⁸. According to the derepression mechanism, the effects of these metal ions binding between CsoR and its operator-promoter DNA was reflected in the expression of the *csor-cueA* divergon. Our findings confirmed that the role of metal ions (Cu²⁺, Zn²⁺ and Cd²⁺) in the inhibition of DNA binding activity (Fig. 5B). We also find Zn²⁺-induced expression of *csor-cueA* divergon was less than Cu²⁺ or Cu⁺ (Figs 2A and 5), in agreement with previous study in *C. glutamicum*²⁷. Additionally, the CsoR-box (5'-TATACCCnnnnGGGTATA-3') of *B. liaoningense* showed higher similarity to *C. glutamicum* (5'-ATACCCnnnGGGTAT-3'), suggesting a similar coordination property of metal cations by the two CsoRs. *T. thermophilus* CsoR can coordinate multiple metal ions including divalent Cu²⁺/Zn²⁺/Cd²⁺ and monovalent Cu⁺⁹. However, the metal-binding motif of *T. thermophilus* CsoR is H-C-H-H, instead of specific C-H-C being conserved in known CsoRs from other bacteria⁹. Sakamoto K *et al.* thought that the low selectivity of *T. thermophilus* CsoR to metal ions is due to the presence of His(70) and His(5) residues⁹. Likewise, we did not find the additional His residue in Bln0360 CsoR corresponding to the His(5) residue in *T. thermophilus* CsoR. Therefore, further studies need to elucidate the critical role of *B. liaoningense* CsoR in the Zn²⁺- and Cd²⁺-dependent expression of *csor-cueA* divergon.

In this study, expression of *cueA* was found to be up-regulated by Cu²⁺/Zn²⁺/Cd²⁺ stress (Figs 2 and 5), suggesting that *cueA* was involved in protecting cells against these metals toxicity. This is validated by complementation studies in Δ *cueA* mutant and Zn/Cd sensitive *E. coli* (Fig. 7). Very few studies so far have addressed the effect of P_{1B1}-type ATPases on resistance to divalent metal ions. Intriguingly, an increase in Cu/Zn/Cd tolerance by overexpression of *cueA* in *csor* deficient mutant was observed in this study (Fig. 6). CueA features two His-rich stretches at its N-terminus, which is fundamentally distinct from the typical GXXCXXC motif in P_{1B}-type ATPase (Fig. 3A). P_{1B}-type ATPases with N-terminal His-rich stretch tend to transport divalent metal ion (Cu²⁺, Zn²⁺, Cd²⁺ and Pb²⁺), and the His-rich stretch is proposed to be a putative MBD^{24,25}. In our complementation studies, deletion of the two His-rich MBD of CueA entirely abolished its function conferring Zn/Cd resistance in *E. coli* GG48 (Fig. 7B), showing the essential role of these amino acids in Zn/Cd resistance to the transporter. There were no previous reports describing the role of His-rich N-MBD of P_{1B2}-type ATPase. However, the His-rich stretch in the C-terminus of plant Zn²⁺/Cd²⁺ transporting ATPases (TcHMA4 and AtHMA4) has been demonstrated experimentally and was shown to be essential for Zn²⁺/Cd²⁺ binding or in the regulation of the enzyme³⁹. Lack of Zn/Cd sensitive phenotype in Δ *cueA* mutant may due to functional redundancy of CueA for the metals, in that multiple Zn²⁺/Cd²⁺ transporters systems are present in the genome of *B. liaoningense* CCNWSX0360 (LUKO0000000.1). This interpretation was supported by an unsuccessful screening for Zn sensitive mutant from the Tn5-induced mutant libraries.

Metal resistance might facilitate the survival of rhizobia in both free-living and symbiotic states in heavy metal-rich conditions. In our study, the *cueA*-deficient mutant showed the same nodulation capacity as the wild-type strain in the absence of Cu, suggesting that CueA is not essential for nodulation in normal conditions. However, in Cu rich conditions, the nodulation occupancy of the mutant decreased significantly (Fig. 8A). This is similar to the previous report that the nodulation of a Ni²⁺-sensitive mutant of *B. japonicum* was affected by the presence of nickel in soil⁴⁰. Similarly, deletion of *dmeRF*, a Ni²⁺/Co²⁺ transport system, resulted in symbiotic performance defects of *Rhizobium leguminosarum* bv. *viciae* in high-cobalt conditions³⁵. In addition, the decreased nodulation occupancy of the *cueA*-deficient mutant was Cu concentration-dependent, as it was less efficient in

the presence of 500 mg/kg Cu²⁺ than 200 mg/kg Cu²⁺ (Fig. 8A). On the basis of these results, it is established that metal resistance determinants gives strains a competitive advantage in the establishment of symbiotic association with their host plants when faced with metal stress. The establishment of a symbiotic system is an intricate process, and how the metal(s) affect(s) the symbiosis remains unknown. With regard to the observed impairment in nodule occupancy efficiency in our Cu sensitive mutant, it is at least clear that Cu resistance is important for the symbiosis between *B. liaoningense* and *V. unguiculata* in Cu polluted soil. Further studies are required to elucidate the Zn/Cd resistance mechanism of CueA and the actual role of the Cu tolerance system in the symbiotic processes of these endosymbiotic bacteria under Cu stress.

Methods

Bacterial strains, media and growth conditions. Bacterial strains used in this study are listed in Table S1. Screening and identification of Cu resistant rhizobia were carried out as described in Supplementary Methods. *E. coli* strains including GG48 ($\Delta zitB::Cm zntA::Km$)⁴¹ were routinely cultured in LB medium at 37 °C. *B. liaoningense* strains were cultured in Tryptone-Yeast⁴² or YMA (Yeast-Mannitol Agar) medium at 28 °C⁴³. Antibiotics were supplemented as required: streptomycin (Sm), 50 µg/ml; neomycin²⁴, 100 µg/ml; kanamycin (Km), 50 µg/ml; ampicillin (Amp), 100 µg/ml; gentamicin (Gm), 25 µg/ml.

Screening Cu sensitive Tn5 mutant and determining the insertion sites. Transposon Tn5 mutagenesis of Bln0360 was carried out using suicide plasmid pRL1063a according to the protocol for mutagenesis of *Rhizobium tropici* CIAT899^{44,45}. Appropriate dilutions of the mating mixture were plated on YMA plates containing Sm and Amp. Individual colonies from YMA medium were picked and streaked onto TY plates supplemented with 0 and 0.8 mM CuSO₄. Mutant clones that grew weakly or not at all on 0.8 mM Cu²⁺ plates but that grew well on control plates were selected for determination of Tn5 insertion sites. To map the sequences contiguous with the inserted transposon, procedures including genomic DNA digestion, self-circularization, transformation, and sequencing were performed following published protocols¹⁵. Homology searches of nucleotide and deduced amino sequences were performed with the BLAST program (<http://www.ncbi.nlm.nih.gov/blast>).

Deletion mutation and complementation of *csor* and *cueA*. Plasmids used in the study were constructed as outlined in Supplementary Methods. Plasmid constructs carrying genes to be deleted (pK18mobsacB- $\Delta cueA$ and pK18mobsacB- $\Delta csor$) were introduced into Bln0360 by conjugal transfer using the *E. coli* S17-1 mobilizing strain according to the method of Simon *et al.*⁴⁶. Gene deletion was achieved by sequential double crossover recombinants selection using YMA plate containing Neo/Amp and 10% sucrose as described previously⁴⁷. To complement the $\Delta csor$ and $\Delta cueA$ mutants, plasmids pBBR5-*csor* and pBBR5-*cueA* respectively were electroporated into the $\Delta csor$ and $\Delta cueA$ mutants. To analyze the role of the two N-terminal His-rich stretches of CueA, plasmid constructs pJQ-1, pJQ-2 and pJQ-12 were transformed into Cu sensitive $\Delta cueA$ mutant and Zn sensitive *E. coli* GG48, respectively.

Construction of chromosomal fusion reporter strains and β -galactosidase assays. The *lacZ* fusion reporter vectors (pK18mobSacB-P_{*cueA*}::*lacZ* and pK18mobSacB-P_{*csor*}::*lacZ*; Table S1) were introduced into *E. coli* S17-1 λ pir and mated with Bln0360 and $\Delta csor$ mutant strain, respectively⁴⁶. Transconjugants were selected by plating on YMA plates supplemented with Neo and Amp and confirmed by PCR and sequencing. The *lacZ* fusion reporter strains were grown to mid-log phase in TY broth and then indicated inducers were added. After 2 h incubation, β -Galactosidase activity was measured using o-nitrophenyl- β -D-galactopyranoside as the substrate and was expressed in Miller units⁴⁸.

Quantitative real-time PCR and rapid amplification of cDNA ends. Total RNA was extracted from exponentially growing cell cultures using the hot phenol method⁴⁹. First-strand cDNA was reverse transcribed using the PrimeScriptTM RT reagent kit with gDNA Eraser (Takara, Dalian, China). qRT-PCR was performed in a CFX96 Real-Time PCR Detection System (Bio-Rad, Hercules, CA USA) with a SYBR[®] Premix Ex TaqTM II (Tli RNaseH Plus) kit (Takara). For all primer sets (Table S2), the following cycling parameters were used: 95 °C for 30 s, followed by 40 cycles of 94 °C for 15 s and 50 °C for 30 s. The relative abundance of 16S rRNA was used as the internal standard. To determine the initiating nucleotide for transcripts of *csor* and *cueA*, the 5' end of *csor* and *cueA* mRNA were analyzed with the SMART RACE cDNA amplification kit (Clontech, Mountain View, CA, USA) according to the manufacturer's protocols.

Assays for tolerance to heavy metals. Log phase cultures of *B. liaoningense* or *E. coli* strains were washed twice and then 10-fold serially diluted in PBS buffer. An aliquot (3 µl) of dilution was spotted onto corresponding agar plates (TY or LB) containing varying concentrations of metal ions including Cu²⁺, Zn²⁺, Cd²⁺, Ni²⁺, or Ag⁺. Growth was monitored after 7 d at 28 °C for *B. liaoningense* and 36 h at 37 °C for *E. coli*. Dose-response growth curves describing the action of metal ions on bacterial cells were also performed. Log phase cultures were used to inoculate into parallel cultures containing increasing metal concentrations. The initial optical density of the cell suspension at 600 nm (OD₆₀₀) was adjusted to 0.02. Cells were cultivated 7 d at 28 °C with shaking at 140 rpm for Bln0360 and 24 h at 37 °C with shaking at 200 rpm for *E. coli*, and the optical density was determined at 600 nm. Each experiment was repeated three times.

Competition assay for nodulation. Nodule occupancy of the $\Delta csor$ and $\Delta cueA$ strains in co-inoculations with Bln0360 were carried out as described by Patankar⁵⁰. Briefly, surface-sterilized *V. unguiculata* seedlings were transplanted into pouches containing a sterilized mixture of vermiculite-perlite (2:1, v/v) supplemented with 0, 200 or 500 mg/kg CuSO₄. After 2 days, individual plants were co-inoculated with 1 ml (10⁶ CFU) of inoculant combination containing the parental Bln0360 and either the $\Delta cueA$ or $\Delta csor$ mutant in approximate 9:1, 1:1

and 1:9 ratios, with nine plants for each treatment. Seedlings without inoculation were used as negative controls. Plants were incubated in a controlled growth chamber (humidity: 70%; day condition: 22 °C, 16 h; night condition: 16 °C, 8 h). Fahraeus nitrogen-free nutrient solution was used to replenish the pouches, if required⁵¹. After 25 days, nodule samples from each treatment were surface sterilized and crushed in an appropriate volume of sterilized water as described by Shima⁵². Bacteria released from the crushed nodules were spotted onto YMA plates and the genetic backgrounds of the resultant colonies were confirmed by colony PCR with primer pair cueA-qc1/cueA-qc4. Significant difference analysis between the recovery ratio and initial inoculation ratio of the mutants was used to evaluate the competitive nodule occupancies⁵³.

Statistical analysis. Statistical analyses were carried out using SPSS 19.0 software (SPSS Inc., Chicago, IL, USA). Paired two-tailed Student's *t*-test was performed to determine significant differences among the treatments in transcription and β -galactosidase activity analysis. Statistical significance in competitive nodulation experiments was assessed by chi-square test at the significance level of $P < 0.05$.

References

- Andreini, C., Bertini, I., Cavallaro, G., Holliday, G. L. & Thornton, J. M. Metal ions in biological catalysis: from enzyme databases to general principles. *J. Biol. Inorg. Chem.* **13**, 1205–1218 (2008).
- Grass, G. & Rensing, C. Genes involved in copper homeostasis in *Escherichia coli*. *J. Bacteriol.* **183**, 2145–2147 (2001).
- Soliz, M. & Stoyanov, J. V. Copper homeostasis in *Enterococcus hirae*. *FEMS Microbiol. Rev.* **27**, 183–195 (2003).
- Dupont, C. L., Grass, G. & Rensing, C. Copper toxicity and the origin of bacterial resistance—new insights and applications. *Metallomics* **3**, 1109–1118 (2011).
- Smith, A. T., Smith, K. P. & Rosenzweig, A. C. Diversity of the metal-transporting P_{1B}-type ATPases. *J. Biol. Inorg. Chem.* **19**, 947–960 (2014).
- Rademacher, C. & Masepohl, B. Copper-responsive gene regulation in bacteria. *Microbiology* **158**, 2451–2464 (2012).
- Fan, B. & Rosen, B. P. Biochemical characterization of CopA, the *Escherichia coli* Cu(I)-translocating P-type ATPase. *J. Biol. Chem.* **277**, 46987–46992 (2002).
- Mana-Capelli, S., Mandal, A. K. & Argüello, J. M. *Archaeoglobus fulgidus* CopB is a thermophilic Cu²⁺-ATPase functional role of its histidine-rich N-terminal metal binding domain. *J. Biol. Chem.* **278**, 40534–40541 (2003).
- Sakamoto, K., Agari, Y., Agari, K., Kuramitsu, S. & Shinkai, A. Structural and functional characterization of the transcriptional repressor CsoR from *Thermus thermophilus* HB8. *Microbiology* **156**, 1993–2005 (2010).
- Canfield, D. E., Glazer, A. N. & Falkowski, P. G. The evolution and future of Earth's nitrogen cycle. *Science* **330**, 192–196 (2010).
- Hao, X. *et al.* Copper tolerance mechanisms of *Mesorhizobium amorphae* and its role in aiding phytostabilization by *Robinia pseudoacacia* in copper contaminated soil. *Environ. Sci. Technol.* **49**, 2328–2340 (2015).
- Wani, P. A., Khan, M. S. & Zaidi, A. Effect of metal-tolerant plant growth-promoting *Rhizobium* on the performance of pea grown in metal-amended soil. *Arch. Environ. Contam. Toxicol.* **55**, 33–42 (2008).
- Gaetke, L. M. & Chow, C. K. Copper toxicity, oxidative stress, and antioxidant nutrients. *Toxicology* **189**, 147–163 (2003).
- Fan, L. *et al.* Characterization of a copper-resistant symbiotic bacterium isolated from *Medicago lupulina* growing in mine tailings. *Bioresour. Technol.* **102**, 703–709 (2011).
- Li, Z., Ma, Z., Hao, X., Rensing, C. & Wei, G. Genes conferring copper resistance in *Sinorhizobium meliloti* CCNWSX0020 also promote the growth of *Medicago lupulina* in copper-contaminated soil. *Appl. Environ. Microbiol.* **80**, 1961–1971 (2014).
- Koronakis, V., Sharff, A., Koronakis, E., Luisi, B. & Hughes, C. Crystal structure of the bacterial membrane protein TolC central to multidrug efflux and protein export. *Nature* **405**, 914–919 (2000).
- Kaneko, T. *et al.* Complete genomic sequence of nitrogen-fixing symbiotic bacterium *Bradyrhizobium japonicum* USDA110. *DNA Res.* **9**, 189–197 (2002).
- Dong, J., Signo, K. S., Vanderlinde, E. M., Yost, C. K. & Dahms, T. E. Atomic force microscopy of a *ctpA* mutant in *Rhizobium leguminosarum* reveals surface defects linking CtpA function to biofilm formation. *Microbiology* **157**, 3049–3058 (2011).
- Gilbert, K. B., Vanderlinde, E. M. & Yost, C. K. Mutagenesis of the carboxy terminal protease CtpA decreases desiccation tolerance in *Rhizobium leguminosarum*. *FEMS Microbiol. Lett.* **272**, 65–74. doi: 10.1111/j.1574-6968.2007.00735.x (2007).
- Nawapan, S. *et al.* Functional and expression analyses of the *cop* operon, required for copper resistance in *Agrobacterium tumefaciens*. *J. Bacteriol.* **191**, 5159–5168 (2009).
- Rensing, C., Fan, B., Sharma, R., Mitra, B. & Rosen, B. P. CopA: an *Escherichia coli* Cu(I)-translocating P-type ATPase. *Proc. Natl. Acad. Sci. USA* **97**, 652–656 (2000).
- Gold, B. *et al.* Identification of a copper-binding metallothionein in pathogenic mycobacteria. *Nat. Chem. Biol.* **4**, 609–616 (2008).
- Blindauer, C. A. & Schmid, R. Cytosolic metal handling in plants: determinants for zinc specificity in metal transporters and metallothioneins. *Metallomics* **2**, 510–529 (2010).
- Tong, L., Nakashima, S., Shibasaki, M., Katsuhara, M. & Kasamo, K. A novel histidine-rich CPx-ATPase from the filamentous cyanobacterium *Oscillatoria brevis* related to multiple-heavy-metal cotolerance. *J. Bacteriol.* **184**, 5027–5035 (2002).
- Argüello, J. Identification of ion-selectivity determinants in heavy-metal transport P_{1B}-type ATPases. *J. Membr. Biol.* **195**, 93–108 (2003).
- Liu, T. *et al.* CsoR is a novel *Mycobacterium tuberculosis* copper-sensing transcriptional regulator. *Nat. Chem. Biol.* **3**, 60–68 (2007).
- Teramoto, H., Inui, M. & Yukawa, H. *Corynebacterium glutamicum* CsoR acts as a transcriptional repressor of two copper/zinc-inducible P_{1B}-type ATPase operons. *Biosci., Biotechnol., Biochem.* **76**, 1952–1958 (2012).
- Omotajo, D., Tate, T., Cho, H. & Choudhary, M. Distribution and diversity of ribosome binding sites in prokaryotic genomes. *BMC Genomics* **16**, 1–8 (2015).
- Grass, G. & Rensing, C. CueO is a multi-copper oxidase that confers copper tolerance in *Escherichia coli*. *Biochem. Biophys. Res. Commun.* **286**, 902–908 (2001).
- Bandara, A. B., Sriranganathan, N., Schurig, G. G. & Boyle, S. M. Carboxyl-terminal protease regulates *Brucella suis* morphology in culture and persistence in macrophages and mice. *J. Bacteriol.* **187**, 5767–5775 (2005).
- Odermatt, A., Krapf, R. & Soliz, M. Induction of the putative copper ATPases, CopA and CopB, of *Enterococcus hirae* by Ag⁺ and Cu²⁺, and Ag⁺ extrusion by CopB. *Biochem. Biophys. Res. Commun.* **202**, 44–48 (1994).
- Rae, T., Schmidt, P., Pufahl, R., Culotta, V. & O'halloran, T. Undetectable intracellular free copper: the requirement of a copper chaperone for superoxide dismutase. *Science* **284**, 805–808 (1999).
- Chang, F.-M. J. *et al.* Cu(I)-mediated allosteric switching in a copper-sensing operon repressor (CsoR). *J. Biol. Chem.* **289**, 19204–19217 (2014).
- Christoffersen, C. A., Brickman, T. J. & McIntosh, M. A. Regulatory architecture of the iron-regulated *fepD-ybdA* bidirectional promoter region in *Escherichia coli*. *J. Bacteriol.* **183**, 2059–2070 (2001).
- Rubio-Sanz, L., Prieto, R., Imperial, J., Palacios, J. & Brito, B. Functional and expression analysis of the metal-inducible *dmeRF* system from *Rhizobium leguminosarum* bv. *viciae*. *Appl. Environ. Microbiol.* **79**, 6414–6422 (2013).

36. Corbett, D. *et al.* The combined actions of the copper-responsive repressor CsoR and copper-metallochaperone CopZ modulate CopA-mediated copper efflux in the intracellular pathogen *Listeria monocytogenes*. *Mol. Microbiol.* **81**, 457–472 (2011).
37. Teramoto, H., Yukawa, H. & Inui, M. Copper homeostasis-related genes in three separate transcriptional units regulated by CsoR in *Corynebacterium glutamicum*. *Appl. Microbiol. Biotechnol.* **99**, 3505–3517 (2015).
38. Lutsenko, S., Petrukhin, K., Cooper, M. J., Gilliam, C. T. & Kaplan, J. H. N-terminal domains of human copper-transporting adenosine triphosphatases (the Wilson's and Menkes disease proteins) bind copper selectively *in vivo* and *in vitro* with stoichiometry of one copper per metal-binding repeat. *J. Biol. Chem.* **272**, 18939–18944 (1997).
39. Eren, E., Kennedy, D. C., Maroney, M. J. & Argüello, J. M. A novel regulatory metal binding domain is present in the C terminus of *Arabidopsis* Zn²⁺-ATPase HMA2. *J. Biol. Chem.* **281**, 33881–33891 (2006).
40. Chaintreuil, C. *et al.* Nickel resistance determinants in *Bradyrhizobium* strains from nodules of the endemic New Caledonia legume *Serianthes calycina*. *Appl. Environ. Microbiol.* **73**, 8018–8022 (2007).
41. Grass, G. *et al.* ZitB (YbgR), a Member of the Cation Diffusion Facilitator Family, Is an Additional Zinc Transporter in *Escherichia coli*. *J. Bacteriol.* **183**, 4664–4667 (2001).
42. Banci, L. *et al.* Structural basis for metal binding specificity: the N-terminal cadmium binding domain of the P1-type ATPase CadA. *J. Mol. Biol.* **356**, 638–650 (2006).
43. Vincent, J. *The cultivation, isolation and maintenance of rhizobia*. Blackwell Scientific, Oxford, United Kingdom (1970).
44. Riccillo, P. M. *et al.* Glutathione is involved in environmental stress responses in *Rhizobium tropici*, including acid tolerance. *J. Bacteriol.* **182**, 1748–1753 (2000).
45. Figurski, D. H. & Helinski, D. R. Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in trans. *Proc. Natl. Acad. Sci. USA* **76**, 1648–1652 (1979).
46. Simon, R., O'Connell, M., Labes, M. & Pühler, A. Plasmid vectors for the genetic analysis and manipulation of rhizobia and other gram-negative bacteria. *Methods Enzymol.* **118**, 640–659 (1985).
47. Angulo, V. A. G. *et al.* Identification and characterization of RibN, a novel family of riboflavin transporters from *Rhizobium leguminosarum* and other proteobacteria. *J. Bacteriol.* **195**, 4611–4619 (2013).
48. Miller, J. H. *Experiments in molecular genetics*. Cold Spring Harbor Laboratory Press, New York (1972).
49. Hosoda, N. *et al.* Translation termination factor eRF3 mediates mRNA decay through the regulation of deadenylation. *J. Biol. Chem.* **278**, 38287–38291 (2003).
50. Patankar, A. V. & González, J. E. An orphan LuxR homolog of *Sinorhizobium meliloti* affects stress adaptation and competition for nodulation. *Appl. Environ. Microbiol.* **75**, 946–955 (2009).
51. Fraeus, G. The infection of clover root hairs by nodule bacteria studied by a simple glass slide technique. *Microbiology* **16**, 374–381 (1957).
52. Eda, S., Mitsui, H. & Minamisawa, K. Involvement of the *smeAB* multidrug efflux pump in resistance to plant antimicrobials and contribution to nodulation competitiveness in *Sinorhizobium meliloti*. *Appl. Environ. Microbiol.* **77**, 2855–2862 (2011).
53. Beattie, G., Clayton, M. & Handelsman, J. Quantitative comparison of the laboratory and field competitiveness of *Rhizobium leguminosarum* biovar *phaseoli*. *Appl. Environ. Microbiol.* **55**, 2755–2761 (1989).

Acknowledgements

This work was funded by the National Key Research and Development Program of China (2016YFD0800706) and National Science Foundation of China (31270012 and 31370142).

Author Contributions

G.W., Z.L., M.C. and X.S. conceived of and organized the study. J.L. and M.Z. performed the experiments. J.L. and M.L. wrote the paper. All authors have read and approved the final manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Liang, J. *et al.* Functional characterization of a *csoR-cueA* divergon in *Bradyrhizobium liaoningense* CCNWSX0360, involved in copper, zinc and cadmium cotolerance. *Sci. Rep.* **6**, 35155; doi: 10.1038/srep35155 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016