

SCIENTIFIC REPORTS



OPEN

Novelty during a late postacquisition time window attenuates the persistence of fear memory

Cynthia Katche¹, Micol Tomaiuolo¹, Guido Dorman¹, Jorge H. Medina^{1,2} & Haydee Viola^{1,3}

Received: 20 June 2016

Accepted: 26 September 2016

Published: 13 October 2016

Learning to avoid threats in the environment is highly adaptive. However, sometimes a dysregulation of fear memories processing may underlie fear-related disorders. Despite recent advances, a major question of how to effectively attenuate persistent fear memories in a safe manner remains unresolved. Here we show experiments employing a behavioural tool to target a specific time window after training to limit the persistence of a fear memory in rats. We observed that exposure to a novel environment 11 h after an inhibitory avoidance (IA) training that induces a long-lasting memory, attenuates the durability of IA memory but not its formation. This effect is time-restricted and not seen when the environment is familiar. In addition, novelty-induced attenuation of IA memory durability is prevented by the intrahippocampal infusion of the CaMKs inhibitor KN-93. This new behavioural approach which targets a specific time window during late memory consolidation, might represent a new tool for reducing the durability of persistent fear memories.

Memories, including those which are painful or fear-inducing, are crucial for our lives and encompass the essence of who we are. They are quite important for our day-to-day functioning and hence for our quality of life. However, sometimes fear learning is maladaptive, generating persistent memories with an excessive fear and anxiety. Current therapies for fear-related disorders involve pharmacological or behavioural manipulations of long-lasting fear memories, and rely mainly on reconsolidation and extinction processes. However, extinction-based exposure therapy has limited efficacy¹. For instance, immediate or delayed extinction procedures induce a reduction in aversive memories that are context-dependent and short-lived^{2,3}. In contrast, when extinction training is given after the retrieval of a traumatic experience in healthy volunteers, a reduction of the original fearful memory is found⁴. This finding paralleled those obtained in rodents⁵, (but see ref. 6). However, it will be desirable to induce fear attenuation without the need to submit the individuals to the retrieval of the fear experience.

We and others have demonstrated the existence of a novel phase, happening around 12 h after acquisition, specifically involved in the persistence but not the formation of fear long-term memory (LTM)^{7,8}. This phase depends on *de novo* protein synthesis in the hippocampus and amygdala and is controlled by dopaminergic inputs⁹. These findings might open an opportunity to generate new treatments to attenuate the persistence of fear memories.

We have previously reported that a robust retrograde amnesia of IA memory formation can be induced by novelty¹⁰ when is presented during the early stage of memory consolidation. Thus, on the basis of the above findings we reasoned that subjecting rats to a novel environment late after acquisition of a one trial IA learning task could selectively attenuate the maintenance of the mnemonic trace without interfering with its formation. This behavioural approach that targets a late and specific time window while consolidation is still in process represents a promising non pharmacological procedure for reducing the durability of persistent memories.

¹Instituto de Biología Celular y Neurociencias, Facultad de Medicina “Dr Eduardo De Robertis” (UBA-CONICET), UBA, Paraguay 2155, 3 piso, Buenos Aires (C1121ABG), Argentina. ²Departamento de Fisiología, Facultad de Medicina, UBA, Paraguay 2155, 7 piso, Buenos Aires (C1121ABG), Argentina. ³Departamento de Fisiología, Biología Molecular y Celular Dr Hector Maldonado, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires (C1428EGA), Argentina. Correspondence and requests for materials should be addressed to H.V. (email: hviola@fmed.uba.ar)




Figure 1. Exposure to a novel OF exploration 11 h post-training impairs IA-LTM persistence but not LTM formation. Rats were subjected to IA training (Ctrl) or to IA training plus a novel OF 11 h (OF 11 h) after IA training. Data are expressed as mean \pm SEM of training (TR) or test (TS) session step down-latency at 2 (a) or 7 days (b) after IA training. * $p < 0.05$, Ctrl vs. OF 11 h group; Student's t test, $n = 10$ –12 per group.

Results

Exposure to a novel OF exploration 11 h post-training impairs IA-LTM persistence. The IA is a fear-motivated associative learning task that is hippocampus-dependent and acquired in a single and brief training session¹¹. Exposure to an OF 11 h after IA training did not affect memory retention performance of the avoidance task when evaluated 2 days after training (Fig. 1a). In contrast, when an independent group of animals was tested 7 days after IA training we found an impairment in memory retention (Fig. 1b, $p < 0.05$ in comparison to the Control IA-trained group of rats that were not exposed to OF; Student's t test, $n = 10$ –12). This effect was observed during a critical time window after training because no altered retention scores were seen when IA-trained rats were exposed to a novel environment 8 or 23 h after training and tested 2 or 7 days thereafter (Fig. 2). These findings indicate that exposure to a novel environment late after training selectively attenuates, in a time-dependent manner, the persistence of IA LTM storage without altering memory formation.

OF requires to be novel in order to impair IA-LTM persistence. Is the attenuating effect of the novel environment on memory persistence due to the perception of novelty? To determine whether novelty is a key factor in reducing the durability of IA LTM, rats were exposed for 30 min to a square OF 24 h prior to the IA training. Animals were then exposed again to the square OF (Fam OF group) or to a round shape novel OF (New OF group) 11 h after IA training and tested for memory retention 7 days after training. As shown in Fig. 3a, the Fam OF group did not exhibit any deficit ($p > 0.05$, compared to Ctrl group, $n = 12$ –13). As before, animals subjected to a novel OF at 11 h posttraining had a clear-cut impairment in retention performance 7 days after training (Fig. 3a, $p < 0.001$ compared to Ctrl rats; Newman-Keuls test after ANOVA, $n = 13$ –14). This amnesic effect induced by a novel OF was also seen when rats were exposed to the square OF the day before the IA training (Fig. 3a, $p < 0.001$ New OF 11hs compared to Ctrl rats; Newman-Keuls test after ANOVA, $n = 9$ –13). Figure 3b shows, in addition, that rats subjected to a 2nd exposure of the same OF performed less crossings and rearings than rats that explored the OF for the 1st time or rats that experienced a different OF during the 2nd exposure. ($p < 0.01$ for crossing, $p < 0.05$ for rearings; Fam OF vs. other groups, Newman-Keuls test after ANOVA, $n = 9$ –14). This implies that in the 2nd exposure to the same OF the animals recognized it as being familiar and, as a consequence, they explored it short time¹².

Novel OF impairs IA-LTM persistence through a CaMK-dependent mechanism. What are the molecular events required for attenuating fear LTM storage induced by novelty? Given that the detection of spatial novelty is associated with an activation of CaMKs in the dorsal hippocampus and that CaMKs are required for memory formation of OF habituation^{13,14}, we next determined whether blockade of hippocampal CaMKs activity (Fig. 4a) could prevent the effect of novelty on memory persistence. Bilateral infusions of the CaMKs inhibitor KN-93 intra-CA1 region of the dorsal hippocampus¹⁵, around 11 h after IA training to rats that were not exposed to spatial novelty, did not modify the retention scores in a 7-day test session (Fig. 4b). However, the infusion of KN-93 into the CA1 region of the dorsal hippocampus 15 min before exposing rats to a novel OF 11 h after IA training, blocked the retrograde amnesic effect of OF exposure (Fig. 4b, $p < 0.01$, compared to Veh + OF rats; Newman-Keuls test after ANOVA, $n = 12$ –13).

Discussion

Anxiety disorders are among the most prevalent psychiatric illnesses¹⁶. They are conceptualized as disorders of emotional learning processes¹⁷ or fear regulation, in which behavioural avoidance is a central feature. The idea of modifying memory processing as a treatment for fear-related disorders is not new. The moment when an aversive experience occurs is the first opportunity to interfere with the formation of a fear-motivated memory.




Figure 2. A novel OF exploration performed 8 or 23 h after training do not affect IA-LTM persistence. Rats were subjected to IA training (Ctrl) or to IA training plus a novel OF 8 h (OF 8 h; **a,b**) or 23 h (OF 23 h; **c,d**) after IA training. Data are expressed as mean \pm SEM of training (TR) or test (TS) session step down-latency at 2 (**a,c**) or 7 days after IA training (**b,d**). Student's *t* test, $n = 8$ per group.

For example, modulation of opioid systems has proved to be effective in PTSD¹⁸. On the other hand, blocking memory formation after trauma with β -adrenoceptor blockers has yielded mixed results¹⁹. In addition, early behavioural intervention after a fearful experience in rodents lead to conflicting findings^{20,21}. In recent years there has been some improvement in the attenuation of fear-related memories by manipulating memory reconsolidation and extinction processes. However, extinction-based exposure therapy might have limited efficacy¹. Immediate or delayed extinction procedures induce a reduction in aversive memories that are context-dependent and short-lived^{2,3}. Several reconsolidation experiments revealed that memories become increasingly resistant to pharmacological manipulations as they grow older^{22,23}, or that postreactivation induced-amnesia recovers over time^{24,25}. On the other hand, a successful attempt to attenuate fear memories in rats and humans was obtained using a novel behavioural design involving a mixed reconsolidation-extinction procedure^{4,5}, (but see ref. 6).

Our results demonstrated that a simple behavioural intervention at the critical time window after training attenuates the persistence of fear-motivated memory storage in rats in a time-dependent manner. The amnesic effect of a new learning on previously encoded material is known as retroactive interference (RI). It has been suggested that there are two types of RI that produce forgetting, named diversion and similarity RI. While the former affects consolidation, the latter affects retrieval^{26,27}. Our present findings represent the first example of a behavioural RI paradigm that successfully impairs LTM persistence in a critical time window many hours after a learning experience. Using this protocol, we expanded the time period in which memory formation and storage can be modified. Moreover, we described that the event that induces the effect is a novel spatial exploration that engages the activation of hippocampal CaMKs. Given that high doses of KN-93 could inhibit some CaMK isoforms²⁸, we cannot ascertain which of the isoforms are involved in this effect.

The delayed RI of fear memory persistence caused by novel OF exposure observed in the present study was not due apparently to a deficit in IA memory retrieval. In fact, animals normally expressed fear memory two days




Figure 3. OF requires to be novel in order to impair IA-LTM persistence. (a) Animals were exposed to a single OF session 11 h post IA training (OF 11 h) or to an additional 30 min square OF session the day before IA training (pre-exposed to OF). Rats explored the square familiar OF (Fam OF 11 h) or a round shape novel OF (New OF 11 h) 11 h after IA training. Data are expressed as mean \pm SEM of training (TR) or test (TS) session step down-latency at 7 days after IA training. * $p < 0.05$, vs. Fam OF 11 h group, *** $p < 0.001$ vs. Ctrl; Newman-Keuls test after ANOVA, $n = 9-14$ per group. (b) Bar graph represents the total number of quadrant crossings (Top) or rearings (Bottom) during 5 min in the OF 11 h, Fam OF 11 h and New OF 11 h groups. Data are presented as mean \pm SEM. * $p < 0.05$; ** $p < 0.01$, vs Fam OF 11 h group, Newman-Keuls test after ANOVA, $n = 9-14$ per group.




Figure 4. Novel OF impairs IA-LTM persistence through a CaMK dependent mechanism. (a) Schematic representation of rat brain sections at three rostrocaudal planes (AP: -3.8 , -4.3 , -4.8 mm, from bregma) taken from the atlas of Paxinos and Watson (1997). In stippling, the extension of the area reached by the infusions in the dorsal hippocampus (CA1). (b) Animals not exposed to OF were bilaterally infused with vehicle (Veh) or KN-93 (KN) into CA1 region of the dorsal hippocampus, 11 h after IA training as control groups. Animals exposed to a novel OF 11 h after IA training received intra CA1 infusions of either vehicle (Veh + OF 11 h) or KN-93 (KN + OF 11 h) 15 min before OF. Data are expressed as mean \pm SEM of training (TR) or test session (TS) step down-latency at 7 days after IA training. * $p < 0.05$; ** $p < 0.01$, *** $p < 0.001$, vs Veh + OF 11 h, Newman-Keuls test after ANOVA, $n = 11-15$ per group.

after IA training. Also, exposure to the OF 8 or 23 h after IA training left the expression of 2 and 7 day-old fear memories intact. Moreover, spatial novelty enhances, but does not reduce, IA memory retrieval when novelty is close to the testing session¹². Stress is an unlike factor involved in the deleterious effect of novelty on IA memory

maintenance since it has been recently shown that exposure to stress around 12 h after training did not impair but rather enhanced contextual fear memory persistence²⁹. We cannot totally rule out that novelty-induced impairment of memory retention might represent a type of “behavioural metaplastic” change³⁰. However, behavioural metaplasticity refers to a modification of a behavioural process (in our case the IA training session) by a prior event and not by a subsequent experience as it occurs in our study (an exposure to an OF several hours after training).

Our results provide further evidence showing that memory is highly dynamic and influenced by other events occurring around and beyond the learning to be remembered³¹. In that sense, the strength of training and the type of -and time when- behavioural manipulations are experienced before or after acquisition, will determine the promoting or interfering effects on memory processing^{32–44}. A possible molecular mechanism underlying this effect is based on the behavioural tagging hypothesis which proposes that memory formation and persistence depend on synthesis of new proteins that will be used at specific substrates (tagged sites) in order to establish the memory trace³¹. On this regard, we have recently found that a weak IA training that generates a short-lasting LTM of a couple of days would create a maintenance-specific tag while proteins necessary for memory persistence would be provided by a close-in-time novel experience⁴⁴. This mechanism could explain interference between two or more memory traces. When the amount of proteins is insufficient for capture at multiple tagged sites, a competition for these resources would develop and, as a consequence, one of the memory traces might be impaired. Some studies showed that plasticity at specific synapses and memory formation for particular experiences are, at least in part, the result of competition for available resources^{37,43,45–48}. Therefore, we suggest that the deleterious effect of the exposure to a spatial novelty on IA-memory persistence is probably due to the “competition maintenance” between tagged sites for the available proteins induced by the two learning tasks. Recently it was showed that dopamine D1/D5 receptor regulates synaptic cooperation and competition in hippocampal neurons⁴⁹. The authors suggested that both, the modulating effect of dopamine on protein synthesis and the number of sites tagged by different stimuli, will interact so that the synapses will cooperate or compete during formation of stable memory traces. Moreover, the authors demonstrated that ERKs are involved in the effect of dopamine D1/D5 receptor agonist on synapse cooperation. However, there is less information regarding possible molecular substrates for the tag. In our study, we showed that CaMKs are involved in the competitive effect of spatial novelty over the persistence of IA memory. We speculate that inactivation of CaMKs may impair the setting of tagged sites induced by OF exposure.

Our finding is utterly promising because it opens a new avenue of research in the control of fear memories. It would be rather difficult to know when a traumatic event is going to take place in our daily life. However, targeting a treatment for fear-related memory disorder many hours after the aversive experience occurred would be a possible intervention to dampen memory. In conclusion, we show that it is possible to attenuate the durability of fear memories by acting on a permissive and restrictive late memory consolidation window (12 h post training). At that time point, and without the need administer any drug^{50,51} or to subject animals to a retrieval session⁵, the exploration to spatial novelty is sufficient to reduce a long-lasting fear memory.

Methods

Subjects. Experiments were conducted in male Wistar rats from the vivarium of the University of Buenos Aires (Buenos Aires, Argentina) weighting 230–260 g and 2–2.5 months old. Animals were housed five to a cage and kept at a constant temperature of 22 °C, with water and food *ad libitum*, under a 12 h light/dark cycle (lights on at 7:00 A.M.). Each animal was used only for one experiment. Experimental procedures followed the guidelines of the USA National Institutes of Health Guide for the Care and Use of Laboratory Animals and were approved by the Animal Care and Use Committees of the University Buenos Aires (CICUAL).

Inhibitory avoidance. Animals were handled once a day for two days and then trained in inhibitory avoidance (IA) as described previously⁷. Briefly the apparatus was a 50 × 25 × 25 cm opaque acrylic box whose floor was a grid made of 1 mm caliber stainless steel bars. The left end of the grid was covered by a 7 cm wide, 5.0 cm high platform. During the handling session animals were manipulated in the same way they were during intracerebral infusions (see below). Briefly, they were grasped by hand and slightly restrained in the lap or the arm of the investigator. During the second day of this manipulation in most animals there were no evident signs of stress. For training, animals were gently placed on the platform and, as they stepped down onto the grid, received a single 3 sec, 0.7 mA scrambled foot-shock. Latency for stepping down onto the grid with all four paws was measured. Rats were tested for retention either at 2 days or 7 days after training, depending on the experiment. The test session was similar in all respects to the training session except that the footshock was not given, and the latency was evaluated for a maximum of 300 seconds. All animals were tested only once. Training was always performed between 8:30–9:30 AM. For each experiment the number of animals in each group is detailed in the Results section.

Open Field. The open field (OF) was a 50 cm high, 50 cm wide, and 39 cm deep arena with black plywood walls and a brown floor divided into nine squares by black lines. The number of line crossings and rearings was measured manually during each minute, in a 5 min test session. The decrease of these parameters is considered an index of spatial habituation⁵².

Surgery. Sixty rats were bilaterally implanted under deep ketamine/xylazine anesthesia (80 and 5 mg/kg, respectively) with 22-g guide cannulae aimed to dorsal CA1 region of the hippocampus (AP −4.3 mm, LL ± 3.0 mm, DV 3 mm). Coordinates were based on Paxinos and Watson atlas⁵³. Cannulae were fixed to the skull with dental acrylic. Obturators were then inserted into the cannulae to prevent blockage, with the same or less

length of the cannulae. At the end of surgery, animals were injected with a single dose of meloxicam (0.2 mg/kg) as analgesic and gentamicin (2.5 mg/Kg) as antibiotic.

Drug infusions. After recovery from surgery (5–7 days), rats were trained in IA and 10.45 h after training received a bilateral infusion of either saline, or the CaMK inhibitor, KN-93 (6 µg/side)¹⁵. The volume infused was 0.8 µl/side and the infusion rate was 1 µl/min. Injectors were left in place for an additional minute following infusion before they were removed carefully to minimize backflow. Thus, the entire infusion procedure took ~4 min. For intracerebral infusions, 30-Gauge needles connected to Hamilton syringes were used. Infusions were delivered through a needle extending 1 mm beyond the tip of the guide cannula. During the procedure, the animals were slightly restrained with the hands, without provoking any evident stress.

Cannula placement. To check cannula placement, 24 h after the end of the behavioural procedures, animals were deeply anesthetized and killed by decapitation 15 min later, and histological localization of the infusion sites was established using a binocular magnifying glasses. Coordinates were based on Paxinos and Watson atlas⁵³. Schematic representation of rat brain sections showing the approximated extension of the area (gray) reached by the infusions of 0.8 µl of methylene blue in the CA1 region of the dorsal hippocampus is shown in Fig. 4a. Infusions spread about 1.5 mm³ and were found to be correct in 53 out of 60 animals. Cannula placements were exactly as in several previous papers^{7,9,44,47}.

Data analysis. In all behavioural experiments statistical analysis was performed by unpaired Student's t test or, when required, one-way ANOVA followed by Newman–Keuls multiple comparison test, comparing mean step-down latencies of the OF/drug-treated groups and control/vehicle-treated groups at each time point studied. Data in the bar graphs are presented as mean ± SEM.

References

- Maren, S. Seeking a spotless mind: extinction, deconsolidation, and erasure of fear memory. *Neuron* **70**, 830–845, doi: 10.1016/j.neuron.2011.04.023 (2011).
- Hermans, D., Craske, M. G., Mineka, S. & Lovibond, P. F. Extinction in human fear conditioning. *Biological psychiatry* **60**, 361–368, doi: 10.1016/j.biopsych.2005.10.006 (2006).
- Woods, A. M. & Bouton, M. E. Immediate extinction causes a less durable loss of performance than delayed extinction following either fear or appetitive conditioning. *Learning & memory* **15**, 909–920, doi: 10.1101/lm.1078508 (2008).
- Schiller, D. *et al.* Preventing the return of fear in humans using reconsolidation update mechanisms. *Nature* **463**, 49–53, doi: 10.1038/nature08637 (2010).
- Monfils, M. H., Cowansage, K. K., Klann, E. & LeDoux, J. E. Extinction-reconsolidation boundaries: key to persistent attenuation of fear memories. *Science* **324**, 951–955, doi: 10.1126/science.1167975 (2009).
- Chan, W. Y., Leung, H. T., Westbrook, R. F. & McNally, G. P. Effects of recent exposure to a conditioned stimulus on extinction of Pavlovian fear conditioning. *Learning & memory* **17**, 512–521, doi: 10.1101/lm.1912510 (2010).
- Bekinschtein, P. *et al.* Persistence of long-term memory storage requires a late protein synthesis- and BDNF- dependent phase in the hippocampus. *Neuron* **53**, 261–277, doi: 10.1016/j.neuron.2006.11.025 (2007).
- Bekinschtein, P. *et al.* BDNF is essential to promote persistence of long-term memory storage. *Proc Natl Acad Sci USA* **105**, 2711–2716 (2008).
- Rossato, J. I., Bevilacqua, L. R., Izquierdo, I., Medina, J. H. & Cammarota, M. Dopamine controls persistence of long-term memory storage. *Science* **325**, 1017–1020, doi: 10.1126/science.1172545 (2009).
- Netto, C. A., Dias, R. D. & Izquierdo, I. Interaction between consecutive learnings: inhibitory avoidance and habituation. *Behavioral and neural biology* **44**, 515–520 (1985).
- Izquierdo, I. & Medina, J. H. Memory formation: the sequence of biochemical events in the hippocampus and its connection to activity in other brain structures. *Neurobiology of learning and memory* **68**, 285–316, doi: 10.1006/nlme.1997.3799 (1997).
- Izquierdo, I. & McGaugh, J. L. Effect of novel experiences on retention of inhibitory avoidance behavior in mice: the influence of previous exposure to the same or another experience. *Behavioral and neural biology* **47**, 109–115 (1987).
- Wolfman, C., Izquierdo, L. A., Schroder, N. & Izquierdo, I. Intra-hippocampal KN-62 hinders the memory of habituation acquired alone, but not simultaneously with a water-finding task. *Behavioural pharmacology* **10**, 99–104 (1999).
- Vianna, M. R. *et al.* Role of hippocampal signaling pathways in long-term memory formation of a nonassociative learning task in the rat. *Learning & memory* **7**, 333–340 (2000).
- Bevilacqua, L. R., Medina, J. H., Izquierdo, I. & Cammarota, M. Memory consolidation induces N-methyl-D-aspartic acid-receptor- and Ca²⁺/calmodulin-dependent protein kinase II-dependent modifications in alpha-amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptor properties. *Neuroscience* **136**, 397–403, doi: 10.1016/j.neuroscience.2005.08.007 (2005).
- Greenberg, P. E. *et al.* The economic burden of anxiety disorders in the 1990s. *The Journal of clinical psychiatry* **60**, 427–435 (1999).
- Gillespie, C. F. & Ressler, K. J. Emotional learning and glutamate: translational perspectives. *CNS spectrums* **10**, 831–839 (2005).
- Holbrook, T. L., Galarneau, M. R., Dye, J. L., Quinn, K. & Dougherty, A. L. Morphine use after combat injury in Iraq and post-traumatic stress disorder. *The New England journal of medicine* **362**, 110–117, doi: 10.1056/NEJMoa0903326 (2010).
- Parsons, R. G. & Ressler, K. J. Implications of memory modulation for post-traumatic stress and fear disorders. *Nature neuroscience* **16**, 146–153, doi: 10.1038/nn.3296 (2013).
- Myers, K. M., Ressler, K. J. & Davis, M. Different mechanisms of fear extinction dependent on length of time since fear acquisition. *Learning & memory* **13**, 216–223, doi: 10.1101/lm.119806 (2006).
- Maren, S. & Chang, C. H. Recent fear is resistant to extinction. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 18020–18025, doi: 10.1073/pnas.0608398103 (2006).
- Milekic, M. H. & Alberini, C. M. Temporally graded requirement for protein synthesis following memory reactivation. *Neuron* **36**, 521–525 (2002).
- Suzuki, A. *et al.* Memory reconsolidation and extinction have distinct temporal and biochemical signatures. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **24**, 4787–4795, doi: 10.1523/JNEUROSCI.5491-03.2004 (2004).
- Lattal, K. M. & Abel, T. Behavioral impairments caused by injections of the protein synthesis inhibitor anisomycin after contextual retrieval reverse with time. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 4667–4672, doi: 10.1073/pnas.0306546101 (2004).
- Prado-Alcala, R. A. *et al.* Amygdala or hippocampus inactivation after retrieval induces temporary memory deficit. *Neurobiol Learn Mem* **86**, 144–149, doi: 10.1016/j.nlm.2006.01.006 (2006).

26. Wixted, J. T. On Common Ground: Jost's (1897) law of forgetting and Ribot's (1881) law of retrograde amnesia. *Psychological review* **111**, 864–879, doi: 10.1037/0033-295X.111.4.864 (2004).
27. Dewar, M. T., Cowan, N. & Sala, S. D. Forgetting due to retroactive interference: a fusion of Muller and Pilzecker's (1900) early insights into everyday forgetting and recent research on anterograde amnesia. *Cortex; a journal devoted to the study of the nervous system and behavior* **43**, 616–634 (2007).
28. Redondo, R. L. *et al.* Synaptic tagging and capture: differential role of distinct calcium/calmodulin kinases in protein synthesis-dependent long-term potentiation. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **30**, 4981–4989, doi: 10.1523/JNEUROSCI.3140-09.2010 (2010).
29. Yang, C. *et al.* Stress within a restricted time window selectively affects the persistence of long-term memory. *PLoS one* **8**, e59075, doi: 10.1371/journal.pone.0059075 (2013).
30. Schmidt, M. V., Abraham, W. C., Maroun, M., Stork, O. & Richter-Levin, G. Stress-induced metaplasticity: from synapses to behavior. *Neuroscience* **250**, 112–120, doi: 10.1016/j.neuroscience.2013.06.059 (2013).
31. Moncada, D., Ballarini, F. & Viola, H. Behavioral Tagging: A Translation of the Synaptic Tagging and Capture Hypothesis. *Neural plasticity* **2015**, 650780, doi: 10.1155/2015/650780 (2015).
32. Moncada, D. & Viola, H. Induction of long-term memory by exposure to novelty requires protein synthesis: evidence for a behavioral tagging. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **27**, 7476–7481, doi: 10.1523/JNEUROSCI.1083-07.2007 (2007).
33. Ballarini, F., Moncada, D., Martinez, M. C., Alen, N. & Viola, H. Behavioral tagging is a general mechanism of long-term memory formation. *Proc Natl Acad Sci USA* **106**, 14599–14604, doi: 10.1073/pnas.0907078106 (2009).
34. Wang, S. H., Redondo, R. L. & Morris, R. G. Relevance of synaptic tagging and capture to the persistence of long-term potentiation and everyday spatial memory. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 19537–19542, doi: 10.1073/pnas.1008638107 (2010).
35. Lu, Y. *et al.* TrkB as a potential synaptic and behavioral tag. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **31**, 11762–11771, doi: 10.1523/JNEUROSCI.2707-11.2011 (2011).
36. Moncada, D., Ballarini, F., Martinez, M. C., Frey, J. U. & Viola, H. Identification of transmitter systems and learning tag molecules involved in behavioral tagging during memory formation. *Proc Natl Acad Sci USA* **108**, 12931–12936, doi: 10.1073/pnas.1104495108 (2011).
37. Redondo, R. L. & Morris, R. G. Making memories last: the synaptic tagging and capture hypothesis. *Nature reviews. Neuroscience* **12**, 17–30, doi: 10.1038/nrn2963 (2011).
38. Almaguer-Melian, W. *et al.* Novelty exposure overcomes foot shock-induced spatial-memory impairment by processes of synaptic-tagging in rats. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 953–958, doi: 10.1073/pnas.1114198109 (2012).
39. Dong, Z. *et al.* Mechanisms of hippocampal long-term depression are required for memory enhancement by novelty exploration. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **32**, 11980–11990, doi: 10.1523/JNEUROSCI.0984-12.2012 (2012).
40. Cassini, L. F. *et al.* Memory reconsolidation allows the consolidation of a concomitant weak learning through a synaptic tagging and capture mechanism. *Hippocampus* **23**, 931–941, doi: 10.1002/hipo.22149 (2013).
41. de Carvalho Myskiw, J., Furini, C. R., Benetti, F. & Izquierdo, I. Hippocampal molecular mechanisms involved in the enhancement of fear extinction caused by exposure to novelty. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 4572–4577, doi: 10.1073/pnas.1400423111 (2014).
42. Salvetti, B., Morris, R. G. & Wang, S. H. The role of rewarding and novel events in facilitating memory persistence in a separate spatial memory task. *Learning & memory* **21**, 61–72, doi: 10.1101/lm.032177.113 (2014).
43. Martinez, M. C., Villar, M. E., Ballarini, F. & Viola, H. Retroactive interference of object-in-context long-term memory: role of dorsal hippocampus and medial prefrontal cortex. *Hippocampus* **24**, 1482–1492, doi: 10.1002/hipo.22328 (2014).
44. Tomaiuolo, M., Kathe, C., Viola, H. & Medina, J. H. Evidence of Maintenance Tagging in the Hippocampus for the Persistence of Long-Lasting Memory Storage. *Neural plasticity* **2015**, 603672, doi: 10.1155/2015/603672 (2015).
45. Fonseca, R., Nagerl, U. V., Morris, R. G. & Bonhoeffer, T. Competing for memory: hippocampal LTP under regimes of reduced protein synthesis. *Neuron* **44**, 1011–1020, doi: 10.1016/j.neuron.2004.10.033 (2004).
46. Govindarajan, A., Israely, I., Huang, S. Y. & Tonegawa, S. The dendritic branch is the preferred integrative unit for protein synthesis-dependent LTP. *Neuron* **69**, 132–146, doi: 10.1016/j.neuron.2010.12.008 (2011).
47. Martinez, M. C., Alen, N., Ballarini, F., Moncada, D. & Viola, H. Memory traces compete under regimes of limited Arc protein synthesis: implications for memory interference. *Neurobiology of learning and memory* **98**, 165–173, doi: 10.1016/j.nlm.2012.05.007 (2012).
48. Sajikumar, S., Morris, R. G. & Korte, M. Competition between recently potentiated synaptic inputs reveals a winner-take-all phase of synaptic tagging and capture. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 12217–12221, doi: 10.1073/pnas.1403643111 (2014).
49. Shivarama Shetty, M., Gopinadhan, S. & Sajikumar, S. Dopamine D1/D5 receptor signaling regulates synaptic cooperation and competition in hippocampal CA1 pyramidal neurons via sustained ERK1/2 activation. *Hippocampus* **26**, 137–150, doi: 10.1002/hipo.22497 (2016).
50. Slipczuk, L. *et al.* Attenuating the persistence of fear memory storage using a single dose of antidepressant. *Molecular psychiatry* **18**, 7–8, doi: 10.1038/mp.2012.4 (2013).
51. Giustino, T. F., Fitzgerald, P. J. & Maren, S. Revisiting propranolol and PTSD: Memory erasure or extinction enhancement? *Neurobiology of learning and memory* **130**, 26–33, doi: 10.1016/j.nlm.2016.01.009 (2016).
52. Winograd, M. & Viola, H. Detection of novelty, but not memory of spatial habituation, is associated with an increase in phosphorylated cAMP response element-binding protein levels in the hippocampus. *Hippocampus* **14**, 117–123, doi: 10.1002/hipo.10153 (2004).
53. Paxinos, G. & Watson, C. Vol. 1 33–37 (Elsevier Ltd, 1997).

Acknowledgements

This work was supported by grants of the National Agency of Scientific and Technological Promotion of Argentina (ANPCyT, Argentina), University of Buenos Aires, (UBACyT, Argentina) and the Argentina National Research Council (CONICET).

Author Contributions

C.K., M.T., J.H.M. and H.V. designed research; C.K., M.T. and G.D. performed research; M.T. and C.K. analyzed data; and J.H.M. and H.V. wrote the paper. The authors declare no conflict of interest.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Katche, C. *et al.* Novelty during a late postacquisition time window attenuates the persistence of fear memory. *Sci. Rep.* **6**, 35220; doi: 10.1038/srep35220 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016