

SCIENTIFIC REPORTS



OPEN

Fold formation at the compartment boundary of *Drosophila* wing requires Yki signaling to suppress JNK dependent apoptosis

Received: 05 September 2016

Accepted: 02 November 2016

Published: 29 November 2016

Suning Liu¹, Jie Sun¹, Dan Wang¹, Gert O. Pflugfelder² & Jie Shen¹

Compartment boundaries prevent cell populations of different lineage from intermingling. In many cases, compartment boundaries are associated with morphological folds. However, in the *Drosophila* wing imaginal disc, fold formation at the anterior/posterior (A/P) compartment boundary is suppressed, probably as a prerequisite for the formation of a flat wing surface. Fold suppression depends on *optomotor-blind* (*omb*). *Omb* mutant animals develop a deep apical fold at the A/P boundary of the larval wing disc and an A/P cleft in the adult wing. A/P fold formation is controlled by different signaling pathways. Jun N-terminal kinase (JNK) and Yorkie (Yki) signaling are activated in cells along the fold and are necessary for the A/P fold to develop. While JNK promotes cell shape changes and cell death, Yki target genes are required to antagonize apoptosis, explaining why both pathways need to be active for the formation of a stable fold.

Epithelial folds contribute to morphogenetic movements and the separation of different cell groups, thus shaping the animal body¹. For example, invagination of the *Drosophila* mesoderm is initiated by ventral fold formation^{2,3}. Segmental and parasegmental grooves transiently appear in the trunk region of the *Drosophila* embryonic epithelium separating fields of cells of different fate^{4,5}. A common but not the only mechanism of epithelial fold formation involves apical cell constriction and the acquisition of a bottle-like cell morphology⁶. By this mechanism tubes can be formed from epithelial sheets as in the development of tracheal and salivary gland primordia⁷. Folds also arise in postembryonic epithelia. In the *Drosophila* eye disc, differentiation depends on the progression of the morphogenetic furrow, a Hedgehog-dependent apical indentation of the eye field^{8,9}. Apical and basal folds can also form at the borders which separate cell groups of different fate in the other imaginal discs of *Drosophila* larvae^{10–12}. Folds can arise in the epithelia of all metazoa. In sea urchins, bottle cells have been shown to be required for invagination of the ectoderm¹³. In vertebrates, classical examples include the formation of the neural tube in chick¹⁴ and of the blastopore lip in amphibians^{15,16}. Neuroectodermal grooves are also found during brain development in mouse and zebrafish^{17,18}.

As outlined above, folds occur in many aspects of *Drosophila* development, making this species an excellent model organism to study mechanisms of fold formation. *Drosophila* has provided information on the molecular underpinnings of the required cell shape changes in various developmental paradigms. The best studied system is gastrulation which was investigated at the levels of genetics, cell biology and biophysics and which, therefore, can serve as a benchmark for studies in other systems^{19–22}. In *Drosophila* gastrulation, the secreted protein Folded gastrulation (Fog) is central to inducing apical constriction of the invaginating cells^{23,24} but is dispensable in several other epithelial folding processes. In the *Drosophila* embryonic ectoderm, the formation of segmental grooves was shown to be controlled by *engrailed* (*en*) expression in boundary cells as well as by Hedgehog (Hh) and Wingless (Wg) signaling⁴. These segment polarity genes are not involved in specifying the position of nearby parasegmental grooves; Wg signaling is required, however, in a non-instructive, permissive role⁵. Formation and progression of the morphogenetic furrow in the larval eye disc is controlled by Hh and Decapentaplegic (Dpp). Induction of the cell adhesion molecule Cad86C by Hh and Dpp may be one of the mechanisms that effects cell shape changes in this tissue²⁵. The way folds form, thus, can differ with regard to molecular and biomechanical

¹Department of Entomology, China Agricultural University, 100193 Beijing, China. ²Institute of Genetics, Johannes Gutenberg-University, 55128 Mainz, Germany. Correspondence and requests for materials should be addressed to J.S. (email: shenjie@cau.edu.cn)

requirements even within one epithelium. For example, in *Drosophila* gastrulation, the ventral furrow forms by apical constriction whereas the dorsal folds arise by a basal shift of the adherens junctions²⁶.

In the larval wing disc pouch (the future wing blade), the normal, graded expression of Dpp and Wg does not instruct folding but rather is involved in maintaining the appropriate position-specific cell shape²⁷. In the columnar main epithelium, loss of Dpp signaling causes extrusion of cells correlated with loss of the apical microtubule web^{28,29}. Similarly, loss of Dpp targets Optomotor-blind (Omb) or Spalt lead to retraction of cells toward basal membrane^{30,31}. Dpp signaling cell-autonomously promotes and maintains the elongated columnar shape of wing disc cells by regulating Rho1 and the regulatory light chain of non-muscle myosin II³². Wg signaling cell-autonomously promotes and maintains the columnar shape of wing disc cells through maintaining Vestigial (Vg) expression³³. The Wg gradient, centered on the D/V boundary, instructs similarly shaped gradients of DE-cadherin concentration and apical cell circumference (high and constricted, respectively, close to the D/V boundary)³⁴. The loss of Adenomatous polyposis coli (APC), a negative regulator of Wg signaling, leads to apical constriction and invagination independent of its effect on the DE-cadherin level. Wg, too, acts by activation of Rho1 and Myosin II³⁵.

The folds which separate parasegments in the *Drosophila* embryonic ectoderm separate fields of cells that are related by lineage (compartments)³⁶. But even in the absence of lineage restriction, groups of epithelial cells differing in gene expression and fated to develop into different structures tend to be separated by a fold. For instance, in the wing imaginal disc, which gives rise to adult notum, hinge, and wing blade, several folds orthogonal to the proximo-distal axis separate gene expression domains without being lineage restriction boundaries^{37,38}. The most distal of these, the blade/hinge fold develops under control of the Omb-related T-box transcription factors Dorsocross (Doc)³⁹. The proximal notum/hinge fold requires the complementary expression of Omb in hinge and Iroquois complex (Iro-C) in notum⁴⁰. In contrast, the A/P compartment boundary is not associated with a fold and remains morphologically inconspicuous throughout development⁴¹, even though it derives from the corresponding infolded parasegmental boundary in the embryonic ectoderm⁴². It is conceivable that fold formation was selected against because of the structural requirement for the adult wing as a flight appendage. Indeed, fold formation is actively suppressed by a genetic program. Omb which is expressed in most of the pouch⁴³ is required to maintain the normal epithelial structure at the A/P boundary. Reduction of Omb level in the pouch causes an apical morphogenetic defect at the A/P boundary due to contraction of cells along their apical-basal axis⁴⁴. We here investigate the mechanisms of boundary fold formation elicited by Omb loss. We found that A/P fold formation is dependent on activation of JNK signaling induced by loss of *omb*. Loss of *omb* also induced Yki activity which promoted cell survival and attenuated the pro-apoptotic activity of JNK. Our results reveal a network of signaling pathways induced by loss of *omb* that controls cell shape and ensures cell survival of folded cells at the A/P boundary.

Results and Discussion

We and others have shown before that Omb prevents aberrant apical fold formation at the A/P boundary^{44,45}. When Omb is directly or indirectly repressed in the P compartment, the A/P boundary of the wing develops a deep apical fold in the larval wing disc and a cleft in the adult wing⁴⁴. We here use this model system to investigate the mechanisms of boundary fold formation under three genetic manipulations, *ptc*-Gal4 UAS-*tkv*, *en*-Gal4 UAS-*omb*-RNAi, and *nub*-Gal4 UAS-*omb*-RNAi (Fig. S1).

JNK is ectopically activated to initiate A/P fold formation. To investigate potential effectors downstream of Omb, we monitored the expression or activity of candidate targets. JNK signaling is an important pathway in the regulation of wing disc morphogenesis^{46,47}. We monitored JNK pathway activity by monitoring transcription of the JNK target gene *puckered* (*puc*)⁴⁸. In both *ptc*>*tkv* and *en*>*omb*-RNAi wing discs, *puc* was activated along the A/P fold (Fig. 1A,B). *puc* was initially activated in cells adjacent to the fold in early-mid L3, then its expression extended further into the A and P compartments (Fig. S2). These data indicate that JNK signaling is activated in the process of A/P fold formation.

We next asked whether JNK activation is required for A/P fold generation. Co-expressing *omb*-RNAi and a dominant negative form of JNK, *bsk*^{DN}⁴⁹, was sufficient to suppress A/P fold formation (Fig. 1C).

This suggests that activation of JNK signaling is required in this process. To test for sufficiency of JNK activation for A/P fold formation, we activated JNK by expressing *hep*^{CA} (encoding a constitutively active form of JNKK⁵⁰) for a short duration controlled by *dpp*-Gal4 and *tub*-Gal80^{ts} (continuous activation of *hep*^{CA} induced severe apoptosis thereby disturbing observation of cell morphology). Under these conditions folds occurred throughout the *dpp*-Gal4 expression domain (Fig. 1D). The broad anterior activation of JNK signaling did not lead, however, to a discrete A/P fold.

The matrix metalloproteinase 1 (*Mmp1*) is induced by ectopic activation of JNK during morphological reorganization of epithelia^{51–55}. When the JNK pathway was activated for 24 h in the *dpp*-Gal4 expression domain, *Mmp1* was broadly induced anterior to the A/P boundary, similar to the plexus of epithelial folds observed under these conditions (Fig. 1E). However, in *en*>*omb*-RNAi wing discs, *Mmp1* accumulated in a discrete stripe of A/P fold cells on both sides of the fold (Fig. 1F and F'). Co-expression of *omb*-RNAi and *Mmp1*-RNAi with *nub*-Gal4 rescued the A/P fold with full penetrance (Fig. 1G). Uniform reduction of *omb* expression on both sides of the A/P boundary, like posterior *omb* reduction, leads to A/P fold formation (Fig. S1E and F).

However, expression of *Mmp1* with *dpp*-Gal4 along the A/P boundary did not generate a fold (Fig. S3). These data suggests that either additional gene expression changes, induced by the loss of *omb*, are necessary for A/P fold formation or that *Mmp1* must be induced on both sides of the A/P boundary.

Yki-Diap1 signaling is activated parallel to the JNK pathway. It has been reported that Dpp and Wg repress JNK signaling to maintain survival of wing pouch cells. Loss of either Dpp or Wg signaling activates




Figure 1. JNK signaling is necessary for A/P fold formation. (A,B) *puc-lacZ* was ectopically activated at the A/P fold. (C) Repression of JNK signaling by co-expressing a dominant negative form of JNK (*bsk^{DN}*) suppressed the A/P fold formation. Anti-Omb staining (green) demonstrates the efficiency of the posterior knock-down. Activation of JNK signaling by *hep^{CA}* for a short duration induced extended folding (D) and *Mmp1* expression (E) in the *dpp-Gal4* domain. (F,F') Focused induction of *Mmp1* expression symmetrically on both sides of the A/P fold in *en>omb-RNAi* wing disc. (G) Co-expressing *Mmp1-RNAi* suppressed A/P fold formation. Arrowheads point at the position of the A/P boundary.

JNK-dependent apoptosis in the wing pouch⁵⁰. However, *omb* knock-down did not induce apparent apoptosis at the A/P boundary⁴⁴, although JNK signaling was activated (Fig. 1A and B). We assume that the apoptosis pathway is repressed in this case. Yki signaling can be induced by the JNK pathway⁵⁶. Yki targets such as *Death-associated inhibitor of apoptosis 1 (Diap1)* and the microRNA *bantam (ban)* can repress apoptosis^{57–59}. We analyzed transcription of the Yki target *expanded (ex⁶⁰)* and observed that *ex-lacZ* was up-regulated at the A/P fold generated by *en>omb-RNAi* (Fig. 2B), suggesting an activation of Yki signaling during A/P fold formation. This was confirmed in *nub>omb-RNAi* wing discs (Fig. 2C). Upregulation at the A/P fold was also observed for *Diap1* (Fig. 2E). The Yki target *ban* is suppressed by Omb in the medial wing discs⁶¹. Consistently, *ban* was up-regulated in the medial wing disc of *nub>omb-RNAi* larvae, with the strongest enhancement along the A/P boundary (Fig. 2G). Therefore, during A/P fold generation, *ban* and *Diap1* were both activated and could suppress potential apoptosis induced by changes in cell shape and JNK activation.

In order to determine whether the Yki targets were induced as a consequence of JNK signaling, we co-expressed *bsk^{DN}* and *omb-RNAi* in the *nub-Gal4* domain. When JNK pathway and A/P fold were suppressed, *ex-lacZ* expression was still activated at the A/P boundary (Fig. S4A). This also held for *Diap1* and *ban* expression (Fig. S4B and C). This suggests that in A/P fold formation Yki can be activated even when the JNK pathway is blocked. Yki activation, thus, appears to occur parallel to JNK signaling and is not sufficient for fold formation.

To test whether the suppression of cell death by Yki signaling is required for *omb*-loss induced A/P fold formation, *yki-RNAi* was co-expressed with *omb-RNAi* in the *nub-Gal4* domain. As shown in Fig. 3A–A', co-expressing *yki-RNAi* was sufficient to suppress the formation of A/P fold. Severe cell death occurred in this double knock-down. When p35 was co-expressed with *omb-RNAi* and *yki-RNAi* in the *nub-Gal4* domain to inhibit apoptosis, cell death was effectively suppressed (Fig. 3B and B') and the A/P fold appeared again (Fig. 3B and C). These data indicate that Yki signaling is required for A/P fold by ensuring cell survival.

Generally, abnormal activation of the JNK pathway induces apoptosis. For instance, expression of activated tumor genes or mutation in tumor suppressor genes lead to JNK-induced cell invasion and apoptosis^{55,62–64}. However, in *omb*-knocked-down wing discs, the activation of JNK pathway did not cause cell death along the A/P




Figure 2. Activation of Yki target genes in *omb* hypomorphic wing discs. (A) Control experiment to show the normal *ex-lacZ* expression. (B,C) *ex-lacZ* was upregulated along the A/P fold in discs in which *omb* was knocked down in the posterior compartment (B) or in the entire pouch (C). (D) Control experiment to show the normal *Diap1-lacZ* expression. (E) *Diap1-lacZ* was upregulated along the A/P fold. (F) Control experiment to show the normal *ban-lacZ* expression. (G) *ban-lacZ* was ectopically activated at the A/P fold.

fold⁴⁴. We suggest that cell death is suppressed by the simultaneous induction of a cell survival pathway. Yki has an important role in promoting cell survival by driving the expression of downstream genes such as *Diap1* and *ban*⁶⁵. We found these genes upregulated along the A/P fold (Fig. 2). This suggests that Yki antagonizes apoptosis along the fold. Previous studies identified JNK as a promoter of Yki activity in the wing disc^{56,63,66}. But this regulatory relationship is not absolute^{62,63,67}. We found that co-expression of *omb*-RNAi and *bsk*^{DN} had no effect on *ex*, *Diap1*, and *ban* expression (Fig. S4). This indicates that, at the A/B boundary, Yki is activated parallel to JNK signaling.

Methods

Drosophila stocks. The transgenes used were as follows: *en-Gal4*, *nub-Gal4*⁶⁸, *dpp-Gal4*, *ptc-Gal4*, *UAS-tkv*, *UAS-CD8-GFP*, *UAS-GFP*, *UAS-omb-RNAi*⁴⁴, *tubP-Gal80^{ts}*⁶⁹, *UAS-MMP1*⁷⁰, *UAS-MMP1-RNAi*⁵⁵, *UAS-p35*, *UAS-hep^{CA}*, *UAS-bsk^{DN}*, *UAS-yki-RNAi* (TsingHua Fly Center). Enhancer trap lines were *hh-lacZ*⁷¹, *puc-lacZ*, *ex-lacZ*, *diap1-lacZ*⁶⁰, and *ban-lacZ*⁷². Stocks, if not mentioned otherwise, were obtained from the Bloomington stock center.




Figure 3. Suppression of apoptosis is required for fold formation. (A–A'') Co-expressing omb-RNAi and yki-RNAi by nub-Gal4 suppressed the A/P fold formation but induced severe apoptosis. (B,B') When apoptosis was suppressed by co-expressing p35 an A/P fold was generated. (C) The fold generated in nub>[omb-RNAi + yki-RNAi + p35] wing discs extended along the A/P boundary (anterior compartment marked by Cubitus interruptus (Ci) expression, green).

Larvae were raised at 25 °C. For efficient expression of RNAi transgenes, larvae were raised at 29 °C. Larvae containing Gal80^{ts}-Gal4 combinations were raised at 18 °C and then were shifted to 29 °C that allows GAL4 to function and activate transcription of UAS controlled transgenes.

Immunohistochemistry. Dissected wing imaginal discs were fixed and stained with antibodies according to the standard procedures. The primary antibodies used were: rabbit anti-Omb, 1:1000; mouse and rabbit anti-beta-galactosidase, 1:2000 (Promega); rabbit anti-caspase3, 1:200 (Santa Cruz), rat anti-2A1 (Ci), 1:200 (Developmental Studies Hybridoma Bank, DSHB); and mouse anti-Mmp1, 1:200 (DSHB). Secondary antibodies used were goat anti-rabbit DyLight 488, goat anti-rat DyLight 488, goat anti-rabbit DyLight 549, goat anti-mouse DyLight 488, goat anti-rabbit Cy3, and goat anti-rabbit Cy5, were diluted 1:200 (Agrisera). Actin was visualized with Rhodamine-phalloidin, 1:1000 (Cytoskeleton). Images were collected using a Leica TCS SP2 AOBS confocal microscope.

References

- Pilot, F. & Lecuit, T. Compartmentalized morphogenesis in epithelia: from cell to tissue shape. *Dev. Dyn* **232**, 685–694 (2005).
- Oda, H. & Tsukita, S. Real-time imaging of cell-cell adherens junctions reveals that *Drosophila* mesoderm invagination begins with two phases of apical constriction of cells. *J. Cell Sci* **114**, 493–501 (2001).
- Kam, Z., Minden, J. S., Agard, D. A., Sedat, J. W. & Leptin, M. *Drosophila* gastrulation: analysis of cell shape changes in living embryos by three-dimensional fluorescence microscopy. *Development* **112**, 365–370 (1991).
- Larsen, C. W., Hirst, E., Alexandre, C. & Vincent, J. P. Segment boundary formation in *Drosophila* embryos. *Development* **130**, 5625–5635 (2003).
- Larsen, C., Bardet, P. L., Vincent, J. P. & Alexandre, C. Specification and positioning of parasegment grooves in *Drosophila*. *Dev. Biol* **321**, 310–318 (2008).
- Sawyer, J. M. *et al.* Apical constriction: a cell shape change that can drive morphogenesis. *Dev. Biol* **341**, 5–19 (2010).
- Maruyama, R. & Andrew, D. J. *Drosophila* as a model for epithelial tube formation. *Dev. Dyn* **241**, 119–135 (2012).
- Corrigan, D., Walther, R. F., Rodriguez, L., Fichelson, P. & Pichaud, F. Hedgehog signaling is a principal inducer of Myosin-II-driven cell ingression in *Drosophila* epithelia. *Dev. Cell* **13**, 730–742 (2007).
- Wolff, T. & Ready, D. F. The beginning of pattern formation in the *Drosophila* compound eye: the morphogenetic furrow and the second mitotic wave. *Development* **113**, 841–850 (1991).
- Dominguez, M. & Casares, F. Organ specification-growth control connection: new in-sights from the *Drosophila* eye-antennal disc. *Dev. Dyn* **232**, 673–684 (2005).
- Greenberg, L. & Hatini, V. Essential roles for lines in mediating leg and antennal proximodistal patterning and generating a stable Notch signaling interface at segment borders. *Dev. Biol* **330**, 93–104 (2009).
- Zirin, J. D. & Mann, R. S. Nubbin and Teashirt mark barriers to clonal growth along the proximal-distal axis of the *Drosophila* wing. *Dev. Biol* **304**, 745–758 (2007).
- Kimberly, E. L. & Hardin, J. Bottle cells are required for the initiation of primary invagination in the sea urchin embryo. *Dev. Biol* **204**, 235–250 (1998).
- Schoenwolf, G. C. & Franks, M. V. Quantitative analyses of changes in cell shapes during bending of the avian neural plate. *Dev. Biol* **105**, 257–272 (1984).
- Hardin, J. & Keller, R. The behaviour and function of bottle cells during gastrulation of *Xenopus laevis*. *Development* **103**, 211–230 (1988).

16. Lee, J. Y. & Harland, R. M. Endocytosis is required for efficient apical constriction during *Xenopus* gastrulation. *Curr. Biol* **20**, 253–258 (2010).
17. Inoue, T. *et al.* Role of cadherins in maintaining the compartment boundary between the cortex and striatum during development. *Development* **128**, 561–569 (2001).
18. Langenberg, T. & Brand, M. Lineage restriction maintains a stable organizer cell population at the zebrafish midbrain-hindbrain boundary. *Development* **132**, 3209–3216 (2005).
19. Dawes-Hoang, R. E. *et al.* Folded gastrulation, cell shape change and the control of myosin localization. *Development* **132**, 4165–4178 (2005).
20. Fox, D. T. & Peifer, M. Abelson kinase (Abl) and RhoGEF2 regulate actin organization during cell constriction in *Drosophila*. *Development* **134**, 567–578 (2007).
21. Kolsch, V., Seher, T., Fernandez-Ballester, G. J., Serrano, L. & Leptin, M. Control of *Drosophila* gastrulation by apical localization of adherens junctions and RhoGEF2. *Science* **315**, 384–386 (2007).
22. Polyakov, O. *et al.* Passive mechanical forces control cell-shape change during *Drosophila* ventral furrow formation. *Biophys. J* **107**, 998–1010 (2014).
23. Costa, M., Wilson, E. T. & Wieschaus, E. A putative cell signal encoded by the folded gastrulation gene coordinates cell shape changes during *Drosophila* gastrulation. *Cell* **76**, 1075–1089 (1994).
24. Fuse, N., Yu, F. & Hirose, S. Gprk2 adjusts Fog signaling to organize cell movements in *Drosophila* gastrulation. *Development* **140**, 4246–4255 (2013).
25. Schlichting, K. & Dahmann, C. Hedgehog and Dpp signaling induce cadherin Cad86C expression in the morphogenetic furrow during *Drosophila* eye development. *Mech. Dev* **125**, 712–728 (2008).
26. Wang, Y. C., Khan, Z., Kaschube, M. & Wieschaus, E. F. Differential positioning of adherens junctions is associated with initiation of epithelial folding. *Nature* **484**, 390–393 (2012).
27. McClure, K. D. & Schubiger, G. Developmental analysis and squamous morphogenesis of the peripodial epithelium in *Drosophila* imaginal discs. *Development* **132**, 5033–5042 (2005).
28. Gibson, M. C. & Perrimon, N. Extrusion and death of DPP/BMP-compromised epithelial cells in the developing *Drosophila* wing. *Science* **307**, 1785–1789 (2005).
29. Shen, J. & Dahmann, C. Extrusion of cells with inappropriate Dpp signaling from *Drosophila* wing disc epithelia. *Science* **307**, 1789–1790 (2005).
30. Shen, J., Dahmann, C. & Pflugfelder, G. O. Spatial discontinuity of optomotor-blind expression in the *Drosophila* wing imaginal disc disrupts epithelial architecture and promotes cell sorting. *BMC Dev. Biol* **10**, 23 (2010).
31. Tang, W., Wang, D. & Shen, J. Asymmetric distribution of Spalt in *Drosophila* wing squamous and columnar epithelia ensures correct cell morphogenesis. *Sci. Rep* **6**, 30236 (2016).
32. Widmann, T. J. & Dahmann, C. Dpp signaling promotes the cuboidal-to-columnar shape transition of *Drosophila* wing disc epithelia by regulating Rho1. *J. Cell Sci* **122**, 1362–1373 (2009a).
33. Widmann, T. J. & Dahmann, C. Wingless signaling and the control of cell shape in *Drosophila* wing imaginal discs. *Dev. Biol* **334**, 161–173 (2009b).
34. Jaiswal, M., Agrawal, N. & Sinha, P. Fat and Wingless signaling oppositely regulate epithelial cell-cell adhesion and distal wing development in *Drosophila*. *Development* **133**, 925–935 (2006).
35. Zimmerman, S. G., Thorpe, L. M., Medrano, V. R., Mallozzi, C. A. & McCartney, B. M. Apical constriction and invagination downstream of the canonical Wnt signaling pathway require Rho1 and Myosin II. *Dev. Biol* **340**, 54–66 (2010).
36. Blair, S. S. Lineage compartments in *Drosophila*. *Curr. Biol* **13**, R548–551 (2003).
37. Villa-Cuesta, E., Gonzalez-Perez, E. & Modolell, J. Apposition of iroquois expressing and non-expressing cells leads to cell sorting and fold formation in the *Drosophila* imaginal wing disc. *BMC Dev. Biol* **7**, 106 (2007).
38. Villa-Cuesta, E. & Modolell, J. Mutual repression between msh and Iro-C is an essential component of the boundary between body wall and wing in *Drosophila*. *Development* **132**, 4087–4096 (2005).
39. Sui, L., Pflugfelder, G. O. & Shen, J. The Dorsocross T-box transcription factors promote tissue morphogenesis in the *Drosophila* wing imaginal disc. *Development* **139**, 2773–2782 (2012).
40. Wang, D., Li, L., Lu, J., Liu, S. & Shen, J. Complementary expression of optomotor-blind and the Iroquois complex promotes fold formation to separate wing notum and hinge territories. *Dev. Biol* **416**, 225–234 (2016).
41. Brower, D. L., Smith, R. J. & Wilcox, M. Cell shapes on the surface of the *Drosophila* wing imaginal disc. *J. Embryol. Exp. Morphol* **67**, 137–151 (1982).
42. Vincent, J. P. Compartment boundaries: where, why and how? *Int. J. Dev. Biol* **42**, 311–315 (1998).
43. Grimm, S. & Pflugfelder, G. O. Control of the gene optomotor-blind in *Drosophila* wing development by decapentaplegic and wingless. *Science* **271**, 1601–1604 (1996).
44. Shen, J., Dorner, C., Bahlo, A. & Pflugfelder, G. O. optomotor-blind suppresses instability at the A/P compartment boundary of the *Drosophila* wing. *Mech. Dev* **125**, 233–246 (2008).
45. Umemori, M., Takemura, M., Maeda, K., Ohba, K. & Adachi-Yamada, T. *Drosophila* T-box transcription factor Optomotor-blind prevents pathological folding and local overgrowth in wing epithelium through confining Hh signal. *Dev. Biol* **308**, 68–81 (2007).
46. Agnes, F., Suzanne, M. & Noselli, S. The *Drosophila* JNK pathway controls the morphogenesis of imaginal discs during metamorphosis. *Development* **126**, 5453–5462 (1999).
47. Tripura, C., Chandrika, N. P., Susmitha, V. N., Noselli, S. & Shashidhara, L. S. Regulation and activity of JNK signaling in the wing disc peripodial membrane during adult morphogenesis in *Drosophila*. *Int. J. Dev. Biol* **55**, 583–590 (2011).
48. Martin-Blanco, E. *et al.* Puckered encodes a phosphatase that mediates a feedback loop regulating JNK activity during dorsal closure in *Drosophila*. *Genes Dev* **12**, 557–570 (1998).
49. Weber, U., Paricio, N. & Mlodzik, M. Jun mediates Frizzled-induced R3/R4 cell fate distinction and planar polarity determination in the *Drosophila* eye. *Development* **127**, 3619–3629 (2000).
50. Adachi-Yamada, T., Fujimura-Kamada, K., Nishida, Y. & Matsumoto, K. Distortion of proximodistal information causes JNK-dependent apoptosis in *Drosophila* wing. *Nature* **400**, 166–169 (1999).
51. Jezowska, B. *et al.* A dual function of *Drosophila* capping protein on DE-cadherin maintains epithelial integrity and prevents JNK-mediated apoptosis. *Dev. Biol* **360**, 143–159 (2011).
52. Leong, G. R., Goulding, K. R., Amin, N., Richardson, H. E. & Brumby, A. M. Scribble mutants promote aPKC and JNK-dependent epithelial neoplasia independently of Crumbs. *BMC Biol* **7**, 62 (2009).
53. Srivastava, A., Pastor-Pareja, J. C., Igaki, T., Pagliarini, R. & Xu, T. Basement membrane remodeling is essential for *Drosophila* disc eversion and tumor invasion. *Proc. Natl. Acad. Sci. USA* **104**, 2721–2726 (2007).
54. Stevens, L. J. & Page-McCaw, A. A secreted MMP is required for reepithelialization during wound healing. *Mol. Biol. Cell* **23**, 1068–1079 (2012).
55. Uhlirova, M. & Bohmann, D. JNK- and Fos-regulated Mmp1 expression cooperates with Ras to induce invasive tumors in *Drosophila*. *EMBO J* **25**, 5294–5304 (2006).
56. Sun, G. & Irvine, K. D. Regulation of Hippo signaling by Jun kinase signaling during compensatory cell proliferation and regeneration, and in neoplastic tumors. *Dev. Biol* **350**, 139–151 (2011).

57. Huang, J., Wu, S., Barrera, J., Matthews, K. & Pan, D. The Hippo signaling pathway coordinately regulates cell proliferation and apoptosis by inactivating Yorkie, the *Drosophila* Homolog of YAP. *Cell* **122**, 421–434 (2005).
58. Nolo, R., Morrison, C. M., Tao, C., Zhang, X. & Halder, G. The bantam microRNA is a target of the hippo tumor-suppressor pathway. *Curr. Biol* **16**, 1895–1904 (2006).
59. Thompson, B. J. & Cohen, S. M. The Hippo pathway regulates the bantam microRNA to control cell proliferation and apoptosis in *Drosophila*. *Cell* **126**, 767–774 (2006).
60. Graves, H. K., Woodfield, S. E., Yang, C. C., Halder, G. & Bergmann, A. Notch signaling activates Yorkie non-cell autonomously in *Drosophila*. *PLoS One* **7**, e37615 (2012).
61. Zhang, X., Luo, D., Pflugfelder, G. O. & Shen, J. Dpp signaling inhibits proliferation in the *Drosophila* wing by Omb-dependent regional control of bantam. *Development* **140**, 2917–2922 (2013).
62. Chen, C. L., Schroeder, M. C., Kango-Singh, M., Tao, C. & Halder, G. Tumor suppression by cell competition through regulation of the Hippo pathway. *Proc. Natl. Acad. Sci. USA* **109**, 484–489 (2012).
63. Enomoto, M. & Igaki, T. Src controls tumorigenesis via JNK-dependent regulation of the Hippo pathway in *Drosophila*. *EMBO Rep* **14**, 65–72 (2013).
64. Igaki, T., Pagliarini, R. A. & Xu, T. Loss of cell polarity drives tumor growth and invasion through JNK activation in *Drosophila*. *Curr. Biol* **16**, 1139–1146 (2006).
65. Zhao, B., Tumaneng, K. & Guan, K. L. The Hippo pathway in organ size control, tissue regeneration and stem cell self-renewal. *Nat. Cell. Biol* **13**, 877–883 (2011).
66. Willsey, H. R. *et al.* Localized JNK signaling regulates organ size during development. *Elife* **5** (2016).
67. Doggett, K., Grusche, F. A., Richardson, H. E. & Brumby, A. M. Loss of the *Drosophila* cell polarity regulator Scribbled promotes epithelial tissue overgrowth and cooperation with oncogenic Ras-Raf through impaired Hippo pathway signaling. *BMC Dev. Biol* **11**, 57 (2011).
68. Calleja, M., Moreno, E., Pelaz, S. & Morata, G. Visualization of gene expression in living adult *Drosophila*. *Science* **274**, 252–255 (1996).
69. McGuire, S. E., Le, P. T., Osborn, A. J., Matsumoto, K. & Davis, R. L. Spatiotemporal rescue of memory dysfunction in *Drosophila*. *Science* **302**, 1765–1768 (2003).
70. Page-McCaw, A., Serano, J., Sante, J. M. & Rubin, G. M. *Drosophila* matrix metalloproteinases are required for tissue remodeling, but not embryonic development. *Dev. Cell* **4**, 95–106 (2003).
71. Lee, J. J., von Kessler, D. P., Parks, S. & Beachy, P. A. Secretion and localized transcription suggest a role in positional signaling for products of the segmentation gene hedgehog. *Cell* **71**, 33–50 (1992).
72. Oh, H. & Irvine, K. D. Cooperative regulation of growth by Yorkie and Mad through bantam. *Dev. Cell* **20**, 109–122 (2011).

Acknowledgements

We thank the Bloomington stock center and TsingHua Fly Center for fly stocks. This research was supported by the National Natural Science Foundation of China [NSFC31372255] and the 973 Program [2013CB127603].

Author Contributions

J.S. developed the concept and designed the experiments. S.L. performed the experiments. D.W., G.P. and J.S. analysed the data and wrote the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Liu, S. *et al.* Fold formation at the compartment boundary of *Drosophila* wing requires Yki signaling to suppress JNK dependent apoptosis. *Sci. Rep.* **6**, 38003; doi: 10.1038/srep38003 (2016).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016