

SCIENTIFIC REPORTS



OPEN

Collisional cross-section of water molecules in vapour studied by means of ^1H relaxation in NMR

Received: 10 February 2016
Accepted: 11 November 2016
Published: 23 December 2016

Daniele Mammoli¹, Estel Canet^{1,2,3}, Roberto Buratto¹, Pascal Miéville¹, Lothar Helm¹ & Geoffrey Bodenhausen^{2,3}

In gas phase, collisions that affect the rotational angular momentum lead to the return of the magnetization to its equilibrium (relaxation) in Nuclear Magnetic Resonance (NMR). To the best of our knowledge, the longitudinal relaxation rates $R_1 = 1/T_1$ of protons in H_2O and HDO have never been measured in gas phase. We report R_1 in gas phase in a field of 18.8 T, i.e., at a proton Larmor frequency $\nu_0 = 800$ MHz, at temperatures between 353 and 373 K and pressures between 9 and 101 kPa. By assuming that spin rotation is the dominant relaxation mechanism, we estimated the effective cross-section σ_j for the transfer of angular momentum due to H_2O - H_2O and HDO - D_2O collisions. Our results allow one to test theoretical predictions of the intermolecular potential of water in gas phase.

Water is the most extensively studied molecule on Earth. A precise determination of its intermolecular potential would allow accurate predictions of its properties in gas, liquid and solid phase. However, despite huge theoretical efforts^{1,2}, a full agreement with experiments³⁻⁶ has not yet been achieved. Nuclear magnetic resonance (NMR) of molecules in gas phase⁷ has some unique features. The coupling between nuclear spins and magnetic moments induced by molecular rotation implies that collisions between molecules lead to a relaxation, i.e. to the return of the longitudinal magnetization M_z to its equilibrium after a perturbation, through a mechanism known as spin-rotation. If spin-rotation is the dominant mechanism, cross-sections for the transfer of angular momentum can be obtained from NMR relaxation rates in gas phase. Such relaxation rates have been measured over a wide range of pressures and temperatures⁸⁻¹⁰. Experimentally determined cross-sections can be used to refine intermolecular potentials¹¹⁻¹³. In methane, isotopic substitution¹⁴⁻¹⁶ affects relaxation rates associated with the different isotopomers such as CH_4 , CH_3D , CH_2D_2 , and CHD_3 . In supercritical water¹⁷⁻¹⁹, spin-rotation significantly contributes to NMR relaxation despite the high density. In the context of our attempts to prepare *para*-water²⁰⁻²⁶ and related spin states in other molecules²⁷⁻³², we have measured longitudinal relaxation rates $R_1 = 1/T_1$ of gaseous H_2O and HDO at different temperatures and pressures. To the best of our knowledge, this is the first time that such observations are reported. Our measurements are useful to refine intermolecular potentials for water vapour. These may be compared with water confined in matrices^{33,34} or in fullerene cages^{21,35} where a gas-phase like behaviour can be observed.

Theory

Collisions between molecules can induce transitions between rotational quantum states. As a result, spin-dependent interactions vary as a function of time and, if the fluctuations occur at frequencies in the vicinity of the nuclear Larmor frequency ω_0 , longitudinal NMR relaxation takes place. Comprehensive theoretical treatments of NMR relaxation can be found elsewhere³⁶⁻⁴⁴. In this article, we shall only mention some aspects of spin-rotation and dipole-dipole relaxation mechanisms that are relevant to longitudinal relaxation in gas phase.

Spin-rotation (SR) relaxation is due to collisions that modulate local fields induced at the sites of the nuclei by the rotation of the surrounding electronic cloud.

¹Institut des Sciences et Ingénierie Chimiques, Ecole Polytechnique Fédérale de Lausanne, 1015 Lausanne, Switzerland. ²Département de Chimie, Ecole Normale Supérieure, PSL Research University, UPMC Univ Paris 06, CNRS, Laboratoire des Biomolécules (LBM), 24 rue Lhomond, 75005 Paris, France. ³Sorbonne Universités, UPMC Univ Paris 06, Ecole Normale Supérieure, CNRS, Laboratoire des Biomolécules (LBM), Paris, France. Correspondence and requests for materials should be addressed to D.M. (email: daniele.mammoli@epfl.ch)

Samples	H ₂ O in H ₂ O								HDO in D ₂ O	
	1		2		3		4		5	
T (K)	p (kPa)	R ₁ (s ⁻¹)	p (kPa)	R ₁ (s ⁻¹)	p (kPa)	R ₁ (s ⁻¹)	p (kPa)	R ₁ (s ⁻¹)	p (kPa)	R ₁ (s ⁻¹)
353	9	73 ± 9	15	79 ± 9	34	57 ± 5	47	54 ± 7	47	57 ± 8
363	10	72 ± 9	18	80 ± 7	42	52 ± 6	69	43 ± 6	69	42 ± 5
373	11	79 ± 9	21	80 ± 7	60	40 ± 6	101	36 ± 7	101	31 ± 3

Table 1. Longitudinal relaxation rates R_1 for gaseous H₂O (samples 1 to 4) and gaseous HDO (sample 5) at 800 MHz and at different temperatures and pressures.




Figure 1. (Points) Experimental rates R_1 of gaseous H₂O at 800 MHz and at pressures $9 < p < 101$ kPa. (Lines) Estimates of R_1 arising from spin-rotation, using Eq. 1 with the parameters in Table 2.

Relaxation induced by spin-rotation can be described by ref. 45:

$$R_1^{SR} = \frac{4\pi^2}{\alpha} C_{eff}^2 \frac{\tau_J}{1 + (\omega_0 - \omega_J)^2 \tau_J^2} \quad (1)$$

where:

$$\tau_J = \frac{1}{\rho \langle v \sigma_J \rangle} \quad \alpha = \frac{\hbar^2}{2I_0 k_B T} \quad \omega_J = \frac{g_{rot} \mu_N H}{\hbar} \nu = \sqrt{\frac{8k_B T}{\pi \mu}} \quad \mu = \frac{m_1 m_2}{(m_1 + m_2)}$$

τ_J is the spin-rotation correlation time, C_{eff} (in Hz) the spin-rotation constant, ω_J the rotational frequency (in rad/s)⁴⁶, ρ is the number density of molecules, ν is the average thermal velocity, σ_J is the collisional cross-section for the transfer of angular momentum, I_0 is the moment of inertia, g_{rot} is the g-factor, μ_N is the nuclear magneton, H is the magnetic field and μ is the reduced mass of the two colliding particles. The correlation time τ_J is related to the lifetime of the rotational quantum states. The relaxation process can be described by characterizing the cross-section for the transfer of angular momentum. Intermolecular potentials used to model the interaction mostly consist of an isotropic part, usually a radial function, depending only on the distance between particles (e.g., Lennard-Jones potential) and an anisotropic part, depending also on the orientation of the molecules with respect to each other. The intermolecular potential of a molecule can be written by considering its axial symmetry⁴⁷ and can be linked to relaxation rates via the Bloom – Oppenheim theory⁴⁸.

Dipole-dipole (DD) relaxation is due to fluctuations of the interaction between magnetic dipoles, which are induced by physical rotation. The DD interactions are described by a correlation time τ_C that is related to the mean time needed for the molecule to undergo a rotation through one radian. DD relaxation can occur between spins in the same molecule (intramolecular DD) or between spins in different molecules (intermolecular DD). Relaxation by the intramolecular DD interaction between the two protons of water is described by ref. 36:

$$R_1^{DD} = \left(\frac{\mu_0}{4\pi} \right)^2 \frac{3\hbar^2 \gamma_H^4}{10r^6} \left[\frac{\tau_c}{1 + (\omega_0 \tau_c)^2} + \frac{4\tau_c}{1 + 4(\omega_0 \tau_c)^2} \right] \quad (2)$$

	C_{eff} (kHz)	ω_0 (rad/s)	ω_J (rad/s)	μ (kg)	I_0 (kg·m ²)
HDO	42.8 ± 0.1 ^(a)	5 · 10 ⁹	6 · 10 ⁸ (b)	1.64 · 10 ⁻²⁶	2.91 · 10 ⁻⁴⁷
H ₂ O	32.2 ± 0.5 ^(a)	5 · 10 ⁹	6 · 10 ⁸ (b)	1.51 · 10 ⁻²⁶	1.94 · 10 ⁻⁴⁷

Table 2. Parameters used to calculate cross-sections via Eq. 1. ^(a)Ref. 53 ^(b)Ref. 66.

T (K)	Cross-section σ_J (Å ²)		Correlation time τ_J (ps)	
	H ₂ O:H ₂ O collisions	HDO:D ₂ O collisions	H ₂ O:H ₂ O collisions	HDO:D ₂ O collisions
353 K	140 ± 26	378 ± 49	82 ± 15	32 ± 5
363 K	142 ± 26	367 ± 42	56 ± 10	22 ± 3
373 K	144 ± 27	354 ± 31	38 ± 7	16 ± 2

Table 3. Correlation times τ_J and cross-sections σ_J for the angular momentum transfer in H₂O:H₂O and HDO:D₂O collisions, calculated with Eq. 1 and parameters in Table 2.

where r is the distance between the protons, γ_H is the gyromagnetic ratio of protons and μ_0 is the magnetic permeability in vacuum.

In liquid phase, the rotational correlation time τ_C is linked to τ_J by the Hubbard relation⁴⁹ $\tau_C/\tau_J = 1/6$. For dilute gases ($\tau_J \rightarrow \infty$), the ratio of correlation times τ_C/τ_J varies from 5/4 (Ivanov model) to 1/4 (extended diffusion model) or 1/24.4 (Langevin model)^{14,15}.

Results

We measured longitudinal relaxation rates R_1 by the conventional inversion-recovery method. Experiments were carried out at temperatures $T = 353, 363,$ and 373 K and pressures $9 < p < 101$ kPa. The translational diffusion of water molecules does not affect our measurements of longitudinal relaxation rates R_1 (see Methods), although it might interfere with measurements of transverse relaxation rates R_2 . The rates R_1 observed in neat water (samples 1–4, H₂O-H₂O collisions) and in a mixture of HDO and D₂O (sample 5, HDO-D₂O collisions) are reported in Table 1.

We shall initially consider spin-rotation to be the dominant relaxation mechanism, neglecting dipole-dipole relaxation. Under our experimental conditions, water vapour is mainly monomeric^{50–52} and the extreme narrowing regime ($(\omega_0 - \omega_J)^2 \tau_J^2 \ll 1$) is not fulfilled: R_1 shows a maximum at a pressure p^{\max} where $\tau_J = 1/(\omega_0 - \omega_J)$ (see Fig. 1).

The number density ρ at pressure p can be estimated via the ideal gas law (see Methods) yielding $\tau_J = RT/(p\nu\sigma_J)$. Hence, it is possible to calculate the cross-section $\sigma_J^{H_2O}$ for H₂O:H₂O collisions at p^{\max} as $\sigma_J^{H_2O}|^{\max} = RT(\omega_0 - \omega_J)/(p\nu^{\max})$. If we assume $\sigma_J^{H_2O}$ to be independent of p (i.e. $\sigma_J^{H_2O} = \sigma_J^{H_2O}|^{\max}$) at the low pressures used in our experiments, we can substitute $\sigma_J^{H_2O} = RT(\omega_0 - \omega_J)/(p\nu^{\max})$ in $\tau_J^{H_2O} = RT/(p\nu\sigma_J^{H_2O})$ to find $\tau_J^{H_2O} = p^{\max}/[p(\omega_0 - \omega_J)]$. This last relationship can be used to predict the dependence of R_1 on p , at a given T , by using Eq. 1 and the parameters in Table 2. All three curves result from fitting a single parameter (p^{\max}), all the other parameters being fixed to the values given in Table 2. The fitted value $p^{\max} = (17 \pm 3)$ kPa provides a fair agreement between experimental relaxation rates (points) and predicted rates (lines) (Fig. 1). The spin-rotation tensor depends on the symmetry of the molecule: in our approximation we take into account only the isotropic constant $C_{eff}^{35,53,54}$ which we consider to be independent of both pressure and temperature. The fitted value of p^{\max} is constant while C_{eff} is fixed to values comprised in its confidence range.

In a more refined analysis we included contributions R_1^{DD} due to the intramolecular dipole-dipole interaction (Eq. 2). We fixed τ_C to values predicted by the Ivanov model ($\tau_C = 5/4\tau_J$), extended diffusion model ($\tau_C = 1/4\tau_J$) or Langevin model ($\tau_C = 1/24.4\tau_J$)^{14,15}. Our experimental data are compatible with a negligible dipole-dipole contribution or with the Langevin model ($\tau_C \ll \tau_J$), according to which significant contributions of R_1^{DD} only occur at low pressures $p < 10$ kPa.

For HDO-D₂O mixtures, experimental relaxation rates R_1 (sample 5) are reported in Table 1. In this case, we can safely neglect DD contributions. The experimental rates R_1 in Table 1 and the parameters in Table 2 are substituted into Eq. 1 to calculate the collision cross-sections.

Cross sections and correlation times for the transfer of the angular momentum in H₂O:H₂O and HDO:D₂O collisions are reported in Table 3.

Discussion

Our analysis provides information about H₂O-H₂O and HDO-D₂O collisions at pressures below 101 kPa and temperatures between 353 and 373 K. The ratio of cross-sections $\sigma_J^{H_2O}/\sigma_J^{HDO} = 0.4 \pm 0.1$ differs from the ratio of the moments of inertia $I_0^{H_2O}/I_0^{HDO} = 0.66$. This discrepancy suggests that there must be appreciable differences between the intermolecular potentials for HDO:D₂O and H₂O:H₂O collisions. This hypothesis is compatible with the fact that H₂O and D₂O have almost equal electric dipole moments⁵⁵ while the electric dipole moment of HDO differs in intensity and orientation from those of H₂O and D₂O⁵⁵. NMR relaxation studies on the influence of hydrogen/deuterium isotopic substitution on collisional cross-sections have been reported for methane^{15,16}.




Figure 2. Schematic view of the coaxial tubes: outer tube (10 mm outer diameter) filled with toluene-d8 and inner tube (5 mm outer diameter, held by Teflon spacers) containing water vapour sealed under vacuum.

However, by isotopic substitution on methane only the moment of inertia is markedly altered. A direct comparison with isotopic substitution on the highly polar H₂O is therefore not possible.

The collisional cross-sections calculated from our NMR data can be used to refine the anisotropic part of the intermolecular potentials for collisions in gas phase^{56–58} via the Bloom – Oppenheim theory⁵⁹. However, such calculations are beyond the scope of this work.

Our findings may be relevant for Dissolution Dynamic Nuclear Polarization (D-DNP)⁶⁰ where a frozen sample is rapidly heated by injecting a burst of superheated D₂O ($T > 373$ K) into the cryostat, and the liquid HDO ‘bolus’, usually containing a hyperpolarized solute, is pushed by pressurized helium gas (typically at 1 MPa) through a polyethylene tube with a 1 mm inner diameter running through a “magnetic tunnel”⁶¹, with a length of ca. 4 m between the polarizer and the NMR or MRI system. Attempts to monitor the speed of the bolus moving through the tube by optical means have shown that it tends to break up into small droplets during the transfer. This increases the surface area where water molecules can exchange between the liquid and gaseous phases. If the liquid/gas exchange is fast, the *averaged* longitudinal relaxation rates are likely to be much shorter than those in liquid water. The shortening of T_1 would lead to a rapid loss of hyperpolarization during the transfer between the polarizer and the NMR magnet. Note that the viscosity and surface tension of the transferred liquid are difficult to control, since it consists of an aqueous solution containing analytes, polarizing agents like TEMPOL and glass-forming agents such as glycerol.

To summarize, we reported NMR relaxation rates due to binary H₂O:H₂O and HDO:D₂O collisions in the gas phase and evaluated the cross-sections for the transfer of the angular momentum which can be used to refine the intermolecular potentials.

Methods

Our experimental setup consisted of a pair of coaxial glass tubes (Fig. 2).

The inner tube with 5 mm outer diameter was held in the center of a 10 mm tube by holders made of PTFE (Teflon). The outer tube contained about 2 mL of deuterated toluene-d8 (boiling point $T_{bp} = 384$ K). Its deuterium signal allows one to lock the static field and to shim its homogeneity. The inner tube contained water that was frozen and flame-sealed under vacuum ($p = 1$ kPa). Four tubes of 3.5 to 4 cm length, labeled as samples 1, 2, 3 and 4, were filled with ca 0.1, 0.2, 0.3 and 4.5 mg H₂O, determined with a precision balance (± 0.1 mg, max. tara 31 g). A fifth tube (sample 5) was filled with 2 mg of 98% D₂O and 2% H₂O (*v:v*), hence containing ca. 2% HDO. The inner tube was completely immersed in the solvent contained in the outer tube (Fig. 2) in order to have a homogeneous temperature and to avoid condensation of water on the walls of the inner tube in regions outside the area where the temperature is accurately controlled. Before and after inserting the samples into the spectrometer, the temperature in the probe was determined with a platinum PT-100 resistance thermometer (“iTRON 08” by JUMO)⁶² using a similar set of two concentric tubes with toluene-d8 in the outer tube. After each experiment the maximum temperature variations were ± 1 K. Two typical ¹H NMR spectra are shown in Fig. 3: the peak near 3.2 ppm (w.r.t. TMS) is due to water in the gas phase at $T = 363$ K.

NMR instrumentation. All NMR experiments have been performed on a Bruker Avance-II 800 MHz spectrometer equipped with a 10 mm BBO probe.

Evaluation of pressure and density. To determine the pressure p and the number density ρ of the water in samples 1 to 5 we measured the mass of water and estimated the volume of the inner tubes. Samples 4 and 5 contain saturated vapour ($p = p^{sat}$). In that case the pressure p^{sat} can be calculated using Antoine’s equation⁶³:

$$\log_{10}(p^{sat}) = A - \frac{B}{C + T} \quad (3)$$

where T is the temperature and A , B and C are sample-specific constants. When expressing the pressure in bar and the temperature in K, we assumed⁶⁴ $A = 5.08354$, $B = 1663.125$ and $C = -45.622$ for both H₂O and D₂O, since their vapour pressures are similar within 1% over our range of temperatures⁶⁵. The number density ρ at a pressure p can be estimated provided that the equation of state of the gas is known *a priori*. We have compared ρ^{ideal} predicted by the ideal gas law with ρ^{virial} obtained from a second-order virial expansion. The deviation $(\rho^{ideal} - \rho^{virial})/\rho^{virial}$ is always below 2% in the range of pressures and temperatures under investigation, so that the use of the ideal gas law is legitimate.




Figure 3. Proton NMR spectra of samples described in Fig. 2 at 800 MHz and at 300 K (top) or 363 K (bottom). The signals at 2.1 and 7 ppm are attributed to residual protons in incompletely deuterated toluene-d₈. The signal at 3.2 stems from water in gas phase. Small peaks between 0.3 and 2 ppm are due to impurities in toluene-d₈. We diluted TMS in toluene-d₈ to use its resonance at 0 ppm as chemical shift reference.

The quantity of water vapour in samples 1 to 3 has been determined by integration of the relevant signals in the NMR spectra. As a reference for integration we added 1,1,2,2-tetrachloroethane (C₂H₂Cl₄, 0.2% v:v) to the toluene-d₈ in the outer sample tube. We calibrated the integral of the C₂H₂Cl₄ reference peak (near ~6 ppm) with respect to the number density of sample 4 (saturated vapour). The pressures in samples 1 to 3 are then determined by scaling the peak intensities of the vapour peak with respect to sample 4. The error on the pressures is assumed to be ±10%. The active volume of the 5 mm inner tube has been estimated from documentation by the manufacturer (Wilmad) to be 0.4 cm³.

Translational diffusion and convection. Translational diffusion of water molecules in gas phase is very fast. Translational motion of water molecules between the active volume of the ¹H NMR coil and the space outside the coil can affect inversion-recovery measurements of *T*₁ relaxation. Indeed, molecules that carry inverted magnetization $-M_z = -M_z^{eq}$ within the active volume may be contaminated with molecules that come from areas outside the rf coil that carry magnetization in equilibrium M_z^{eq} that has not been inverted. To ascertain the relevance of these effects on the time scale of the *T*₁ measurement (max. $5 \cdot T_1 = 140$ ms) we performed the following test. The inner tubes were only a few mm longer than the active region of the ¹H coil of the 10 mm probe which is about 3 cm long. We measured *R*₁ at the highest temperature *T* = 373 K (where the effects of diffusion are most pronounced) in two arrangements. First, we centered the inner tube with respect to the active region of the ¹H coil. In this configuration, molecules can diffuse to and from the volumes above and below the active region. Secondly, we moved the inner tube up so that its bottom end was aligned with the lower end of the active region of the coil. In this manner, only molecules that cross the limit of the active region of the rf coil from above can influence the NMR signal. Any difference in *R*₁ observed with these two configurations should be due to diffusion or convection. We found the *R*₁ values to be identical within their errors, suggesting that contributions from diffusion can be neglected. Since we immersed the inner tube completely in a liquid with a controlled temperature, we assumed that there was no significant temperature gradient, so that convection due to differences in density should be negligible. Nevertheless, the experimental errors of the relaxation rates were doubled to take into account uncertainties stemming from diffusion and convection.

References

- Bukowski, R., Szalewicz, K., Groenenboom, G. C. & van der Avoird, A. Predictions of the Properties of Water from First Principles. *Science* **315**, 1249–1252 (2007).
- Nilsson, A. & Pettersson, L. G. M. The structural origin of anomalous properties of liquid water. *Nat. Commun.* **6**, 8998 (2015).
- Astrath, N. G. C., Malacarne, L. C., Baesso, M. L., Lukaszewicz, G. V. B. & Bialkowski, S. E. Unravelling the effects of radiation forces in water. *Nat. Commun.* **5**, (2014).
- Russo, J. & Tanaka, H. Understanding water's anomalies with locally favoured structures. *Nat. Commun.* **5**, (2014).
- Elgabarty, H., Khaliullin, R. Z. & Kühne, T. D. Covalency of hydrogen bonds in liquid water can be probed by proton nuclear magnetic resonance experiments. *Nat. Commun.* **6**, 8318 (2015).
- Ramasesha, K., De Marco, L., Mandal, A. & Tokmakoff, A. Water vibrations have strongly mixed intra- and intermolecular character. *Nat. Chem.* **5**, 935–940 (2013).
- Jameson, C. J. Gas-phase NMR spectroscopy. *Chem. Rev.* **91**, 1375–1395 (1991).
- Armstrong, R. L., Kisman, K. E. & Kalechstein, W. Longitudinal Relaxation Time Measurements in Hydrogen Gas Mixtures at Low Densities. *Can. J. Phys.* **53**, 1–4 (1975).
- Jameson, C. J., Jameson, A. K., Smith, N. C. & Jackowski, K. Cross sections for transfer of rotational angular momentum in CO₂ from ¹³C spin relaxation studies in the gas phase. *J. Chem. Phys.* **86**, 2717 (1987).
- Jameson, C. J. & Jameson, A. K. Effective collision cross sections for SF₆ from nuclear magnetic relaxation. *J. Chem. Phys.* **88**, 7448 (1988).
- Riehl, J. W. Spin-lattice relaxation and the anisotropic part of the H[sub 2][Single Bond]He and H[sub 2][Single Bond]Ne intermolecular potential. *J. Chem. Phys.* **58**, 4571 (1973).

12. Zarur, G. Effective potential formulation of molecule-molecule collisions with application to H₂[Single Bond]H₂. *J. Chem. Phys.* **60**, 2057 (1974).
13. Jameson, C. J. In *New Developments in NMR - Chapter 1. Fundamental Intramolecular and Intermolecular Information from NMR in the Gas Phase* (eds. Jackowski, K. & Jaszunski, M.) 1–51 (Royal Society of Chemistry, 2016).
14. Jameson, C. J., Jameson, A. K., Smith, N. C., Hwang, J. K. & Zia, T. Carbon-13 and proton spin relaxation in methane in the gas phase. *J. Phys. Chem.* **95**, 1092–1098 (1991).
15. ter Horst, M. A., Jameson, C. J. & Jameson, A. K. Molecular reorientation of CD₄ in gas-phase mixtures. *Magn. Reson. Chem.* **44**, 241–248 (2006).
16. Bloom, M., Bridges, F. & Hardy, W. N. Nuclear spin relaxation in gaseous methane and its deuterated modifications. *Can. J. Phys.* **45**, 3533–3554 (1967).
17. Lamb, W. J. NMR study of compressed supercritical water. *J. Chem. Phys.* **74**, 913 (1981).
18. Jonas, J., DeFries, T. & Lamb, W. J. NMR proton relaxation in compressed supercritical water. *J. Chem. Phys.* **68**, 2988 (1978).
19. Lamb, W. J. Self-diffusion in compressed supercritical water. *J. Chem. Phys.* **74**, 6875 (1981).
20. Mammoli, D. *et al.* Challenges in preparing, preserving and detecting para-water in bulk: overcoming proton exchange and other hurdles. *Phys Chem Chem Phys* (2015).
21. Beduz, C. *et al.* Quantum rotation of ortho and para-water encapsulated in a fullerene cage. *Proc. Natl. Acad. Sci.* **109**, 12894–12898 (2012).
22. Mamone, S. *et al.* Nuclear spin conversion of water inside fullerene cages detected by low-temperature nuclear magnetic resonance. *J. Chem. Phys.* **140**, 194306 (2014).
23. Horke, D. A., Chang, Y.-P., Długołęcki, K. & Küpper, J. Separating Para and Ortho Water. *Angew. Chem. Int. Ed.* **53**, 11965–11968 (2014).
24. Kravchuk, T. *et al.* A Magnetically Focused Molecular Beam of Ortho-Water. *Science* **331**, 319–321 (2011).
25. Tikhonov, V. I. Separation of Water into Its Ortho and Para Isomers. *Science* **296**, 2363–2363 (2002).
26. Meier, B. *et al.* Electrical detection of ortho–para conversion in fullerene-encapsulated water. *Nat. Commun.* **6**, 8112 (2015).
27. Mammoli, D. *et al.* Hyperpolarized para -Ethanol. *J. Phys. Chem. B* **119**, 4048–4052 (2015).
28. Bornet, A. *et al.* Long-Lived States of Magnetically Equivalent Spins Populated by Dissolution-DNP and Revealed by Enzymatic Reactions. *Chem. - Eur. J* **20**, 17113–17118 (2014).
29. Zhang, Y., Soon, P. C., Jerschow, A. & Canary, J. W. Long-Lived ¹H Nuclear Spin Singlet in Dimethyl Maleate Revealed by Addition of Thiols. *Angew. Chem. Int. Ed.* **53**, 3396–3399 (2014).
30. Tayler, M. C. D. *et al.* Direct Enhancement of Nuclear Singlet Order by Dynamic Nuclear Polarization. *J. Am. Chem. Soc.* **134**, 7668–7671 (2012).
31. Stevanato, G. *et al.* A Nuclear Singlet Lifetime of More than One Hour in Room-Temperature Solution. *Angew. Chem. Int. Ed.* **54**, 3740–3743 (2015).
32. Kovtunov, K. V. *et al.* Long-Lived Spin States for Low-Field Hyperpolarized Gas MRI. *Chem. - Eur. J* **20**, 14629–14632 (2014).
33. Fajardo, M. E., Tam, S. & DeRose, M. E. Matrix isolation spectroscopy of H₂O, D₂O, and HDO in solid parahydrogen. *J. Mol. Struct.* **695–696**, 111–127 (2004).
34. Redington, R. L. & Milligan, D. E. Infrared Spectroscopic Evidence for the Rotation of the Water Molecule in Solid Argon. *J. Chem. Phys.* **37**, 2162 (1962).
35. Li, Y. *et al.* Comparison of Nuclear Spin Relaxation of H₂O@C₆₀ and H₂@C₆₀ and Their Nitroxide Derivatives. *J. Phys. Chem. Lett.* **3**, 1165–1168 (2012).
36. Kowalewski, J. & Mäler, L. *Nuclear spin relaxation in liquids: theory, experiments, and applications* (Taylor & Francis, 2006).
37. Abragam, A. *The principles of nuclear magnetism* (Oxford Univ. Press, 2006).
38. Bransden, B. H. & Joachain, C. J. *Physics of atoms and molecules* (Prentice Hall, 2003).
39. Callaghan, P. T. *Principles of nuclear magnetic resonance microscopy* (Clarendon Press, 2007).
40. *Protein NMR spectroscopy: principles and practice* (Academic Press, 2007).
41. Keeler, J. *Understanding NMR spectroscopy* (John Wiley and Sons, 2010).
42. Levitt, M. H. *Spin dynamics: basics of nuclear magnetic resonance* (John Wiley & Sons, 2008).
43. Ernst, R. R., Bodenhausen, G. & Wokaun, A. *Principles of Nuclear Magnetic Resonance in One and Two Dimensions* (Clarendon Press, Oxford, 1987).
44. Slichter, C. P. *Principles of magnetic resonance* (Springer-Verlag, 1992).
45. Courtney, J. A. & Armstrong, R. L. A Nuclear Spin Relaxation Study of the Spin–Rotation Interaction in Spherical Top Molecules. *Can. J. Phys.* **50**, 1252–1261 (1972).
46. Anderson, C. H. & Ramsey, N. F. Magnetic Resonance Molecular-Beam Spectra of Methane. *Phys. Rev.* **149**, 14–24 (1966).
47. Gray, C. G. On the theory of multipole interactions. *Can. J. Phys.* **46**, 135–139 (1967).
48. Bloom, M. & Oppenheim, I. In *Advances in Chemical Physics* (ed. Hirschfelder, J. O.) **12**, 549–599 (John Wiley & Sons, Inc., 1967).
49. Hubbard, P. S. Theory of Nuclear Magnetic Relaxation by Spin-Rotational Interactions in Liquids. *Phys. Rev.* **131**, 1155–1165 (1963).
50. Slanina, Z. Computational studies of water clusters: temperature, pressure, and saturation effects on cluster fractions within the RRHO MCY-B/EPEN steam. *J. Mol. Struct.* **237**, 81–92 (1990).
51. Mhin, B. J., Lee, S. J. & Kim, K. S. Water-cluster distribution with respect to pressure and temperature in the gas phase. *Phys. Rev. A* **48**, 3764–3770 (1993).
52. Johansson, E., Bolton, K. & Ahlström, P. Simulations of vapor water clusters at vapor–liquid equilibrium. *J. Chem. Phys.* **123**, 24504 (2005).
53. Bluyssen, H., Dymanus, A., Reuss, J. & Verhoeven, J. Spin-rotation constants in H₂O, HDO and D₂O. *Phys. Lett. A* **25**, 584–585 (1967).
54. Puzzarini, C., Cazzoli, G., Harding, M. E., Vázquez, J. & Gauss, J. The hyperfine structure in the rotational spectra of D₂17O and HD17O: Confirmation of the absolute nuclear magnetic shielding scale for oxygen. *J. Chem. Phys.* **142**, 124308 (2015).
55. Clough, S. A. Dipole moment of water from Stark measurements of H₂O, HDO, and D₂O. *J. Chem. Phys.* **59**, 2254 (1973).
56. Rajan, S., Lalita, K. & Babu, S. V. Intermolecular potentials from nmr data. 1. CH₄-N₂ AND CH₄-CO₂. *Can. J. Phys.* **53**, 1624–1630 (1975).
57. Pandey, L., Reddy, C. P. K. & Sarkar, K. L. Intermolecular potentials from nmr data - H₂-N₂O AND H₂-CO₂. *Can. J. Phys.* **61**, 664–670 (1983).
58. Coroiu, I., Demco, D. E. & Bogdan, N. Anisotropic intermolecular potential from nuclear spin-lattice relaxation in hexafluoride gases. *Appl. Magn. Reson.* **14**, 9–17 (1998).
59. Bloom, M. & Oppenheim, I. In *Advances in Chemical Physics* **12**, 549–599 (Hirschfelder J.O.).
60. Ardenkjaer-Larsen, J. H. *et al.* Increase in signal-to-noise ratio of >10,000 times in liquid-state NMR. *Proc. Natl. Acad. Sci. USA* **100**, 10158–10163 (2003).
61. Milani, J. *et al.* A magnetic tunnel to shelter hyperpolarized fluids. *Rev. Sci. Instrum.* **86**, 24101 (2015).
62. Ammann, C., Meier, P. & Merbach, A. A simple multinuclear NMR thermometer. *J. Magn. Reson.* **1969** **46**, 319–321 (1982).
63. Antoine, C. [Tensions des vapeurs; nouvelle relation entre les tensions et les températures] [Vapor Pressure: a new relationship between pressure and temperature]. *Comptes Rendus des Séances de l'Académie des Sciences (in French)* **107**, 681–684, 778–780, 836–837 (1888).
64. NIST database. <http://webbook.nist.gov/cgi/cbook.cgi?ID=C7732185&Mask=4&Type=ANTOINE&Plot=on>.
65. Jones, W. M. Vapor Pressures of Tritium Oxide and Deuterium Oxide. Interpretation of the Isotope Effects. *J. Chem. Phys.* **48**, 207 (1968).
66. Schwenke, D. W. Beyond the Potential Energy Surface: Ab initio Corrections to the Born–Oppenheimer Approximation for H₂O⁺. *J. Phys. Chem. A* **105**, 2352–2360 (2001).

Acknowledgements

We are indebted to the Swiss National Science Foundation (FNRS), the École polytechnique fédérale de Lausanne (EPFL), the French Centre national de la recherche scientifique (CNRS) and the European Research Council (ERC) for providing financial support.

Author Contributions

D.M., E.C. and R.B. designed and performed the experiment. P.M., L.H., G.B. conceived the experiment. D.M. analysed the data and wrote the paper.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Mammoli, D. *et al.* Collisional cross-section of water molecules in vapour studied by means of ^1H relaxation in NMR. *Sci. Rep.* **6**, 38492; doi: 10.1038/srep38492 (2016).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016