SCIENTIFIC REPORTS

OPEN

Positive association between physical activity and *PER3* expression in older adults

Masaki Takahashi¹, Atsushi Haraguchi², Yu Tahara³, Natsumi Aoki¹, Mayuko Fukazawa², Kumpei Tanisawa⁴, Tomoko Ito⁴, Takashi Nakaoka⁵, Mitsuru Higuchi^{4,6} & Shigenobu Shibata^{1,6}

Received: 05 September 2016

Accepted: 25 November 2016

Published: 03 January 2017

The circadian clock regulates many physiological functions including physical activity and feeding patterns. In addition, scheduled exercise and feeding themselves can affect the circadian clock. The purpose of the present study was to investigate the relationship between physical/feeding activity and expression of clock genes in hair follicle cells in older adults. Twenty adult men (age, 68 ± 7 years, mean $\pm$ SE) were examined in this cross-sectional study. Prior to hair follicle cell collection, the participants were asked to wear a uniaxial accelerometer for one week. The timings of breakfast, lunch, and dinner were also recorded. Hair follicle cells were then collected over a 24 h period at 4 h intervals. The amplitude of *PER3* expression was positively correlated with moderate and vigorous physical activity ($r = 0.582$, $p = 0.007$) and peak oxygen uptake ($r = 0.481$, $p = 0.032$), but these correlations were not observed for *NR1D1* or *NR1D2*. No association was noted between meal times and the amplitude or the acrophase for any of these three clock genes. These findings suggest that rhythmic expression of the circadian clock gene *PER3* is associated with the amount of daily physical activity and physical fitness in older adults.

The circadian clock system consists of a central clock located in the suprachiasmatic nucleus of the hypothalamus and peripheral clocks located in peripheral tissues, all of which have an approximately 24 h rhythm in mammals¹. Circadian clock genes regulate the daily rhythms of physiology and behavior such as locomotor activity, feeding time, sleep-wake cycle, body temperature, metabolism, alertness, and cognition in all mammals including humans^{1–4}. Scheduled feeding and exercise in mice can entrain peripheral circadian rhythms under light-dark conditions⁵. In addition, scheduled feeding and exercise in mice can restore to normal the amplitude reduction and arrhythmicity of clock gene expression created under constant light conditions⁶. Recently, one study in humans has reported that clock gene expression in subcutaneous fat is regulated by body weight changes and is associated with BMI, serum cholesterol levels and expression of metabolic and inflammatory genes². Another study in humans has shown that heavy physical exercise might influence the circadian phase of clock gene expression in hair follicle cells⁷. Thus, there are strong relationships between physical/feeding activities and clock gene expression in the peripheral organs of mice and human.

In aging, it has been reported that body temperature amplitude is progressively blunted in humans⁸. In mice, aging also causes a functional deterioration in circadian systems including the expression of circadian clock genes^{9–11}. In fact, a relationship between abnormal circadian clock functions and age-related cognitive deficits and Alzheimer's disease has been demonstrated^{12–14}. One study reported that the circadian rhythmicity of core body temperature is correlated with a higher level of physical activity in older adults¹⁵. However, little is known about the relationship between daily physical activity and/or aerobic capacity and the rhythmic expression of circadian clock genes in older adults. From the viewpoint of abnormal circadian rhythm and prevention of various diseases with aging, it may be important to elevate the amplitude of clock gene expression by increasing physical activity.

In order to evaluate clock gene expression in humans, several different sampling methods have been used including measurement of blood metabolites and, collection of white blood cells or oral mucosa cells^{16–18}. To

¹Faculty of Science and Engineering, Waseda University, Shinjuku, 162-8480, Japan. ²Graduate School of Advanced Science and Engineering, Waseda University, Shinjuku, 162-8480, Japan. ³Waseda Institute for Advanced Study, Waseda University, Shinjuku, 162-8480, Japan. ⁴Faculty of Sports Sciences, Waseda University, Tokorozawa, 359-1192, Japan. ⁵Department of Medicine, Tokyo Women's Medical University, Medical Center East, Arakawa, Tokyo, 116-8567, Japan. ⁶Institute of Advanced Active Aging Research, Waseda University, Tokorozawa, 359-1192, Japan. Correspondence and requests for materials should be addressed to S.S. (email: shibatasa@waseda.jp)

Figure 1. The diurnal *PER3* (a), *NR1D1* (b), and *NR1D2* (c) expression in hair follicle cells and pattern of physical activity (d) (n = 20). Data are means $\pm$ SE. Main effect of time (*PER3*, $P = 0.001$; *NR1D1*, $P = 0.001$, *NR1D2*, $P = 0.001$; physical activity, $P = 0.001$; one-way ANOVA).

solve the problem of obtaining peripheral tissues in a non-invasive way, several studies have reported the use of hair follicle cells as a quick and convenient means for evaluating human circadian clock genes^{19,20}. Moreover, Akashi *et al.* have reported that circadian fluctuations in *PER3*, *NR1D1*, and *NR1D2* in hair follicle cells could be used to evaluate circadian rhythms in humans^{19,21}. *NR1D1* and *NR1D2* (also known as *Rev-erb* alpha and beta) are nuclear receptors that regulate a number of physiological functions including circadian rhythm, lipid metabolism, immune function, and cellular differentiation^{22–24}. Although it is difficult to clarify the physiological function of *PER3*, some studies have indicated that a *PER3* variant is correlated with the circadian phenotype (Morningness-Eveningness score; MEQ), sleep phase disorder, and sleep homeostasis^{25–27}. Recently, one study has reported that *PER3* variant causes a circadian phenotype, and is associated with a seasonal mood trait²⁸. The purpose of the present study was to investigate the relationship between physical/feeding activity and expression of clock genes in hair follicle cells in older adults. We hypothesized that an increase in daily physical activity and time of breakfast would both be correlated with clock gene expression patterns.

Results

The rhythm of clock gene expression in hair follicle cells. The expression rhythms of three clock genes, namely *PER3*, *NR1D1*, and *NR1D2*, as well as the house keeping gene *18s-rRNA*, were investigated in this study. One-way ANOVA revealed a significant main effect of time for *PER3*, *NR1D1*, and *NR1D2* expression ($p = 0.001$ for each) (Fig. 1a,b,c). However, there are no significant main effect of time for *18s-rRNA* (Figure S1). The results of cosinor analysis, including the amplitude and the acrophase values, are presented in Table 1. The individual amplitude and acrophase data are shown in Table S1.

Relationship between physical activity and the amplitude or the acrophase of clock gene expression rhythm. The amount of moderate to vigorous physical activity (MVPA) and step counts detected by the accelerometer were 170 ± 26 min/week (mean $\pm$ SE, range: 23–423 min/week) and 8389 ± 802 step/count (mean $\pm$ SE, range: 2847–15782 step/day) respectively. The $\dot{V}O_2$ peak was 25 ± 1 ml/min (mean $\pm$ SE, range: 16–36 ml/min).

Patterns of physical activity were also investigated in this study. One-way ANOVA revealed a significant main effect of time for patterns of physical activity (Fig. 1d). The patterns of physical activity were not associated with the amplitude of clock gene expression (*PER3*, *NR1D1*, and *NR1D2*) (Fig. 2a) (Figure S2a,b).

The amplitude of *PER3* expression was positively correlated with MVPA ($r = 0.582$, $p = 0.007$) (Fig. 2b, Table S2) and tended to be positively correlated with step counts ($r = 0.383$, $p = 0.096$) (Fig. 2c). The acrophase of *PER3* expression was positively associated with MVPA, but not with step counts (Table S3). However, MVPA and step counts were not associated with either the amplitude or the acrophase of *NR1D1* and *NR1D2* expression (Tables S2 and S3).

Gene	Forward	Reverse	Probe
18S-rRNA	CGCCGCTAGAGGTGAAATTC	CGAACCTCCGACTTTCGTTCT	CCGGCGCAAGACGGACCAGA
PER3	CTACCTGCACCCTGAAGATCGTTCTC	CTGGAATCCAGTATGATGTAGTCTCCGTTT	CTCTGATGGTTGCCATAC
NR1D1	GCTCAGTGCCATGTTTCGACTTC	AAGTCTCCAAGGGCCGGTTC	AAGCTCAACTCCCTGGC
NR1D2	TCCAGTACAAGAAGTGCCTGAAGAATGAAA	CACGCTTAGGAATACGACCAAAACCGA	ATGTCAGCAATGTCTG

Table 1. Primer sequences for real-time RT-PCR analysis and TaqMan MGB probes.

Figure 2. The relationship across all participants between amplitude of physical activity (a), moderate to vigorous physical activity (MVPA) levels (b), step count (c), $\dot{V}O_2$ peak (d) and expression of *PER3* ($n = 20$). The correlation coefficient ((a), $P = 0.150$, (b), $P = 0.007$, (c), $P = 0.096$, (d), $P = 0.032$; Pearson's correlation coefficient).

The amplitude of *PER3* expression was found to be higher in the active group compared to the inactive group ($p = 0.039$) (Figure S2a), but this was not the case for the amplitude of *NR1D1* and *NR1D2* expression (Figure S2b,c).

Relationship between $\dot{V}O_2$ peak and the amplitude or the acrophase of clock gene expression.

The amplitude but not the acrophase of *PER3* expression was positively correlated with $\dot{V}O_2$ peak ($r = 0.481$, $p = 0.032$) (Fig. 2d, Tables S2 and S3). In contrast, the $\dot{V}O_2$ peak was not associated with either the amplitude or the acrophase of *NR1D1* or *NR1D2* expression (Tables S2 and S3).

Timing of diet and MEQ score and its effect on the amplitude or the acrophase of clock gene expression. The effect of meal times was also evaluated in this study. Breakfast time (range; 5:30–10:00), lunch time (range; 11:30–13:30), and dinner time (range; 17:00–20:00) were not correlated with the amplitude or acrophase of *PER3*, *NR1D1*, or *NR1D2* expression (Tables S2 and S3). MEQ scores (age-adjusted MEQ score range; 38–60) were not correlated with the amplitude or acrophase of *PER3*, *NR1D1*, or *NR1D2* expression (Tables S2 and S3).

Discussion

To the best of our knowledge, the present study is the first to examine the relationship between the amount of daily physical activity/ $\dot{V}O_2$ peak and the expression of clock genes in humans. The main findings are that, in older adults, there was a positive correlation between MVPA and the amplitude/acrophase of *PER3*, and a positive

correlation between $\dot{V}O_2$ peak and *PER3* amplitude. These findings indicate that elevating the aerobic capacity by increasing daily physical activity may affect the internal circadian clock system driven by *PER3*.

It is well documented that circadian rhythms regulate physiological and biological processes involved in the sleep-wake cycle, body temperature, metabolism, alertness, and cognition^{1–4}. Akashi *et al.* have reported that assessing peripheral clock gene expression in hair follicle cells could be an effective tool for evaluating human circadian rhythms^{19,21}. These studies showed that the peak times of expression of *PER3*, *NR1D1*, and *NR1D2* were correlated with the average wake time. Moreover, Watanabe *et al.* have demonstrated that the phase of *PER3* expression in leukocytes significantly with that seen in beard hair follicle cells²⁰. In the present study, the expression patterns of *PER3*, *NR1D1*, and *NR1D2* observed are consistent with those reported in previous studies^{19,20}. Our findings convincingly demonstrate that examining clock gene expression using hair follicle cells is a convenient method in old adults.

Several studies have reported that the circadian system can be influenced by exercise and physical activity^{29–31}. One study has shown that circadian clock gene expression was phase-delayed following high-intensity exercise training in the evening⁷. In addition, amplitudes in body temperature are higher in active adults than in sedentary adults^{31,32}. Recently, it has been shown that a high level of aerobic capacity may be related to the elevation of rhythmicity in body temperature in older adults¹⁵. However, few studies are available addressing the relationship between daily physical activity status/physical fitness and circadian expression of clock genes in older adults. The amplitude of circadian clock gene expression is thought to indicate the strength of the circadian rhythm, which decreases with aging⁸. In the present study, the amplitude of *PER3* expression was positively correlated with the amount of moderate and vigorous intensity physical activity in older adults. Interestingly, *PER3* expression was higher in active older adults who adhered to physical activity guidelines for health (i.e. >150 min/week) than in inactive adults.

Although the mechanism is unclear at present, the circadian clocks in peripheral tissues might be induced by exercise or physical activity^{33,34}. A previous study has reported that scheduled exercise in mice altered the daily rhythm of clock gene expression³³. Increasing physical activity by scheduled exercise elevated the amplitude and phase of both heart rate and body temperature rhythms. Additionally, the amplitude of *PER2:LUCIFERASE* expression rhythms in the SCN and peripheral tissues of *Per2:Luc* knock-in mice were increased by exercise. Consistent with the previous study in mice, the present study found that the amplitude of *PER3* expression was positively correlated with the amount of physical activity in older adults.

The amplitude of *PER3* expression was also positively correlated with the $\dot{V}O_2$ peak in older adults. It is worth noting that the mean values observed in the present study are consistent with data previously observed in sedentary older adults³⁵. The $\dot{V}O_2$ peak expresses the aerobic capacity, which declines with aging³⁶. A previous study has reported that $\dot{V}O_{2\max}$, which is one of the aerobic capacity parameters, declines by approximately 10% per decade³⁶. Moreover, especially in older adults, overall physical fitness and aerobic capacity is dependent on daily physical activity. In fact, the $\dot{V}O_2$ peak in the present study was positively correlated with the amount of moderate and vigorous physical activity. Thus, it may be important to elevate aerobic fitness and capacity or increasing daily physical activity for elevating circadian rhythm amplitude.

We also assessed the pattern of physical activity and its relation to circadian clock gene expression. Although the peak time of physical activity was 3:00 PM in all participants, there was no association between the peak time of daily physical activity and the acrophase of *PER3*, *NR1D1*, and *NR1D2* expression. Therefore, regular physical activity may increase the amplitude of clock genes without influencing the time of physical activity. On the other hand, the acrophase of *PER3* expression was positively association with MVPA. In mice, high intensity treadmill exercise could entrain peripheral clocks⁵. MVPA may therefore be sufficient physical activity for entrainment of human peripheral clocks. In the present study, there was no association between amplitude/acrophase of clock gene expression and the timing of breakfast, lunch, and dinner. Although feeding pattern is well known to be associated with clock phase in murine peripheral clocks³⁷, feeding time appears not to be an important factor in setting the amplitude/acrophase in human peripheral clocks.

Watanabe *et al.* have reported that the circadian rhythmicity of *PER3* in hair follicle cells assessed by the cosinor method program was not observed in half of the subjects²⁰. On the other hand, Akashi *et al.*, again using the cosinor method program, have shown that the peak time of *PER3*, *NR1D1*, and *NR1D2* expression in hair follicle cells correlated with the average wakefulness time¹⁹. Therefore, the acrophase calculated using the cosinor method program may still be considered valid.

The present study has, however, some limitations. Firstly, it is difficult to separate cause and effect, because this study was a cross-sectional study. Thus, an intervention study would be required to examine the effect of increased physical activity on clock gene expression. Secondly, it is unclear if there is an effect of other external synchronizers such as light on clock gene expression in this study. Although active participants may spend more time outside, and therefore get more exposure to sunlight, further investigation will be needed to take light exposure into consideration, by using a device that could accurately measure light exposure. Thirdly, it is unclear whether the effects of physical activity on other clock genes such as *PER2* and *BMAL-1* and clock genes in other tissues is similar or different. Additional research will be required to examine the effects of physical activity on circadian clock genes obtained from the blood samples, or oral mucosa cells compared to hair follicle cells. Regarding the expression of *PER2* and *BMAL-1* in hair follicle cells, a previous study has shown that the expression of *PER2* oscillated with lower amplitude, and only slight oscillations were detected for *BMAL-1*¹⁹. Thus, monitoring these clock genes in hair follicle cells may be less useful in evaluating circadian rhythms in human. Finally, it has been difficult to clarify the physiological and biological function of *PER3*. In fact, understanding the role of *PER3* in humans has been a focus for the last several years. In this regard, several studies have indicated that polymorphisms in the *PER3* gene in humans may be related to the pathogenesis of delayed sleep phase syndrome^{26,38}. Recently, one genome-wide association analysis with 89,283 subjects reported that *PER3* variants

	Total (n = 20)	
	Amplitude	Acrophase
<i>PER3</i>	1.06 ± 0.08	6.84 ± 0.47
<i>NR1D1</i>	0.70 ± 0.08	4.20 ± 0.80
<i>NR1D2</i>	0.76 ± 0.07	4.76 ± 0.55

Table 2. Amplitude and acrophase of clock gene expression rhythms.

are associated with self-reporting of being a ‘morning person’³⁹. Another study has also demonstrated that a *PER3* variant causes a circadian phenotype and was associated with a seasonal mood trait²⁸. Taken together these reports indicate that *PER3* may have an important role in the development of chronotype, sleep phenotype, and depression and this will likely guide future research.

In conclusion, our study has revealed an association between *PER3* expression and daily physical activity in a sample of older adults. We have also demonstrated higher levels of *PER3* in older men who have higher aerobic fitness.

Methods

Participants. Twenty-eight participants were recruited from the general population from two local communities in the Tokorozawa metropolitan area. All participants in this study were men. The participants in the present study did not perform any shift-work or travel for one week prior to hair follicle cell collection. None of the participants were involved in athletics although some were recreationally active. Eight participants were subsequently excluded from the data analysis. The exclusion criteria included yielding an insufficient total amount of RNA (<40 ng/μL), and having insufficient accelerometer wear time (see “Physical activity measurement” section below). The physical characteristics of the participants are shown in Table 2. This study was conducted according to the guidelines laid down in the Declaration of Helsinki and was approved by the ethics committees of Waseda University (2014-G003). Informed consent was obtained from all participants following a detailed description of the experiment.

Baseline measurements of anthropometry, diet time, sleep profile, chronotype. For all participants, anthropometric variables were measured before physical activity measurements were performed. Body mass was measured to the nearest 0.1 kg using a digital scale (Karada Scan HBF-701, Omron Co., Japan). Height was measured to the nearest 0.1 cm using a wall-mounted stadiometer (YL-65, Yagami, Co., Japan). Body mass index (BMI) was calculated as weight in kilograms divided by the square of height in meters. Waist circumference was measured to the nearest 0.1 cm at the level of the umbilicus using a flexible plastic tape.

Participants were asked to respond to questions relating to diet and sleep such as “What time do you usually eat breakfast, lunch, and dinner?”, “During the last month, what time do you usually go to sleep and wake up?”, “How many hours do you usually sleep per day during the last month?”.

To assess chronotype, we used the Japanese version of the Horne-Ostberg Morningness-Eveningness Questionnaire (MEQ)^{40,41}. This tool has been validated in a Japanese population^{41,42}. The MEQ consisted of 19 questions related to preferred sleep time and daily performance (e.g. What would be the best time to perform hard physical work?). Because diurnal preference changes with age, MEQ scores were adjusted by age (age-adjusted MEQ score, MEQ score + 0.3512 × (39.212–age))⁴². The scores ranged from 16 to 86. Based on age-adjusted MEQ scores, participants were classified into three chronotype groups; either morningness (score 59–86), intermediate (score 42–58), or eveningness (score 16–41). The participants in the present study were classified as follows: 1 morningness, 19 intermediate.

Baseline measurements and determination of $\dot{V}O_2$ peak. At the first laboratory visit, the subjects’ height, weight, and percent body fat were measured by the same experimenter. After these measurements were taken, peak oxygen uptake ($\dot{V}O_2$ peak) was measured using a maximal graded exercise test on a cycle ergometer (Ergonomic 828E-Monark, Varberg, Sweden). The graded cycle exercise began at a workload of 30 W, which was increased by 15 W every minute until the subject could no longer maintain the required pedaling frequency of 60 rpm. During exercise, a 12-lead electrocardiogram was electronically recorded (Stress Test System ML-6500; Fukuda Denshi, Japan), and heart rate was derived from the R-R interval. In addition, ratings of perceived exertion (RPE) were determined at the end of each minute of exercise. Pulmonary gas exchange (oxygen uptake [$\dot{V}O_2$], carbon dioxide output [$\dot{V}CO_2$], minute ventilation [$\dot{V}E$], and respiratory exchange ratio [RER]) were determined breath-by-breath using a gas analyzer (AE-300S; Minato Medical Science, Japan).

To ascertain that $\dot{V}O_2$ peak was attained, at least two of the following four criteria had to be met: (1) $\dot{V}O_2$ plateaued despite increasing exercise intensity, (2) the highest HR measured during the last minute of exercise was >90% of the predicted maximal heart rate (using the formula 220 – age [in years]), (3) the highest RER was >1.10 during the final stage of incremental exercise, and (4) the subject achieved an RPE of ≥18.

Physical activity measurement. In order to determine daily physical activity levels, participants were asked to wear a uniaxial accelerometer (Lifecorder-EX; Suzuken Co. Ltd., Nagoya, Japan) for one week prior to hair follicle cell collection. A number of studies have reported that the Lifecorder-EX is considered as a validated accelerometer to evaluate and monitor step counts and physical activity levels^{43,44}. The device reports the intensity of the activity based on an 11-point scale (0, 0.5, 1–9; 0 being the lowest-intensity activity and 9 being the highest).

	Total (n = 20)
Age (years)	71 ± 1
Body mass (kg)	64 ± 2
Body mass index	23 ± 1
Time of breakfast (24 h)	8 ± 1
Time of lunch (24 h)	12 ± 1
Time of dinner (24 h)	19 ± 1
MEQ score	52 ± 1
$\dot{V}O_2$ peak (ml/min)	25 ± 1
Step count (step/day)	8389 ± 802
MVPA (min/week)	170 ± 26

Table 3. Physical characteristics of the participants. Data are means ± SE.

The intensity of activity is determined by measuring the magnitude and frequency of acceleration every 4 s. Data from participants who had worn the accelerometer for at least 10 h (600 min) a day for at least four weekdays and one weekend day were considered valid. The main physical activity variable used in this study was the duration of moderate to vigorous physical activity (MVPA). All minutes of recording with an activity level ≥ 4 were classified as MVPA. The activity level threshold of 4 was derived from a calibration study and corresponded to approximately three metabolic equivalents⁴¹. To establish whether physical activity had an effect on clock gene expression, we divided the participants into two groups namely, an active group and an inactive group. The active group included participants who performed MVPA for 150 min or more per week. This cut-off was chosen because this amount of physical activity is the minimum recommended by current physical activity health guidelines. The inactive group included participants who performed MVPA for less than 150 min per week.

Hair follicle cells collection and laboratory assays. Following the physical activity measurements, hair follicle cells were collected from all participants. The participants were asked to refrain from consuming excess alcohol, and to maintain normal eating and sleeping habits during the sampling day. Hairs were collected over a 24 h period at 4 h intervals by firmly holding and pulling the facial hair root, and then quickly soaking the hair in dissolution buffer (RNeasy Micro Kit; QIAGEN, Hilden, Germany). More than 10 hairs were collected at each sampling point. All samples were stored at -80°C until RNA purification. The RNeasy Micro Kit was used with frozen cytolysis solution to purify total RNA. Total RNA was reverse-transcribed using an Advanced cDNA Synthesis Kit for RT-qPCR (Bio-Rad Laboratories, Inc., CA, USA), and real-time PCR was performed using a TaqMan MGB probe (Applied Biosystems, CA, USA), and 1/20 volume of the reverse-transcription product. Data were obtained using the 7500 fast (Applied Biosystems, CA, USA) and expression was normalized to 18 S rRNA. The sequences of the primers and probes used are shown in Table 3. These primers and probe are consistent with those used in previous study²¹. To evaluate the circadian rhythm of clock gene expression and physical activity pattern, we used the cosinor method. The peak phase, amplitude, and rhythmicity of normalized data were determined using the single cosinor method program (Acro.exe version 3.5)⁴⁵.

Statistical analysis. Data was analyzed using Predictive Analytics Software (PASW) version 23.0 for Windows (SPSS Japan Inc., Tokyo, Japan). The Kolmogorov-Smirnov test was used to check for normal blood marker distribution. The distribution of these markers did not differ significantly from the normal. Using a one-way repeated measures ANOVA, we analyzed clock gene expression and physical activity patterns over a 24 h period. When significant main effects of time (24 h variation) were detected, we evaluated the amplitude and acrophase using the aforementioned cosinor method. Student's t-test was used to compare differences between the active and inactive groups. Partial correlation was used to examine the relationship between step count, the amount of physical activity, $\dot{V}O_2$ peak and expression of clock genes. Statistical significance was accepted at the 5% level. Results are presented as means ± standard error (SE).

References

1. Bass, J. & Takahashi, J. S. Circadian integration of metabolism and energetics. *Science* **330**, 1349–1354 (2010).
2. Pivovarov, O. *et al.* Regulation of the clock gene expression in human adipose tissue by weight loss. *Int J Obes (Lond)* **40**, 899–906 (2016).
3. Fang, M. Z. *et al.* Sleep interruption associated with house staff work schedules alters circadian gene expression. *Sleep Med* **16**, 1388–1394 (2015).
4. Kyriacou, C. P. & Hastings, M. H. Circadian clocks: genes, sleep, and cognition. *Trends Cogn Sci* **14**, 259–267 (2010).
5. Sasaki, H. *et al.* Forced rather than voluntary exercise entrains peripheral clocks via a corticosterone/noradrenaline increase in PER2::LUC mice. *Sci Rep* **6**, 27607 (2016).
6. Hamaguchi, Y., Tahara, Y., Hitosugi, M. & Shibata, S. Impairment of Circadian Rhythms in Peripheral Clocks by Constant Light Is Partially Reversed by Scheduled Feeding or Exercise. *J Biol Rhythms* **30**, 533–542 (2015).
7. Okamoto, A., Yamamoto, T., Matsumura, R., Node, K. & Akashi, M. An out-of-lab trial: a case example for the effect of intensive exercise on rhythms of human clock gene expression. *J Circadian Rhythms* **11**, 10 (2013).
8. Hofman, M. A. & Swaab, D. F. Living by the clock: the circadian pacemaker in older people. *Ageing Res Rev* **5**, 33–51 (2006).
9. Kondratova, A. A. & Kondratov, R. V. The circadian clock and pathology of the ageing brain. *Nat Rev Neurosci* **13**, 325–335 (2012).
10. Brown, S. A., Schmitt, K. & Eckert, A. Aging and circadian disruption: causes and effects. *Aging (Albany NY)* **3**, 813–817 (2011).
11. Nakamura, T. J. *et al.* Age-related decline in circadian output. *J Neurosci* **31**, 10201–10205 (2011).
12. Oosterman, J. M., van Someren, E. J., Vogels, R. L., Van Harten, B. & Scherder, E. J. Fragmentation of the rest-activity rhythm correlates with age-related cognitive deficits. *J Sleep Res* **18**, 129–135 (2009).

13. Naismith, S. L. *et al.* Circadian misalignment and sleep disruption in mild cognitive impairment. *J Alzheimers Dis* **38**, 857–866 (2014).
14. Coogan, A. N. *et al.* The circadian system in Alzheimer's disease: disturbances, mechanisms, and opportunities. *Biol Psychiatry* **74**, 333–339 (2013).
15. Dupont Rocher, S., Bessot, N., Sesboue, B., Bulla, J. & Davenne, D. Circadian Characteristics of Older Adults and Aerobic Capacity. *J Gerontol A Biol Sci Med Sci* **71**, 817–822 (2016).
16. Kasukawa, T. *et al.* Human blood metabolite timetable indicates internal body time. *Proc Natl Acad Sci USA* **109**, 15036–15041 (2012).
17. Boivin, D. B. *et al.* Circadian clock genes oscillate in human peripheral blood mononuclear cells. *Blood* **102**, 4143–4145 (2003).
18. Cajochen, C. *et al.* Evening exposure to blue light stimulates the expression of the clock gene PER2 in humans. *Eur J Neurosci* **23**, 1082–1086 (2006).
19. Akashi, M. *et al.* Noninvasive method for assessing the human circadian clock using hair follicle cells. *Proc Natl Acad Sci USA* **107**, 15643–15648 (2010).
20. Watanabe, M. *et al.* Rhythmic expression of circadian clock genes in human leukocytes and beard hair follicle cells. *Biochem Biophys Res Commun* **425**, 902–907 (2012).
21. Sato, M., Matsuo, T., Atmore, H. & Akashi, M. Possible contribution of chronobiology to cardiovascular health. *Front Physiol* **4**, 409 (2013).
22. Burris, T. P. Nuclear hormone receptors for heme: REV-ERBalpha and REV-ERBbeta are ligand-regulated components of the mammalian clock. *Mol Endocrinol* **22**, 1509–1520 (2008).
23. Zhao, X. *et al.* Nuclear receptors rock around the clock. *EMBO Rep* **15**, 518–528 (2014).
24. Zhang, Y. *et al.* GENE REGULATION. Discrete functions of nuclear receptor Rev-erbalpha couple metabolism to the clock. *Science* **348**, 1488–1492 (2015).
25. Hida, A. *et al.* Screening of clock gene polymorphisms demonstrates association of a PER3 polymorphism with morningness-eveningness preference and circadian rhythm sleep disorder. *Sci Rep* **4**, 6309 (2014).
26. Ebisawa, T. *et al.* Association of structural polymorphisms in the human period3 gene with delayed sleep phase syndrome. *EMBO Rep* **2**, 342–346 (2001).
27. Viola, A. U. *et al.* PER3 polymorphism predicts sleep structure and waking performance. *Curr Biol* **17**, 613–618 (2007).
28. Zhang, L. *et al.* A PERIOD3 variant causes a circadian phenotype and is associated with a seasonal mood trait. *Proc Natl Acad Sci USA* **113**, E1536–1544 (2016).
29. Facer-Childs, E. & Brandstaetter, R. The impact of circadian phenotype and time since awakening on diurnal performance in athletes. *Curr Biol* **25**, 518–522 (2015).
30. Schroder, E. A. & Esser, K. A. Circadian rhythms, skeletal muscle molecular clocks, and exercise. *Exerc Sport Sci Rev* **41**, 224–229 (2013).
31. Atkinson, G., Coldwells, A., Reilly, T. & Waterhouse, J. A comparison of circadian rhythms in work performance between physically active and inactive subjects. *Ergonomics* **36**, 273–281 (1993).
32. Weinert, D. & Waterhouse, J. The circadian rhythm of core temperature: effects of physical activity and aging. *Physiol Behav* **90**, 246–256 (2007).
33. Schroeder, A. M. *et al.* Voluntary scheduled exercise alters diurnal rhythms of behaviour, physiology and gene expression in wild-type and vasoactive intestinal peptide-deficient mice. *J Physiol* **590**, 6213–6226 (2012).
34. Castillo, C., Molyneux, P., Carlson, R. & Harrington, M. E. Restricted wheel access following a light cycle inversion slows re-entrainment without internal desynchrony as measured in Per2Luc mice. *Neuroscience* **182**, 169–176 (2011).
35. Ogawa, T. *et al.* Effects of aging, sex, and physical training on cardiovascular responses to exercise. *Circulation* **86**, 494–503 (1992).
36. Fleg, J. L. *et al.* Accelerated longitudinal decline of aerobic capacity in healthy older adults. *Circulation* **112**, 674–682 (2005).
37. Tahara, Y. & Shibata, S. Circadian rhythms of liver physiology and disease: experimental and clinical evidence. *Nat Rev Gastroenterol Hepatol* **13**, 217–226 (2016).
38. Archer, S. N. *et al.* A length polymorphism in the circadian clock gene Per3 is linked to delayed sleep phase syndrome and extreme diurnal preference. *Sleep* **26**, 413–415 (2003).
39. Hu, Y. *et al.* GWAS of 89,283 individuals identifies genetic variants associated with self-reporting of being a morning person. *Nat Commun* **7**, 10448 (2016).
40. Horne, J. A. & Ostberg, O. A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *Int J Chronobiol* **4**, 97–110 (1976).
41. Ishihara, K. *et al.* [The results of investigation by the Japanese version of Morningness-Eveningness Questionnaire]. *Shinrigaku Kenkyu* **57**, 87–91 (1986).
42. Hida, A. *et al.* Individual traits and environmental factors influencing sleep timing: a study of 225 Japanese couples. *Chronobiol Int* **29**, 220–226 (2012).
43. Ayabe, M. *et al.* Accumulation of short bouts of non-exercise daily physical activity is associated with lower visceral fat in Japanese female adults. *Int J Sports Med* **34**, 62–67 (2013).
44. Kumahara, H. *et al.* The use of uniaxial accelerometry for the assessment of physical-activity-related energy expenditure: a validation study against whole-body indirect calorimetry. *Br J Nutr* **91**, 235–243 (2004).
45. Refinetti, R., Lissen, G. C. & Halberg, F. Procedures for numerical analysis of circadian rhythms. *Biol Rhythm Res* **38**, 275–325 (2007).

Acknowledgements

This work was supported by the council for Science, Technology and Innovation, SIP, “Technologies for creating next-generation agriculture, forestry and fisheries” (funding agency: Bio-oriented Technology Research Advancement Institution, NARO) (S. S.).

Author Contributions

M.T., A.H., Y.T., K. T., M.H., T.N., S.S. designed the research, analyzed the data, and wrote the manuscript, M.T., A.H., N.A., M.F., K.T., and T.I. performed the experiments.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Takahashi, M. *et al.* Positive association between physical activity and *PER3* expression in older adults. *Sci. Rep.* **7**, 39771; doi: 10.1038/srep39771 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017