SCIENTIFIC REPORTS

OPEN

Therapeutic Effects of Human Umbilical Cord-Derived Mesenchymal Stem Cells in Acute Lung Injury Mice

Received: 06 July 2016
Accepted: 29 November 2016
Published: 04 January 2017

Hua Zhu^{1,2,*}, Yi Xiong^{1,*}, Yunqiu Xia^{1,3}, Rong Zhang¹, Daiyin Tian^{1,3}, Ting Wang^{1,3}, Jihong Dai^{1,3}, Lijia Wang¹, Hongbing Yao^{1,4}, Hong Jiang², Ke Yang^{1,5}, Enmei Liu^{1,3}, Yujun Shi⁶, Zhou Fu^{1,3,5}, Li Gao^{1,4,5} & Lin Zou^{1,5,7}

The incidence and mortality of acute lung injury (ALI)/acute respiratory distress syndrome (ARDS) are still very high, but stem cells show some promise for its treatment. Here we found that intratracheal administration of human umbilical cord-mesenchymal stem cells (UC-MSCs) significantly improved survival and attenuated the lung inflammation in lipopolysaccharide (LPS)-induced ALI mice. We also used the proteins-chip and bioinformatics to analyze interactions between UC-MSCs treatment and immune-response alternations of ALI mice. Then we demonstrated that UC-MSCs could inhibit the inflammatory response of mouse macrophage in ALI mice, as well as enhance its IL-10 expression. We provide data to support the concept that the therapeutic capacity of UC-MSCs for ALI was primarily through paracrine secretion, particularly of prostaglandin-E2 (PGE2). Furthermore, we showed that UC-MSCs might secrete a panel of factors including GM-CSF, IL-6 and IL-13 to ameliorate ALI. Our study suggested that UC-MSCs could protect LPS-induced ALI model by immune regulation and paracrine factors, indicating that UC-MSCs should be a promising strategy for ALI/ARDS.

Acute lung injury (ALI), as the early and basic pathophysiologic change in acute respiratory distress syndrome (ARDS), is a challenging disease in clinical critical care medicine¹. ALI/ARDS is a form of severe respiratory disorders which are most commonly caused by sepsis, pneumonia, trauma and aspiration². Although many ALI/ARDS treatments have been investigated, the mortality is between 36–44% with little improvement³. In the United States, it is estimated that there are 190,600 cases and 74,000 deaths annually from ALI that results in a high cost of health-care⁴. However, the current therapeutic strategies for ALI are mainly focused on supportive treatment, and a novel effective therapy is required.

Stem cells, especially mesenchymal stem cells (MSCs), have demonstrated therapeutic potential in lung diseases⁵. MSCs are collected from bone marrow (BM), umbilical cord (UC), adipose tissue, amniotic fluid, skeletal muscle, synovial, gingiva and other tissues⁶. The bone marrow-derived mesenchymal stem cells (BM-MSCs) have presented ability in resolving lipopolysaccharide (LPS)-induced ALI, hyperoxia, pneumonia and systemic sepsis in animal models⁷. However, the clinical application of BM-MSCs is limited due to invasive sample collection, reduced cell number, proliferation, and differentiation capacity in aging donors⁸. Human umbilical cord-derived mesenchymal stem cells (UC-MSCs) are considered a better choice of MSCs for clinical application because of their easy collection, high cell vitality⁹, low immunogenicity¹⁰, and high paracrine potential for accelerating injury tissue repair processes¹¹. UC-MSCs have potential efficacy in the prevention or treatment of lung injury^{12,13}.

¹Pediatrics Research Institute, Ministry of Education Key Laboratory of Child Development and Disorders, Children's Hospital of Chongqing Medical University, Chongqing 400014, China. ²Department of Pediatrics, First Affiliated Hospital of China Medical University, Shenyang 110001, China. ³Department of Respiratory Medicine, Children's Hospital of Chongqing Medical University, Chongqing 400014, China. ⁴Department of Otorhinolaryngology, Children's Hospital of Chongqing Medical University, Chongqing 400014, China. ⁵Chongqing Engineering Research Center of Stem Cell Therapy, Chongqing 400014, China. ⁶Laboratory of Pathology, West China Hospital, Sichuan University, Chengdu 610041, China. ⁷Center for Clinical Molecular Medicine, Children's Hospital of Chongqing Medical University, Chongqing 400014, China. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to L.G. (email: 416875725@qq.com) or L.Z. (email: zoulin74@126.com)

However, the therapeutic effects and potential molecular mechanism of UC-MSCs in an ALI model remain unclear. In this study, we found therapeutic effects of UC-MSCs in classical LPS-induced ALI mice by secreting factors.

Methods

Animal. All the protocols were carried out in accordance with the approved guidelines. Eight- to ten-week-old BALB/c and C57BL/6 female mice were obtained from the Experimental Animal Center of Chongqing Medical University. All the experiments were approved by the Ethics Committee of Chongqing Medical University.

Harvesting and concentrating of UC-MSCs conditional medium. When the cell count of UC-MSCs in T25 culture flasks reached 0.5×10^6 , the cells were washed 3 times with PBS and re-cultured with 5 ml new serum-free medium for 24 h. Subsequent serum-free medium was used as UC-MSCs conditional medium. For *in vivo* experiments, 2 ml of this original medium was concentrated to an 80 μ l volume using a 3000 Da centrifugal filter (Millipore, Massachusetts, USA) by a relative slower centrifugal speed 4500 g for 35–40 minutes, which also could avoid an excessive loss of the small molecule PGE2 in UC-MSCs concentrated medium (Suppl. Fig. 4g).

Treatment of LPS-induced ALI mice with UC-MSCs or UC-MSCs conditional medium. After 1 hour of LPS exposure, BALB/c or C57/BL6 mice were intravenously administered UC-MSCs (0.5×10^6), or concentrated conditional medium (80 μ l, harvested from 0.5×10^6 UC-MSCs), or PBS (80 μ l) respectively. Mice were then monitored for post-exposure LPS for 120 h.

Protein-array of bronchoalveolar lavage fluid (BALF). Protein-array analysis of BALF was performed using the Raybiotech™ 308 biotin label-based mouse antibody array (Raybiotech, Georgia, USA) according to the manufacturer's protocol, as the previous report¹⁴. The expression levels for each protein were normalized to the mean intensity of positive control in all samples and were further analyzed using a fold change ≥ 2.0 (t-test, two-tailed, $p < 0.05$). Differentially expressed proteins were determined by Hierarchical Clustering (Gene Cluster 3.0, Tokyo, Japan) and gene-ontology (GO) classifications analysis using the DAVID online database.

Celecoxib inhibited PGE2 synthesis in UC-MSCs. To identify the role of UC-MSCs paracrine factor PGE2, we used celecoxib (Pfizer, New York, USA) to block the PGE2 synthesis in cells as previously described¹⁵. UC-MSCs were cultured with different concentrations of celecoxib or control (0.1% DMSO) in 10% FBS DMEM/F12 for 48 h. The cells were then washed with PBS three times and were harvested as celecoxib-pretreated UC-MSCs (UC-MSCs^{-PGE2}), in which PGE2 synthesis was inhibited. Alternatively, the UC-MSCs^{-PGE2} were re-cultured with new serum-free medium for 24 h, and the subsequent serum-free medium was used as the PGE2-reduced UC-MSCs conditional medium (CM^{-PGE2}). ELISA test was used to analyze the efficacy of celecoxib to inhibit the PGE2 paracrine of UC-MSCs, and MTT assay was used to investigate the effect of celecoxib on the growth of UC-MSCs.

The investigations of paracrine activities of UC-MSCs. 0.5×10^6 UC-MSCs were cultured in 2 ml ALI mice bronchoalveolar lavage fluid (BALF), added with 2 ml DMEM/F12 for 24 h, and cytokines/chemokines secreted by UC-MSCs in culture supernatant were tested by Multiplex-ELISA arrays (Millipore, Massachusetts, USA) according to the manufacturers' instructions¹⁶.

Data and statistical analysis. The values are expressed as the mean $\pm$ SE. Multiple comparisons of parametric data were performed using one-way ANOVA at specific time-points (72 h) and two-way ANOVA at different time-points (0 h, 24 h, 72 h, 120 h), followed by Bonferroni multiple test between groups. The survival curves were compared with a log-rank test (Graphpad Prism5.0, California, USA). $p < 0.05$ was considered statistically significant.

(The other detailed Supplementary Information for Methods is presented online).

Results

UC-MSCs improve survival and attenuate lung inflammation in LPS-induced ALI mice. We successfully isolated UC-MSCs from human Wharton's Jelly (Suppl. Fig. 1) and constructed a stable ALI animal model by intratracheal injection of 5 mg/kg LPS (Suppl. Fig. 2). The ALI mice were then transtracheally injected with UC-MSCs (0.5×10^6) after being administered LPS for 1 h (Suppl. Fig. 2a). The survival and weight improvement of LPS-induced ALI mice treated with UC-MSCs, both in BALB/c and C57/BL6 mice (Fig. 1a,b), indicated the significant therapeutic effects of UC-MSCs in ALI. Then we used BALB/c mice in the following experiments. The pathological damage of lung tissue from LPS-induced ALI mice treated with UC-MSCs was significantly decreased (Fig. 1c,d), especially 72 h post-treatment. The lung injury was evidently declined, meanwhile the myeloperoxidase (MPO) activity, total protein concentration, total cell and neutrophil counts in bronchoalveolar lavage fluid (BALF) were significantly reduced, confirming that UC-MSCs attenuate the lung inflammation of LPS-induced ALI mice (Fig. 1e–h).

UC-MSCs regulate the lung immune response of LPS-induced ALI mice. To understand the molecules involved in UC-MSCs treatment for LPS-induced ALI mice, we used a 308 protein array to analyze in the BALF (Suppl. Table 1). Only 32 proteins were significantly decreased, while 5 proteins were significantly increased in the LPS-induced ALI mice treated with UC-MSCs for 72 h (Fig. 2a). The gene-ontology (GO) classifications of these 37 proteins (Suppl. Table 2) showed that 17 proteins (45.95%) had a correlation with the host immune response. Among these 17 proteins, eleven (64.71%) were related to defense response and particularly 6 (35.30%)

Figure 1. Human umbilical cord-derived mesenchymal stem cells (UC-MSCs) alleviate lipopolysaccharide (LPS)-induced acute lung injury (ALI) mice by intratracheal delivery. Kaplan-Meier survival curves (a) and body weight alterations (b) of LPS-induced ALI mice after intratracheal delivery of 0.5×10^6 UC-MSCs or fibroblasts (Fb) for 120 h. Green line, BALB/C mice treated with PBS (Ctrl); Red line, BALB/C mice induced by LPS; Black line, LPS-induced BALB/C mice treated with fibroblasts; Orange line, BALB/C mice treated with UC-MSCs; Blue line, LPS-induced BALB/C mice treated with UC-MSCs; Purple line, LPS-induced C57BL mice treated with UC-MSCs. (c) Hematoxylin-eosin (HE) staining of lung sections of BALB/C mice with intratracheal LPS administration or with intratracheal LPS and UC-MSCs at different post-injection times. The lungs from LPS-induced mice treated with UC-MSCs exhibited decreased intra-alveolar edema. Magnifications $\times 100$ or $\times 400$. In (a) to (c), $n = 24$ for each group. (d) Lung injury scores of LPS-induced ALI mice with or without UC-MSCs treatment. The detection of myeloperoxidase (MPO) activity (e), total protein concentration (f), total cell count (g) and neutrophil count (h) in bronchoalveolar lavage fluid (BALF) in LPS-induced ALI mice with UC-MSCs treatment. In (d) to (h), data are the mean $\pm$ SEM, $n = 8-10$ at each time point. * $p < 0.05$ or ** $p < 0.01$ vs. control mice (Ctrl); # $p < 0.05$ or ## $p < 0.01$ vs. LPS-induced ALI mice (ALI).

Figure 2. UC-MSCs regulate the protein expression profiles of BALF in LPS-induced ALI mice. (a) The heat map of the altered protein files over twofold ($p < 0.05$) in 308 mouse proteins from BALF in LPS-induced ALI mice treated with UC-MSCs for 72 h by cluster analysis ($n = 3$ each group). Only 32 proteins were significantly decreased and 5 related proteins were significantly increased. The increased proteins are shown in progressively brighter shades of red, and decreased proteins are shown in progressively darker shades of green. (b) The analysis of gene-ontology (GO) classifications on 37 differentially expressed proteins. It showed that 17 proteins (45.95%) had a correlation with immune response, and among these 17 proteins, 11 proteins (64.71%) were related to defense response, particularly 6 proteins (35.30%) including TNF- α , IL-1 β , IL-6, IL-10, CCL17 and CCL22, were associated with the immune subtypes of macrophages. M1 macrophage, classically activated macrophage. M2 macrophage, alternatively activated macrophage. In (c) to (d), the concentration of TNF- α and IL-10 in BALF, detected by ELISA from LPS-induced ALI mice treated with UC-MSCs. In (b) to (d), values are the means $\pm$ SE. $n = 8$ for each group. * $p < 0.05$ or ** $p < 0.01$ vs. control mice (Ctrl); # $p < 0.05$ or ## $p < 0.01$ vs. LPS-induced ALI mice (ALI).

were associated with the immune subtypes of macrophages (Suppl. Fig. 3a), indicating that UC-MSCs regulate the lung immune response of LPS-induced ALI mice (Fig. 2b).

We also found that the markers of classically activated macrophages (M1, CM), TNF- α , IL-1 β and IL-6 were significantly down-regulated by UC-MSCs treatment¹⁷. More interestingly, in 5 up-regulated proteins, 3 proteins including IL-10, CCL17 and CCL22 were reported to the markers of alternatively activated macrophages (M2, AM)¹⁸, especially IL-10 had been proved to be a potent anti-inflammatory cytokine to protect lung injury¹⁹. The results of the q-PCR (Suppl. Fig. 3b) and ELISA assays (Fig. 2c,d) were consistent with those of the protein microarray, suggesting that UC-MSCs treatment would polarize the lung macrophages into M2 macrophages in ALI mice.

UC-MSCs modulate polarization of mouse macrophages *in vivo* and *in vitro*. To explore the effects of UC-MSCs on macrophage transition, we examined the lung macrophages in UC-MSCs-treated ALI mice by flow cytometry. The results demonstrated that the total number of lung macrophages (Marker, F4/80 and CD11b) was significantly reduced in UC-MSCs-treated ALI mice, but the IL-10 positive lung macrophages (Marker, IL-10) were up-regulated (Fig. 3a,b, $p < 0.05$), indicating that UC-MSCs modulate the immune response of lung macrophages in ALI mice by promoting lung macrophages to secrete more IL-10 to protect ALI mice. Therefore, UC-MSCs might alleviate the inflammatory response in ALI mice by mediating macrophages transition.

We then co-cultured LPS-stimulated mouse macrophage RAW264.7 cells with UC-MSCs to verify their relationship *in vitro*. The morphology of macrophages was significantly altered by LPS stimulation, including an enlarged cell body, irregular shape, and a large number of pseudopodia. Interestingly, when co-cultured with UC-MSCs, the morphology of LPS-stimulated RAW264.7 cells was presented as normal RAW264.7 cells (Fig. 3c,d). In addition, TNF- α expression in LPS-stimulated macrophages was significantly decreased by UC-MSCs co-culture, while IL-10 expression was increased (Fig. 3e,f), indicating that UC-MSCs promote the M2 polarization of macrophages.

UC-MSCs protect LPS-induced ALI mice *via* paracrine activity. Given that MSCs generate their therapeutic action through cell engraftment differentiation to directly repair injured tissues or through paracrine to facilitate self-healing of tissues²⁰, we labeled UC-MSCs with DAPI *in vitro* and then tracked the labeled UC-MSCs in ALI mice post-injection at different times. The results showed that UC-MSCs were concentrated in the trachea 0 h post-injection, continuously infiltrated the lung parenchyma through the tracheal wall 12 h post-injection, and populated the whole lung parenchyma 24 h post-injection. However, UC-MSCs were still localized in the lung parenchyma 48 h post-injection, but the intensity of fluorescence was much lower than that of 24 h post-injection. Additionally, the DAPI-labeled UC-MSCs almost disappeared in the fluorescent microscope 72 h post-injection (Fig. 4a,b), which indicated that the time of UC-MSCs engraftment and therapeutic effects was 24 h–48 h post-injection, suggesting that UC-MSCs secrete soluble factors responsible for relieving ALI mice, rather than differentiating target cells and replacing.

To further confirm the paracrine activity of UC-MSCs in LPS-induced ALI mice, we applied the concentrated conditioned medium from UC-MSCs to treat LPS-induced ALI mice. The results demonstrated that the conditional medium of UC-MSCs could improve the survival and body weight (Fig. 4c,d), and ameliorate the lung injury of LPS-induced ALI mice (Fig. 4e,f), indicating that UC-MSCs secrete soluble factors responsible for ALI mice treatment.

UC-MSCs protect ALI mice by secretion of PGE2. MSCs are reported to release multiple paracrine factors, especially prostaglandin-E2 (PGE2), secreted from BM-MSCs, with powerful immunological regulation functions²¹, which cause macrophages to secrete IL-10 to accelerate recovery of damaged tissues^{22,23}. However, little is known about the ability of paracrine PGE2 in UC-MSCs. We analyzed the data of the gene expression array (GSE48022) from the GEO database and found that the gene expression of prostaglandin-endoperoxide synthase 2 (PTGS2/Cox-2), the rate-limiting enzyme in the biosynthesis of PGE2, is much higher in UC-MSCs than that in BM-MSCs (Suppl. Fig. 4a), indicating the powerful ability of synthesis and secretion of PGE2 in UC-MSCs. To further verify the secretion ability of PGE2 and its biological function, we analyzed the concentration of the small molecules in the UC-MSCs conditional medium, and found that the expression of PGE2 was several folds higher than that in LPS-stimulated macrophage medium. In addition, when co-cultured with LPS-stimulated macrophages, the production of PGE2 in UC-MSCs was significantly enhanced (Suppl. Fig. 4b), suggesting that the key role of PGE2 secreted by UC-MSCs for macrophages transition.

To confirm the effect of PGE2 secreted by UC-MSCs on macrophages, we applied celecoxib, a selective inhibitor of PGE2 biosynthesis, to approve the role of PGE2 in UC-MSCs-treated macrophages and mice. The results showed that a 1 $\mu\text{mol/L}$ concentration of celecoxib significantly reduced PGE2 expression in UC-MSCs without drug toxicity (Suppl. Fig. 4c,d). Then we harvested the 1 $\mu\text{mol/L}$ celecoxib-pretreated UC-MSCs (UC-MSCs^{-PGE2}), which PGE2 synthesis in cells was inhibited, and the conditional medium from celecoxib-pretreated UC-MSCs (UC-MSCs^{-PGE2}), which PGE2 concentrations was reduced. And we added them into LPS-stimulated macrophages and measured the TNF- α and IL-10 concentrations in the culture medium. The results showed that both the celecoxib-pretreated UC-MSCs (UC-MSCs^{-PGE2}) and the conditional medium from celecoxib-pretreated UC-MSCs (CM^{-PGE2}) had a weaker capacity for inhibiting TNF- α (Suppl. Fig. 4e) and stimulating IL-10 (Suppl. Fig. 4f) in LPS-stimulated macrophages, indicating that UC-MSCs increased PGE2 synthesis and secretion when interacting with LPS-stimulated macrophages, resulting in the inhibitory effects on TNF- α and stimulatory effects on IL-10.

To further investigate the importance of PGE2 expression in the treatment of UC-MSCs in ALI mice, we treated mice with the celecoxib-pretreated UC-MSCs (UC-MSCs^{-PGE2}) or the conditional medium from celecoxib-pretreated UC-MSCs (CM^{-PGE2}) and observed the lung pathological alterations and the concentration of soluble factors in the BALF of these mice. The results (Fig. 5a–f) showed that the therapeutic effect of UC-MSCs on LPS-induced ALI mice declined with the decrease of PGE2 production, demonstrating that the rapid high expression of PGE2 secreted by UC-MSCs mediated the protection of UC-MSCs on LPS-induced ALI mice.

The paracrine molecules of UC-MSCs in relief of ALI. To further investigate the paracrine secretions of UC-MSCs in this mouse model of ALI, we found that a panel of the molecules secreted by the stem cells was increased significantly when cultured in BALF from ALI mouse compared with control mice (Fig. 6a). This indicated that UC-MSCs could secrete a group of paracrine (including GM-CSF, IL-6 and IL-13) responsible for ameliorating ALI in our mouse model. All the paracrine molecules are summarized in Fig. 6b.

Figure 3. UC-MSCs modulate the immune response of macrophages in LPS-induced ALI mice. (a) The representative flow cytometric figures of lung CD11b⁺ F4/80⁺ and IL10⁺ macrophages. The lung tissues from LPS-induced ALI mice were harvested, grinded, filtered, and washed to obtain the monotypes. The monotypes were then incubated with anti-CD11b, anti-F4/80 or anti-IL-10 and were analyzed by flow cytometry. The cells were gated on anti-CD11b and anti-F4/80 simultaneously, and followed by gating on anti-IL-10. (b) The statistical analysis of lung CD11b⁺ F4/80⁺ macrophages and IL10⁺ macrophages were performed by flow cytometry. (c) The morphologic alterations of mouse macrophage RAW264.7 cells were stimulated with LPS and were co-cultured with UC-MSCs, resulting in enlarged cell body, irregular shape, and a large number of pseudopodia. (d) The statistical average surface area of a single macrophage was analyzed by Image Pro-Plus (IPP) 6.0 software, and the cell surface area in macrophage Raw264.7 cells was taken as 1.0 fold. The secretion concentration of TNF-α (e) and IL-10 (f) was measured by ELISA. Values are the means ± SE. n = 6–8 for each group. **p* < 0.05 or ***p* < 0.01 vs. control mice (Ctrl) or macrophages. #*p* < 0.05 or ##*p* < 0.01 vs. LPS-induced ALI mice (ALI) or LPS-stimulated macrophages. Mac., macrophages.

Figure 4. UC-MSCs ameliorate the lung injury of LPS-induced ALI mice through paracrine role. (a) Representative images of UC-MSCs stained with DAPI and imaged by fluorescence microscopy (left panel) or by light microscopy (right panel). (b) The fluorescence images of DAPI-stained UC-MSCs in lung cryosections from LPS-induced ALI mice at different post-delivery times. After UC-MSCs were intratracheally delivered, DAPI-labeled UC-MSCs were presented along the bronchial wall at post-deliver 0 h, they were accumulated in lung tissue 12 h–24 h post-delivery, rapidly decreasing in number 48 h post-delivery, and becoming rare 72 h to 120 h post-delivers. Magnification, $\times 100$. $n = 24$ each group. (c–f) Injection of conditional medium (CM) of UC-MSCs effectively protects LPS-induced ALI mice. Kaplan-Meier survival curves (c) and body weight alteration (d) of mice, $n = 24$ for each group. HE staining of lung sections (e) and lung injury scores (f) in LPS-induced ALI mice treated with CM from the same amount of UC-MSCs or fibroblasts (Fb). Magnification, $\times 100$. In (f), Values are the means $\pm$ SE. $n = 7$ –9 for each group. $**p < 0.01$, LPS-induced ALI mice vs. LPS-induced ALI mice treated with conditional medium from UC-MSCs. CM, conditional medium.

Figure 5. PGE2 secreted by UC-MSCs ameliorate the lung injury of LPS-induced ALI. Kaplan-Meier survival curves (a), body weight change (b), HE staining of lung sections (c), and lung injury scores (d) of LPS-induced ALI mice with a variety of treatments. The protection of UC-MSCs and UC-MSCs conditional medium (CM) on mice was weakened by reduced PGE2 levels. $n = 24$ for each group. Magnification, $\times 100$. The concentration of TNF- α (e) and IL-10 (f) detected by ELISA in the lung BALF of mice with several treatments. In (c–e), values are the means $\pm$ SE, $n = 7$ –9 at each point. * $p < 0.05$ or ** $p < 0.01$, LPS-induced ALI mice treated with normal UC-MSCs (ALI + UC-MSCs) vs. LPS-induced ALI mice treated with celecoxib-pretreated UC-MSCs (ALI + UC-MSCs^{-PGE2}). # $p < 0.05$ or ## $p < 0.01$, LPS-induced ALI mice treated with conditional medium from normal UC-MSCs (ALI + CM) vs. LPS-induced ALI mice treated with conditional medium from celecoxib-pretreated UC-MSCs (ALI + CM^{-PGE2}).

Figure 6. The multiple factors secreted by UC-MSCs may contribute to lung injury amelioration. **(a)** The paracrine factors of UC-MSCs were analyzed by using Multiple-ELISA assay and expressed as a heatmap. The increased proteins are shown in progressively brighter shades of red, and decreased proteins are shown in progressively darker shades of green. $^*p < 0.05$ or $^{**}p < 0.01$, UC-MSCs + BALF (ALI) vs. UC-MSCs + BALF (Ctrl), $n = 8-9$. **(b)** The scheme of factors secreted by UC-MSCs in ALI amelioration.

Discussion

In this study, we illustrated that intratracheal injection of UC-MSCs increased the survival rate of LPS-induced ALI mice and significantly reduced pulmonary inflammation. The soluble factors secreted by UC-MSCs, in particular, PGE2, GM-CSF, IL-6 and IL-13 may be the therapeutic basis of UC-MSCs for ALI.

ALI usually develops in patients with predisposing conditions that induce systemic inflammatory response, such as sepsis, pneumonia, major trauma, multiple transfusions, aspiration, and acute pancreatitis²⁴. Among them, sepsis is the major etiology for ALI development, wherein Gram-negative bacteria are a prominent cause²⁵. LPS, the main component of the outer membranes of Gram-negative bacteria, is the most important antigen that promotes the development of ALI²⁶. Therefore, the LPS-induced animal models could highlight ways to explore mechanisms of multiple diseases and provide useful information on the discovery of novel biomarkers and drug targets²⁷. In our study, the animal model induced by intra-tracheal administration of LPS successfully duplicated human ALI. The LPS-induced ALI mouse model had a classical pulmonary inflammatory state of the lungs between 48 h and 120 h, but after the administration of UC-MSCs, a decreased lung pathological lesion and inflammatory response was observed at 72 h after UC-MSCs injection, including the lung MPO activity, total protein concentration, lung neutrophil infiltration intensity, the expression of various pro-inflammation cytokines in BALF and the immune response of LPS-induced ALI mice. The results were consistent with previous studies on the beneficial effects of BM-MSCs in resolving ALI^{5,7,28,29}, proving that the intra-tracheal injection of UC-MSCs had protective effects in attenuating lung injury and inhibiting the lung inflammation of LPS-induced ALI mice.

Although there are therapeutic effects of MSCs for different ALI models²⁸, the precise molecular mechanisms of MSCs action in ALI therapy are still unclear. Some studies found that BM-MSCs could mobilize into circulation in mice after LPS exposure, accumulate within the inflammatory site and differentiate to form endothelial and epithelium-like phenotype cells, like ACE1 and ACE2^{29,30}. Some studies postulated that soluble factors secreted by BM-MSCs played a vital role in wound repair, most likely through their anti-inflammatory, anti-apoptotic, angiogenic and immunomodulatory properties^{31,32}. To determine the UC-MSCs' location and their potential action principles, we located the DAPI-labeled UC-MSCs in ALI mice and found that UC-MSCs concentrated in the mouse lung within 24 h–48 h. Such a short period of UC-MSCs action suggested that paracrine is the dominant mechanism rather than cell differentiation and replacement in ALI repair. This hypothesis was further confirmed by UC-MSCs conditional medium ameliorating LPS-induced ALI mice. Our results revealed that the paracrine

capacity of UC-MSCs had a primary functional and survival benefit in LPS-induced ALI. The secreted soluble factors in the UC-MSCs conditional medium are key for relieving ALI.

PGE₂, one of the UC-MSCs paracrine factors, has demonstrated beneficial effects in LPS-induced ALI mice, which could modulate macrophage responses and attenuate the LPS-induced inflammation. Previous studies demonstrated that an intravenous injection of BM-MSCs can beneficially modulate the host immune response, by increasing the release of PGE₂ from the BM-derived MSCs, acting on the EP₂ and EP₄ receptors of the macrophages, increasing the production of IL-10 in the macrophages and reducing inflammation^{22,23,33}. Here, we found that UC-MSCs secrete a large amount of PGE₂; however, with the decline of the PGE₂ level in UC-MSCs and UC-MSCs conditional medium, both showed a lower ability to promote the production of IL-10 in LPS-stimulated RAW264.7 cells and in LPS-induced ALI mice. The increasing IL-10 level in the lungs suppresses LPS-induced ALI by reducing induction of pro-inflammatory mediators and chemokines, reducing mast-cell activation and NF- κ B activation, and down-modulating pathways generating oxidative stress³⁴. Activation of MSCs impacts their therapeutic abilities in lung injury by increasing IL-10 and IL-1 α levels. Thus, our data suggested that UC-MSCs-secreted PGE₂ mediates the protective function of UC-MSCs in lung injury.

Moreover, we observed that UC-MSCs cultured in BALF from ALI mouse could have a high-level expression of some paracrine including GM-CSF, G-CSF, IL-6, IL-13 and others, as shown in Fig. 6a. Studies found Both GM-CSF and G-CSF belong to the family of colony-stimulating factors, playing a key role in host defense by regulating proliferation and differentiation of granulocytes and mononuclear phagocytes³⁵. In response to pulmonary infection, local delivery of GM-CSF could increase the number of alveolus exudate macrophages derived from blood monocytes, to improve the host defense against microbial pathogens³⁶. During recently years, the benefits of GM-CSF application have been demonstrated to protect lung injury in sepsis³⁷. As for IL-6, it is a multifunctional cytokine involved in the regulation of complex cellular processes such as proliferation and differentiation, and act as a key player during immune and inflammation response³⁸. Recently, IL-6 is found to promote survival of human CD11b⁺ peripheral blood mononuclear cells and induce M2-type macrophage polarization³⁹. In addition to IL-6, the capacity of IL-13 on the transformation of macrophages into M2 macrophages had been confirmed in many previous studies⁴⁰. Importantly, M2 macrophages are considered to play an essential role in regulation of immune response and resolution of lung inflammation⁴¹. Furthermore, studies also found that MSCs could induce M2 macrophages polarization through other soluble factors, such as TGF- β ¹⁴², TSG-6⁴³ and IDO⁴⁴, which could get together to promote disease amelioration⁴⁵. Therefore, we deduced that UC-MSC could modulate immune responses of macrophages against LPS-induced lung injury not just by PGE₂, but rather *via* a panel of secretory factors, such as GM-CSF, IL-6 and IL-13 also contributed to the treatment of ALI mice.

In conclusion, our results demonstrated that UC-MSCs significantly attenuate inflammatory responses and lung injury in LPS-induced ALI mice. The protective effect on ALI was due to secreted a panel of secretory factors, suggesting that UC-MSCs could be a novel therapeutic choice for ALI.

References

- Force, A. D. T. *et al.* Acute respiratory distress syndrome: the Berlin Definition. *JAMA* **307**, 2526–2533 (2012).
- Han, S. *et al.* The acute respiratory distress syndrome: from mechanism to translation. *J Immunol* **194**, 855–860 (2015).
- Phua, J. *et al.* Has mortality from acute respiratory distress syndrome decreased over time?: A systematic review. *Am J Respir Crit Care Med* **179**, 220–227 (2009).
- Rubenfeld, G. D. *et al.* Incidence and outcomes of acute lung injury. *N. Engl. J. Med.* **353**, 1685–1693 (2005).
- Antunes, M. A. *et al.* Mesenchymal stem cell trials for pulmonary diseases. *J Cell Biochem* **115**, 1023–1032 (2014).
- Lv, F. J. *et al.* Concise review: the surface markers and identity of human mesenchymal stem cells. *Stem cells* **32**, 1408–1419 (2014).
- Akram, K. M. *et al.* Mesenchymal stem cell therapy and lung diseases. *Adv Biochem Eng Biotechnol* **130**, 105–129 (2013).
- Panepucci, R. A. *et al.* Comparison of gene expression of umbilical cord vein and bone marrow-derived mesenchymal stem cells. *Stem cells* **22**, 1263–1278 (2004).
- Li, G. *et al.* Comparative proteomic analysis of mesenchymal stem cells derived from human bone marrow, umbilical cord, and placenta: implication in the migration. *Proteomics* **9**, 20–30 (2009).
- Lee, M. *et al.* Low immunogenicity of allogeneic human umbilical cord blood-derived mesenchymal stem cells *in vitro* and *in vivo*. *Biochem Biophys Res Commun* **446**, 983–989 (2014).
- El Omar, R. *et al.* Umbilical cord mesenchymal stem cells: the new gold standard for mesenchymal stem cell-based therapies? *Tissue Eng Part B Rev* **20**, 523–544 (2014).
- Moodley, Y. *et al.* Human umbilical cord mesenchymal stem cells reduce fibrosis of bleomycin-induced lung injury. *Am J Pathol* **175**, 303–313 (2009).
- Liu, L. *et al.* Intranasal versus intraperitoneal delivery of human umbilical cord tissue-derived cultured mesenchymal stromal cells in a murine model of neonatal lung injury. *Am J Pathol* **184**, 3344–3358 (2014).
- Tang, Y. L. *et al.* Hypoxic preconditioning enhances the benefit of cardiac progenitor cell therapy for treatment of myocardial infarction by inducing CXCR4 expression. *Circ Res* **104**, 1209–1216 (2009).
- Bowman, C. C. *et al.* Cyclooxygenase-2-Mediated Prostaglandin E₂ Production in Mesenteric Lymph Nodes and in Cultured Macrophages and Dendritic Cells after Infection with Salmonella. *J Immunol* **172**, 2469–2475 (2004).
- Trune, D. R. *et al.* Simultaneous measurement of multiple ear proteins with multiplex ELISA assays. *Hear Res* **275**, 1–7 (2011).
- Xie, G. *et al.* p-Cymene protects mice against lipopolysaccharide-induced acute lung injury by inhibiting inflammatory cell activation. *Molecules* **17**, 8159–8173 (2012).
- Rajaiah, R. *et al.* Dissociation of Endotoxin Tolerance and Differentiation of Alternatively Activated Macrophages. *J Immunol* **190**, 4763–4772 (2013).
- Hoegl S, B. K. *et al.* Inhaled IL-10 reduces biotrauma and mortality in a model of ventilator-induced lung injury. *Respir Med* **103**, 463–470 (2009).
- Inamdar, A. C. *et al.* Mesenchymal stem cell therapy in lung disorders: pathogenesis of lung diseases and mechanism of action of mesenchymal stem cell. *Exp Lung Res* **39**, 315–327 (2013).
- Kalinski, P. Regulation of immune responses by prostaglandin E₂. *J Immunol* **188**, 21–28 (2012).
- Ylostalo, J. H. *et al.* Human mesenchymal stem/stromal cells cultured as spheroids are self-activated to produce prostaglandin E₂ that directs stimulated macrophages into an anti-inflammatory phenotype. *Stem cells* **30**, 2283–2296 (2012).
- Nemeth, K. *et al.* Bone marrow stromal cells attenuate sepsis via prostaglandin E₂-dependent reprogramming of host macrophages to increase their interleukin-10 production. *Nat Med* **15**, 42–49 (2009).

24. Arthur, P. & Wheeler, G. R. B. Acute lung injury and the acute respiratory distress syndrome: a clinical review. *Lancet* **369**, 1553–1565 (2007).
25. Sheu, C. C. *et al.* Clinical characteristics and outcomes of sepsis-related vs non-sepsis-related ARDS. *Chest* **138**, 559–567 (2010).
26. Shi, D. *et al.* Protective effects and mechanisms of mogroside V on LPS-induced acute lung injury in mice. *Pharm Biol* **52**, 729–734 (2014).
27. Chen, H., B. C. *et al.* The value of the lipopolysaccharide induced acute lung injury. *Expert Rev Respir Med* **4**, 773–783 (2010).
28. Monsel, A., Z. Y. *et al.* Therapeutic Effects of Human Mesenchymal Stem Cell-derived Microvesicles in Severe Pneumonia in Mice. *Am J Respir Crit Care Med* **192**, 324–336 (2015).
29. Rojas, M. *et al.* Bone marrow-derived mesenchymal stem cells in repair of the injured lung. *Am J Respir Cell Mol Biol* **33**, 145–152 (2005).
30. Kotton, D. N., M., B. *et al.* Bone marrow-derived cells as progenitors of lung alveolar epithelium. *Development* **128**, 5181–5188 (2001).
31. Kawai, T. *et al.* Secretomes from bone marrow-derived mesenchymal stromal cells enhance periodontal tissue regeneration. *Cytotherapy* **17**, 369–381 (2015).
32. Waszak, P. *et al.* Preconditioning enhances the paracrine effect of mesenchymal stem cells in preventing oxygen-induced neonatal lung injury in rats. *Stem Cells and Dev* **21**, 2789–2797 (2012).
33. Zhang, Y. *et al.* Inhibition of the prostaglandin-degrading enzyme 15-PGDH potentiates tissue regeneration. *Science* **348**, aaa2340 (2015).
34. Hiroshima, Y. *et al.* S100A8 induces IL-10 and protects against acute lung injury. *J Immunol* **192**, 2800–2811 (2014).
35. Hamilton, J. A. Colony-stimulating factors in inflammation and autoimmunity. *Nat Rev Immunol* **8**, 533–544 (2008).
36. Steinwede, K. *et al.* Local delivery of GM-CSF protects mice from lethal pneumococcal pneumonia. *J Immunol* **187**, 5346–5356 (2011).
37. Mathias, B. *et al.* A Review of GM-CSF Therapy in Sepsis. *Medicine (Baltimore)* **94**, e2044 (2015).
38. Dittrich, A. *et al.* Systems biology of IL-6, IL-12 family cytokines. *Cytokine Growth Factor Rev* **26**, 595–602 (2015).
39. Zhang, Q. Z. *et al.* Human gingiva-derived mesenchymal stem cells elicit polarization of m2 macrophages and enhance cutaneous wound healing. *Stem cells* **28**, 1856–1868 (2010).
40. Martinez, F. O. *et al.* Alternative activation of macrophages: an immunologic functional perspective. *Annu Rev Immunol* **27**, 451–483 (2009).
41. Aggarwal, N. R. *et al.* Diverse macrophage populations mediate acute lung inflammation and resolution. *Am J Physiol Lung Cell Mol Physiol* **306**, L709–725 (2014).
42. Salazar, K. D. *et al.* Mesenchymal stem cells produce Wnt isoforms and TGF-beta1 that mediate proliferation and procollagen expression by lung fibroblasts. *Am J Physiol Lung Cell Mol Physiol* **297**, L1002–1011 (2009).
43. Choi, H. *et al.* Anti-inflammatory protein TSG-6 secreted by activated MSCs attenuates zymosan-induced mouse peritonitis by decreasing TLR2/NF-kappaB signaling in resident macrophages. *Blood* **118**, 330–338 (2011).
44. Francois, M. *et al.* Human MSC suppression correlates with cytokine induction of indoleamine 2,3-dioxygenase and bystander M2 macrophage differentiation. *Mol Ther* **20**, 187–195 (2012).
45. Zheng, G. *et al.* Mesenchymal Stromal Cells Affect Disease Outcomes via Macrophage Polarization. *Stem Cells Int* **2015**, 989473 (2015).

Acknowledgements

We appreciated Dangsheng Li, Ph.D. for revising and editing this manuscript. Prof. Lin Zou was funded from National Natural Science Foundation Committee of China (81373444, 81570142). Prof. Zhou Fu was funded by the key grant from Chongqing Science and Technology Commission (Cstc2014yykfc10003). Dr. Daiyin Tian was supported from the National Natural Science Foundation Committee of China (81000002).

Author Contributions

H.Z. contributed to study design, performance of all experiments, data analysis and interpretation, and writing of the manuscript. Y.X. and L.G. contributed to study design, performance of the majority of experiments, data analysis and interpretation, and editing of the manuscript. Y.X., R.Z. and L.W. contributed to performance of partial experiments. D.T., T.W., J.D., H.Y., H.J., K.Y. and E.L. contributed to revising of the manuscript. Y.S. contributed to study design, and editing of the manuscript. Z.F. contributed to study design, financial support, data analysis and interpretation, editing and revising of the manuscript, and final approval. L.Z. contributed to study design, financial support, data analysis and interpretation, writing, editing and revising of the manuscript, and final approval.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhu, H. *et al.* Therapeutic Effects of Human Umbilical Cord-Derived Mesenchymal Stem Cells in Acute Lung Injury Mice. *Sci. Rep.* **7**, 39889; doi: 10.1038/srep39889 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017