

SCIENTIFIC REPORTS



OPEN

Diverse strategies of O₂ usage for preventing photo-oxidative damage under CO₂ limitation during algal photosynthesis

Received: 29 April 2016

Accepted: 15 December 2016

Published: 20 January 2017

Ginga Shimakawa¹, Yusuke Matsuda², Kensuke Nakajima², Masahiro Tamoi³, Shigeru Shigeoka³ & Chikahiro Miyake¹

Photosynthesis produces chemical energy from photon energy in the photosynthetic electron transport and assimilates CO₂ using the chemical energy. Thus, CO₂ limitation causes an accumulation of excess energy, resulting in reactive oxygen species (ROS) which can cause oxidative damage to cells. O₂ can be used as an alternative energy sink when oxygenic phototrophs are exposed to high light. Here, we examined the responses to CO₂ limitation and O₂ dependency of two secondary algae, *Euglena gracilis* and *Phaeodactylum tricornutum*. In *E. gracilis*, approximately half of the relative electron transport rate (ETR) of CO₂-saturated photosynthesis was maintained and was uncoupled from photosynthesis under CO₂ limitation. The ETR showed biphasic dependencies on O₂ at high and low O₂ concentrations. Conversely, in *P. tricornutum*, most relative ETR decreased in parallel with the photosynthetic O₂ evolution rate in response to CO₂ limitation. Instead, non-photochemical quenching was strongly activated under CO₂ limitation in *P. tricornutum*. The results indicate that these secondary algae adopt different strategies to acclimatize to CO₂ limitation, and that both strategies differ from those utilized by cyanobacteria and green algae. We summarize the diversity of strategies for prevention of photo-oxidative damage under CO₂ limitation in cyanobacterial and algal photosynthesis.

Air consists of 21% O₂, the concentration of which increased during the evolution of oxygenic phototrophs, in particular the oceanic cyanobacteria, around 2.3 billion years ago¹. Due to its electron configuration, O₂ has a very high oxidizing potential and is the final electron acceptor in aerobic respiratory electron transport.

Oxygenic photosynthesis uses photon energy to produce sugar from CO₂ and H₂O, and releases O₂ as a waste product. Two photosystems, PSI and PSII, play central roles in this process, which involves an electron transport system located on thylakoid membranes. The reaction centers, P700 and P680, are photo-oxidized via light-harvesting pigments such as chlorophyll (Chl). The oxidized P700 in PSI accepts electrons from PSII via plastoquinone, the cytochrome b₆/f complex, and plastocyanin (or cytochrome c₆). This electron transport is accompanied by the generation of a proton gradient across the membranes (ΔpH), allowing the production of ATP by ATP synthase². At the acceptor side of PSI, NADP⁺ is reduced to NADPH by accepting electrons from P700 through ferredoxin and ferredoxin-NADP⁺ reductase. O₂ is produced via the oxidation of H₂O by oxidized P680 in the luminal side of PSII. Together, these reactions are termed ‘photosynthetic electron transport’, and are the source of chemical energy in the forms NADPH and ATP, which are used for CO₂ assimilation in the Calvin-Benson cycle³.

The production and consumption of energy by photosynthetic electron transport and the Calvin-Benson cycle becomes unbalanced without sufficient CO₂ (CO₂-limited photosynthesis; Fig. 1). Excess photon energy causes the production of reactive oxygen species (ROS), which trigger oxidative damage to PSII and PSI, so-called photoinhibition^{4–7}.

¹Department of Biological and Environmental Science, Faculty of Agriculture, Graduate School of Agricultural Science, Kobe University, 1-1 Rokkodai, Nada, Kobe 657-8501, Japan. ²Research Center for the Development of Intelligent Self-Organized Biomaterials, Research Center for Environmental Bioscience, Department of Bioscience, Kwansei-Gakuin University, 2-1 Gakuen, Sanda, Hyogo 669-1337, Japan. ³Department of Advanced Bioscience, Faculty of Agriculture, Kinki University, 3327-204 Nakamachi, Nara 631-8505, Japan. Correspondence and requests for materials should be addressed to C.M. (email: cmiyake@hawk.kobe-u.ac.jp)




Figure 1. A simple model of photosynthesis. Orange line shows photon energy and red line shows photosynthetic activity. Photosynthesis actively occurs when sufficient CO_2 is available (CO_2 -saturated photosynthesis, double-headed black arrow). Under CO_2 -limited photosynthesis (double-headed blue arrow), excess photon energy accumulates in a photosynthetic electron transport system located on the thylakoid membranes, which causes the production of reactive oxygen species.

It is broadly accepted that O_2 is essential, not only as the respiratory electron acceptor, but also as an electron sink for various reactions of photosynthesis: O_2 -dependent alternative electron flow (AEF)⁸, including the Mehler-ascorbate peroxidase (MAP) pathway^{9,10}, singlet O_2 production in PSII⁵, flavodiroxin protein (FLV) reactions^{11–13}, mitochondrial respiration¹⁴, and plastidial (or cyanobacterial) respiration^{15,16}. Also, photorespiration can be explained as an O_2 -dependent AEF in the broad sense^{17–19}. A large O_2 -dependent AEF that replaces photosynthesis can alleviate photoinhibition by dissipating excess energy to O_2 ^{11–13,17–22}. Recently, we showed that an O_2 -dependent AEF is essential for the oxidation of P700 under CO_2 limitation to protect PSI against photo-oxidative damage in the cyanobacterium *Synechococcus* sp. PCC 7002 (S. 7002)^{22,23}. That is, oxygenic phototrophs accessed O_2 to prevent photo-oxidative damage derived from O_2 . However, both the magnitude and the molecular mechanisms of O_2 -dependent AEF vary across species in oxygenic phototrophs^{12,13,18,22,23}.

There are alternative mechanisms, which do not depend on O_2 , that dissipate excess photon energy in oxygenic phototrophs. First, during Chl fluorescence measurements, non-photochemical quenching (NPQ) is observed as a decrease in maximum Chl fluorescence yields (F_m or F_m'). Simply put, NPQ is a process of heat dissipation of photon energy around PSII. The molecular mechanisms of NPQ in various oxygenic phototrophs are diverse and include the xanthophyll cycle, light-harvesting complex II aggregation, and state transition, some of which are activated by ΔpH ^{24,25}. The degree of induced NPQ varies widely depending on growth and measurement conditions^{24,25}. Second, the electron transport in the cytochrome b_6/f complex has suppressed sensitivity to ΔpH ²⁶ or reduced plastoquinone pool²⁷, which is expected to oxidize PSI to alleviate the production of ROS by PSI owing to P700 quenching. Finally, cyclic electron transport (CET) around PSI supports the formation of ΔpH to induce NPQ and down-regulate the electron transport in the cytochrome b_6/f complex²⁸. We note that CET is defined as an AEF but does not require O_2 . In prokaryotic and eukaryotic algae, CET around PSI is suggested to be driven in several pathways, including chloroplast NADPH dehydrogenase (NDH) 1 and 2, and an elusive ferredoxin-plastoquinone reductase²⁹. Further, CET around PSII has been found in the green alga *Chlorella pyrenoidosa*^{30,31}.

In this study, we measured responses to CO_2 limitation of the cyanobacterium *Synechococcus elongatus* PCC 7942 (S. 7942) and two secondary algae, the Euglenoid *Euglena gracilis* and the pennate marine diatom *Phaeodactylum tricornutum*. We aimed to elucidate the diversity of mechanisms to utilize O_2 as an alternative electron acceptor in photosynthetic electron transport to CO_2 in cyanobacteria and algae. Cyanobacteria are known as the ancestors of chloroplasts, and have evolved into the chloroplasts of various photosynthetic eukaryotes via endosymbiosis. In contrast, the secondary algae are known to be the products of two endosymbiotic events and to have evolved from cyanobacteria along different lineages from that of land plants³². Chloroplasts of *E. gracilis* are possibly derived from a green plastid-containing organism and are surrounded by a triple, rather than a double, membrane as found in vascular plants and green algae³², which possess Chl *a* and *b* as light-harvesting pigments. However, the relative content of Chl *b* to Chl *a* in *E. gracilis* is less than that in vascular plants³³. Conversely, *P. tricornutum* harbors chloroplasts that possibly originated from red algae, and are surrounded by a quadruple membrane³². The light-harvesting complex of *P. tricornutum* has fucoxanthin-Chl *a/c* binding proteins containing the carotenoids diadinoxanthin and diatoxanthin, which are involved in NPQ³⁴.

Results

Responses of photosynthetic electron transport to CO_2 limitation in S. 7942, E. gracilis, and P. tricornutum. We measured O_2 and relative Chl fluorescence in S. 7942, *E. gracilis*, and *P. tricornutum* using an O_2 electrode and a PAM fluorometer to evaluate the responses of photosynthetic electron transport.

We estimated AEF activities in S. 7942, *E. gracilis*, and *P. tricornutum* using the relationship between photosynthetic O₂ evolution rates and the relative electron transport rate (ETR) at PSII. Photosynthetic O₂ evolution rate reflects the activity of photosynthesis (the Calvin-Benson cycle), whereas relative ETR is related to total electron transport activity, including AEF. Actually, we have found that the deletion of FLV2 and 4 (FLV2/4), which is the molecular mechanism of AEF under CO₂ limitation, gives the proportional linear relationship between photosynthetic O₂ evolution rates and relative ETR in the cyanobacterium *Synechocystis* sp. PCC 6803 (S. 6803) (Supplemental Fig. S1)¹³. These rates showed proportional linearity in CO₂-saturated conditions in the two secondary algae, but not in S. 7942 (Supplemental Figs S2–S4), which suggests that electron transports at PSII were strongly coupled to photosynthesis in *E. gracilis* and *P. tricornutum* when sufficient CO₂ was available. In S. 7942, relative ETR was already partially uncoupled from photosynthetic O₂ evolution rate during CO₂-saturated photosynthesis at supersaturated photon flux densities (Supplemental Fig. S2), indicating that cyanobacterial AEF functions in such situations³⁵. We note that AEF can be reflected in the relative ETR only when the AEF has a high activity level comparable to photosynthesis.

Responses of algal photosynthesis to CO₂ limitation were measured by following the method previously described^{12,13}. The responses of photosynthetic parameters to CO₂ limitation are shown in Fig. 2, and the original traces used to estimate these parameters are presented in Supplemental Figs S5–S7. The cyanobacterial and algal cells in fresh media were applied to an O₂ electrode chamber without adding inorganic carbon sources, and then illuminated with white actinic light (AL). Illumination with AL stimulated photosynthesis, which was accompanied by an increase in O₂ in the reaction medium (Supplemental Figs S5–S7). However, CO₂ in the medium was gradually removed by algal photosynthesis, as the diffusion of CO₂ from the atmosphere into the reaction medium was very slow, compared with the consumption by photosynthetic CO₂ assimilation in the experimental system. O₂ in the reaction medium began to decrease when CO₂ was depleted (Supplemental Figs S5–S7), indicating that photosynthesis was suppressed during the transition to CO₂ limitation. The addition of CO₂ (as NaHCO₃) to the reaction medium restored photosynthetic activity (Fig. 2, Supplemental Figs S5–S7).

During the measurements, the top of the chamber remained open, which enabled O₂ and CO₂ to diffuse into or out of the reaction medium. This open system relieved excessive increases in O₂ in the reaction mixture, which enabled O₂ to be measured for longer without passing over an undetectable point of the O₂ electrode. We temporarily closed the chamber to exclude the effects of diffusion of O₂ for determination of photosynthetic O₂ evolution rates (Supplemental Figs S5–S7)¹³.

In several earlier studies, it was observed that S. 7942 induced little AEF or NPQ in response to CO₂ limitation^{12,23,36}. Thus, we used S. 7942 as a control to compare the responses of *E. gracilis* and *P. tricornutum* in this study. In S. 7942, the photosynthetic O₂ evolution rate decreased in the transition to CO₂-limited photosynthesis, which was paralleled by the decrease in the relative ETR without NPQ induction (Fig. 2A, Supplemental Fig. S5). The proportional linear relationship between gross photosynthetic activity and relative ETR indicated that S. 7942 hardly drives AEF in the transition to CO₂ limitation (Supplemental Fig. S5). The increase in NPQ after adding NaHCO₃ has been observed in a previous study³⁶, although the molecular mechanism was unclear.

In both secondary algae, particularly in *E. gracilis*, some relative ETR was uncoupled from the O₂ evolution rates during CO₂-limited photosynthesis (Fig. 2B and C, Supplemental Figs S6 and S7), indicating that an AEF partially replaced photosynthesis during CO₂-limited photosynthesis in these algae. Compared with S. 7942, *E. gracilis* maintained relative ETR uncoupled from the photosynthetic O₂ evolution rate during CO₂-limited photosynthesis, which reached approximately half that during CO₂-saturated photosynthesis (Supplemental Fig. S6). Conversely, NPQ was slightly induced in the transition to CO₂ limitation in *E. gracilis*, which was not alleviated after at least 5 min after NaHCO₃ was added (Fig. 2B). These results concurred with those of a previous study³⁴.

In the diatom *P. tricornutum*, a dramatic induction of NPQ was observed in the transition to CO₂ limitation with the suppression of photosynthesis, although the AEF activity was much less, compared with *E. gracilis* (Fig. 2C). The relaxation of NPQ after adding NaHCO₃ was faster than that in *E. gracilis* (Fig. 2C), which is in agreement with a number of studies of diatomaceous NPQ^{24,25,34}. These data suggest that the NPQ in *E. gracilis* and *P. tricornutum* are derived from different molecular mechanisms.

Dependences of relative ETR on O₂ under CO₂ limitation in S. 7942, *E. gracilis*, and *P. tricornutum*. Diverse responses of photosynthetic electron transport to CO₂ limitation in S. 7942, *E. gracilis*, and *P. tricornutum* (Fig. 2) suggest different strategies of O₂ usage when photosynthesis is suppressed. To compare the O₂ usage of photosynthetic electron transport in these cyanobacterium and algae, we investigated the dependencies of relative ETR on O₂ during CO₂-limited photosynthesis. We eliminated O₂ in the medium by adding glucose, catalase, and glucose oxidase during CO₂-limited photosynthesis using the method described by Shimakawa *et al.*²³. We confirmed in advance that the addition of exogenous glucose during illumination did not affect the O₂ evolution rates and relative ETR in these species (Supplemental Table S1)²³. The top of the O₂ electrode chamber was closed to exclude the effects of diffusion of O₂ and CO₂ into or out of the reaction medium. It should be noted that there may have been some unintended consequences of using anaerobic conditions. However, removing O₂ did not affect relative ETR, at least during CO₂-saturated photosynthesis, in *E. gracilis* or *P. tricornutum* (Supplemental Fig. S8).

Compared with S. 7942 and *P. tricornutum*, which required little O₂ to drive AEF during CO₂-limited photosynthesis (Fig. 3A and C)^{12,23}, *E. gracilis* showed a biphasic dependence on O₂ (Fig. 3B). This indicated that more than two molecular mechanisms functioned as the O₂-dependent AEF in *E. gracilis*. Conversely, *P. tricornutum* was unlikely to rely on O₂-dependent AEF to alleviate photo-oxidative damage under CO₂ limitation, compared with *E. gracilis*. There was residual relative ETR under anaerobic conditions in *E. gracilis* and *P. tricornutum*, which might be derived from an O₂-insensitive AEF, including CET around PSI and PSII^{28–31}.




Figure 2. Responses of photosynthesis to CO₂ limitation in *S. 7942* (A), *Euglena gracilis* (B), and *Phaeodactylum tricornutum* (C). The graphs show the time course of O₂ evolution rate (red circles), relative electron transport rate (ETR) (blue triangles), and non-photochemical quenching (NPQ) (orange diamonds) in the fresh media containing the cells (10 μg Chl *a* mL⁻¹). Illumination with white actinic light (AL) (300 μmol photons m⁻² s⁻¹ for *S. 7942* and *E. gracilis*; 400 μmol photons m⁻² s⁻¹ for *P. tricornutum*) began at 0. NaHCO₃ (final concentration 10 mM) was added at the times indicated by red arrows. The double-headed black and blue arrows show CO₂-saturated and CO₂-limited photosynthesis, respectively.

Discussion

Figure 4 is a summary diagram of our previous and present results^{12,13,22,23} that presents the diverse O₂ usage strategies of photosynthetic electron transport to dissipate excess energy under CO₂ limitation in cyanobacteria, green algae, and two classes of algae with secondary plastids. Oxygenic phototrophs possess a number of molecular mechanisms that protect their cells against photo-oxidative damage by ROS. In this study, we focused on the physiological significance of O₂ as an alternative ‘safety valve’ in photosynthetic electron transport, and compared responses of photosynthesis to CO₂ limitation in genetically, phylogenetically, biologically, and ecologically different cyanobacteria and two classes of algae with secondary plastids. These organisms had different pigment compositions (Supplemental Table S2), which made it difficult to quantitatively compare photosynthetic O₂ evolution rates and relative ETR. Therefore, we simply defined the ratio of relative ETR during CO₂-limited photosynthesis to that during CO₂-saturated photosynthesis as residual relative ETR under CO₂ limitation in each species, and summarized the dependencies on O₂ as shown in Fig. 4. The cyanobacterium *S. 6803*, harboring FLV2/4, showed the activation of an O₂-dependent AEF during CO₂-limited photosynthesis¹³. This was different from *S. 7942*, which does not possess FLV2/4 (Fig. 2A)¹². Conversely, the marine species *S. 7002*, which does




Figure 3. Dependence of relative electron transport rate (ETR) on O_2 during CO_2 -limited photosynthesis in *S. 7942* (A), *Euglena gracilis* (B), and *Phaeodactylum tricornutum* (C). To remove dissolved O_2 , D-glucose (5 mM), catalase (250 units mL^{-1}), and glucose oxidase (5 units mL^{-1}) was added to the fresh media containing the cells (10 μg Chl a mL^{-1}). Photon flux densities of white actinic light (AL) were 300 μmol photons $m^{-2} s^{-1}$ for *S. 7942* and *E. gracilis*; 400 μmol photons $m^{-2} s^{-1}$ for *P. tricornutum*. Triangles, inverse triangles, and squares represent three independent measurements, respectively.

not harbor FLV2/4, maintained a relatively high electron flux to O_2 during CO_2 -limited photosynthesis owing to the higher contribution of FLV1 and 3 homologs (FLV1/3) to AEF, compared with *S. 7942* and *S. 6803*^{22,23}. The green alga *Chlamydomonas reinhardtii* drives an O_2 -dependent AEF in the transition from CO_2 -saturated to CO_2 -limited photosynthesis, similar to *S. 7002*²³. The dependences of relative ETR on O_2 under CO_2 limitation in *S. 6803*, *S. 7942*, *S. 7002*, and *C. reinhardtii* have already been reported in Shimakawa *et al.*²³. In addition, in *E. gracilis*, the electron flux to O_2 partially replaced photosynthesis under CO_2 limitation, while the dependency on O_2 was different from that in *S. 6803*, *S. 7002*, and *C. reinhardtii* (Figs 3B and 4). The biphasic O_2 dependency of relative ETR in *E. gracilis* indicated that this alga might drive the other AEF, which has low affinity for O_2 (e.g.




Figure 4. The diversity of O₂ usage strategies under CO₂ limitation in cyanobacterial and algal photosynthesis. Shown are cyanobacteria: *Synechocystis* sp. PCC 6803 (S. 6803), *S. elongatus* PCC 7942 (S. 7942), and *Synechococcus* sp. PCC 7002 (S. 7002); the green alga *C. reinhardtii*; the Euglenoid *Euglena gracilis*; and the diatom *Phaeodactylum tricornutum*. Cyanobacterial and algal cells were grown under ambient CO₂. Residual relative electron transport rate (ETR) under CO₂-limited photosynthesis indicates the ratio of relative ETR during CO₂-limited photosynthesis to that during CO₂-saturated photosynthesis, which is shown with the dependency on O₂ in reference to data in this and previous studies^{12,13,22,23}.

photorespiration)^{37,38}, in addition to the AEF that has high O₂ affinity. Conversely, the diatom *P. tricornutum* hardly showed O₂ usage (Figs 3C and 4). Compared with cyanobacteria and algae, C₃ plants mainly drive photorespiration as an O₂-dependent AEF that replaces photosynthesis at the CO₂ compensation point^{18,19}, whereas this is not observed in the C₄ plant maize¹⁸. Overall, there appear to be a number of diverse strategies of O₂ utilization that prevent photo-oxidative damage under CO₂ limitation, irrespective of the species of oxygenic phototroph, and O₂ is essential for some oxygenic phototrophs to protect cells against excess photon energy^{21,22}.

Overall, dissipating photon energy to O₂ is not necessarily a universal strategy in oxygenic phototrophs during CO₂-limited photosynthesis (Fig. 4). In many oxygenic phototrophs, the MAP pathway and respiratory terminal oxidases reduce O₂ at low concentrations, but in most cases, the rates estimated are less than 10% of CO₂-saturated gross photosynthesis^{39–41}, whereas some species show high activity of MAP pathway *in vivo*⁴². In this study, we measured the activities of the MAP pathway under CO₂ limitation in S. 7942, *E. gracilis*, and *P. tricornutum* by adding exogenous H₂O₂⁴³. The maximum activities reached approximately 70%, 25%, and 10% of the gross photosynthetic O₂-evolution rates in S. 7942, *E. gracilis*, and *P. tricornutum*, respectively (Supplemental Fig. S9). These estimates can be applied to the dependence of relative ETR on O₂ under CO₂ limitation in *E. gracilis*, but not in S. 7942 or *P. tricornutum* (Fig. 3). That is, both S. 7942 and *P. tricornutum* probably suppressed the MAP pathway under CO₂ limitation, and relied upon alternative strategies to dissipate excess photon energy.

There are mechanisms other than O₂-dependent AEF that function in the protection of cells against photo-oxidative damage, which would explain why there are diverse strategies of O₂ usage in oxygenic phototrophs. Increased NPQ is broadly used in many oxygenic phototrophs to dissipate excess photon energy, but the molecular mechanisms are unlikely to have the same origin. In the cyanobacterium S. 6803, it is observed that the orange carotenoid protein is expressed and functions in NPQ in response to high light levels⁴⁴. However, in the transition to CO₂ limitation, no induction of NPQ was observed in S. 7942 or S. 6803 (Fig. 2A)^{12,23,36}. The strategy to enhance NPQ under CO₂ limitation might not have been widespread in oxygenic phototrophs during their early evolution. The diatom *P. tricornutum* showed a large increase in NPQ under CO₂ limitation (Fig. 2C), which would be strictly related to ΔpH, some carotenoids, and the gene product of *lhcx5*^{24,25,45}. Conversely, the suppression of electron transport in the cytochrome *b6/f* complex is stimulated by ΔpH²⁶ and reduced the plastoquinone pool²⁷, both of which can cause the oxidation of P700 to dissipate excess photon energy²². Additionally, O₂-insensitive AEF, including CET around PSI^{28,29} and PSII^{30,31} may function to alleviate photo-oxidative damage. Furthermore, phototaxis possibly functions as a main strategy to avoid excess photon energy under CO₂ limitation in motile algae, such as *E. gracilis*⁴⁶. These O₂-insensitive strategies to alleviate photo-oxidative damage would enable various oxygenic phototrophs to be independent of O₂-dependent AEF. Nevertheless, the questions of the benefit (or cost) of O₂ usage to dissipate excess photon energy remains. There are still many questions over the diverse strategies that oxygenic phototrophs use to counter the detrimental effects of sunlight.

Methods

Growth conditions and determination of Chlorophyll. Cyanobacteria and algae were cultured in baffled shake flasks on a rotary shaker (100 rpm) under ambient CO₂. For all measurements, cells from the exponential growth phase were used.

S. 7942 was cultured in BG-11 medium⁴⁷ under light:dark conditions (25 °C, 16 h, 150 μmol photons m⁻² s⁻¹, fluorescent lamp; 23 °C, 8 h, dark). To quantify Chl, cells were centrifugally harvested and re-suspended by vortexing in 1 mL 100% (v/v) methanol. After subsequent incubation at room temperature for 5 min, the suspension was centrifuged at 10,000 × g for 2 min. Total Chl was determined from the supernatant⁴⁸.

E. gracilis Z (NIES-48) was photoautotrophically cultured in Cramer-Myers medium⁴⁹ under light:dark conditions (25 °C, 16 h, 150 μmol photons m⁻² s⁻¹, fluorescent lamp: 23 °C, 8 h, dark). Both Chl *a* and Chl *b* were quantified following the above-mentioned method⁴⁸.

P. tricornutum (UTEX642) was photoautotrophically cultured in the artificial seawater medium described previously, with the addition of 0.31% half-strength Guillard's 'F' solution^{50,51}, under light:dark conditions (22 °C, 14 h, 100 μmol photons m⁻² s⁻¹, fluorescent lamp: 20 °C, 10 h, dark). Both Chl *a* and Chl *c* were quantified as described above, except that the cells were re-suspended in a 1 mL mixture (10% [v/v] distilled water, 10% [v/v] dimethyl sulfoxide, and 80% [v/v] acetone)⁵².

Measurement of O₂ and Chl fluorescence. Net uptake and evolution of O₂ was measured simultaneously with Chl fluorescence. Cell samples in freshly prepared media (2 mL, 10 μg Chl *a* mL⁻¹) were stirred with a magnetic microstirrer and illuminated with white actinic light (AL) at 25 °C (for S. 7942 and *E. gracilis*) or 20 °C (for *P. tricornutum*). A halogen lamp (Xenophot HLX 64625, Osram, München, Germany) from the LS2 light source (Hansatech, King's Lynn, UK) was used as the white AL source. O₂ was monitored continuously using an O₂ electrode (Hansatech, King's Lynn, UK) while the measuring cuvette remained open to allow diffusion of O₂ and CO₂ between the medium and the air^{12,13}. The top of the cuvette was temporarily closed (1–3 min) while the O₂ evolution rate was determined^{12,13}. Representative raw traces of O₂ and relative Chl fluorescence in S. 7942, *E. gracilis*, and *P. tricornutum* are shown in Supplemental Figs S5A, S6A and S7A, respectively.

The relative Chl fluorescence originating from Chl *a* was measured using a PAM-Chl fluorometer (PAM-101; Walz, Effeltrich, Germany)^{53,54}. Pulse-modulated excitation was achieved using an LED lamp with a peak emission at 650 nm. Modulated fluorescence was measured at λ > 710 nm (Schott RG9 long-pass filter). The minimum Chl fluorescence (F₀) was determined from illumination using a measuring light (ML). The steady-state fluorescence (F_s) was monitored under AL, and 1,000-ms pulses of saturated light (10,000 μmol photons m⁻² s⁻¹) were supplied at arbitrary intervals to determine the maximum variable fluorescence (F_{m'}). The fluorescence terminology used in this study follows that of van Kooten and Snel (1990)⁵⁵. The effective quantum yield of PSII, Y(II), was defined as (F_{m'} - F₀)/F_{m'}. Relative ETR at PSII was estimated as the product of Y(II) and photon flux density of white AL. NPQ was calculated as (F_m - F_{m'})/F_{m'}⁵⁶. For S. 7942, F_m was determined in the presence of 3-(3,4-dichlorophenyl)-1,1-dimethylurea to exclude effects of state transition³⁵.

To measure the dependence of relative ETR on O₂ (Fig. 3, Supplemental Fig. S8), we added glucose (5 mM), catalase (250 units mL⁻¹, Wako, from bovine liver), and glucose oxidase (5 units mL⁻¹, Wako, from *Aspergillus niger*) to the medium with the chamber closed to block the diffusion of air to the medium. After photosynthetic O₂ evolution rates decreased to 0, we added these agents and evaluated the relative ETR²³.

Activity of the Mehler-ascorbate peroxidase pathway in cyanobacterial and algal cells was estimated from H₂O₂-dependent O₂ evolution rates during CO₂-limited photosynthesis⁴³. To exclude the effects of catalase, we added hydroxylamine (HA) to the reaction medium at 0.1 mM (for S. 7942 and *P. tricornutum*) or 0.5 mM (for *E. gracilis*).

Measurement of nitrogen. Cyanobacterial and algal cells were centrifugally harvested and dried overnight at 60 °C. Dried pellets were digested using the Kjeldahl method with sulfuric acid and H₂O₂. Total N content was determined using Nessler's reagent after adding sodium potassium tartrate and NaOH⁵⁷.

References

1. Kasting, J. F. Theoretical constraints on oxygen and carbon dioxide concentrations in the Precambrian atmosphere. *Precambrian Res.* **34**, 205–229 (1987).
2. Mitchell, P. Chemiosmotic coupling in oxidative and photosynthetic phosphorylation. *Biol. Rev.* **41**, 445–501 (1966).
3. Trebst, A. Energy conservation in photosynthetic electron transport of chloroplasts. *Annu. Rev. Plant Physiol.* **25**, 423–458 (1974).
4. Murata, N., Takahashi, S., Nishiyama, Y. & Allakhverdiyev, S. I. Photoinhibition of photosystem II under environmental stress. *Biophys. Biochim. Acta Bioenerg.* **1767**, 414–421 (2007).
5. Roach, T., Na, C. S. & Krieger-Liszka, A. High light-induced hydrogen peroxide production in *Chlamydomonas reinhardtii* is increased by high CO₂ availability. *Plant J.* **81**, 759–766 (2015).
6. Sonoike, K. Photoinhibition of photosystem I. *Physiol. Plant.* **142**, 56–64 (2011).
7. Sejima, T., Takagi, D., Fukayama, H., Makino, A. & Miyake, C. Repetitive short-pulse light mainly inactivates photosystem I in sunflower leaves. *Plant Cell Physiol.* **55**, 1184–1193 (2014).
8. Miyake, C. Alternative electron flows (water-water cycle and cyclic electron flow around PSI) in photosynthesis: molecular mechanisms and physiological functions. *Plant Cell Physiol.* **51**, 1951–1963 (2010).
9. Mehler, A. H. Studies on reactivities of illuminated chloroplasts. I. Mechanism of the reduction of oxygen and other Hill reagents. *Arch. Biochem. Biophys.* **33**, 65–77 (1951).
10. Miyake, C., Schreiber, U., Hormann, H., Sano, S. & Asada, K. The FAD-enzyme monodehydroascorbate radical reductase mediates photoproduction of superoxide radicals in spinach thylakoid membranes. *Plant Cell Physiol.* **39**, 821–829 (1998).
11. Helman, Y. *et al.* Genes encoding A-type flavoproteins are essential for photoreduction of O₂ in cyanobacteria. *Curr. Biol.* **13**, 230–235 (2003).
12. Hayashi, R. *et al.* O₂-dependent large electron flow functioned as an electron sink, replacing the steady-state electron flux in photosynthesis in the cyanobacterium *Synechocystis* sp. PCC 6803, but not in the cyanobacterium *Synechococcus* sp. PCC 7942. *Biosci. Biotechnol. Biochem.* **78**, 384–393 (2014).
13. Shimakawa, G. *et al.* FLAVODIIRON2 and FLAVODIIRON4 proteins mediate and oxygen-dependent alternative electron flow in *Synechocystis* sp. PCC 6803 under CO₂-limited conditions. *Plant Physiol.* **167**, 472–480 (2015).
14. Noguchi, K. & Yoshida, K. Interaction between photosynthesis and respiration in illuminated leaves. *Mitochondrion* **8**, 87–99 (2008).
15. Joët, T. *et al.* Involvement of a plastid terminal oxidase in plastoquinone oxidation as evidenced by expression of the *Arabidopsis thaliana* enzyme in tobacco. *J. Biol. Chem.* **277**, 31623–31630 (2002).
16. Lea-Smith, D. J. *et al.* Thylakoid terminal oxidases are essential for the cyanobacterium *Synechocystis* sp. PCC 6803 to survive rapidly changing light intensities. *Plant Physiol.* **162**, 484–495 (2013).
17. Kozaki, A. & Takeba, G. Photorespiration protects C3 plants from photooxidation. *Nature* **384**, 557–560 (1996).
18. Sejima, T. *et al.* Post-illumination transient O₂-uptake is driven by photorespiration in tobacco leaves. *Physiol. Plant.* **156**, 227–238 (2016).

19. Takagi, D., Hashiguchi, M., Sejima, T., Makino, A. & Miyake, C. Photorespiration provides the chance of cyclic electron flow to operate for the redox-regulation of P700 in photosynthetic electron transport system of sunflower leaves. *Photosynth. Res.* doi: 10.1007/s11120-016-0267-5 (2016).
20. Zhang, P., Allahverdiyeva, Y., Eisenhut, M. & Aro, E. M. Flavodiiron proteins in oxygenic photosynthetic organisms: photoprotection of photosystem II by Flv2 and Flv4 in *Synechocystis* sp. PCC 6803. *PLoS One* **4**, e5331 (2009).
21. Allahverdiyeva, Y. *et al.* Flavodiiron proteins Flv1 and Flv3 enable cyanobacterial growth and photosynthesis under fluctuating light. *Proc. Natl. Acad. Sci. USA* **110**, 4111–4116 (2013).
22. Shimakawa, G., Shaku, K. & Miyake, C. Oxidation of P700 in photosystem I is essential for the growth of cyanobacteria. *Plant Physiol.* doi: http://dx.doi.org/10.1104/pp.16.382 012277 (2016).
23. Shimakawa, G. *et al.* Diversity in photosynthetic electron transport under [CO₂]-limitation: the cyanobacterium *Synechococcus* sp. PCC 7002 and green alga *Chlamydomonas reinhardtii* drive an O₂-dependent alternative electron flow and non-photochemical quenching of chlorophyll fluorescence during CO₂-limited photosynthesis. *Photosynth. Res.* **130**, 293–305 (2016).
24. Goss, R. & Lepetit, B. Biodiversity of NPQ. *J. Plant Physiol.* **172**, 13–32 (2015).
25. Derkx, A., Schavén, K. & Bruce, D. Diverse mechanisms for photoprotection in photosynthesis. Dynamic regulation of photosystem II excitation in response to rapid environmental change. *Biochim. Biophys. Acta Bioenerg.* **1847**, 468–485 (2015).
26. Hope, A. B., Valente, P. & Matthews, D. B. Effects of pH on the kinetics of redox reactions in and around the cytochrome *bf* complex in an isolated system. *Photosynth. Res.* **42**, 111–120 (1994).
27. Shaku, K., Shimakawa, G., Hashiguchi, M. & Miyake, C. Reduction-induced suppression of electron flow (RISE) in the photosynthetic electron transport system of *Synechococcus elongatus* PCC 7942. *Plant Cell Physiol.* **57**, 1443–1453 (2016).
28. Allen, J. F. Cyclic, pseudocyclic and noncyclic photophosphorylation: new links in the chain. *Trends Plant Sci.* **8**, 15–19 (2003).
29. Peltier, G., Tolleter, D., Billon, E. & Cournac, L. Auxiliary electron transport pathways in chloroplasts of microalgae. *Photosynth. Res.* **106**, 19–31 (2010).
30. Falkowski, P. G., Fujita, Y., Ley, A. & Mauzerall, D. Evidence for cyclic electron flow around photosystem II in *Chlorella pyrenoidosa*. *Plant Physiol.* **81**, 310–312 (1986).
31. Miyake, C. & Yokota, A. Cyclic flow of electrons within PSII in thylakoid membranes. *Plant Cell Physiol.* **42**, 508–515 (2001).
32. Falkowski, P. G. *et al.* The evolution of modern eukaryotic phytoplankton. *Science* **305**, 354–360 (2004).
33. Cunningham, F. X. Jr. & Schiff, J. A. Chlorophyll-protein complexes from *Euglena gracilis* and mutants deficient in chlorophyll *b*. *Plant Physiol.* **80**, 231–238 (1986).
34. Casper-Lindley, C. & Björkman, O. Fluorescence quenching in four unicellular algae with different light-harvesting and xanthophyll-cycle pigments. *Photosynth. Res.* **56**, 277–289 (1998).
35. Campbell, D., Hurry, V., Clarke, A. K., Gustafsson, P. & Öquist, G. Chlorophyll fluorescence analysis of cyanobacterial photosynthesis and acclimation. *Microbiol. Mol. Biol. Rev.* **62**, 667–683 (1998).
36. Miller, A. G., Espie, G. S. & Bruce, D. Characterization of the non-photochemical quenching of chlorophyll fluorescence that occurs during the active accumulation of inorganic carbon in the cyanobacterium *Synechococcus* PCC 7942. *Photosynth. Res.* **49**, 251–262 (1996).
37. Jordan, D. B. & Ogren, W. L. Species variation in the specificity of ribulose bisphosphate carboxylase/oxygenase. *Nature* **291**, 513–515 (1981).
38. Yokota, A. & Kitaoka, S. Rates of glycolate synthesis and metabolism during photosynthesis of *Euglena* and microalgae grown on low CO₂. *Planta* **170**, 181–189 (1987).
39. Trimborn, S., Thoms, S., Petrou, K., Kranz, S. A. & Rost, B. Photophysiological responses of Southern Ocean phytoplankton to changes in CO₂ concentrations: Short-term versus acclimation effects. *J. Exp. Mar. Biol. Ecol.* **451**, 44–54 (2014).
40. Bailleul, B. *et al.* Energetic coupling between plastids and mitochondria drives CO₂ assimilation in diatoms. *Nature* **524**, 366–369 (2015).
41. Driever, S. M. & Baker, N. R. The water-water cycle in leaves is not a major alternative electron sink for dissipation of excess excitation energy when CO₂ assimilation is restricted. *Plant Cell Environ.* **34**, 837–846 (2011).
42. Waring, J., Klenell, M., Bechtold, U., Underwood, G. J. C. & Baker, N. R. Light-induced responses of oxygen photoreduction, reactive oxygen species production and scavenging in two diatom species. *J. Phycol.* **46**, 1206–1217 (2010).
43. Miyake, C., Michihata, F. & Asada, K. Scavenging of hydrogen peroxide in prokaryotic and eukaryotic algae: acquisition of ascorbate peroxidase during the evolution of cyanobacteria. *Plant Cell Physiol.* **32**, 33–43 (1991).
44. Wilson, A. *et al.* A soluble carotenoid protein involved in phycobilisome-related energy dissipation in cyanobacteria. *Plant Cell* **18**, 992–1007 (2006).
45. Bailleul, B. *et al.* An atypical member of the light-harvesting complex stress-related protein family modulates diatom responses to light. *Proc. Natl. Acad. Sci. USA* **107**, 18214–18219 (2010).
46. Richter, P. R. *et al.* High light exposure leads to a sign change of gravitaxis in the flagellate *Euglena gracilis*. *Acta Protozool.* **41**, 343–351 (2002).
47. Allen, M. M. Simple conditions for growth of unicellular blue-green algae on plates. *J. Phycol.* **4**, 1–4 (1968).
48. Grimmel, L. H. & Boardman, N. K. Photochemical activities of a particle fraction P₁ obtained from the green alga *Chlorella fusca*. *Biochem. Biophys. Res. Commun.* **49**, 1617–1623 (1972).
49. Cramer, M. & Myers, J. Growth and photosynthetic characteristics of *Euglena gracilis*. *Arch. Mikrobiol.* **17**, 384–402 (1952).
50. Guillard, R. R. L. & Ryther, J. H. Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustadt and *Detonula confervacea* (Cleve) Gran. *Can. J. Microbiol.* **8**, 229–239 (1962).
51. Guillard, R. R. L. *Culture of phytoplankton for feeding marine invertebrates*. In: Smith, W. L. & Chanley, M. H. (eds) *Culture of Marine Invertebrate Animals*, Plenum Press, New York, pp 29–60 (1975).
52. Jeffrey, S. W. & Humphrey, G. F. New spectrophotometric equations for determining chlorophylls *a*, *b*, *c*₁ and *c*₂ in higher plants, algae and natural phytoplankton. *Biochem. Physiol. Pflanz.* **167**, 191–194 (1975).
53. Schreiber, U., Schlüter, U. & Bilger, W. Continuous recording of photochemical and non-photochemical chlorophyll fluorescence quenching with a new type of modulation fluorometer. *Photosynth. Res.* **10**, 51–62 (1986).
54. Genty, B., Briantais, J. M. & Baker, N. R. The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim. Biophys. Acta Gen. Subj.* **990**, 87–92 (1989).
55. van Kooten, O. & Snel, J. F. H. The use of chlorophyll fluorescence nomenclature in plant stress physiology. *Photosynth. Res.* **25**, 147–150 (1990).
56. Baker, N. R. Chlorophyll fluorescence: a probe of photosynthesis *in vivo*. *Annu. Rev. Plant Biol.* **59**, 89–113 (2008).
57. Shimakawa, G. *et al.* Respiration accumulates Calvin cycle intermediates for the rapid start of photosynthesis in *Synechocystis* sp. PCC 6803. *Biosci. Biotechnol. Biochem.* **78**, 1997–2007 (2014).

Acknowledgements

This work was supported by the Japan Society for the Promotion of Science (JSPS, Scientific Research Grant No. 26450079 to C.M.) and the Ministry of Education, Culture, Sports, Science, and Technology, Japan (Scientific Research on Innovative Area No. 22114512 to C.M.). G.S. is supported as a JSPS research fellow (grant no. 16J03443). We would like to thank Editage (www.editage.jp) for English language editing.

Author Contributions

C.M. conceived the original screening and research plans; C.M. and Y.M. supervised the experiments; G.S. performed most of the experiments; Y.M., K.N., M.T., S.S., and C.M. provided technical assistance to G.S.; C.M. and G.S. designed the experiments and analyzed the data; C.M. and G.S. conceived the project and wrote the article with contributions from all the authors; C.M. supervised and complemented the writing.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Shimakawa, G. *et al.* Diverse strategies of O₂ usage for preventing photo-oxidative damage under CO₂ limitation during algal photosynthesis. *Sci. Rep.* 7, 41022; doi: 10.1038/srep41022 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017