SCIENTIFIC REPORTS

OPEN

Received: 23 August 2016
Accepted: 19 January 2017
Published: 24 March 2017

Renin-angiotensin system acting on reactive oxygen species in paraventricular nucleus induces sympathetic activation *via* AT1R/PKC γ /Rac1 pathway in salt-induced hypertension

Qing Su¹, Chan-Juan Huo¹, Hong-Bao Li¹, Kai-Li Liu¹, Xiang Li¹, Qing Yang¹, Xin-Ai Song¹, Wen-Sheng Chen², Wei Cui³, Guo-Qing Zhu⁴, Xiao-Lian Shi^{1,5}, Jin-Jun Liu¹ & Yu-Ming Kang¹

Brain renin-angiotensin system (RAS) could regulate oxidative stress in the paraventricular nucleus (PVN) in the development of hypertension. This study was designed to explore the precise mechanisms of RAS acting on reactive oxygen species (ROS) in salt-induced hypertension. Male Wistar rats were administered with a high-salt diet (HS, 8.0% NaCl) for 8 weeks to induced hypertension. Those rats were received PVN infusion of AT1R antagonist losartan (LOS, 10 μ g/h) or microinjection of small interfering RNAs for protein kinase C γ (PKC γ siRNA) once a day for 2 weeks. High salt intake resulted in higher levels of AT1R, PKC γ , Rac1 activity, superoxide and malondialdehyde (MDA) activity, but lower levels of copper/zinc superoxide dismutase (Cu/Zn-SOD), superoxide dismutase (SOD) and glutathione (GSH) in PVN than control animals. PVN infusion of LOS not only attenuated the PVN levels of AT1R, PKC γ , Rac1 activity, superoxide and decreased the arterial pressure, but also increased the PVN antioxidant capacity in hypertension. PVN microinjection of PKC γ siRNA had the same effect on LOS above responses to hypertension but no effect on PVN level of AT1R. These results, for the first time, identified that the precise signaling pathway of RAS regulating ROS in PVN is *via* AT1R/PKC γ /Rac1 in salt-induced hypertension.

High salt intake is a significant environmental factor, which strongly associated with blood pressure and the effects of salt loading on the renin-angiotensin system (RAS) in development of salt-induced hypertension^{1,2}. It is reported that high salt intake elevated the concentration of [Na⁺] in cerebrospinal fluid (CSF) and led to the activation of brain RAS and the increase of angiotensin II (Ang II) production in Dahl sensitive rat^{1,3}. As the primary effector peptide of the RAS, Ang II binding to the angiotensin II type-1 receptor (AT1R) contributes to the regulation of blood pressure and the development and/or the maintenance of metabolic syndrome⁴.

It is well known that paraventricular nucleus of the hypothalamus (PVN) in the brain is an important central site for integration of sympathetic nerve activity and regulation of cardiovascular function^{5,6}. Abundant evidences suggest that the elevated AT1R leads to reactive oxygen species (ROS) accumulation such as superoxide radical,

¹Department of Physiology and Pathophysiology, Xi'an Jiaotong University School of Basic Medical Sciences, Key Laboratory of Environment and Genes Related to Diseases (Xi'an Jiaotong University), Ministry of Education, Xi'an 710061, China. ²Department of Cardiovascular Surgery, Xijing Hospital, Fourth Military Medical University, Xi'an 710032, China. ³Department of Endocrinology and Metabolism, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an Jiaotong University Health Science Center, Xi'an 710061, China. ⁴Key Laboratory of Cardiovascular Disease and Molecular Intervention, Department of Physiology, Nanjing Medical University, Nanjing 210029, China. ⁵Department of Pharmacology, School of Basic Medical Sciences, Xi'an Jiaotong University Health Science Center, Xi'an 710061, China. Correspondence and requests for materials should be addressed to J.-J.L. (email: jupet@163.com) or Y.-M.K. (email: ykang@mail.xjtu.edu.cn)

Figure 1. Effects of AT1R antagonist (LOS) on mean arterial pressure (MAP) in salt-induced hypertensive rats. MAP is increased in high salt (HS) intake rats. Bilateral PVN infusion of LOS for 14 days attenuated high salt-induced presser response on the 5th day. Values are expressed as means $\pm$ SE. * $P < 0.05$ vs NS groups (NS + PVN LOS or NS + aCSF); † $P < 0.05$ HS + LOS vs HS + aCSF.

hydroxyl ion, and hydrogen peroxide in the PVN, which contribute to the overactivity of pre-autonomic PVN neurons and result in sympathoexcitation and metabolic disorders in the pathophysiology of hypertension^{5,7–9}. Furthermore, data from our laboratory suggest that Ang II increased the PVN levels of ROS, and bilateral PVN infusion of AT1R blocker decreased the production of ROS in PVN, as well as sympathetic nerve activity and blood pressure^{10–12}. Therefore, the activation of AT1R augments the intracellular level of ROS and sympathetic nerve activity. However, the precise signaling pathway of RAS modulating the oxidative stress in PVN is still unknown in hypertension.

Superoxide generated by NAD(P)H oxidase has emerged as a key intermediary in the central and peripheral effects of Ang II¹³. NAD(P)H oxidase is composed of membrane-bound subunits, gp91^{phox} and p22^{phox}, cytoplasmic subunits, p47^{phox}, p40^{phox}, p67^{phox}, and Rac1 and/or Rac2¹⁴. Rac1 (Ras-related C3 botulinum toxin substrate1) is a small GTPase which is essential for the assembly and activation of NAD(P)H oxidase and the production of superoxide¹⁵. A growing body of evidence indicate that the increased Ang II in brain induces the NAD(P)H-dependent ROS production by the protein kinase C (PKC) which is a critical step in Rac1 activation and subsequent enzyme assembly^{13,16}. PKC is a family of phospholipids serine/threonine protein kinases, which have been classified into conventional PKCs (α , β and γ), novel PKCs (δ , ϵ , η and θ), and atypical PKCs (ζ and λ/ι)¹⁷. PKC γ , unlike other PKC isozymes, is expressed only in the central nervous system and eye tissues¹⁷. Aronowski and colleagues (2000) found that PKC γ is activated in the period of brain ischemia/reperfusion which is closely related to the oxidative stress^{18,19}. These findings suggest that the linking of AT1R and NAD(P)H oxidase-derived ROS is involved in the effects of PKC γ /Rac1/NAD(P)H pathway. Furthermore, it is unclear whether AT1R/PKC γ /Rac1/NAD(P)H pathway in PVN mediates ROS production in the development of hypertension. Therefore, the aim of the present study was to determine whether the precise mechanism of RAS modulating oxidative stress in PVN is through AT1R/PKC γ /Rac1/NAD(P)H signaling pathway during the development of high-salt hypertension.

Results

Effect of losartan on mean arterial pressure (MAP). High salt diet induced a significant increase in mean arterial pressure (MAP) compared with control rats after 8 weeks (139 ± 4.7 mmHg vs. 94.7 ± 3.6 mmHg, $P < 0.05$) of high salt diet before bilateral PVN infusion of LOS or microinjections of PKC γ siRNA. In Fig. 1, HS + PVN aCSF exhibited a significant increase in MAP when compared with Wistar normal salt diet (NS) rats (at Day 14, 163.3 ± 5.6 vs. 95.6 ± 3.0 mmHg, $P < 0.05$). Whereas HS + PVN LOS rats exhibited significantly reduced MAP from Day 5, and it remained lower for the duration of the study (at Day 14, 119.2 ± 3.5 vs. 165.4 ± 5.3 mmHg, $P < 0.05$) when compared with HS + PVN aCSF rats.

Effect of losartan on sympathetic nerve activity. In Fig. 2, HS + PVN aCSF rats exhibited a 7.1-fold increase in renal sympathetic nerve activity (RSNA, % of max) and 3.3-fold increase in plasma noradrenaline (NE) when compared with control rats ($P < 0.05$). Bilateral PVN infusion of LOS declined to a 64.5% decrease RSNA and 54.7% plasma NE in high salt-induced hypertensive rats ($P < 0.05$). NS + PVN LOS did not attenuate RSNA when compared with NS + PVN aCSF.

Effect of losartan on PKC γ and AT1R positive neurons in PVN. In Fig. 3, immunohistochemistry and immunofluorescence results revealed that high salt diet respectively induced a 4.8-fold increase and 2.6-fold increase in the number of AT1R and PKC γ positive neurons compared with control rats ($P < 0.05$). Bilateral PVN infusion of LOS respectively declined to a 51.2% decrease and 60% decrease in the number of AT1R and PKC γ positive neurons in hypertensive rats ($P < 0.05$). NS + PVN LOS did not attenuate the number of AT1R and PKC γ positive neurons when compared with NS + PVN aCSF.

Figure 2. Effects of LOS on renal sympathetic nerve activity (RSNA) and norephedrine (NE) in plasma in salt-induced hypertensive rats. RSNA was increased in high salt (HS) intake rats. Bilateral PVN infusion of LOS decreased RSNA in HS rats. (A) A representative renal sympathetic nerve activity in different groups. (B) Bar graph comparing renal sympathetic nerve activity in different groups. (C) Bar graph comparing NE in plasma in different groups. Values are expressed as means $\pm$ SE. * $P < 0.05$ vs NS groups (NS + PVN LOS or NS + aCSF); † $P < 0.05$ HS + LOS vs HS + aCSF.

Effect of losartan on p-Rac1 positive neurons and the level of superoxide in PVN. In Fig. 4, immunohistochemistry and immunofluorescence results revealed that high salt diet respectively induced a 2.3-fold increase and 3.5-fold increase in the number of p-Rac1 positive neurons and superoxide compared with control rats ($P < 0.05$). Bilateral PVN infusion of LOS respectively declined to a 56.0% decrease and 53.1% decrease in the number of p-Rac1 positive neurons and superoxide in hypertensive rats ($P < 0.05$). NS + PVN LOS did not attenuate the number of p-Rac1 positive neurons and superoxide when compared with NS + PVN aCSF.

Effect of losartan on protein expression levels of AT1R, PKC γ , p-Rac1 and Cu/Zn-SOD in PVN. In Fig. 5A–C, western blot results indicated that high salt-induced rats respectively induced a 10.3-fold (AT1R), 8.9-fold (PKC γ) and 9.7-fold (GTP-Rac1, Pull-down Assay) increase, but 5.6-fold decrease (Cu/Zn-SOD) in the protein expression level ($P < 0.05$) in PVN compared with control rats. Bilateral PVN infusion of LOS respectively declined to 38% (AT1R), 37% (PKC γ), and 64% (GTP-Rac1, Pull-down Assay) protein expression level, but increased Cu/Zn-SOD protein level to 46% compared with high salt-induced rats ($P < 0.05$).

Effect of losartan on mRNA levels of AT1R, PKC γ , and Cu/Zn-SOD in PVN. In Fig. 5D, the high salt diet respectively induced a 3.1-fold (AT1R) and 4.2-fold (PKC γ) but 3.0-fold decrease (Cu/Zn-SOD) in the mRNA level in PVN compared with control rats ($P < 0.05$). Bilateral PVN infusion of LOS respectively declined to 42.8% (AT1R) and 43.2% (PKC γ) mRNA level, but increased Cu/Zn-SOD mRNA level to 36.5% compared with high salt-induced rats ($P < 0.05$).

Effect of PKC γ (PKC γ siRNA) on mean arterial pressure (MAP). In Fig. 6, the MAP of HS + PVN scrambled siRNA group is also much higher than control animals (at Day 14, 160.6 ± 4.9 vs. 95.1 ± 3.1 mmHg, $P < 0.05$) at the end of experiment. Bilateral PVN microinjection of PKC γ siRNA significantly decreased the MAP compared to HS + PVN scrambled siRNA rat (at Day 14, 132.6 ± 3.4 vs. 163.5 ± 5.6 mmHg, $P < 0.05$). Normal salt diet groups did not show any significant change in MAP at the end of experiment.

Effect of PKC γ (PKC γ siRNA) on sympathetic nerve activity. In Fig. 7, HS + PVN scrambled siRNA rats exhibited a 4.1-fold increase in RSNA and a 2.5-fold increase in plasma NE when compared with control rats ($P < 0.05$). Bilateral PVN infusion of PKC γ siRNA declined to a 65.4% RSNA and 72% plasma NE in high salt-induced hypertensive rats ($P < 0.05$). NS + PVN PKC γ siRNA did not show any significant change in RSNA and plasma NE when compared with NS + PVN scrambled siRNA at the end of experiment.

Figure 3. Effects of LOS on AT1R and PKC γ expression in the PVN of salt-induced hypertensive rats. Expressions of AT1R and PKC γ in high salt intake (HS) rats were higher than in control rats. Treatment with LOS attenuated the AT1R and PKC γ expression in the PVN of HS rats. (A) Immunofluorescence for AT1R (bright green) and immunohistochemistry for PKC γ (bright brown) in the PVN in different groups. (B) and (C) Column diagram showing the numbers of AT1R and PKC γ positive neurons in the PVN in different groups. * $P < 0.05$ vs NS groups (NS + PVN LOS or NS + aCSF); $^{\dagger}P < 0.05$ HS + LOS vs HS + aCSF.

Effect of PKC γ (PKC γ siRNA) on PKC γ and AT1R positive neurons in PVN. In Fig. 8, high salt diet also induced respectively induced a 6.7-fold increase and 3.2-fold increase in the number of AT1R and PKC γ positive neurons compared with control rats ($P < 0.05$). However, bilateral PVN microinjection of PKC γ siRNA declined to 60% PKC γ positive neurons, and had no effect on the AT1R positive neurons in PVN in hypertensive rats at the end of experiment.

Effect of PKC γ (PKC γ siRNA) on p-Rac1 positive neurons and the level of Superoxide in PVN. In Fig. 9, high salt diet also induced respectively induced a 1.3-fold increase and 3.7-fold increase in the number of p-Rac1 positive neurons and superoxide compared with control rats ($P < 0.05$). However, bilateral PVN microinjection of PKC γ siRNA respectively declined to 34% the number of p-Rac1 positive neurons and 70% superoxide in PVN in hypertensive rats at the end of experiment ($P < 0.05$).

Effect of PKC γ (PKC γ siRNA) on protein expression levels of AT1R, PKC γ , p-Rac1 and Cu/Zn-SOD in PVN. In Fig. 10A–C, western blot results indicated that high salt-induced rats respectively induced a 9.6-fold (AT1R), 1.8-fold (PKC γ) and 7.3-fold (GTP-Rac1, Pull-down Assay) increase, but 7.8-fold decrease (Cu/Zn-SOD) in the protein expression level ($P < 0.05$) in PVN compared with control rats. Bilateral PVN infusion of PKC γ siRNA in PVN respectively declined to 51% (AT1R), 31% (PKC γ), and 49% (GTP-Rac1, Pull-down Assay) protein expression level, but increased Cu/Zn-SOD protein level to 32% compared with high salt-induced rats ($P < 0.05$). And administration of PKC γ siRNA in PVN had no effect on PVN levels of AT1R in salt-induced hypertensive rats. Bilateral PVN infusion of PKC γ siRNA in normal rat downregulated a 55% protein expression compared with normal rats with scrambled siRNA.

Effect of PKC γ (PKC γ siRNA) on mRNA levels of AT1R, PKC γ , and Cu/Zn-SOD in PVN. In Fig. 10D, RT-PCR results indicated that high salt-induced rats respectively induced a 10.1-fold (AT1R) and 8.4-fold (PKC γ), but 7.2-fold decrease (Cu/Zn-SOD) in the protein expression level ($P < 0.05$) in PVN compared with control rats. Bilateral PVN infusion of PKC γ siRNA in PVN respectively declined to 31% (PKC γ) mRNA level, but increased Cu/Zn-SOD mRNA level to 36% compared with high salt-induced rats ($P < 0.05$). And administration of PKC γ siRNA in PVN had no effect on PVN mRNA levels of AT1R in salt-induced hypertensive rats. Bilateral PVN infusion of PKC γ siRNA in normal rat downregulated a 52% mRNA level compared with normal rats with scrambled siRNA.

Figure 4. Effects of LOS on p-Rac1 expression and ROS activity in the PVN of salt-induced hypertensive rats. ROS activity was measured by fluorescent-labeled dihydroethidium (DHE) staining. High salt (HS) intake rats had higher level of p-Rac1 expression and immunofluorescent intensity of DHE when compared with control rats. Bilateral PVN infusion of LOS for 14 days reduced the levels of superoxide and p-Rac1 expression in the PVN of HS rats. (A) Immunohistochemistry for p-Rac1 and the immunofluorescence for superoxide in the PVN in different groups. (B) Column diagram showing the effects of LOS on the p-Rac1 positive neurons in the PVN in different groups. (C) Column diagram showing immunofluorescent intensity of DHE in the PVN in different groups. Values are expressed as means $\pm$ SE. * $P < 0.05$ vs NS groups (NS + PVN LOS or NS + aCSF); $^{\dagger}P < 0.05$ HS + LOS vs HS + aCSF.

Effect of losartan and PKC γ (PKC γ siRNA) on arterial pressure and heart rate (HR). The mean arterial pressure (MAP) and heart rate (HR) were measured with a pressure transducer (MLT0380, AD Instruments, Australia) via a catheter in the right carotid artery. The MAP and HR in high salt-induced hypertensive rats were significantly higher than that in normal salt diet rats. There was no significant difference in the body weight between hypertensive rats and control rats. Chronic PVN infusion of LOS and PKC γ siRNA attenuated MAP and HR in hypertensive rats, but not in normal salt diet rats (Table 1).

Effect of losartan and PKC γ (PKC γ siRNA) on oxidative stress in PVN. Compared with NS rats, HS + PVN aCSF and HS + PVN scrambled siRNA groups had higher level of malondialdehyde (MAD) and NAD(P)H oxidase activity, but lower levels of superoxide dismutase (SOD) and glutathione (GSH) in PVN. Infusion of LOS or microinjection of PKC γ siRNA into the PVN attenuated the increased levels of MAD and NAD(P)H oxidase activity, but decreased levels of SOD activity and GSH in PVN of HS groups (Table 2).

Discussion

The novel finding of this study is that both AT1R and PKC γ play an important role in modulating the salt-induced central RAS activation and oxidative stress responses. And these results, for the first time, identified that the precise signaling pathway of RAS regulating ROS in PVN may be through AT1R/PKC γ /Rac1 pathway, which induces overproduction of ROS from Rac1-dependent NAD(P)H and finally increase sympathetic nerve activity and blood pressure. Therefore, PKC γ signaling pathway contributes to RAS modulating oxidative stress in PVN in the development of salt-induced hypertension.

PVN is a central integration site for the regulation of cardiovascular functions². Substantial findings have confirmed high salt intake is a significant environmental factor, which strongly associated with RAS and ROS in the PVN^{3,20}. Our study also showed that the activated RAS could regulate the reactive oxygen species overproduction in PVN in the progression of hypertension. As the primary effector peptide of the RAS, Ang II binding to AT1R contributes to the regulation of blood pressure and the development and/or the maintenance of metabolic syndrome. There is also increasing evidence showed that the augmented Ang II in brain induces the NAD(P)

Figure 5. Effects of LOS on AT1R, PKC γ , GTP-Rac1 (Pull-down Assay) and Cu/Zn-SOD proteins expression, and effects of LOS on AT1R, PKC γ and Cu/Zn-SOD mRNA level in the PVN of salt-induced hypertensive rats. High salt (HS) intake rats had higher levels of AT1R, PKC γ and GTP-Rac1, but lower level of Cu/Zn-SOD compare with control rats. Bilateral PVN infusion of LOS for 14 days attenuated expression of AT1R, PKC γ and GTP-Rac1, and augmented expression of Cu/Zn-SOD in hypertensive rats. (A) A representative immunoblot; (B) PVN tissues were analyzed for Rac1 activity (Rac1-GTP) as described in METHODS. As controls, protein levels of Rac1 (total-Rac1) in PVN were measured. (C) Densitometric analysis of protein expression of AT1R, PKC γ and Cu/Zn-SOD in the PVN in different groups; (D) Densitometric analysis of mRNA expression of AT1R, PKC γ and Cu/Zn-SOD in the PVN in different groups. Values are expressed as means $\pm$ SE. * $P < 0.05$ vs NS groups (NS + PVN LOS or NS + aCSF); † $P < 0.05$ HS + LOS vs HS + aCSF.

Figure 6. Effects of PKC γ siRNA on mean arterial pressure (MAP) in salt-induced hypertensive rats. MAP is increased in high salt (HS) intake rats. Bilateral PVN microinjections of PKC γ for 14 days attenuated high salt-induced pressor response on the 8th day. Values are expressed as means $\pm$ SE. * $P < 0.05$ vs NS groups (NS + PKC γ siRNA or NS + scrambled siRNA); † $P < 0.05$ HS + PKC γ siRNA vs HS + scrambled siRNA.

H-dependent ROS production by PKC. Members of the PKC family play a key role in the regulation of cellular functions in the nervous system²¹. Meanwhile, previous work by Griendling K and coworkers (2009) reviewed that Ang II activated by NAD(P)H oxidase is mainly dependent on the PKC and other signaling pathways in the central nervous system¹³. Lin D. and Takemoto D. J. (2005) had showed that PKC γ could activate the oxidative

Figure 7. Effects of PKC γ siRNA on RSNA and NE in plasma in salt-induced hypertensive rats. RSNA was increased in high salt (HS) intake rats. Bilateral PVN infusion of PKC γ siRNA decreased RSNA in HS rats. (A) A representative renal sympathetic nerve activity in different groups. (B) Bar graph comparing renal sympathetic nerve activity in different groups. (C) Bar graph comparing NE in plasma in different groups. Values are expressed as means $\pm$ SE. * $P < 0.05$ vs NS groups (NS + PKC γ siRNA or NS + scrambled siRNA); $^{\dagger}P < 0.05$ HS + PKC γ siRNA vs HS + scrambled siRNA.

stress in the cardiovascular and neurodegenerative diseases²². And PKC γ is a unique isoform of PKC that is found in neurons and eye tissues. Rac1 (Ras-related C3 botulinum toxin substrate1) is a small GTPase which is essential for the assembly and activation of NAD(P)H oxidase²³. Growing evidence indicate that NOX/Rac1 activation is a main pathway in cardiovascular and cerebrovascular disorders such as myocardial infarction, hypertension, atherosclerosis and stroke¹⁴. Zimmerman M. C. (2004) had, for the first time, identified a Rac1-dependent NAD(P)H oxidase as the source of central Ang II-induced O₂ production, and implicated this oxidase in cardiovascular diseases associated with dysregulation of brain Ang II signaling, including hypertension¹⁴. Moreover, increasing evidences demonstrate that increased Ang II in brain induces the NAD(P)H-dependent ROS production by PKC is a critical step in Rac1 activation and subsequent enzyme assembly.

In this study, PVN microinjection of PKC γ siRNA could decrease the sympathetic nervous activity and mean arterial pressure, which indicates that PKC γ plays a significant role in the brain Ang II regulation of oxidative stress in the pathophysiology of high salt-induced hypertension. And our results also show that high salt diet not only elevated the PVN level of AT1R, PKC γ , Rac1, and NADPH oxidase-dependent superoxide in PVN, but also decreased the antioxidant capacity in PVN, therefore, increased the sympathetic nervous activity and mean arterial pressure. Chronic infusion of AT1R antagonist (LOS) into PVN decreased the PVN expression of AT1R, PKC γ , Rac1 and the level of superoxide in hypertensive rats, but increased the antioxidant capacity in PVN. Campese VM (2000) found LOS could stimulate the production of nitric oxide in PVN, and the nitric oxide could regulate the sympathetic outflow and reduces blood pressure levels²⁴. This response is also consistent with our results. However, PVN microinjection of PKC γ siRNA suppressed NAD(P)H oxidase-dependent oxidative stress in PVN, decreased the PVN levels of PKC γ , Rac1 and NAD(P)H oxidase subunits but did not affect AT1R. So we would only speculate that AT1R could act on PKC γ and Rac1-dependent NAD(P)H oxidase and regulates the NAD(P)H oxidase-dependent oxidative stress in PVN, which implicated RAS regulating ROS in PVN may be through the AT1R/PKC γ /Rac1 signaling pathway and increase the sympathetic nervous activity and mean arterial pressure in the progression of salt-induced hypertension, which is presented in Fig. 11.

Material and Methods

Animals. Male Wistar rats weighing 150 g to 200 g were purchased from Experimental Animal Center of Xi'an Jiaotong University. All animals were housed in temperature ($23 \pm 2^{\circ}\text{C}$) and light-controlled (12 h light/dark cycle) animal quarters and were provided with rat chow *ad libitum*. All experimental procedures were reviewed and approved by the National Institutes of Health Guide for the Care and Use of Laboratory Animals (the US National Institutes of Health Publication No. 85-23, revised 1996). All of the animal procedures were conducted according to the Animal Care and Use Committees of Xi'an Jiaotong University.

Figure 8. Effects of PKC γ siRNA on AT1R and PKC γ expression in the PVN of salt-induced hypertensive rats. Expressions of AT1R and PKC γ in high salt intake (HS) rats were higher than in control rats. Microinjections of PKC γ siRNA attenuated the PKC γ expression and had no effect on the AT1R positive neurons in the PVN of HS rats. (A) Immunofluorescence for AT1R (bright green) and immunohistochemistry for PKC γ (bright brown) in the PVN in different groups. (B) and (C) Column diagram showing the numbers of AT1R and PKC γ positive neurons in the PVN in different groups. Values are expressed as means $\pm$ SE. * $P < 0.05$ vs NS groups (NS + PKC γ siRNA or NS + scrambled siRNA); [†] $P < 0.05$ HS + PKC γ siRNA vs HS + scrambled siRNA.

PKC λ siRNA preparation and transfection in PVN. PKC λ siRNA (siPKC) oligonucleotides and the scrambled siRNA were purchased from GenePharma (Shanghai, China). We selected three potential siRNA sequences with high knock-down probability values. The siRNA sequences in *Prkcg*-Rat started at bases 364, 1277, and 1537 (364: 5'-AAGUCGGUACAGUGACUGCTT-3'; 1277: 5'-UUUCCCUAGAACCAUGAGGTT-3'; 1537: 5'-AUGUGGUACAUUAAAUCGCTT-3'). To test the silence ability of these siRNA sequences target genes in the PVN, 15 nM of each siRNA carried out with Lipofectamine TM 2000 (Invitrogen) was infusion of PVN for two weeks in the high salt-induced hypertensive rats according to the manufacturer's instructions. After 14 days, the rats were euthanized to collect PVN tissue. Western blot was used for measurement of PKC proteins expression in PVN. Finally, we choose PKC γ siRNA sequences 1537 as follows: 5'-AUGUGGUACAUUAAAUCGCTT-3'² in our further study.

General experimental protocol. Rats were fed on the normal salt diet containing 0.3% NaCl (NS) or high salt diet containing 8% NaCl (HS) for 8 weeks to induce hypertension. After 8 weeks, animals from NS group and HS group are received bilateral PVN infusion of AT1R antagonist losartan (LOS, 10 μ g/h) or artificial cerebrospinal fluid (aCSF), and bilateral PVN microinjection of PKC γ small interfering RNA (PKC γ siRNA), or scrambled siRNA for 2 weeks with high salt diet respectively²⁵.

Implantation of bilateral PVN osmotic minipump for chronic infusion. The method for PVN minipump has been previously described^{18,26,27}. Each rat head was placed into a stereotaxic apparatus after anesthetized with ketamine (90 mg/kg) and xylazine (10 mg/kg) intraperitoneally (i.p). A skin incision was made, and the skull was opened. The location was 1.8 mm caudal to the bregma, 0.4 mm lateral to central line, and 7.9 mm below the skull surface (Chen *et al.*, 2011; Zhu *et al.*, 2004). The skull was then opened, and the minipumps (ALZET Osmotic Pumps, 2002 Model, 0.5 μ l/h) connected with losartan (LOS, 10 μ g/h) or artificial cerebrospinal fluid (aCSF) for the continuous infusion were implanted subcutaneously in the back of the neck. The infusion was continued for 2 weeks. Rats received buprenorphine (0.01 mg/kg, sc) immediately following surgery. The success rate of bilateral PVN microinjection was around 68%.

Figure 9. Effects of PKC γ siRNA on p-Rac1 expression and ROS activity in the PVN of salt-induced hypertensive rats. ROS activity was measured by fluorescent-labeled dihydroethidium (DHE) staining. High salt (HS) intake rats had higher level of p-Rac1 expression and immunofluorescent intensity of DHE when compared with control rats. Bilateral PVN microinjections of PKC γ siRNA for 14 days reduced the levels of superoxide and p-Rac1 expression in the PVN of HS rats. (A) Immunohistochemistry for p-Rac1 and the immunofluorescence for superoxide in the PVN in different groups. (B) Column diagram showing the effects of PKC γ siRNA on the p-Rac1 positive neurons in the PVN in different groups. (C) Column diagram showing immunofluorescent intensity of DHE in the PVN in different groups. Values are expressed as means $\pm$ SE. * $P < 0.05$ vs NS groups (NS + PKC γ siRNA or NS + scrambled siRNA); $^{\dagger}P < 0.05$ HS + PKC γ siRNA vs HS + scrambled siRNA.

Bilateral PVN cannulae implantation for chronic infusion studies. The method for implantation of bilateral PVN cannulae has been described previously. Briefly, under anesthesia, rat head was placed into a stereotaxic apparatus. A stainless steel double cannula (Plastics One, Inc.) with a center-to-center distance of 0.5 mm, was implanted into the PVN using an introducer, according to stereotaxic coordinates (1.8 mm caudal to the bregma, 0.4 mm lateral to central line, and 7.9 mm below the skull surface)¹⁰. The cannula was fixed to the cranium using dental acrylic and two stainless steel screws. 50 nL of scrambled siRNA or small interfering RNAs for PKC γ (PKC γ siRNA) were microinjected into the bilateral PVN each side, which were completed within 1 min once per day. The success rate of bilateral PVN cannulation is about 65%.

Blood pressure measurement. Arterial pressure was measured noninvasively *via* tail-cuff instrument and their Recording System. Conscious rats from each group were warmed to an ambient temperature of 30 °C by placing them in a holding device mounted on a thermostatically controlled warming plate. Each rat was allowed to accommodate the cuff for 10 minutes before blood pressure measurement. The rat arterial pressure was measured every week during the 8 weeks and every day after LOS or siRNA infusion. Mean arterial pressure and heart rate data were consisted of 20 times, which were collected for 40 min within the same 2-hour time window each day and then averaged those data until the end of this study²⁸.

Sympathetic neural recordings. Rats were anaesthetized with a ketamine (90 mg/kg) and xylazine (10 mg/kg) mixture (ip). After retroperitoneal laparotomy, the left renal nerves were isolated *via* glass microelectrode technique under an inversion microscope. The renal nerve was hung by a platinum electrode which is connected to the recording system. In order to moisturize the nerves and isolate electrical disturbance, the nerve should be covered by paraffin oil tampons. Maximum renal sympathetic nerve activity (RSNA) was detected using an intravenous bolus administration of sodium nitroprusside (SNP, 10 mg). The recordings of rectified and integrated RSNA were analyzed using methods described as previously^{8,26}.

Figure 10. Effects of PKC γ siRNA on AT1R, PKC γ , GTP-Rac1 (Pull-down Assay) and Cu/Zn-SOD proteins expression, and effects of LOS on AT1R, PKC γ and Cu/Zn-SOD mRNA level in the PVN of salt-induced hypertensive rats. High salt (HS) intake rats had higher levels of AT1R, PKC γ and GTP-Rac1, but lower level of Cu/Zn-SOD compare with control rats. Bilateral PVN infusion of PKC γ siRNA for 14 days attenuated expression of AT1R, PKC γ and GTP-Rac1, and augmented expression of Cu/Zn-SOD in hypertensive rats. However, administration of PKC γ siRNA in PVN had no effect on PVN levels of AT1R in salt-induced hypertension. **(A)** A representative immunoblot; **(B)** PVN tissues were analyzed for Rac1 activity (Rac1-GTP) as described in METHODS. As controls, protein levels of Rac1 (total-Rac1) in PVN were measured. **(C)** densitometric analysis of protein expression of AT1R, PKC γ and Cu/Zn-SOD in the PVN in different groups; **(D)** Densitometric analysis of mRNA expression of AT1R, PKC γ and Cu/Zn-SOD in the PVN in different groups. Values are expressed as means $\pm$ SE. * $P < 0.05$ vs NS groups (NS + PKC γ siRNA or NS + scrambled siRNA); † $P < 0.05$ HS + PKC γ siRNA vs HS + scrambled siRNA. # $P < 0.05$ NS + PKC γ siRNA vs NS + scrambled siRNA.

Figure 11. The schematic showing the proposed signaling pathways of renin-angiotensin system (RAS) regulating reactive oxygen species (ROS) within the hypothalamic paraventricular nucleus (PVN) via AT1R/PKC γ /Rac1 in high salt-induced hypertension.

PVN	Body weight (g)	MAP (mmHg)	HR (bpm)
HS + PVN LOS	367 ± 9	119 ± 5 ^{†,*}	342 ± 10 [†]
HS + PVN aCSF	355 ± 7	165 ± 8*	384 ± 13*
NS + PVN LOS	359 ± 8	100 ± 6	354 ± 8
NE + PVN aCSF	347 ± 8	98 ± 4	346 ± 9
HS + PVN PKC siRNA	358 ± 9	132 ± 7 ^{†,*}	351 ± 11 [†]
HS + PVN scrambled siRNA	366 ± 10	162 ± 6*	394 ± 13*
NS + PVN PKC siRNA	351 ± 8	103 ± 4	345 ± 7
NE + PVN scrambled siRNA	359 ± 9	97 ± 5	349 ± 8

Table 1. Changes of body weight, MAP and HR at the end of the 2nd week of the experiment (n = 7, for each group). Values are expressed as means SE. *P < 0.05 versus NS groups; [†]P < 0.05 HS + PVN LOS or HS + PVN PKC siRNA versus HS control groups.

PVN	MDA (mmol/mgp)	SOD (U/mgp)	GSH (umol/mgp)	NAD(P)H oxidase (RLU/mg/sec)
HS + PVN LOS	4.87 ± 0.19 ^{†,*}	6.26 ± 0.30 ^{†,*}	2.85 ± 0.24 ^{†,*}	1.91 ± 1.20 ^{†,*}
HS + PVN aCSF	7.01 ± 0.24*	2.67 ± 0.10*	0.98 ± 0.13*	3.69 ± 1.41*
NS + PVN LOS	1.68 ± 0.22	9.61 ± 0.40	4.76 ± 0.15	0.93 ± 0.34
NE + PVN aCSF	1.25 ± 0.21	10.00 ± 1.40	5.18 ± 0.37	0.87 ± 0.29
HS + PVN PKC siRNA	3.82 ± 0.14 ^{†,*}	4.57 ± 0.17 ^{†,*}	1.99 ± 0.18 ^{†,*}	2.80 ± 1.16 ^{†,*}
HS + PVN scrambled siRNA	6.28 ± 0.24*	3.67 ± 0.15*	0.98 ± 0.13*	4.27 ± 1.78*
NS + PVN PKC siRNA	2.55 ± 0.13	7.11 ± 0.37	3.24 ± 0.12	1.33 ± 0.32
NE + PVN scrambled siRNA	2.68 ± 0.14	7.85 ± 0.44	2.97 ± 0.11	1.18 ± 0.26

Table 2. Effect of LOS or PKC γ siRNA on MDA, SOD, GSH levels and NAD(P)H oxidase activity in PVN of salt-induced hypertensive and control groups (n = 7, for each group). Values are expressed as means SE. *P < 0.05 versus NS groups; [†]P < 0.05 HS + PVN LOS or HS + PVN PKC siRNA versus HS control groups.

Immunohistochemistry and immunofluorescence staining. Rats were anaesthetized and received a thoracotomy and were perfused through the left ventricle first with 300 mL of 0.01 M phosphate-buffered solution (PBS) at pH 7.4 and then with 300 mL of 4% paraformaldehyde. The brains were immediately removed and immersed in 4% paraformaldehyde, and then immersed in 30% sucrose for at least 2 days. Samples were embedded in OCT and microdissection procedures were used to isolate the PVN tissue. The tissues were collected from both sides of the PVN of individual rat and sectioned into several 18 mm transverse sections at about 1.80 mm from bregma and stored at −80 °C for future use for immunohistochemistry and immunofluorescence staining⁵.

Immunohistochemistry and immunofluorescence were performed as described previously². The primary antibodies (AT1R (sc-160811), PKC γ (sc-211), p-Rac1 (sc-12924-R), and Cu/Zn-SOD (sc-8637) were purchased from Santa Cruz Biotechnology. Superoxide anion levels in PVN were determined by fluorescent-labeled dihydroethidium (DHE; Molecular Probes) staining. Coronal sections (14 μ m) were incubated for 10 minutes with DHE (1 μ mol/L, Sigma) at 37 °C, as previously described⁵.

Rac1 activity assay. Rac1 activity was assayed using a pull-down assay (Millipore kit, Cat. #17-283) as described previously^{29,30}. Briefly, the PVN tissue lysate was prepared by homogenization in ice-cold 1 $\times$ lysis buffer (diluted from 5 $\times$ Mg²⁺ Lysis Buffer, MLB, Cat. #20-168, Millipore) and placed on ice for 30 min. After 30 min, the lysate was centrifuged at 14,000 g for 15 min and the supernatant was collected. The sample concentrations were determined using the BCA Protein Assay Kit (Beyotime Institute of Biotechnology). Protein p21-activated protein kinase 1 (Pak1) is an effect protein that can bind active Rac1. One mg of total protein was incubated with 10 μ l PAK-1 PBD (a GST fusion protein corresponding to the p21-binding domain (PBD) of human PAK-1) agarose beads for 1 h; the beads were then washed 3 times. The agarose beads were resuspended in 40 μ l of 2 $\times$ reducing sample buffer and boiled for 5 min. The bound proteins were separated by 12% SDS-PAGE. GTP-bound Rac1 was detected by immunoblotting with an anti-Rac1 antibody (Cat. #05-389, Millipore).

Western blotting. Rats were sacrificed and decapitated. The PVN were quickly collected on ice and then submerged in liquid nitrogen. Tissues were stored at −80 °C until further testing. Protein extracted from PVN tissue was used for measuring the expression. PVN tissue samples were lysed in a RIPA buffer with protease inhibitor and phosphatase inhibitor cocktail. Sonication the protein content of the resulting samples was determined by a modified BCA protein assay. The samples were then subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis using Electrophoresis and Blotting Apparatus (Bio-Rad). Protein products were separated by SDS-PAGE electrophoresis and transferred to nitrocellulose membranes. Then membranes were blocked with 3.0% BSA in TBST buffer (TBS plus 0.1% Tween-20) for 1.5 h at room temperature and incubated with primary antibodies overnight at 4 °C.

Rat genes	Forward (5'-3')	Reverse (5'-3')
AT1R	CAAAGGAGATGGGAGGTCA	TGACAAGCAGTTTGGCTTTG
PKC γ	AGGTGCTGAGAGCGAAGCTCCGC	CTTGCCCTGTCTTCCTATCTC
Cu/Zn-SOD	GGTGGGCCAAAGGATGAAGAG	CCACAAGCCAAACGACTTCC
GAPDH	AGACAGCCGCATCTTCTGT	CTTGCCGTGGGTAGAGTCAT

Table 3. Rat primers used for real-time PCR. AT1R, angiotensin II type 1 receptor; PKC γ , protein kinase C γ ; Cu/Zn-SOD, copper/zinc superoxide dismutase; GAPDH, Glyceraldehyde 3-phosphate dehydrogenase.

The primary antibodies for AT1R (sc-160811), PKC γ (sc-211), and Cu/Zn-SOD (sc-8637) were from Santa Cruz Biotechnology. Protein loading was controlled by probing all western blots with β -actin antibody which was from Santa Cruz Biotechnology and normalizing AT1R, PKC γ , and Cu/Zn-SOD protein intensities to that of β -actin. Band densities were analyzed using NIH Image J software¹².

Real-time PCR. The hypothalamic tissue containing PVN was dissected as described previously. Briefly, rat brains were isolated and cut into a coronal segment (-0.92 mm to -2.13 mm posterior to bregma). A block of the hypothalamus containing PVN was excised from the coronal section. Total RNA was extracted from micro-dissected PVN using Tri-Zol reagent (Invitrogen) and reverse transcribed using oligo (dT) with conditions at 23°C for 10 min, 37°C for 60 min, and 95°C for 5 min. The cDNA used for real-time PCR with specific primers for AT1R, PKC γ and Cu/Zn-SOD and GAPDH were shown in Table 3. The quantitative fold changes in mRNA expression were determined relative to glyceraldehyde-phosphate dehydrogenase (GAPDH) mRNA levels in each corresponding group^{6,15}.

ELISA studies. MDA, SOD, GSH and NAD(P)H oxidase activity in PVN were quantified using commercially available rat ELISA kits (Invitrogen Corporation, CA, USA) according to the manufacturer's instructions^{29–32}. NE in plasma were quantified using commercially available rat ELISA kits (Invitrogen Corporation, CA, USA) according to the manufacturer's instructions⁶.

Statistical analysis. All data are expressed as mean $\pm$ SE. The significance of differences between mean values was analyzed by ANOVA followed by Tukey's test. A probability value of $P < 0.05$ was considered to be statistically significant.

References

- Drenjancevic-Peric, I. *et al.* High-salt diet and hypertension: focus on the renin-angiotensin system. *Kidney & blood pressure research* **34**, 1–11, doi: 10.1159/000320387 (2011).
- Huang, B. S. *et al.* Activation of brain renin-angiotensin-aldosterone system by central sodium in Wistar rats. *American journal of physiology. Heart and circulatory physiology* **291**, H1109–H1117, doi: 10.1152/ajpheart.00024.2006 (2006).
- Huang, B. S., White, R. A., Bi, L. & Leenen, F. H. Central infusion of aliskiren prevents sympathetic hyperactivity and hypertension in Dahl salt-sensitive rats on high salt intake. *American journal of physiology. Regulatory, integrative and comparative physiology* **302**, R825–R832, doi: 10.1152/ajpregu.00368.2011 (2012).
- Sriramula, S., Cardinale, J. P., Lazartigues, E. & Francis, J. ACE2 overexpression in the paraventricular nucleus attenuates angiotensin II-induced hypertension. *Cardiovascular research* **92**, 401–408, doi: 10.1093/cvr/cvr242 (2011).
- Su, Q. *et al.* Alpha lipoic acid supplementation attenuates reactive oxygen species in hypothalamic paraventricular nucleus and sympathoexcitation in high salt-induced hypertension. *Toxicology letters* **241**, 152–158, doi: 10.1016/j.toxlet.2015.10.019 (2015).
- Li, H. B. *et al.* Central blockade of salusin beta attenuates hypertension and hypothalamic inflammation in spontaneously hypertensive rats. *Scientific reports* **5**, 11162, doi: 10.1038/srep11162 (2015).
- Zhang, M. *et al.* Endogenous hydrogen peroxide in the hypothalamic paraventricular nucleus regulates neurohormonal excitation in high salt-induced hypertension. *Toxicology letters* **235**, 206–215, doi: 10.1016/j.toxlet.2015.04.008 (2015).
- Kang, Y. M. *et al.* Paraventricular nucleus corticotrophin releasing hormone contributes to sympathoexcitation via interaction with neurotransmitters in heart failure. *Basic research in cardiology* **106**, 473–483, doi: 10.1007/s00395-011-0155-2 (2011).
- Huber, M. J. *et al.* Activation of the (pro)renin receptor in the paraventricular nucleus increases sympathetic outflow in anesthetized rats. *American journal of physiology. Heart and circulatory physiology* **309**, H880–H887, doi: 10.1152/ajpheart.00095.2015 (2015).
- Yu, X. J. *et al.* Interaction between AT1 receptor and NF- κ B in hypothalamic paraventricular nucleus contributes to oxidative stress and sympathoexcitation by modulating neurotransmitters in heart failure. *Cardiovascular toxicology* **13**, 381–390, doi: 10.1007/s12012-013-9219-x (2013).
- Gregg, D., Rauscher, F. M. & Goldschmidt-Clermont, P. J. Rac regulates cardiovascular superoxide through diverse molecular interactions: more than a binary GTP switch. *American journal of physiology. Cell physiology* **285**, C723–C734, doi: 10.1152/ajpcell.00230.2003 (2003).
- Su, Q. *et al.* Inhibition of reactive oxygen species in hypothalamic paraventricular nucleus attenuates the renin-angiotensin system and proinflammatory cytokines in hypertension. *Toxicology and applied pharmacology* **276**, 115–120, doi: 10.1016/j.taap.2014.02.002 (2014).
- Garrido, A. M. & Griendling, K. K. NADPH oxidases and angiotensin II receptor signaling. *Molecular and cellular endocrinology* **302**, 148–158, doi: 10.1016/j.mce.2008.11.003 (2009).
- Zimmerman, M. C. *et al.* Requirement for Rac1-dependent NADPH oxidase in the cardiovascular and dipsogenic actions of angiotensin II in the brain. *Circulation research* **95**, 532–539, doi: 10.1161/01.RES.0000139957.22530.b9 (2004).
- Xiang, L. *et al.* Mixture of Peanut Skin Extract and Fish Oil Improves Memory in Mice via Modulation of Anti-Oxidative Stress and Regulation of BDNF/ERK/CREB Signaling Pathways. *Nutrients* **8**, doi: 10.3390/nu8050256 (2016).
- Zhang, H. M. & Su, Q. PKC in developmental hypothyroid rat brain. *Neurological sciences: official journal of the Italian Neurological Society and of the Italian Society of Clinical Neurophysiology* **35**, 1161–1166, doi: 10.1007/s10072-014-1716-6 (2014).
- Chou, W. H. & Messing, R. O. Protein kinase C isozymes in stroke. *Trends in cardiovascular medicine* **15**, 47–51, doi: 10.1016/j.tcm.2005.01.003 (2005).

18. Aronowski, J. & Labiche, L. A. Perspectives on reperfusion-induced damage in rodent models of experimental focal ischemia and role of gamma-protein kinase C. *ILAR journal/National Research Council, Institute of Laboratory Animal Resources* **44**, 105–109 (2003).
19. Aronowski, J., Grotta, J. C., Strong, R. & Waxham, M. N. Interplay between the gamma isoform of PKC and calcineurin in regulation of vulnerability to focal cerebral ischemia. *Journal of cerebral blood flow and metabolism: official journal of the International Society of Cerebral Blood Flow and Metabolism* **20**, 343–349, doi: 10.1097/00004647-200002000-00016 (2000).
20. Koga, Y. *et al.* High salt intake enhances blood pressure increase during development of hypertension via oxidative stress in rostral ventrolateral medulla of spontaneously hypertensive rats. *Hypertension research: official journal of the Japanese Society of Hypertension* **31**, 2075–2083, doi: 10.1291/hypres.31.2075 (2008).
21. Lin, D. & Takemoto, D. J. Oxidative activation of protein kinase Cgamma through the C1 domain. Effects on gap junctions. *The Journal of biological chemistry* **280**, 13682–13693, doi: 10.1074/jbc.M407762200 (2005).
22. Carrizzo, A. *et al.* Rac-1 as a new therapeutic target in cerebro- and cardio-vascular diseases. *Current drug targets* **15**, 1231–1246 (2014).
23. Madamanchi, N. R., Vendrov, A. & Runge, M. S. Oxidative stress and vascular disease. *Arteriosclerosis, thrombosis, and vascular biology* **25**, 29–38, doi: 10.1161/01.ATV.0000150649.39934.13 (2005).
24. Campese, V. M., Ye, S., Truong, R. H. & Gamburd, M. Losartan reduces sympathetic nerve outflow from the brain of rats with chronic renal failure. *Journal of the renin-angiotensin-aldosterone system: JRAAS* **1**, 202–208, doi: 10.3317/jraas.2000.026 (2000).
25. Yu, Y. *et al.* Early interference with p44/42 mitogen-activated protein kinase signaling in hypothalamic paraventricular nucleus attenuates angiotensin II-induced hypertension. *Hypertension* **61**, 842–849, doi: 10.1161/HYPERTENSIONAHA.111.00080 (2013).
26. Kang, Y. M. *et al.* TNF-alpha in hypothalamic paraventricular nucleus contributes to sympathoexcitation in heart failure by modulating AT1 receptor and neurotransmitters. *The Tohoku journal of experimental medicine* **222**, 251–263 (2010).
27. Dange, R. B. *et al.* Central blockade of TLR4 improves cardiac function and attenuates myocardial inflammation in angiotensin II-induced hypertension. *Cardiovascular research* **103**, 17–27, doi: 10.1093/cvr/cvu067 (2014).
28. Elks, C. M. *et al.* A blueberry-enriched diet attenuates nephropathy in a rat model of hypertension via reduction in oxidative stress. *PloS one* **6**, e24028, doi: 10.1371/journal.pone.0024028 (2011).
29. Kamataki, T., Naminohira, S., Sugita, O. & Kitagawa, H. Lipid peroxidation activity mediated by NADPH-cytochrome C reductase purified from rabbit liver microsomes. *Japanese journal of pharmacology* **28**, 819–827 (1978).
30. Kono, Y. Generation of superoxide radical during autooxidation of hydroxylamine and an assay for superoxide dismutase. *Archives of biochemistry and biophysics* **186**, 189–195 (1978).
31. Cheng, P. W. *et al.* Resveratrol Inhibition of Rac1-Derived Reactive Oxygen Species by AMPK Decreases Blood Pressure in a Fructose-Induced Rat Model of Hypertension. *Scientific reports* **6**, 25342, doi: 10.1038/srep25342 (2016).
32. Sedlak, J. & Lindsay, R. H. Estimation of total, protein-bound, and nonprotein sulphhydryl groups in tissue with Ellman's reagent. *Analytical biochemistry* **25**, 192–205 (1968).

Acknowledgements

This work was supported by National Natural Science Foundation of China (Nos 81370356, 91439120, 81471471 and 31271213) and China Postdoctoral Science Foundation (grant numbers 2016M592802).

Author Contributions

Yu-Ming Kang, Wen-Sheng Chen, Jin-Jun Liu and Xiao-Lian Shi designed the study. Qing Su, Guo-Qing Zhu, Wei Cui and Hong-Bao Li performed the data analysis and drafted the manuscript. Qing Su, Kai-Li Liu, Xiang Li, Chan-Juan Huo, Xin-Ai Song and Qing Yang performed all experiments and participated in data analysis. All authors reviewed the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Su, Q. *et al.* Renin-angiotensin system acting on reactive oxygen species in paraventricular nucleus induces sympathetic activation via AT1R/PKCγ/Rac1 pathway in salt-induced hypertension. *Sci. Rep.* **7**, 43107; doi: 10.1038/srep43107 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017