

SCIENTIFIC REPORTS



OPEN

OGG1-DNA interactions facilitate NF- κ B binding to DNA targets

Lang Pan^{1,2}, Wenjing Hao^{1,2}, Xu Zheng^{1,2}, Xianlu Zeng^{1,2}, Adeel Ahmed Abbasi¹, Istvan Boldogh³ & Xueqing Ba^{1,2}

Received: 10 October 2016

Accepted: 20 January 2017

Published: 07 March 2017

DNA repair protein counteracting oxidative promoter lesions may modulate gene expression. Oxidative DNA bases modified by reactive oxygen species (ROS), primarily as 7, 8-dihydro-8-oxo-2'-deoxyguanosine (8-oxoG), which is repaired by 8-oxoguanine DNA glycosylase1 (OGG1) during base excision repair (BER) pathway. Because cellular response to oxidative challenge is accompanied by DNA damage repair, we tested whether the repair by OGG1 is compatible with transcription factor binding and gene expression. We performed electrophoretic mobility shift assay (EMSA) using wild-type sequence deriving from *Cxcl2* gene promoter and the same sequence bearing a single synthetic 8-oxoG at defined 5' or 3' guanine in runs of guanines to mimic oxidative effects. We showed that DNA occupancy of NF- κ B present in nuclear extracts from tumour necrosis factor alpha (TNF α) exposed cells is OGG1 and 8-oxoG position dependent, importantly, OGG1 counteracting 8-oxoG outside consensus motif had a profound influence on purified NF- κ B binding to DNA. Furthermore, OGG1 is essential for NF- κ B dependent gene expression, prior to 8-oxoG excised from DNA. These observations imply that pre-excision step(s) during OGG1 initiated BER evoked by ROS facilitates NF- κ B DNA occupancy and gene expression.

Arising through a plethora of biological matrices and a variety of settings, reactive oxygen species (ROS) are well recognized for representing a double-edged sword as both deleterious and beneficial species¹. ROS cause predominantly base damages that do not occur randomly, and exhibit a strong bias distribution in genome^{2,3}. Due to their lowest ionization potential among DNA components⁴, guanine bases, especially two- and more-base runs of guanines, e.g., 5'G < 5'GG < 5'GGG, are preferentially targeted by oxidants, and readily oxidizes to 7, 8-dihydro-8-oxo-2'-deoxyguanosine (8-oxoG)^{5,6}. 8-OxoG is often used as a cellular biomarker to indicate the extent of oxidative stress^{7,8}. Differing from normal guanine only by a single atom change, 8-oxoG does neither distort DNA helix structure nor perturb DNA or RNA polymerases function⁹. Such damage usually recognized precisely by 8-oxoguanine DNA glycosylase1 (OGG1) and repaired through base excision repair (BER) pathway^{10,11}. Several studies have documented that the key searching step for damaged site in BER involves nonspecific DNA binding with induced destabilization by bending, and subsequent flipping of the lesion into the enzyme active site pockets^{12–14}. OGG1 is a bifunctional DNA glycosylase endowed with an AP lyase activity. Following oxidized base removed, OGG1 cleaves DNA phosphodiester backbone at the 3' side of the abasic site, producing a nick in the DNA backbone^{15,16}.

Both DNA glycosylases and BER pathway are generally well conserved in evolution, tailoring to deal with the special features of base lesions. Most notably, the principal substrate for OGG1 recognizing is 8-oxoG paired with C, while a diverse array of oxidized DNA bases are repaired in the enzymic system, for example, hydantoin lesions are the best substrates for *Nei*-like DNA glycosylases (NEIL1–2)^{17–19}. Additional studies reported that *Neil1*^{-/-} mice showed enhanced inflammatory responses upon chronic UV exposure relative to wild-type (Wt) controls²⁰. By contrast, *Ogg1*^{-/-} mice exhibited lower expression of pro-inflammatory cytokines and inflammatory cells infiltration, even though resulting high levels of genomic 8-oxoG, embryonic development or life span were not affected^{21–24}. The non-random distribution of 8-oxoG in genome and substrate specificity of OGG1 suggest that 8-oxoG and OGG1 initiated BER may play an important role in physiological and pathophysiological processes.

In addition to the extensive studies devoted to the role of oxidative nuclear DNA damage in coding sequences, the outcome of oxidative base lesions in non-coding regions remains to be elucidated. Intriguingly,

¹The Key Laboratory of Molecular Epigenetics of the Ministry of Education, Northeast Normal University, Changchun, Jilin 130024, China. ²Institute of Genetics and Cytology, Northeast Normal University, Changchun, Jilin 130024, China. ³Department of Microbiology and Immunology, Sealy Center for Molecular Medicine, University of Texas Medical Branch at Galveston, Galveston, Texas 77555, USA. Correspondence and requests for materials should be addressed to X.B. (email: baxq755@nenu.edu.cn)

vertebrate genome evolutionarily has a high GC content in the promoter regions of RNA polymerase II (RNA Pol II)-transcribed genes. Moreover, several transcription factors (TFs) (NF- κ B, SP1, AP1 and NRF1) binding sites are rich in guanines, for example, the κ B site recognized by NF- κ B with a conserved sequence of 5'-GGGRNYYYCC-3' (R is an unspecified purine; Y is an unspecified pyrimidine; and N is any nucleotide²⁵), which are putatively vulnerable to oxidants. It is well documented that changes in *cis* elements result from oxidative modification may modulate the pattern of TFs binding^{26,27}. In contrast to that *cis* effects, however, how DNA repair protein acting on oxidative bases affects promoter function remains elusive.

Here, we propose OGG1 counteracting 8-oxoG produced on key nucleotide in *Cxcl2* promoter benefits NF- κ B binding to its consensus motif. Previously, we have shown that OGG1 and NF- κ B/RelA coimmunoprecipitated from *Cxcl2* promoter²⁸. But due to the major impediment to the ChIP-based evaluation is identifying genomic regions that interact with TFs but not individual binding sites²⁹, it prompts us to address the issue in site resolution. In the present study, we generated several duplex oligonucleotides corresponding to *Cxcl2* gene promoter containing putative κ B motif, one without (Wt) and the rest with single 8-oxoG situated at defined 5' or 3' guanine in runs of guanines to mimic ROS-induced oxidative damage occurrence. We performed electrophoretic mobility shift assay (EMSA) using nuclear extracts from mock and TNF α treated MLE-12 cells and recombinant proteins. We found that OGG1 recognizing 8-oxoG introduced outside cognate site facilitates NF- κ B-DNA association robustly. The biological consequences of oxidants attack within *Cxcl2* promoter and the contribution of OGG1 counteracting 8-oxoG are facilitating NF- κ B transcriptional activation. The results presented here provide novel insight into the ways in which organisms utilize oxidized guanines to regulate gene expression in a time-frame of DNA damage response.

Results

OGG1 plays a key role in providing the transcriptional activity from the *Cxcl2* promoter. Our previous work has established that OGG1 can augment a subset of cytokine and chemokine gene expression in mouse lung^{28,30,31}, including CXCLs, CCLs, ILs and TNF α . To confirm that OGG1 facilitates gene expression through promoter activation, we carried out small interfering RNA (siRNA)-mediated *Ogg1* knockdown and monitored how it affects luciferase activity. We were able to attain ~70% siOgg1 knockdown compared with control non-specific siRNA (Fig. 1a, insert). The activation of *Cxcl2* promoter revealed ~4 fold increase after 6 h of TNF α treatment, while in OGG1 deficient cells, inducible luciferase activity was significantly decreased (Fig. 1a). To confirm these results, we used *Ogg1*^{-/-} and *Ogg1*^{+/+} MEFs and showed that TNF α induced significantly less luciferase expression in *Ogg1*^{-/-} than wild type ones (Fig. 1b). To address the impact of OGG1 at transcription level, luciferase mRNA level was determined from reporter plasmids transfected cells over one hour TNF α exposure. RT-qPCR analyses showed a considerable increase of luciferase transcript in OGG1 expressing cells (~8 fold in MLE-12 cells, Fig. 1c; ~5 fold in MEF cells, Fig. 1d), while OGG1 depletion dampened transcript levels. Pre-treatment of cells with ROS scavenger N-acetyl-L-cysteine (NAC, GSH precursor) abrogated luciferase expression, suggesting oxidative DNA bases (e.g., 8-oxoG) were important for efficient *Cxcl2* gene expression in OGG1 expressing cells (Fig. 1e).

OGG1 is essential for NF- κ B-mediated transcriptional regulation. Based on previous work, there are two NF- κ B (-66 to -57, GGGAATTTCC, designated as κ B2; -47 to -38, GGGCTTTTCC, designated as κ B1) and one SP1 (-109 to -100, GGGGCGGGGC) putative binding sites immediately adjacent to TATA box in *Cxcl2* promoter²⁸. To understand the functional role of potential *cis* regulatory elements in promoter activation, luciferase reporter gene constructs containing *Cxcl2* promoter fragments (depicted in Fig. 2a) and with one κ B site mutation at each time were transfected into MLE-12 cells following with or without TNF α treatment. MLE-12 cells carrying the intact *Cxcl2* gene promoter (Wt-Luc) conferred strong TNF α responsiveness (Fig. 2b). The mutation of proximal κ B site to a null sequence hardly affected inducible luciferase activity (Fig. 2b, κ B2-Luc), while mutation of distal κ B site abrogated inducible luciferase activity significantly (Fig. 2b, κ B1-Luc), implying that κ B2 site is indispensable for *Cxcl2* promoter activation. When both two κ B sites were mutated to null sites, no inducible luciferase activity showed following TNF α treatment (Fig. 2b, Null-Luc), suggesting that NF- κ B binding sites are crucial for *Cxcl2* promoter-driven gene expression. These reporter plasmid variants were also transfected to MEF cells, and the patterns were similar to those in MLE-12 cells (Fig. 2c).

To examine whether OGG1 acts on these κ B sites differently, we carried out siRNA-mediated *Ogg1* knockdown and monitored how it affects luciferase activity from mutated κ B site reporter plasmids expressing cells. It showed that lacking of OGG1 decreased inducible luciferase activity both in κ B2-Luc and κ B1-Luc transfected cells (Fig. 2d,e). Of note, compared with non-treated situation, inducible luciferase activity in κ B1-Luc transfected cells increased ~4-fold, implying other transcriptional factors, like SP1 participates in gene expression; OGG1 knockdown represses this induction to basal level, indicating that OGG1 may also regulate the performance of other *trans* acting factors.

NF- κ B-DNA associations are increased in nuclear extracts from OGG1 expressing cells. To examine whether NF- κ B binds to the putative sites from *Cxcl2* promoter, we carried out electrophoretic mobility shift assay (EMSA) using seven double-stranded 33 bp oligonucleotides containing κ B2 site (5'-GGAATTTCC-3') and four double-stranded 31 bp oligonucleotides containing κ B1 site (5'-GGCTTTTCC-3'). An efficient inducible activity was observed as strong retarded complex was detected using nuclear extracts (NE) from TNF α -exposed cells with κ B2-Wt oligo, while NF- κ B binding to same oligo was under detectable level in NE prepared from mock-treated cells (Fig. 3a, lanes 1 and 2). Compared with NF- κ B-Wt oligo complex, 8-oxoG modified at G1 showed nearly equal binding, while κ B2-G2 showed no retarded complex, implying that OGG1 recognizing 8-oxoG within κ B site competes with NF- κ B on motif binding (Fig. 3a, lanes 3 and 4). The level of protein-DNA complex formed with 8-oxoG modified upstream of κ B site was far greater than Wt, ~12-fold using κ B2-G3 and




Figure 1. *Ogg1* depletion decreases *Cxcl2* promoter driven luciferase activity. (a,b) Luciferase expression driven by *Cxcl2* promoter is dependent on *Ogg1* in MLE-12 (a), and MEF cells (b). All cells were exposed to $\text{TNF}\alpha$ for six hours. Insert: *Ogg1*-siRNA mediated *Ogg1* knockdown in MLE-12 cells (WB analysis). (c,d) Decreased luciferase mRNA levels in *Ogg1*-siRNA transfected MLE-12 cells (c) and *Ogg1*^{-/-} MEF cells (d) upon $\text{TNF}\alpha$ exposure. Whole RNAs were isolated at 1 h post-exposure and luciferase mRNA levels were determined by RT-qPCR. (e) $\text{TNF}\alpha$ -induced promoter activation is inhibited by antioxidant pre-treatment in MLE-12 cells. RLU, relative luciferase unit. ** $p < 0.01$, *** $p < 0.001$ (n = 3).

G4; G6 8-oxoG modified at reverse strand bind $\text{NF-}\kappa\text{B}$ ~10-fold to Wt (Fig. 3a, lanes 5, 6 and 8). On the contrary, $\kappa\text{B2-G5}$ 8-oxoG, just adjacent κB site modified at reverse strand, showed slightly decreased $\text{NF-}\kappa\text{B}$ binding over that of Wt (Fig. 3a, lane 7). In order to test if *OGG1* in NE was responsible for altered $\text{NF-}\kappa\text{B}$ binding, we carried out siRNA-mediated *Ogg1* knockdown and tested the effects on $\text{NF-}\kappa\text{B}$ -DNA association. We found that the inducible $\text{NF-}\kappa\text{B}$ -DNA association significantly decreased ($\kappa\text{B2-G3}$, G4 and G6 oligoes) to those of *OGG1* expressing ones (Fig. 3a, lanes 12, 13 and 15). Whether *OGG1* depleted or not, $\text{NF-}\kappa\text{B}$ binding to $\kappa\text{B2-G2}$ is not affected (Fig. 3a, lane 11). These extensive retarded complexes were specific $\text{NF-}\kappa\text{B}$ contained, as in each case, binding was blocked by unlabelled probe competitor (Fig. 3a, lanes 20–23).

In the case of κB1 oligoes, Wt oligo shifted obvious retarded complex with $\text{TNF}\alpha$ -treated NE; 8-oxoG introduced at G1 decreased that association slightly, while G2 and G3 8-oxoG introduced outside κB site shifted ~17-fold $\text{NF-}\kappa\text{B}$ to Wt (Fig. 3b, lanes 1–9). Competition assay showed those protein-DNA complexes were competed out by cold probe (Fig. 3b, lanes 14–16).

In case that deficient $\text{NF-}\kappa\text{B}$ binding was due to the impairment of p65 translocation, we tested p65 protein level in the cytoplasmic and nuclear fraction from *Ogg1* knockdown cells. Westernblotting revealed that cytoplasmic levels of p65 decreased and nuclear levels increased after induction, suggesting that *OGG1* does not govern p65 nuclear importation (Fig. 3e). Taken together, these observations suggest a dominant role of *OGG1* in modulating $\text{NF-}\kappa\text{B}$ binding to its DNA targets.

OGG1-DNA interactions enhance $\text{NF-}\kappa\text{B}$ binding synergistically. To provide additional evidence for *OGG1* modulating $\text{NF-}\kappa\text{B}$ binding through interaction with DNA, we utilized recombinant $\text{NF-}\kappa\text{B}$ subunits and performed EMSA either singly or in the presence of purified *OGG1*. Purified p65 and p50 proteins were pre-incubated at 37 °C for 60 min, so that p65/p50 heterodimer and p50/p50 homodimer were allowed to form³², after that EMSA was carried out with κB2 -Wt oligo added at RT within 10 min time course. Results showed that p50/p50 homodimer binds the probe from 1 min on, and gradually increase until 10 min, while obvious p65/p50 heterodimer-DNA complex appeared at 5 min and then approached an equilibrium until 10 min (Fig. 4a). Similar data were obtained using κB1 -Wt oligo (data not shown). Therefore, we chose 10 min as the incubation time for analysing the binding properties of purified protein in EMSA.




Figure 2. OGG1 is essential for NF- κ B directed promoter activation. (a) Graphical depiction of *Cxcl2* promoter constructs: a wild-type and three mutated *Cxcl2* promoter fragments were cloned to drive luciferase expression as described in Materials and Methods. (b,c) Luciferase expression driven by Wt and mutated *Cxcl2* promoter. Constructs were transfected into MLE-12 (b) and MEF (c) cells. TNF α treated for 6 h and dual luciferase assays were performed as described in Materials and Methods. (d,e) Effect of Ogg1 on mutated Luc expression. κB2-Luc and κB1-Luc constructs were transfected into Ogg1 expression and siRNA-mediated knockdown MLE-12 cells. TNF α treated for 6 h and dual luciferase assays were performed as described above. RLU, relative luciferase unit. ***p < 0.001 (n = 3).

Figure 4b shows that p50/p50 homodimer bound to κB2 oligos with varying efficiency, and p65/p50 heterodimer recognizes oligos with bias, preferring κB2-G3 and G4 rather than κB2-G2 (Fig. 4b, lanes 1–7). After adding purified OGG1, the association of both homo- and heterodimer with oligos (except κB2-G2) was stronger than NF-κB alone (Fig. 4b, lanes 8–14). Especially, NF-κB binding to κB2-G3, G4 and G6 exhibited higher level than Wt, implying OGG1 encountering 8-oxoG outside κB motif increases NF-κB binding to DNA. In the presence of OGG1, NF-κB binding to κB2-G2 was decreased compared to Wt, suggesting 8-oxoG introduced within κB motif and recognized by OGG1 would compromise NF-κB binding. NF-κB alone could recognize four κB1 oligos (Fig. 4c, lanes 1–4), after adding OGG1, both homo- and heterodimer binding were increased, among which κB1-G3 showed highest shifted amount (Fig. 4c, lanes 5–8).

As we showed above, OGG1 significantly affects NF-κB binding to κB2-G6 oligo and efficiently bound to that oligo itself, we chose κB2-G6 for further studies. Purified p65 and p50 were pre-incubated, followed by mixing with increasing amounts of OGG1 and EMSA was performed. It showed that the occupancies of both hetero- and homodimer on κB2-G6 oligo were increased in OGG1 concentration dependent manner (Fig. 4d, lanes 1–3). Amazingly, OGG1-DNA complex was significantly less in absence of NF-κB (Fig. 4d, lanes 4–6). We can conclude from these results that there is a bidirectional interaction between OGG1 and NF-κB, OGG1 facilitates NF-κB assembly on DNA and NF-κB reciprocally increases OGG1-DNA association. To examine the effect of OGG1 on NF-κB subunit, p50 and p65 were pre-incubated individually and then mixed with purified OGG1. p65 alone bound to oligo only at high concentration (15 ng/sample) (Fig. 4d, lane 9), which may represent p65/p65 homodimer³³. Importantly, DNA occupancy of recombinant p65 increased more than 10-fold in the presence of OGG1 (Fig. 4d, lanes 12 and 15). Interestingly, OGG1 didn't shift κB2-G6 oligo with low concentration of p65 (Fig. 4d, lanes 11 and 12, 13 and 14), only at 15 ng of purified p65 could show obvious OGG1-DNA complex (Fig. 4d, lanes 12 and 15). The level of p50/p50 homodimer-DNA complex was enhanced mildly by the lowest OGG1 concentration (1 ng/sample), which was not further increased by higher amounts (Fig. 4d, lanes 16–19). OGG1-mediated oligo shifts were observed at indicating concentration. Surprisingly, purified OGG1 didn't affect NF-κB-DNA complex migration, instead, independent OGG1 shifts were observed with 8-oxoG containing oligos (Fig. 4b, c and d).

In independent experiments, the ability of purified OGG1 binding to 8-oxoG-introduced κB2 and κB1 oligos were also determined (Fig. 4e). Owing to insufficient flanking bases for OGG1 docking in κB2-G3, G4 and G5 oligos, OGG1 shifts those oligos with higher amounts (data not shown).




Figure 4. OGG1 enhances NF- κ B-DNA association. (a) Kinetic NF- κ B homo- and heterodimer binding to κ B2-Wt oligo. EMSA shows purified p65 (3.75 ng/sample) and p50 (2.75 ng/sample) proteins binding to κ B2-Wt probe over a 10 min time course. (b,c) Effects of OGG1 encountering 8-oxoG on NF- κ B home- and heterodimer binding to κ B2-oligos (b) and κ B1-oligos (c). Purified p65 (3.75 ng/sample) and p50 (2.75 ng/sample) were pre-incubated and oligoes were added with or without purified OGG1 (4 ng/sample) after 10 min incubation. (d) The effect of OGG1 on NF- κ B subunit binding to κ B2-G6 oligo. Lanes 1–3, recombinant p65 (3.75 ng/sample) and p50 (2.75 ng/sample) were incubated with increasing amounts of OGG1 (1, 2 and 4 ng), and then mixed with oligo at RT for 10 min. Lanes 4–6, increasing amounts of OGG1 (1, 2 and 4 ng) binding to κ B2-G6 oligo. Lanes 7–15, pre-incubated p65 (3.75, 7.5 and 15 ng respectively) were mixed alone or with 2 or 4 ng of purified OGG1, followed by adding κ B2-G6 oligo. Lanes 16–19, pre-incubated p50 (2.75 ng/sample) were mixed with increasing amounts of OGG1 (1, 2 and 4 ng respectively), followed by an EMSA with κ B2-G6 oligo. < denotes p65/p50 heterodimer-DNA complex, * denotes p50/p50 homodimer-DNA complex, triangle indicates OGG1-DNA complex, NS denotes non-specific bands. (e) Purified OGG1 binds to 8-oxoG containing DNA. EMSA was performed using increasing concentrations of purified OGG1 (0, 2, 4 and 8 ng respectively) incubated with indicated 8-oxoG containing oligoes at RT for 10 min.

macromolecules, especially DNA base lesions that occur through oxidative reactions. Taking that not only activation of transcription factors and their nuclear translocation, but their access to the binding motifs in the chromatin is actively controlled, 8-oxoG and its repair complex (OGG1-DNA interactosome) perform in gene regulatory regions could provide another insight in gene expression.

What we demonstrate here is simulating one way of mediating transcription through OGG1-DNA interactosome. Implicit in that notion are extensions of three: first, 8-oxoG in response elements themselves can act as *cis* effects that has either no effect, induces a full or partial inhibition or enhances TFs binding; second, OGG1 surveying the genome for damaged bases can act as *trans* effects that facilitates TFs binding to DNA synergistically; third, the combination of 8-oxoG and OGG1 determines the levels of TFs binding, in part by which selective gene transcription may be achieved.

We observed varying NF- κ B binding as a function of 8-oxoG in oligoes, which is not surprising as the *cis* effects are the most studied and offer a good argument for the significant role of oxidative damage interference in promoter functions. 8-OxoG formation in DNA response elements has the capacity to interfere with normal gene regulation on the basis of changes in the binding affinity of TFs. The affinity changes are either increased, unchanged, or diminished that depend on various factors, including the particular TF-promoter system under investigation (e.g., SP1, AP1, or NF- κ B) and the precise site of damage within the consensus binding motif^{26,27}.




Figure 5. OGG1 is required for a subset of pro-inflammatory gene expression upon TNF α stimulation. (a) Changes in global 8-oxoG levels assayed by dot blot analysis. Genomic DNA was isolated at times indicated and immunoblotted as described in Materials and Methods. (b) Changes in OGG1 substrate levels in *Cxcl2*, *Tnf* and *Il-1 β* proximal promoter regions after TNF α challenge. DNAs were isolated at 0 and 30 min after TNF α exposure and promoter regions were amplified by RT-qPCR after mock (–) and OGG1 digestion (+). (c) Time course of gene expression upon TNF α exposure of MLE-12 cells. RNAs were isolated at times indicated and mRNA levels were determined by RT-qPCR. (d,e) *Ogg1* depletion decreases gene expression in MLE12 (d) and MEF (e) cells. Parallel cultures of MLE12 cells were transfected with control or *Ogg1*-siRNA, after which TNF α treated for 30 min. RNAs were isolated at times indicated and mRNA levels were determined by RT-qPCR. (f) TNF α -induced gene activation is inhibited by antioxidant pre-treatment. *** $p < 0.001$ (n = 3).

Although the mechanism(s) of affinity changes are not well defined, they most likely involve 8-oxoG:C pairs showing more increased local and global flexibility than G:C⁴⁴.

Another more intriguing aspect of DNA damage outcome on gene regulation involves *trans* effects from DNA binding proteins. In exposure of oxidative stress, damage-specific DNA glycosylase OGG1 searches oxidized guanines without any delay and initiates BER, through which small base lesions that do not significantly distort DNA double helix are corrected. Previously, we have shown that OGG1 is non-productively bound to oxidative lesions and its excision activity is inhibited in response to TNF α treatment²⁸. In the present study, we showed that OGG1 could facilitate NF- κ B binding to its DNA targets through recognizing 8-oxoG outside consensus motif, the mechanism by which is unknown at present. The crystallographic observation of OGG1-DNA complex shed some light upon the biological effects of OGG1 on modifying the allosteric effect of DNA. Verdine and co-workers have revealed that OGG1 bends DNA sharply when it is bound to an 8-oxoG containing site, giving rise to a pronounced kink in duplex, and more importantly, also showed that OGG1 induces similar bending when it binds to undamaged sites ($\sim 70^\circ$)^{45–47}. Impressively, a universal feature of most DNA repair glycosylases, especially for OGG1, is the free energy cost for DNA bending during base flipping since it takes place in the presence of only protein, DNA, and the cofactor S-adenosylmethionine^{13,48}. It is quite remarkable that bending or kinking of DNA facilitates NF- κ B binding, presumably because flexible duplex structure overcomes the energy barrier upon bending⁴⁹. We suspect that OGG1 bent DNA when searching for damaged sites, lowering the energy cost for NF- κ B twisting DNA and creating a stereo-specific interface that suited NF- κ B recognizing consensus motif. Deficiency in UNG did not decrease pro-inflammatory gene expression upon TNF α exposure encouraged us to test uracil levels deaminated from cytosine in the future. Our results showing OGG1 counteracting 8-oxoG situated outside κ B motif increased NF- κ B-DNA association indicate that much larger segments than consensus recognition sites need to take into account for individual TF function. Note that DNA glycosylases process on genome without sequence bias^{13,45}, we observed OGG1 regulates promoter activity at both canonical and non-canonical κ B sites.

The combination of 8-oxoG and OGG1 determines the levels of TFs binding on DNA. 8-OxoG limits the position of OGG1 where TFs binding would be benefited. In that case, OGG1 recognizing 8-oxoG produced within TFs binding site shoves corresponding TF by competition, while outside motifs facilitates TFs binding robustly. The final outcome for TF binding in response to stimuli might compromise those two parts *in vivo*.

It's still an enigma that purified OGG1 didn't slower the mobility of NF- κ B-DNA complex in EMSA. One possibility is that the high mobility of OGG1 ensures its availability throughout the genome and makes the

Oligo Sequences	Site
5'-TGAG ³ GGG ⁴ ACCCTGAGCTCA G¹GG²AATTTC C CTGG-3'	κB2
3'-ACTCCCCTGGG ⁶ ACTCGAGT CCCTTAAAGG G ⁵ ACC-5' Biotin	
5'-TTCCTGGTCCCC G¹GGCTTTTC C AGACATCG-3'	κB1
3'-AAGGG ³ ACCAGGG ² CCCGAAAAGG TCTGTAGC-5' Biotin	

Table 1. The sequence of oligoes used in this study. Consensus motif is in bold. Wt refers to no 8-oxoG modified oligoes, and the position of 8-oxoG is shown by numbers above. Each oligo contains a single 8-oxoG modification. The κB2 oligoes contain classic κB motif GGGAAATTTC, named κB2-G1 to G6; κB1 oligoes contain GGGCTTTTCC binding site named κB1-G1 to G3, differing only by a cytosine in the position 4 from the canonical sequence GGGRNYYCC (R = Purine, W = A or T, N = Any nucleotide, Y = Pyrimidine)²⁵.

transcription apparatus highly dynamic, which is the common property for nuclear proteins⁵⁰. TFs binding to consensus motifs normally proscribed by the energetic cost of DNA bending and twisting within its persistence length, the function of OGG1 is more like an “architectural” protein that allows TF-DNA interactions. Carey also suggested that architectural proteins can be bypassed if the strength or flexibility of the interactions can absorb the energy cost of DNA distortion⁵¹.

In conclusion, we were alert for the possibility that DNA glycosylases are the first sensor detecting physiological changes and taking rapid responses, beckoning TFs like NF-κB to locate their sites. Our data provided here may be viewed that oxidation of guanines in gene regulatory regions may be utilized by transcriptional machinery. Our future goal is to ascertain the exact DNA conformational changes caused by specific glycosylase that are required for TF binding, and the potential cooperative action(s) of other substrate-specific DNA glycosylases function in gene regulation. Owing to the paucity of our knowledge concerning the function of OGG1 in transcriptional response and the likelihood that increased knowledge of TFs will lead to increased insight into the causes of human diseases, it is of utmost importance that we expand our understanding of how site-specific DNA glycosylases contribute to gene regulation.

Methods and Materials

Cell culture and treatments. MLE-12 (an immortalized type 2 mouse lung epithelial cell line), *Ogg1*^{-/-} and *Ogg1*^{+/+} mouse embryo fibroblasts (MEFs) were kindly provided by Professor Istvan Boldogh (University of Texas Medical Branch at Galveston, Texas). Cells at about 80% of confluence were treated with 20 ng/ml recombinant human TNFα (Cat # H8916; Sigma) for indicated time course at 37 °C. In the case of antioxidant treatment, cells were pre incubated for 1 h in the presence of 10 mM N-acetyl cysteine (NAC) (Cat # A7250; Sigma)⁵² followed by TNFα treatment.

Depletion of target gene expression. For RNA interference experiments, scramble control or smart-pool siRNA against *Ogg1* and *Ung* (Dharmacon) were transfected into MLE-12 using Lipofectamine 2000 (Invitrogen) as previously described²⁸. Cells were used for experiments after 48 h of incubation, and target gene knockdown efficiency was validated by Western blot.

Luciferase reporter assays. Wt-Luc containing mouse *Cxcl2* promoter (−571 to +81) was generate as previously described²⁸ and mutant versions κB2-Luc and κB1-Luc were created using Fast Mutagenesis System (Cat # FM111, TransGen Biotech). MLE-12, wild-type and *Ogg1* deficient MEF were transiently transfected with luciferase reporter DNA and Renilla luciferase expression plasmid was co-transfected as an internal control. The total amount of plasmid DNA was kept constant for all assays. Transient transfections were carried out using Lipofectamine 2000 (Invitrogen) and luciferase activities of total cell lysates were measured using the dual-luciferase reporter assay system (Promega) following the manufacturer’s instructions.

Preparation of cell nuclear extracts (NE). The preparation an extract from nuclei was made using CelLytic™ NuCLEAR™ Extraction Kit (Cat # NXTRACT, Sigma) and total protein was quantified by Pierce BCA Protein Assay Kit (Cat # 23225, Thermo Scientific).

Electrophoretic mobility shift assay (EMSA). Consensus oligonucleotides containing the binding sites for NF-κB were synthesized at Sangon Biotech (Shanghai, China), and site-specific guanine residue within each of the oligonucleotide was replaced with single 8-oxoG modification (Table 1). EMSA was performed using LightShift® Chemiluminescent EMSA Kit (Cat # 20148, Thermo Scientific). For binding reactions, 10 fmol Biotin-labeled probe was mixed with 1 μg NE in a total volume of 10 μl containing 10 mM Tris-HCl (pH 8.0), 5 mM NaCl, 1 mM EDTA, 1 mM DTT, 0.1 mg/ml BSA and 0.1 μg/μl Poly[d(I-C)]⁵³. p50/p65 hetero-complexes were made by adding 2.75 ng of p50 (Cat # AG-40T-0021, Adipogen Corp) and 3.75 ng of p65 (Cat # AG-40T-0020, Adipogen Corp) purified protein, followed by incubation at 37 °C for 60 min³². After that, the mixture was incubated with probe with or without indicated purified OGG1 (Cat # ENZ-253, ProSpec). Binding was performed for 5–10 min at room temperature (RT). The reaction mixtures were applied to a 6% polyacrylamide gels in low-ionic-strength buffer (0.5xTBE) and electrophoresed for 2 h at 4 °C.

Dot blot analysis. Genomic DNA was isolated using a Qiagen Genra Puregene Kit or the phenol-chloroform-isoamyl alcohol method. RNase A digestion was included in the isolation procedure. Isolated genomic DNA (2 μg per sample) was denatured in 0.1 M NaOH for 5 min at 95 °C, and chilled rapidly on ice.

Samples were serially diluted twofold and spotted on a positively charged nitrocellulose membrane using a Bio-Dot Microfiltration Apparatus (Cat # 170-6545, Bio-Rad). The blotted membrane was washed in $2 \times$ SSC buffer, and UV cross-linked. The membrane was then blocked in Odyssey buffer (Li-Cor) diluted 1:1 in PBS for 1 hour at RT. Mouse anti-8-oxoG monoclonal antibody (Cat # MOG100 P, Japan Institute for the Control of Aging) in Odyssey:PBS was added for 1 hour at RT. The membrane was washed for 10 min three times in PBS, and then incubated with HRP-conjugated sheep anti-mouse immunoglobulin-G (IgG) (GE, 1:5,000) secondary antibodies in Odyssey:PBS for 1 h at room temperature. The membrane was then washed for 10 min three times in PBS and visualized by chemiluminescence with GE ECL Plus.

Flare qPCR. Oxidative modified base products in promoter regions were determined by OGG1 digestion-coupled qPCR (Flare-qPCR)²⁸. Briefly, cells were exposed to TNF α for 30 min or not, DNA was isolated and 2 μ g was mock- or OGG1 (5 ng) digested. After phenol/chloroform extraction 20 ng DNA was subjected to qPCR. Primers: *Tnf*: F: 5'-AACTCTCAAGCTGCTCTGCC-3' R: 5'-CAAGGAATCTCCTCCCCGTC-3' (-499 to -138); *Cxcl2*: F: 5'-GCTCAGTACACCGCAGGAAC-3', R: 5'-CTGCCCTTCCACTATGGGAC-3' (-531 to -186), *Il-1 β* : F: 5'-AAGGAAGTGCGTGTCTCTCC-3' R: 5'-TCAAGGGGTGGCAGATAGTG-3' (-567 to -251).

Quantitative RT-PCR (RT-qPCR). Total RNA was isolated from cultured cells using Qiagen RNeasy with on-column DNase I treatment. Complementary DNA (cDNA) was generated from 1 μ g of RNA using random hexamers to prime the reaction. The cDNA was used as template for RT-qPCR. RT-qPCR was performed in combination with the SYBR Green qPCR Master Mix (Cat# 638320, TaKaRa). Relative expression levels of target genes were calculated by $\Delta\Delta$ Ct method. The relative amount of each gene was normalized using β actin and Tubulin housekeeping genes.

Statistical tests. Results were tested for statistical significance using one-way ANOVA to analyse changes at mRNA levels. All data values are presented as mean \pm SD. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

References

- Thannickal, V. J. Oxygen in the evolution of complex life and the price we pay. *Am J Respir Cell Mol Biol* **40**, 507–510 (2009).
- Evans, M. D. & Cooke, M. S. Factors contributing to the outcome of oxidative damage to nucleic acids. *Bioessays* **26**, 533–542 (2004).
- Ming, X. *et al.* Mapping structurally defined guanine oxidation products along DNA duplexes: influence of local sequence context and endogenous cytosine methylation. *J Am Chem Soc* **136**, 4223–4235 (2014).
- Cadet, J., Douki, T. & Ravanat, J. L. Oxidatively generated base damage to cellular DNA. *Free Radic Biol Med* **49**, 9–21 (2010).
- Giese, B. Long-distance electron transfer through DNA. *Annu Rev Biochem* **71**, 51–70 (2002).
- Margolin, Y., Cloutier, J. F., Shafirovich, V., Geacintov, N. E. & Dedon, P. C. Paradoxical hotspots for guanine oxidation by a chemical mediator of inflammation. *Nat Chem Biol* **2**, 365–366 (2006).
- Wallace, S. S. DNA glycosylases search for and remove oxidized DNA bases. *Environ Mol Mutagen* **54**, 691–704 (2013).
- Barnes, D. E. & Lindahl, T. Repair and genetic consequences of endogenous DNA base damage in mammalian cells. *Annu Rev Genet* **38**, 445–476 (2004).
- Breen, A. P. & Murphy, J. A. Reactions of oxyl radicals with DNA. *Free Radic Biol Med* **18**, 1033–1077 (1995).
- David, S. S., O'Shea, V. L. & Kundu, S. Base-excision repair of oxidative DNA damage. *Nature* **447**, 941–950 (2007).
- Krokan, H. E. & Bjoras, M. Base excision repair. *Cold Spring Harb Perspect Biol* **5**, a012583 (2013).
- Huffman, J. L., Sundheim, O. & Tainer, J. A. DNA base damage recognition and removal: new twists and grooves. *Mutat Res* **577**, 55–76 (2005).
- Stivers, J. T. Site-specific DNA damage recognition by enzyme-induced base flipping. *Prog Nucleic Acid Res Mol Biol* **77**, 37–65 (2004).
- Stivers, J. T. & Jiang, Y. L. A mechanistic perspective on the chemistry of DNA repair glycosylases. *Chem Rev* **103**, 2729–2759 (2003).
- Odell, I. D., Wallace, S. S. & Pederson, D. S. Rules of engagement for base excision repair in chromatin. *J Cell Physiol* **228**, 258–266 (2012).
- Hazra, T. K. *et al.* Oxidative DNA damage repair in mammalian cells: a new perspective. *DNA Repair (Amst)* **6**, 470–480 (2007).
- Krishnamurthy, N., Zhao, X., Burrows, C. J. & David, S. S. Superior removal of hydantoin lesions relative to other oxidized bases by the human DNA glycosylase hNEIL1. *Biochemistry* **47**, 7137–7146 (2008).
- Nemec, A. A., Wallace, S. S. & Sweasy, J. B. Variant base excision repair proteins: contributors to genomic instability. *Seminars in cancer biology* **20**, 320–328 (2010).
- Zharkov, D. O. *et al.* Structural analysis of an Escherichia coli endonuclease VIII covalent reaction intermediate. *EMBO J* **21**, 789–800 (2002).
- Calkins, M. J. *et al.* Enhanced sensitivity of Neil1 $^{-/-}$ mice to chronic UVB exposure. *DNA Repair (Amst)* **48**, 43–50 (2016).
- Li, G. *et al.* 8-Oxoguanine-DNA glycosylase 1 deficiency modifies allergic airway inflammation by regulating STAT6 and IL-4 in cells and in mice. *Free Radic Biol Med* **52**, 392–401 (2012).
- Klungland, A. *et al.* Accumulation of premutagenic DNA lesions in mice defective in removal of oxidative base damage. *Proc Natl Acad Sci USA* **96**, 13300–13305 (1999).
- Minowa, O. *et al.* Mmh/Ogg1 gene inactivation results in accumulation of 8-hydroxyguanine in mice. *Proc Natl Acad Sci USA* **97**, 4156–4161 (2000).
- Bacsi, A. *et al.* Down-regulation of 8-oxoguanine DNA glycosylase 1 expression in the airway epithelium ameliorates allergic lung inflammation. *DNA Repair (Amst)* **12**, 18–26 (2013).
- Wang, V. Y. *et al.* The transcriptional specificity of NF-kappaB dimers is coded within the kappaB DNA response elements. *Cell Rep* **2**, 824–839 (2012).
- Ghosh, R. & Mitchell, D. L. Effect of oxidative DNA damage in promoter elements on transcription factor binding. *Nucleic Acids Res* **27**, 3213–3218 (1999).
- Hailer-Morrison, M. K., Kotler, J. M., Martin, B. D. & Sugden, K. D. Oxidized guanine lesions as modulators of gene transcription. Altered p50 binding affinity and repair shielding by 7,8-dihydro-8-oxo-2'-deoxyguanosine lesions in the NF-kappaB promoter element. *Biochemistry* **42**, 9761–9770 (2003).
- Ba, X. *et al.* 8-oxoguanine DNA glycosylase-1 augments proinflammatory gene expression by facilitating the recruitment of site-specific transcription factors. *J Immunol* **192**, 2384–2394 (2014).
- Wong, D. *et al.* Extensive characterization of NF-kappaB binding uncovers non-canonical motifs and advances the interpretation of genetic functional traits. *Genome Biol* **12**, R70 (2011).

30. Aguilera-Aguirre, L. *et al.* Whole transcriptome analysis reveals an 8-oxoguanine DNA glycosylase-1-driven DNA repair-dependent gene expression linked to essential biological processes. *Free Radic Biol Med* **81**, 107–118 (2015).
31. Aguilera-Aguirre, L. *et al.* Whole transcriptome analysis reveals a role for OGG1-initiated DNA repair signaling in airway remodeling. *Free Radic Biol Med* **89**, 20–33 (2015).
32. Fujita, T., Nolan, G. P., Ghosh, S. & Baltimore, D. Independent modes of transcriptional activation by the p50 and p65 subunits of NF-kappa B. *Genes Dev* **6**, 775–787 (1992).
33. Chen, Y. Q., Ghosh, S. & Ghosh, G. A novel DNA recognition mode by the NF-kappa B p65 homodimer. *Nat Struct Biol* **5**, 67–73 (1998).
34. Jamaluddin, M., Wang, S., Boldogh, I., Tian, B. & Brasier, A. R. TNF-alpha-induced NF-kappaB/RelA Ser(276) phosphorylation and enhanceosome formation is mediated by an ROS-dependent PKAc pathway. *Cell Signal* **19**, 1419–1433 (2007).
35. Vlahopoulos, S., Boldogh, I., Casola, A. & Brasier, A. R. Nuclear factor-kappaB-dependent induction of interleukin-8 gene expression by tumor necrosis factor alpha: evidence for an antioxidant sensitive activating pathway distinct from nuclear translocation. *Blood* **94**, 1878–1889 (1999).
36. Pastukh, V., Ruchko, M., Gorodnya, O., Wilson, G. L. & Gillespie, M. N. Sequence-specific oxidative base modifications in hypoxia-inducible genes. *Free Radic Biol Med* **43**, 1616–1626 (2007).
37. Doseth, B. *et al.* Uracil-DNA glycosylase in base excision repair and adaptive immunity: species differences between man and mouse. *J Biol Chem* **286**, 16669–16680 (2011).
38. Bubici, C., Papa, S., Dean, K. & Franzoso, G. Mutual cross-talk between reactive oxygen species and nuclear factor-kappa B: molecular basis and biological significance. *Oncogene* **25**, 6731–6748 (2006).
39. Baeuerle, P. A. & Henkel, T. Function and activation of NF-kappa B in the immune system. *Annu Rev Immunol* **12**, 141–179 (1994).
40. German, P. *et al.* 8-Oxoguanine DNA glycosylase1-driven DNA repair-A paradoxical role in lung aging. *Mech Ageing Dev* (2016).
41. Bacsi, A., Pan, L., Ba, X. & Boldogh, I. Pathophysiology of bronchoconstriction: role of oxidatively damaged DNA repair. *Curr Opin Allergy Clin Immunol* **16**, 59–67 (2016).
42. Brasier, A. R. The NF-kappaB regulatory network. *Cardiovasc Toxicol* **6**, 111–130 (2006).
43. Sen, C. K. & Packer, L. Antioxidant and redox regulation of gene transcription. *Faseb J* **10**, 709–720 (1996).
44. Cheng, X. *et al.* Dynamic behavior of DNA base pairs containing 8-oxoguanine. *J Am Chem Soc* **127**, 13906–13918 (2005).
45. Chen, L., Haushalter, K. A., Lieber, C. M. & Verdine, G. L. Direct visualization of a DNA glycosylase searching for damage. *Chem Biol* **9**, 345–350 (2002).
46. Bruner, S. D., Norman, D. P. & Verdine, G. L. Structural basis for recognition and repair of the endogenous mutagen 8-oxoguanine in DNA. *Nature* **403**, 859–866 (2000).
47. Banerjee, A., Yang, W., Karplus, M. & Verdine, G. L. Structure of a repair enzyme interrogating undamaged DNA elucidates recognition of damaged DNA. *Nature* **434**, 612–618 (2005).
48. Roberts, R. J. On base flipping. *Cell* **82**, 9–12 (1995).
49. Schreck, R., Zorbas, H., Winnacker, E. L. & Baeuerle, P. A. The NF-kappa B transcription factor induces DNA bending which is modulated by its 65-kD subunit. *Nucleic Acids Res* **18**, 6497–6502 (1990).
50. Misteli, T. Protein dynamics: implications for nuclear architecture and gene expression. *Science* **291**, 843–847 (2001).
51. Carey, M. The enhanceosome and transcriptional synergy. *Cell* **92**, 5–8 (1998).
52. Schreck, R., Rieber, P. & Baeuerle, P. A. Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1. *Embo J* **10**, 2247–2258 (1991).
53. Lenardo, M., Pierce, J. W. & Baltimore, D. Protein-binding sites in Ig gene enhancers determine transcriptional activity and inducibility. *Science* **236**, 1573–1577 (1987).

Acknowledgements

This work was supported by grants from the National Nature Science Foundation of China (grant number: 31571339 to X.B), the Programme for Introducing Talent to Universities (Grant no. B07017 to X.B), and United States National Institute of Environmental Health and Sciences (grant number: RO1 ES018948 to IB); United States National Institute of Allergic and Infectious Diseases (grant number: PO1-AI062885, to IB); United States National Institute of Environmental Health and Sciences Centre Grant (grant number: P30 ES006676 to IB).

Author Contributions

L.P., I.B. and X.B. conceived and designed the experiments, L.P., W.H., X.Z. and A.A.A. performed the experiments, L.P., I.B. and X.B. analysed the data, L.P. wrote the manuscript. Manuscript is reviewed, discussed and approved by all the authors.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Pan, L. *et al.* OGG1-DNA interactions facilitate NF- κ B binding to DNA targets. *Sci. Rep.* **7**, 43297; doi: 10.1038/srep43297 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017