

SCIENTIFIC REPORTS

OPEN

Corrigendum: Epileptiform activity in the mouse visual cortex interferes with cortical processing in connected areas

L. Petrucco, E. Pracucci, M. Brondi, G. M. Ratto & S. Landi

Scientific Reports 7:40054; doi: 10.1038/srep40054; published online 10 January 2017; updated on 14 March 2017

This Article contains typographical errors in the Results section under subheading ‘ISs fragment the up states in the opposite hemisphere’.

“The presence of ISs in Hem 2 was associated with a change of the statistics of the slow-wave oscillations in Hem 1; in particular, as shown in Fig. 3c,d, the frequency of up states increased from 0.70 ± 0.06 Hz to 0.81 ± 0.06 Hz (paired t-test, $n = 8$, $p = 0.007$), while their duration decreased from 0.74 ± 0.04 s to 0.64 ± 0.03 s (paired t-test, $n = 8$, $p = 0.003$)”.

should read:

“The presence of ISs in Hem 2 was associated with a change of the statistics of the slow-wave oscillations in Hem 1; in particular, as shown in Fig. 3c,d, the frequency of up states increased from 0.57 ± 0.03 Hz to 0.64 ± 0.02 Hz (paired t-test, $n = 8$, $p = 0.004$), while their duration decreased from 0.74 ± 0.04 s to 0.64 ± 0.03 s (paired t-test, $n = 8$, $p = 0.003$)”.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017