

# SCIENTIFIC REPORTS

OPEN

## Vlasouliolides A-D, four rare C<sub>17</sub>/C<sub>15</sub> sesquiterpene lactone dimers with potential anti-inflammatory activity from *Vladimiria souliei*

Received: 28 September 2016

Accepted: 31 January 2017

Published: 03 March 2017

Li-Ping Chen<sup>1,\*</sup>, Guo-Zhen Wu<sup>1,\*</sup>, Jian-Ping Zhang<sup>1</sup>, Ji Ye<sup>1</sup>, Qing-Xin Liu<sup>1</sup>, Yun-Heng Shen<sup>1</sup>, Hui-Liang Li<sup>1</sup> & Wei-Dong Zhang<sup>1,2</sup>

Vlasouliolides A-D (1–4), four rare sesquiterpene lactone dimers, were isolated from *Vladimiria souliei*. The common structural characteristic of 1–4 is the C<sub>32</sub> skeleton comprising two sesquiterpene lactone units linked by a C11–C13' single bond with one acetyl connected to the C-13 position of one of the two sesquiterpene lactone units. The stereochemistries of 1–4 were assigned by a combination of NOESY correlations and Cu–K $\alpha$  X-ray crystallographic analyses. Compounds 1–4 strongly inhibited the production of NO in LPS-stimulated RAW 264.7 cells. Furthermore, 1 and 2 inhibited the activation of NF- $\kappa$ B in LPS-induced 293T cells.

Naturally occurring sesquiterpene lactone dimers (SLDs) are a type of complex found in natural products with many pharmacological activities<sup>1,2</sup>. Since the first SLD absinthin was isolated from *Artemisia absinthium* in 1953<sup>3</sup>, more than 160 SLDs have been obtained. To the best of our knowledge, most of the published SLDs have C<sub>30</sub> cores derived from two C<sub>15</sub> sesquiterpenoid units<sup>4–6</sup>.


The genus *Vladimiria*, belonging to the family of Asteraceae, comprises approximately 12 species that are mainly distributed in the Sichuan Province, China<sup>7</sup>. Sesquiterpenes, as major constituents isolated from the *Vladimiria* species, possess various types of structures including guaianolide, carabranane, eudesmane and germacranane sesquiterpenes<sup>8–12</sup>. The plant of *Vladimiria souliei*, as a traditional Chinese medicine, has been used for relieving pain and stomach diseases since ancient times<sup>7,13</sup>. Additionally, the chemical constituents from the roots of *V. souliei* exhibited significant antimicrobial, antitumor and inhibitory effects on NO production activities<sup>10,13</sup>. In the course of our investigation on structurally novel SLDs from the family of Asteraceae, four rare SLDs with C<sub>32</sub> cores were discovered from *Vladimiria souliei*, and they were designated vlasouliolides A-D (1–4) (Fig. 1).

### Results and Discussion


**Structure elucidation.** Vlasouliolide A (1), [ $\alpha$ ]<sub>D</sub><sup>25</sup> +3.77 (*c* 0.24, CH<sub>3</sub>COCH<sub>3</sub>), was obtained as a colorless orthorhombic crystal. Its molecular formula was determined to be C<sub>32</sub>H<sub>40</sub>O<sub>5</sub> by positive HRESIMS at *m/z* 527.2763 ([M + Na]<sup>+</sup>, calcd. 527.2773). The <sup>1</sup>H NMR spectrum of 1 (Supplementary Table S1) showed characteristic signals for one methyl singlet at  $\delta_{\text{H}}$  2.16 (3H, s, -Ac), two oxymethines at  $\delta_{\text{H}}$  3.91 (1H, t, *J* = 9.5 Hz, H-6') and  $\delta_{\text{H}}$  4.17 (1H, t, *J* = 9.6 Hz, H-6), and four sets of terminal alkylene ( $\delta_{\text{H}}$  4.83, 1H, s, 4.76, 1H, s, H<sub>2</sub>-14;  $\delta_{\text{H}}$  4.88, 1H, s, 4.78, 1H, s, H<sub>2</sub>-14';  $\delta_{\text{H}}$  5.29, 1H, d, *J* = 1.8 Hz, 5.06, 1H, d, *J* = 1.7 Hz, H<sub>2</sub>-15;  $\delta_{\text{H}}$  5.14, 1H, d, *J* = 1.9 Hz, 5.03, 1H, d, *J* = 1.7 Hz, H<sub>2</sub>-15').

Analysis of the <sup>13</sup>C NMR spectrum revealed the existence of 32 carbons (Supplementary Table S2), including 1  $\times$  CH<sub>3</sub>, 14  $\times$  CH<sub>2</sub> (containing 4 sp<sup>2</sup> alkylenes at  $\delta_{\text{C}}$  109.3, 109.6, 111.6, and 112.0), 9  $\times$  CH (containing 2 oxymethines at  $\delta_{\text{C}}$  82.4 and 84.5), and 8  $\times$  C (containing 3 carbonyl carbons at  $\delta_{\text{C}}$  177.5, 178.3 and 205.2). Detailed analysis of the NMR spectrum of 1 indicated that vlasouliolide A (1) should contain an acetyl due to the resonance signals at  $\delta_{\text{C}}$  205.2, 30.7, and  $\delta_{\text{H}}$  2.16 (3H, s, -Ac). Deducting the carbon resonances for the acetyl, the remaining 30 carbon resonances implied that two similar sesquiterpene lactone units existed.

<sup>1</sup>Department of Phytochemistry, School of Pharmacy, Second Military Medical University, Shanghai 200433, P. R. China. <sup>2</sup>Shanghai Institute of Pharmaceutical Industry, Shanghai 200040, P. R. China. \*These authors contributed equally to this work. Correspondence and requests for materials should be addressed to H.-L.L. (email: faranli@hotmail.com) or W.-D.Z. (email: wdzhangy@hotmail.com)




**Figure 1.** Chemical structures of 1–4.



**Figure 2.** (A) Selected NMR correlations and (B) X-ray crystallographic structure of 1.

The planar structure of vlasouliolide A was constructed by comprehensive analyses of the 2D NMR spectra (Fig. 2A). Two similar long proton-bearing structural fragments, H<sub>2</sub>-3/H<sub>2</sub>-2/H-1/H-5/H-6/H-7/H<sub>2</sub>-8/H<sub>2</sub>-9 and H<sub>2</sub>-3'/H<sub>2</sub>-2'/H-1'/H-5'/H-6'/H-7'/H<sub>2</sub>-8'/H<sub>2</sub>-9', as well as an extra short chain, H-7'/H-11'/H<sub>2</sub>-13', were constructed based on interpretation of the <sup>1</sup>H-<sup>1</sup>H COSY and HSQC-TOCSY spectra. The HMBC spectrum of **1** showed the key correlations of four typical terminal alkylene groups (Fig. 2A), including H<sub>2</sub>-14/C-1 and C-9; H<sub>2</sub>-15/C-3 and C-5; H<sub>2</sub>-14'/C-1' and C-9'; and H<sub>2</sub>-15'/C-3' and C-5'. Thus, subunits A and B were deduced to be two guaianolide moieties similar to dehydrocostus lactone<sup>14,15</sup>. Therefore, the structure of **1** should be composed of two guaianolide moieties and an acetyl moiety. In the HMBC spectrum of **1**, the long correlations from H<sub>3</sub>-17 (δ<sub>H</sub> 2.16, 3H, s) to C-13 (δ<sub>C</sub> 46.4) as well as H<sub>2</sub>-13 (δ<sub>H</sub> 2.86, 2.56, 1H, d, J = 19 Hz, respectively) to C-7 (δ<sub>C</sub> 48.2) and C-12 (δ<sub>C</sub> 178.3) indicated that the acetyl moiety was connected to the C-13 position of one guaianolide unit. Compared with dehydrocostus lactone, the C-11 position of subunit A was a quaternary carbon, while the C-13' position of subunit B was an sp<sup>3</sup> methylene. These changes implied that the two sesquiterpene lactone units should be linked directly *via* a C-C bond between C-11 and C-13', and this assumption was verified by the HMBC correlations of H<sub>2</sub>-13' (δ<sub>H</sub> 2.23, 1.35, 1H, m, respectively) with C-11 (δ<sub>C</sub> 46.1), C-12 (δ<sub>C</sub> 178.3), C-11' (δ<sub>C</sub> 41.9) and C-12' (δ<sub>C</sub> 177.5). Thus, the planar structure of **1** was elucidated as shown in Fig. 1.

The relative configuration of **1** was characterized by interpretation of the NOESY spectrum (Fig. 2A). The similar NOESY correlations of H-1/H-7/H-5 and H-1'/H-7'/H-5' in subunits A and B indicated that they are on the same face. In addition, the large coupling constant between H-6/H-7 and H-6'/H-7' (J = 9.6, 9.5 Hz, respectively) implied that H-6/H-7 and H-6'/H-7' were in the *trans*-form. Consequently, in subunit A, H-7 was arbitrarily assigned as having an α-orientation. The correlation between H-7 and H<sub>2</sub>-13 suggested that the side chain C<sub>13</sub>/C<sub>16</sub>/C<sub>17</sub> should be located below the molecular plane. Thus, the H<sub>2</sub>-13' was assigned as having a β-orientation, as



**Figure 3.** (A) The deshielding effect of the carbonyl group at C-12' and (B) X-ray crystallographic structure of **2**.


indicated by the NOESY correlations of H-6 with H<sub>2</sub>-13'. It should be noted that H<sub>2</sub>-13' exhibited NOESY correlations with H-7'. Combined with the NOESY correlation between H<sub>2</sub>-13 and H-11', the relative configuration of H-11' should be the  $\alpha$ -orientation. Finally, the absolute configuration of **1** was confirmed to be 1*R*, 5*R*, 6*S*, 7*S*, 11*S*, 1'*R*, 5'*R*, 6'*S*, 7'*S*, 11'*S* by Cu-K $\alpha$  X-ray crystallographic analysis (Fig. 2B).

Vlasouliolide B (**2**), [ $\alpha$ ]<sub>D</sub><sup>25</sup> +20.41 (*c* 0.21, CH<sub>3</sub>COCH<sub>3</sub>), possessed the same molecular formula as that of **1** as determined by positive HRESIMS at *m/z* 527.2758 ([*M* + Na]<sup>+</sup>, calcd. 527.2773). Analysis of the NMR data (Supplementary Tables S1 and S2) indicated that compound **2** possesses the same planar structure as that of **1**. The notable NOESY correlation of H<sub>2</sub>-13/H-11'/H-7' in **2** instead of that H-11'/H-6' in **1** implied that **2** was an 11'-epimer of **1**. Comparing the NMR spectroscopic data of **2** with those of **1**, an obvious downfield shift of Ha-13 from  $\delta$ <sub>H</sub> 2.86 in **1** to  $\delta$ <sub>H</sub> 3.36 in the <sup>1</sup>H NMR was observed. As shown in Fig. 3A, the opposite arrangement of the two subunits led to the C-12' carbonyl and H<sub>2</sub>-13 in **1** being far from each other. In contrast, the subunits A and B were arranged in the same direction in **2**, resulting in the H<sub>2</sub>-13 being spatially adjacent to the carbon-oxygen double bond at C-12'. Thus, the intramolecular deshielding effect contributed to the downfield shift of Ha-13 from  $\delta$ <sub>H</sub> 2.86 in **1** to  $\delta$ <sub>H</sub> 3.36 in **2**. Finally, the absolute configuration of C-11' was confirmed as *R* by Cu-K $\alpha$  X-ray diffraction (Fig. 3B).

Vlasouliolide C (**3**) was isolated as a colorless monoclinic crystal with [ $\alpha$ ]<sub>D</sub><sup>25</sup> +30.23 (*c* 0.04, CHCl<sub>3</sub>). Its molecular formula was determined to be C<sub>32</sub>H<sub>42</sub>O<sub>5</sub> by positive HRESIMS at *m/z* 529.2940 ([*M* + Na]<sup>+</sup>, calcd. 529.6728). The 1D NMR data (Supplementary Tables S1 and S2) of **3** disclosed a C<sub>32</sub> skeleton similar to in **1** and **2**. Comprehensive analysis of the 2D NMR spectra indicated that compound **3** possessed the same 13-acetyl-mokkolactone fragment (subunit A) as **1** and **2**. Unlike the structures of **1** and **2**, an eudesmane moiety existed as subunit B in the structure of **3**, as deduced by the <sup>1</sup>H-<sup>1</sup>H COSY correlations of H<sub>2</sub>-3'/H<sub>2</sub>-2'/H-1', H-5'/H-6'/H-7'/H<sub>2</sub>-8'/H<sub>2</sub>-9' and H-7'/H-11'/H<sub>2</sub>-13' together with crucial HMBC correlations from H<sub>2</sub>-15' to C-3' and C-5' as well as from CH<sub>3</sub>-14 to C-1, C-2, and C-9. Therefore, the structure of **3** was formed by an acetyl-substituted guaianolide moiety and eudesmane moiety. The HMBC correlations from H<sub>2</sub>-13' ( $\delta$ <sub>H</sub> 2.30, 1.38) to C-11' ( $\delta$ <sub>C</sub> 41.6), C-12' ( $\delta$ <sub>C</sub> 178.8), C-11 ( $\delta$ <sub>C</sub> 46.2), C-12 ( $\delta$ <sub>C</sub> 178.3) and C-13 ( $\delta$ <sub>C</sub> 47.2) suggested that subunits A and B should also be directly linked by a C-11/13' single bond (Fig. 4A).

The relative configuration of subunit B was deduced by analysis of the NOESY correlations (Fig. 4A). H-11', H-6' and H-14' were assigned as having the  $\alpha$ -orientation, and H-5', H-7' and H-13' were assigned as having the  $\beta$ -orientation, which were consistent with the biosynthetic precursor  $\beta$ -cyclocostunolide<sup>16</sup>. Similar to **1**, the opposite arrangement of the two subunits was verified by the NOESY correlations of H-13'/H-6 and H-11'/H-13. The structure of **3** was finally elucidated as shown in Fig. 1, and the absolute configuration was assigned as 1*R*, 5*R*, 6*S*, 7*S*, 11*S*, 5'*S*, 6'*S*, 7'*S*, 10'*S*, 11'*S* by Cu-K $\alpha$  X-ray crystallographic analysis (Fig. 4B).

Vlasouliolide D (**4**), a colorless orthorhombic crystal with [ $\alpha$ ]<sub>D</sub><sup>25</sup> +55.21 (*c* 0.08, CH<sub>3</sub>COCH<sub>3</sub>), possessed the same molecular formula as that of **3** determined by positive HRESIMS at *m/z* 529.2923 ([*M* + Na]<sup>+</sup> calcd. 529.6722). The 1D NMR data of **4** (Supplementary Tables S1 and S2) revealed that **4** was constructed from one acetyl, one dehydrocostus lactone, and one  $\beta$ -cyclocostunolide moieties, which were identical with those of **3**. However, different from **3**, the acetyl in **4** was located at the C-13 position of  $\beta$ -cyclocostunolide to form 13-acetyl-eudesmenolide (subunit A), which is supported by the key HMBC correlations from H<sub>2</sub>-13 ( $\delta$ <sub>H</sub> 2.79, 2.66, 1H, d, *J* = 17 Hz, respectively) to C-7 ( $\delta$ <sub>C</sub> 51.9), C-11 ( $\delta$ <sub>C</sub> 46.2) and C-12 ( $\delta$ <sub>C</sub> 178.9). Subunit B was also connected to subunit A *via* a C-11/13' single bond. Therefore, the structure of **4** was determined as shown in Fig. 1. The absolute configuration of **4** was determined as 5*S*, 6*S*, 7*S*, 10*R*, 11*S*, 1'*R*, 5'*R*, 6'*S*, 7'*S*, 11'*S* by Cu-K $\alpha$  X-ray crystallographic analysis (Fig. 5).



**Figure 4.** (A) Selected NMR correlations and (B) X-ray crystallographic structure of **3**.


**Biological activity assay.** The roots of *V. souliei* are often used in traditional Chinese medicine for the treatment of digestive disorders and inflammatory diseases. Many sesquiterpenes isolated from *V. souliei* showed inhibitory activity against lipopolysaccharide (LPS)-induced nitric oxide (NO) production in murine RAW264.7 cells<sup>10</sup>. Excessive NO has been implicated in the pathological process of tissue damage following inflammation<sup>17</sup>. The inhibition of the overproduction of NO is an important therapeutic way to treat inflammatory diseases<sup>18</sup>. We investigated the anti-inflammatory activities of compounds **1–4** by a LPS-induced NO production assay in RAW 264.7 macrophages<sup>5</sup>. Compounds **1–4** exhibited significant inhibitory effects against NO production with IC<sub>50</sub> values of 1.14, 2.53, 1.57 and 3.19  $\mu$ M, respectively. Moreover, these compounds were not notably cytotoxic at the concentrations required for inhibiting NO production, as determined by an MTT assay. NF- $\kappa$ B is a transcription factor that controls immune responses and plays a pivotal role in the regulation of NO expression<sup>19</sup>. We conducted an NF- $\kappa$ B luciferase reporter assay in 293T cells to evaluate the impact of compounds **1–4** on the transcriptional activity of NF- $\kappa$ B<sup>20</sup>.

The NF- $\kappa$ B reporter luciferase construct and Renilla luciferase control vector were cotransfected in 293T cells for 24 h. Thereafter, the cells were left untreated or were treated with a 10  $\mu$ M concentration of compound for an additional 1 h before LPS activation for 4 h. Compounds **1** and **2** displayed inhibition towards the NF- $\kappa$ B pathway, while **3** and **4** showed no effects in the reduction of NF- $\kappa$ B luciferase activity (Fig. 6A). We also examined the effects of compounds **1–4** on the expression of I $\kappa$ B $\alpha$  and P65 proteins in the NF- $\kappa$ B pathway<sup>21</sup>. RAW 264.7 cells pretreated with **1–4** at the indicated concentrations for 1 h were subjected to LPS stimulation before Western blot analysis. Compounds **1–4** had no inhibitory effects on the degradation of I $\kappa$ B $\alpha$ , while **1** and **2** can dose-dependently down-regulate the LPS-induced phosphorylation of the NF- $\kappa$ B p65 subunit (Fig. 6B). These data indicated that the NO inhibitory activities of **1** and **2** might be attributed to suppressing NF- $\kappa$ B activation, while those of **3** and **4** might not be due to this mechanism.

In conclusion, sesquiterpene lactone dimers usually have a C<sub>30</sub> framework biosynthetically derived from a Diels-Alder adduct of two homo or hetero sesquiterpene monomers. Vlasouliolides A–D (**1–4**) possessed a C<sub>32</sub> skeleton derived from two sesquiterpene lactone molecules and an acetyl group. Commonly, in the chemical structures of **1–4**, the acetyl was connected to the C-13 of one of the two sesquiterpene lactone units to form a C<sub>17</sub> unit, and the C<sub>17</sub>/C<sub>15</sub> units were further directly linked by a C11–C13' single bond. This is the first report regarding C<sub>17</sub>/C<sub>15</sub> sesquiterpene lactone dimers from nature with significant anti-inflammation activities, which might be potentially useful for the treatment of inflammatory diseases. The discovery of Vlasouliolides A–D may encourage further investigations by natural product chemists, synthetic chemists, and pharmacists.

## Methods

**General experimental procedures.** Column chromatography (CC): silica gel H (10–40  $\mu$ m) and silica gel (200–300 mesh) (Marine Chemical Factory, Qingdao, P. R. China); Sephadex LH-20 (Pharmacia Fine Chemicals, Piscataway, NJ, USA); RP-C18 gel (40–63  $\mu$ m; Daiso, Co., Japan). TLC: silica gel plates (Yantai Jiang You Silicone Development Co., Yantai, P. R. China), visualization by spraying with 10% H<sub>2</sub>SO<sub>4</sub> in EtOH. HPLC: Agilent 1260 series (Agilent Technologies, US) with a Zorbax SB-C18 (5  $\mu$ m, 9.4  $\times$  150 mm) column. NMR: Bruker Avance III-500 and Avance III-600 spectrometers (Bruker, Switzerland). MS: Agilent MSD-Trap-XCT (for ESI) and Agilent-6520 Q-TOF mass spectrometers (for HR-ESI). Melting point: X-4B digital display micro-melting apparatus (Shanghai Jingsong Instrument, Shanghai, P. R. China). Optical rotation: Rudolph Autopo V (Rudolph Research Analytical, Hackettstown, NJ). UV: Agilent 1260 series DAD detector (Agilent Technologies, US). CD: Brighttime Chirascan (Applied Photophysics Ltd, UK). IR: Thermo Scientific Nicolet 6700 (Thermo Scientific, USA). RAW 264.7 cells and 293T cells: ATCC (American type culture collection). Dulbecco's modified Eagle's medium and fetal bovine serum: Gibco Invitrogen (Carlsbad, CA, USA). LPS Griess reagent and MTT: Sigma-Aldrich (St. Louis, MO, USA).




**Figure 5.** X-ray crystallographic structure of **4**.

**Plant material.** The roots of *V. souliei* were collected from the Sichuan province of China in October 2014 and authenticated by professor Bao-Kang Huang, Department of Pharmacognosy, School of Pharmacy, Second Military Medical University. A voucher specimen (No. 201412-VS) is deposited in the Department of Pharmacognosy, Second Military Medical University.

**Extraction and isolation.** The chipped and dried roots of *V. souliei* (20.0 kg) were extracted by maceration with 95% ethanol overnight at room temperature ( $3 \times 60$  L). After removal of the solvent, the ethanol extract (2.12 kg) was successively partitioned between water and petroleum ether (PE)/ethyl acetate (EtOAc) to give PE, EtOAc and water extracts. The EtOAc extract (0.626 kg) was segmented by silica gel column chromatography (PE/EtOAc, 50:1–0:1) to yield 17 fractions (Fr. 1–17). Fraction 7 (31.99 g) underwent chromatography over an RP-C18 medium-pressure column (MeOH/H<sub>2</sub>O, 10:90 to 90:10) to give 12 subfractions (Fr. 7.1–7.12). Subfraction 7.9 (2.5 g) underwent further chromatography over an RP-C18 medium-pressure column (CH<sub>3</sub>CN/H<sub>2</sub>O, 45:55) and was finally purified by semi-preparative RP-C18 HPLC (MeOH/H<sub>2</sub>O, 71:29) to produce **1** (42.9 mg) and **4** (5.2 mg). Subfraction 7.10 (1.89 g) underwent chromatography over an RP-C18 medium-pressure column using MeOH/H<sub>2</sub>O (55:45–100:0) as the elution solvent, and the 70–80% fraction was purified by semi-preparative RP-C18 HPLC (CH<sub>3</sub>CN/H<sub>2</sub>O, 45:55), yielding **2** (22.2 mg). Fraction 8 (5.40 g) underwent chromatography over





**Figure 6. Effects of 1–4 on LPS-stimulated NF-κB activation.** (A) The NF-κB luciferase reporter assay shows that compounds 1 and 2 displayed suppression of LPS-induced NF-κB activation in 293T cells. (Mean ± SD in three separate experiments. \* $p < 0.05$ ; \*\* $p < 0.001$ ). (B) The effects of compounds 1–4 on the phosphorylation of IκBα and p65 in the presence of LPS stimulation in RAW 264.7 cells.

an RP-C18 medium-pressure column using MeOH/H<sub>2</sub>O in a gradient (10:90–100:0) to yield eight subfractions (Fr. 8.1–8.8). Subfraction 8.5 (2.35 g) was purified by an RP-C18 medium-pressure column (MeOH/H<sub>2</sub>O, 55:45 to 100:0) followed by semi-preparative RP-C18 HPLC (CH<sub>3</sub>CN/H<sub>2</sub>O, 63:27) to produce 3 (10.6 mg). Overall, we obtained compounds 1 (42.9 mg), 2 (22.2 mg), 3 (10.6 mg) and 4 (5.2 mg).

**Spectroscopic data.** **Vlasouliolide A (1)** Colorless orthorhombic crystals in EtOH/H<sub>2</sub>O. m.p.: 169–173 °C;  $[\alpha]_D^{25} +3.77$  ( $c$  0.24, CH<sub>3</sub>COCH<sub>3</sub>); UV (CH<sub>3</sub>OH/H<sub>2</sub>O)  $\lambda_{max}$  210 nm; IR (KBr)  $\nu_{max}$  3089, 2931, 2854, 1770, 1714, 1639, 1446, 1400, 1367, 1334, 1209, 1178, 1126, 995, 890 cm<sup>-1</sup>; <sup>1</sup>H- and <sup>13</sup>C-NMR data, see Tables S1 and S2; ESIMS  $m/z$  527.4 ([M + Na]<sup>+</sup>), 503.2 ([M – H]<sup>-</sup>); positive HRESIMS  $m/z$  527.2763 ([M + Na]<sup>+</sup>, calcd. 527.2773).

**Vlasouliolide B (2)** Colorless orthorhombic crystals in EtOH/H<sub>2</sub>O. m.p.: 171–173 °C;  $[\alpha]_D^{25} +20.41$  ( $c$  0.21, CH<sub>3</sub>COCH<sub>3</sub>); UV (CH<sub>3</sub>CN/H<sub>2</sub>O)  $\lambda_{max}$  210, 254, 290 nm; IR (KBr)  $\nu_{max}$  3081, 2931, 2854, 1760, 1712, 1641, 1455, 1365, 1313, 1213, 1164, 995, 885 cm<sup>-1</sup>; <sup>1</sup>H- and <sup>13</sup>C-NMR data, see Tables S1 and S2; ESIMS  $m/z$  527.3 ([M + Na]<sup>+</sup>), 539.3 ([M + Cl]<sup>-</sup>); positive HRESIMS  $m/z$  527.2758 ([M + Na]<sup>+</sup>, calcd. 527.2773).

**Vlasouliolide C (3)** Colorless monoclinic crystals in CHCl<sub>3</sub>/MeOH. m.p.: 211–218 °C;  $[\alpha]_D^{25} +30.23$  ( $c$  0.04 CHCl<sub>3</sub>); UV (CH<sub>3</sub>CN/H<sub>2</sub>O)  $\lambda_{max}$  210 nm; IR (KBr)  $\nu_{max}$  3086, 2929, 2849, 1763, 1719, 1649, 1444, 1380, 1336, 1260, 1195, 1159, 1103, 1019, 991, 911, 891 cm<sup>-1</sup>; <sup>1</sup>H- and <sup>13</sup>C-NMR data, see Tables S1 and S2; ESIMS  $m/z$  529.4 ([M + Na]<sup>+</sup>), 541.5 ([M + Cl]<sup>-</sup>); positive HRESIMS  $m/z$  529.2940 ([M + Na]<sup>+</sup>, calcd. 529.2930).

**Vlasouliolide D (4)** Colorless orthorhombic crystals in EtOH/H<sub>2</sub>O. m.p.: 172–182 °C;  $[\alpha]_D^{25} +55.21$  ( $c$  0.08 CH<sub>3</sub>COCH<sub>3</sub>); UV (CH<sub>3</sub>CN/H<sub>2</sub>O)  $\lambda_{max}$  210 nm; IR (KBr)  $\nu_{max}$  2933, 1772, 1712, 1648, 1457, 1367, 1205, 1176, 1130, 995, 892 cm<sup>-1</sup>; <sup>1</sup>H- and <sup>13</sup>C-NMR data, see Tables S1 and S2; ESIMS  $m/z$  529.4 ([M + Na]<sup>+</sup>), 505.3 ([M – H]<sup>-</sup>); positive HRESIMS  $m/z$  529.2923 ([M + Na]<sup>+</sup>, calcd. 529.2930).

**Measurement of LPS-Induced NO Production.** RAW 264.7 cells were seeded in 96-well culture plates at  $5 \times 10^5$  cells/well at 37 °C for 6 h in DMEM medium. The cells were pretreated with different concentrations of samples for 12 h and then incubated for 16 h with or without 1 μg/mL LPS. The nitrite concentration in the culture supernatant was measured using Griess reagent (1% sulfanilamide, 0.1% N-1-naphthylethylenediamine dihydrochloride and 2.5% phosphoric acid). The absorbance was measured at 540 nm using a microplate reader after incubation for 15 min. The nitrite levels in the samples were calculated from a standard curve created using known concentrations of sodium nitrite.

**MTT assay.** RAW 264.7 cells were seeded in 96-well plates at  $5 \times 10^5$  cells/well for 6 h and treated with different concentrations of compounds for 24 h. Thereafter, MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] was added to each well at a final concentration of 0.5 mg/ml and incubated at 37 °C for 3 h. The amount of MTT formazan was determined by dissolving it in dimethyl sulfoxide (DMSO) and measuring its absorbance at 495 nm using a microplate reader.

**Dual-Luciferase reporter gene assay.** 293T cells were seeded in 24-well plates at  $2 \times 10^6$  cells/well for 6 h. After being cotransfected with expression plasmids for NF-κB firefly luciferase and TK-Renilla luciferase for 24 h, the cells were treated with 10 μM concentrations of compounds 1–4 for an additional 1 h. Thereafter, the cells were stimulated with 1 μg/ml LPS for another 6 h and then lysed in lysis buffer. Luciferase activities were measured by the dual-luciferase reporter gene assay system (Promega). NF-κB firefly luciferase activity was normalized to the Renilla luciferase activity.

**Antibodies and Western blot analysis.** Antibodies for p-p65, p65, p-IκBα, IκBα and Gapdh were purchased from Cell Signaling Technology. For western blot analysis, RAW 264.7 cells were lysed in 2× SDS sample buffer (62.5 mM Tris-HCl, pH = 6.8, 2% SDS, 10% glycerol, 50 mM DTT, and 0.01% bromophenol blue) after

treatment with the compounds and stimulation by LPS. Cell lysates were fractionated by sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–PAGE) after denaturation treatment and subjected to immunoblot analysis with GAPDH as the loading control.

## References

1. Wang, G. W. *et al.* *Inula* sesquiterpenoids: structural diversity, cytotoxicity and anti-tumor activity. *Exper Opin Inv Drug*. **23**, 317–45 (2014).
2. Zhang, J. P. *et al.* The genus *Carpesium*: a review of its ethnopharmacology, phytochemistry and pharmacology. *J Ethnopharmacol*. **163**, 173–91 (2015).
3. Zhan, Z. J., Ying, Y. M., Ma, L. F. & Shan, W. G. Natural disesquiterpenoids. *Nat. Prod. Rep.* **28**, 594–629 (2011).
4. Liao, S. G. & Yue, J. M. *Dimeric Sesquiterpenoids* (eds Kinghorn, A. D., Falk, H., Gibbons, S., Kobayashi) 1–112 (Switzerland, 2016).
5. Qin, J. J. *et al.* Neojaponicone A, a bioactive sesquiterpene lactone dimer with an unprecedented carbon skeleton from *Inula japonica*. *Chem. Commun.* **47**, 1222–1224 (2011).
6. Yang, Y. X. *et al.* Carpedilactones A–D, four new isomeric sesquiterpene lactone dimers with potent cytotoxicity from *Carpesium faberi*. *Org. Lett.* **16**, 4216–4219 (2014).
7. Editorial commission of Traditional Chinese Medicine, State Administration Of Traditional Chinese Medicine (Ed.). *Traditional Chinese Medicine*. *Shanghai Science & Technology Press*, Vol. 7, p. 815–817 (Shanghai, 1999).
8. Tan, R. X., Jakupovic, J., Bohlmann, F., Jia, Z. J. & Schuster, A. Sesquiterpene lactones from *Vladimiria souliei*. *Phytochemistry*. **29**, 1209–1212 (1990).
9. Xu, J. *et al.* Two carabrane-type sesquiterpenes from *Vladimiria souliei*. *Phytochem Lett.* **2**, 204–206 (2009).
10. Xu, J. *et al.* Sesquiterpenes from *Vladimiria souliei* and their inhibitory effects on NO production. *Fitoterapia*. **82**, 508–511 (2011).
11. Chen, J. J. *et al.* Sesquiterpenoids from the roots of *Vladimiria muliensis*. *J. Asian Nat. Prod. Res.* **17**, 1–8 (2015).
12. Wei, H., Ma, G., Peng, Y., He, C. & Xiao, P. Chemical Constituents of the Roots of *Dolomiaea souliei*. *Chem. Nat. Compd.* **37**, 1249–53 (2014).
13. Zhou, L. Z., Jiang, J. H., Li, Y. P. & Chen, Y. G. Chemical and Bioactive Studies on Tibetan Medicines Plants of *Vladimiria Genus*. *Yunnan Chemical Technology* (2010).
14. Rigby, J. H. & Wilson, J. A. Z. Total synthesis of guaianolides: (±)-dehydrocostus lactone and (±)-estafiatin. *J. Am. Chem. Soc.* **106**, 8217–8224 (1984).
15. Saori Yuuya *et al.* Guaianolides as Immunomodulators. Synthesis and Biological Activities of Dehydrocostus Lactone, Mokko Lactone, Eremanthin, and Their Derivatives. *J. Nat. Prod.* **62**, 22–30 (1999).
16. Wei, H. *et al.* Chemical constituents of *Dolomiaea souliei*. *China J. Chin. Mater. Med.* **37**, 1249–1253 (2012).
17. Sakurai, H. *et al.* Nitric oxide production and inducible nitric oxide synthase expression in inflammatory arthritides. *J. Clin. Invest.* **96**, 2357–2363 (1995).
18. Tan-no, K. *et al.* Anti-inflammatory Effect of Propolis through Inhibition of Nitric Oxide Production on Carrageenin-Induced Mouse Paw Edema. *Biol. Pharm. Bull.* **29**, 96–99 (2006).
19. Duarte, J., Francisco, V. & Perez-Vizcaino, F. Modulation of nitric oxide by flavonoids. *Food Funct.* **5**, 1653–1668 (2014).
20. Ono, R., Kaisho, T. & Tanaka, T. PDLIM1 inhibits NF- $\kappa$ B-mediated inflammatory signaling by sequestering the p65 subunit of NF- $\kappa$ B in the cytoplasm. *Sci. Rep.* **5**, 18327 (2015).
21. Yu, C., Qi, D., Sun, J. F., Li, P. & Fan, H. Y. Rhein prevents endotoxin-induced acute kidney injury by inhibiting NF- $\kappa$ B activities. *Sci. Rep.* **5**, 11822 (2015).

## Acknowledgements

This work was supported by the Professor of Chang Jiang Scholars Program, the National Nature Science Foundation of China (81102335, 81230090, 81473109, 81502957), the National High-Tech Research and Development Program of China (863 Program, 2014AA022201-03), the Scientific Foundation of Shanghai China (13401900103), the Shanghai Engineering Research Center for the Preparation of Bioactive Natural Products (16DZ2280200) and the China Postdoctoral Science Foundation funded project (2015M572740).

## Author Contributions

Chen, L.P. and Wu, G.Z. carried out the experimental work. Zhang, J.P., Ye, J., Liu, Q.X., Shen, Y.H. and Zhang, W.D. provided oversight. Li, H.L. conducted the experiments. Chen, L.P., Wu, G.Z. and Li, H.L. conceived the experiments and wrote the manuscript.

## Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing Interests:** The authors declare no competing financial interests.

**How to cite this article:** Chen, L.-P. *et al.* Vlasouliolides A–D, four rare C<sub>17</sub>/C<sub>15</sub> sesquiterpene lactone dimers with potential anti-inflammatory activity from *Vladimiria souliei*. *Sci. Rep.* **7**, 43837; doi: 10.1038/srep43837 (2017).

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017