

SCIENTIFIC REPORTS



OPEN

Identifying the chiral d -wave superconductivity by Josephson φ_0 -states

Jun-Feng Liu¹, Yong Xu¹ & Jun Wang²

Received: 02 December 2016

Accepted: 30 January 2017

Published: 07 March 2017

We propose the Josephson junctions linked by a normal metal between a $d + id$ superconductor and another $d + id$ superconductor, a d -wave superconductor, or a s -wave superconductor for identifying the chiral $d + id$ superconductivity. The time-reversal breaking in the chiral d -wave superconducting state is shown to result in a Josephson φ_0 -junction state where the current-phase relation is shifted by a phase φ_0 from the sinusoidal relation, other than 0 and π . The ground-state phase difference φ_0 and the critical current can be used to definitely confirm and read the information about the $d + id$ superconductivity. A smooth evolution from conventional $0-\pi$ transitions to tunable φ_0 -states can be observed by changing the relative magnitude of two types of d -wave components in the $d + id$ pairing. On the other hand, the Josephson junction involving the $d + id$ superconductor is also the simplest model to realize a φ_0 -junction, which is useful in superconducting electronics and superconducting quantum computation.

Much interest has been attracted to recent theoretical predictions that the superconducting pairing driven by strong electron correlation in materials with three- and sixfold rotational lattice symmetries favors topological chiral $d + id$ symmetry. This symmetry is close to the d -wave superconducting pair in high temperature superconductors, but time-reversal broken. The materials which have been proposed to be chiral d -wave superconductors include graphene and silicene^{1–3}, MoS₂^{4–6}, In₃Cu₂VO₉⁷, SrPtAs^{8,9}, and bilayer SrIrO₃^{10,11}. But the experimental identification is still lacking. One of the experimental method to verify the chiral d -wave superconductivity is detecting the edge supercurrent induced by the topologically nontrivial superconducting order. But so far, no definite evidence of edge supercurrent is observed. Also, even theoretically, whether chiral d -wave supports an edge supercurrent still remains controversial¹². Therefore, how to identify this novel superconducting order in experiments is of significant importance to further develop the microscopic theory of high temperature superconductors. The Josephson effect is another powerful method to identify the superconducting pair, should be promising to definitely identify the chiral d -wave superconductivity.

The efforts to detect the chiral d -wave order by Josephson effect are currently focusing on the signal of the critical current of the Josephson junction which varies with the superconducting order or other junction parameters. But the critical current, the amplitude of supercurrent, includes only indirect information on the pair symmetry, and can not definitely identify the chiral d -wave order. By contrast, the ground-state phase difference of the Josephson current, namely, the so-called Josephson φ_0 -states contains more ample and direct information on the pair potential, thus should be more promising to definitely identify the chiral d -wave pairing. Therefore, the theoretical and experimental investigation in the relation between the ground-state phase difference and the chiral d -wave pairing become important. To our best knowledge, the effort in this way is still blank Fig. 1.

In the Josephson φ_0 -state, or the anomalous Josephson effect, the current-phase relation (CPR) has a phase shift φ_0 compared with the conventional CPR, namely, $I(\varphi) = I_c \sin(\varphi - \varphi_0)$ ^{13–20}. The ground-state phase difference φ_0 is neither 0 nor π in general, and tunable by the junction parameters. Such a φ_0 -state has been predicted in Josephson junctions with coexisting exchange field and spin-orbit coupling^{14–17,19}, multilayer ferromagnets^{18,20}, noncentrosymmetric superconductors^{21–23}, and topological edge or interface states^{24–26}. Very recently, the first experimental demonstration of a φ_0 -junction has been reported in a quantum dot junction by use of a quantum interferometer device²⁷. These φ_0 -junctions with tunable ground-state phase difference may have applications in superconducting computer memory components²⁸, superconducting phase batteries and rectifiers²⁹, as well

¹Department of Physics, South University of Science and Technology of China, Shenzhen 518055, China.

²Department of Physics, Southeast University, Nanjing 210096, China. Correspondence and requests for materials should be addressed to J.-F.L. (email: liujf@sustc.edu.cn) or J.W. (email: jwang@seu.edu.cn)




Figure 1. Schematic diagrams of Josephson junctions linked by a normal metal between a $d + id$ superconductor and (a) another $d + id$ superconductor, (b) a d -wave superconductor, and (c) a s -wave superconductor. The junctions lie in the x - y plane and the transport is along the x -direction.

as flux- or phase-based quantum bits³⁰. Among various ways to achieve a φ_0 -junction, the junctions involving chiral d -wave superconductors should be the simplest one because the time-reversal symmetry is already broken in chiral d -wave pairing.

In this study, we investigate the Josephson junctions linked by a normal metal between a $d + id$ superconductor and another $d + id$ superconductor, a d -wave superconductor, or a s -wave superconductor (see Fig. 1). Anomalous Josephson effect appears as a result of the broken time-reversal symmetry in the $d + id$ superconductor. The ground-state phase difference φ_0 other than 0 and π should be the definite evidence of the $d + id$ pairing. Furthermore, the ground-state phase difference and the critical current can be used to determine the ratio between two types of d -wave components in the $d + id$ superconductor. The demonstration of a smooth evolution from conventional 0 - π transitions to tunable φ_0 -states is also interesting.

The paper is organized as follows. In Sec. II we present the model Hamiltonian and introduce the method to solve the CPR. The numerical results and relevant discussion of three types of junctions will be given in Sec. III. Finally, the conclusion will be given in Sec. IV.

Model and Methods

We begin with the BdG Hamiltonian of a two-dimensional $d + id$ superconductor with parabolic spectrum in the normal state. The particular form of the spectrum may not take much effect in the ground-state phase difference of the Josephson junction, but the phase of the pair potential does. The spin degree of freedom is also not important and ignored here. Then the simplest BdG Hamiltonian of a $d + id$ superconductor can be written as

$$H = \begin{pmatrix} \varepsilon_{\mathbf{k}} & \Delta(\theta) \\ \Delta^*(\theta) & -\varepsilon_{\mathbf{k}} \end{pmatrix} \tag{1}$$

where $\varepsilon_{\mathbf{k}} = \frac{\hbar^2(k_x^2 + k_y^2)}{2m} - \mu = \frac{\hbar^2 k^2}{2m} - \mu$ is the kinetic energy measured from the chemical potential μ . The $d + id$ pair potential is

$$\begin{aligned} \Delta(\theta) &= \Delta_1 \cos [2(\theta - \gamma)] + i\Delta_2 \sin [2(\theta - \gamma)] \\ &= |\Delta(\theta)| e^{i\delta(\theta)} \end{aligned} \tag{2}$$

with

$$\begin{aligned} |\Delta(\theta)| &= \sqrt{\Delta_1^2 \cos^2 [2(\theta - \gamma)] + \Delta_2^2 \sin^2 [2(\theta - \gamma)]}, \\ \delta(\theta) &= \text{sign}[\sin 2(\theta - \gamma)] \arccos \frac{\Delta_1 \cos [2(\theta - \gamma)]}{|\Delta(\theta)|}. \end{aligned} \tag{3}$$

Here θ is the injection angle satisfying $\tan\theta = k_y/k_x$, γ is the angle between the x -direction and the α axis of the superconductor. Δ_1 and Δ_2 are two positive real numbers, denoting the amplitudes of two kinds of d -waves. It is clearly shown that an additional phase $\delta(\theta)$ emerges in the pair potential and depends on the injection angle.

For a Josephson junction between two $d + id$ superconductors, the pair potential can be approximately described by two step functions $\Delta(x) = [\Delta_L \Theta(-x) e^{i\varphi/2} + \Delta_R \Theta(x-L) e^{-i\varphi/2}]$ where L is the length of the normal layer and φ is the macroscopic phase difference between two superconductors. Similar to Eq. (delta), the left (right) pair potential Δ_L (Δ_R) reads $\Delta_\lambda(\theta) = \Delta_{\lambda 1} \cos[2(\theta - \gamma_\lambda)] + i\Delta_{\lambda 2} \sin[2(\theta - \gamma_\lambda)] = |\Delta_\lambda(\theta)| e^{i\delta_\lambda(\theta)}$ with $\lambda = L$ or R denoting the left or right superconductor respectively. For simplicity, we assume that the momentum component in the y -direction is conserved. Then the eigen wavefunctions in two superconductors can be written as

$$\phi_{\lambda e \pm} = \begin{pmatrix} u_{\lambda \pm} e^{i(\frac{1}{4}\lambda\varphi + \frac{1}{2}\delta_{\lambda \pm})} \\ v_{\lambda \pm} e^{-i(\frac{1}{4}\lambda\varphi + \frac{1}{2}\delta_{\lambda \pm})} \end{pmatrix} e^{\pm ik_{\lambda \tau \pm} x} \tag{4}$$

for electron-like quasiparticles and

$$\phi_{\lambda h \mp} = \begin{pmatrix} v_{\lambda \mp} e^{i(\frac{1}{4}\lambda\varphi + \frac{1}{2}\delta_{\lambda \mp})} \\ u_{\lambda \mp} e^{-i(\frac{1}{4}\lambda\varphi + \frac{1}{2}\delta_{\lambda \mp})} \end{pmatrix} e^{\mp ik_{\lambda \tau \mp} x} \tag{5}$$

for hole-like quasiparticles where $k_{\lambda \tau \pm} = \sqrt{k_F^2 \cos^2 \theta + \tau \frac{2m}{\hbar^2} \sqrt{E^2 - |\Delta_{\lambda \pm}|^2}}$, and $u_{\lambda \pm} = \sqrt{(E + \Omega_{\lambda \pm})/2E}$, $v_{\lambda \pm} = \sqrt{(E - \Omega_{\lambda \pm})/2E}$ with $\Omega_{\lambda \pm} = \sqrt{E^2 - |\Delta_{\lambda \pm}|^2}$. Here $k_F = \sqrt{2m\mu}/\hbar$ is the Fermi wave vector, $\lambda = 1$ (-1) for the left (right) superconductor, $\tau = 1$ (-1) for electron-like (hole-like) quasiparticles. $\Delta_{\lambda \pm} = \Delta_\lambda(\pm\theta)$ with $\Delta_{\lambda+}$ ($\Delta_{\lambda-}$) being the pair potential for right-going electron-like and left-going hole-like (left-going electron-like and right-going hole-like) quasiparticles. $\delta_{\lambda \pm} = \delta(\pm\theta, \gamma_\lambda)$ according to Eq. (3).

The eigen wavefunctions can be easily solved for the normal layer. Then the scattering problem can be solved by considering the boundary conditions at two interfaces. Each interface gives a scattering matrix, from which the reflection matrix of the right-going (left-going) incident particles R_1 (R_2) can be abstracted in the normal layer. To calculate the Josephson current, we can work out the Green's function $G(z, z', E)$ in the normal layer which is made of the reflection matrices R_1 and R_2 ³¹. Then the Josephson current in the normal layer can be evaluated by

$$I(\varphi) = \frac{e\hbar k_B T}{2im} \lim_{x' \rightarrow x} \left(\frac{\partial}{\partial x'} - \frac{\partial}{\partial x} \right) \sum_{\omega_n} \text{Tr} G_{\omega_n}(x, x')$$

where $G_{\omega_n}(z, z')$ is the Matsubara-Green's function with the Matsubara frequencies $\omega_n = \pi k_B T(2n + 1)$, $n = 0, \pm 1, \pm 2, \dots$. By integrating over the injection angle γ , the total Josephson current in two dimension is

$$J(\varphi) = \frac{k_F W}{2\pi} \int_{-\pi/2}^{\pi/2} I(\varphi, \theta) \cos \theta d\theta, \tag{6}$$

where W is the transversal width of the junction.

Results and Discussion

Next, we present the numerical results and relevant discussion of three types of junctions involving chiral d -wave superconductors. We focus on the ground-state phase difference and critical current of the CPR. The interface transparency is found to neither change the ground-state phase difference, nor change the relative magnitude of critical current, but the absolute value of critical current. Therefore, the results about the effect of interface barriers is not presented here.

Junction between two $d + id$ superconductors. For the Junction between two $d + id$ superconductors with different directions of the α -axis, the Josephson current is shown in Fig. 2. In the simple case $\Delta_{L1} = \Delta_{L2} = \Delta_{R1} = \Delta_{R2}$, the additional phase in the pairing potential for the left and right superconductor $\delta_\lambda(\theta) = 2(\theta - \gamma_\lambda)$ according to Eq. (3). Then the additional phase difference is $2(\gamma_R - \gamma_L)$, which means the ground-state phase difference is $2\gamma_L$ when $\gamma_R = 0$. It is exactly the case shown in Fig. 2(a) where the nearly sinusoidal CPR just moves to the right with increasing γ_L . The ground-state phase difference increases smoothly, while the shape and amplitude of the CPR keep unchanged. In the other limit $\Delta_{L2} = 0$, $\Delta_L(\theta) = \Delta_{L1} \cos[2(\theta - \gamma_L)]$. The left superconductor becomes a d -wave superconductor and the additional phase is 0. However, $|\Delta_L(\theta)|$ reaches its maximum at $\theta = \gamma_L$, $\gamma_L \pm \frac{\pi}{2}$, or $\gamma_L - \pi$. It implies that the Josephson current is dominated by the components related to these two incident angles. For these two components, the pair potential changes sign for both left and right superconductors. Then the ground-state phase difference is nearly still $2\gamma_L$ because the additional phase for the right superconductor is 2θ with $\gamma_R = 0$. As shown in Fig. 2(b), the ground-state phase difference φ_0 nearly keeps unchanged with varied ratio Δ_{L2}/Δ_{L1} and just equals $2\gamma_L$ with increasing γ_L . The critical current J_c is shown in Fig. 2(c). With decreasing ratio Δ_{L2}/Δ_{L1} , the critical current decreases because $|\Delta_L(\theta)|$ decreases for all the incident angles θ except for the angles at which $|\Delta_L(\theta)|$ takes the maximum. Note that with a fixed Δ_{L2}/Δ_{L1} smaller than 1, the critical current oscillates with a period of $\frac{\pi}{2}$ with increasing γ_L . This oscillation is due to the factor $\cos \theta$ in Eq. (6) and the period $\frac{\pi}{2}$ is attributed to the four-fold rotational symmetry of $d + id$ pairing. With decreasing ratio Δ_{L2}/Δ_{L1} from 1 to 0, the amplitude of the oscillation increases from 0.




Figure 2. Josephson current for the junction between two $d + id$ superconductors with different directions of the α -axis. (a) CPR for various γ_L with fixed $\Delta_{L2}/\Delta_{L1} = 1$. (b) The ground-state phase difference and (c) the critical current as functions of Δ_{L2}/Δ_{L1} and γ_L are shown in the contour plots. The temperature $T = 0.5T_c$ with T_c the critical temperature. $\Delta_{L1} = \Delta_{R1} = \Delta_{R2} = 10^{-3} \mu$, $\gamma_R = 0$. $|\Delta_L|_{max} = \max(\Delta_{L1}, \Delta_{L2})$.

Junction between a $d + id$ superconductor and a d -wave superconductor. Figure 3 shows the Josephson current through the junction between a $d + id$ superconductor and a d -wave superconductor. The d -wave pairing in the right superconductor is chosen to be a d_{xy} -wave pairing by setting $\Delta_{R2} = 0$ and $\gamma_R = \frac{\pi}{4}$. According to the above discussion, the right d_{xy} -wave pairing decides that the dominant components in the Josephson current come from the incident angles $\theta = \pm \frac{\pi}{4}$. When $\Delta_{L2}/\Delta_{L1} = 1$, the additional phases in $\Delta_L(\theta)$ for the two components are $\delta_L(\theta) = \pm \frac{\pi}{2} - 2\gamma_L$, which leads to the ground-state phase difference $\varphi_0 = 2\gamma_L - \frac{\pi}{2}$ by taking into account of the sign change in d_{xy} -wave pairing. The numerical results on φ_0 shown in Fig. 3(a,b) verify this discussion. Because $|\Delta_L(\theta)|$ is independent of θ and γ_L for $\Delta_{L2}/\Delta_{L1} = 1$, the critical current is also independent of γ_L as shown in Fig. 3(a,c). In the other limit $\Delta_{L2} = 0$, the left superconductor reduces to a d -wave superconductor and its additional phase becomes 0. Then the anomalous Josephson effect disappears and φ_0 must be either 0 or π . With increasing γ_L , the junction experiences conventional $0-\pi$ transitions which is accompanied with heavy oscillations in the critical current. The oscillations in the critical current come from the varying angles between the α -axes of two d -wave superconductors. Specifically, the junction is a 0 -junction for $\gamma_L \in [0, \frac{\pi}{2}]$ and turns into a π -junction for $\gamma_L \in [\frac{\pi}{2}, \pi]$, accompanied with minima in J_c at the transition points $\gamma_L = 0, \frac{\pi}{2}, \pi$. It is interesting to note the smooth transition between these two limiting cases $\Delta_{L2}/\Delta_{L1} = 0$ and 1. With the changing of Δ_{L2}/Δ_{L1} from 1 to 0, φ_0 increases more and more nonuniformly with increasing γ_L , at the same time accompanied with more and more heavy oscillations in J_c , and finally evolves into a conventional $0-\pi$ transition.




Figure 3. Josephson current for the Junction between a $d + id$ superconductors and a d -wave superconductor. (a) CPR for various γ_L with fixed $\Delta_{L2}/\Delta_{L1} = 1$. (b) The ground-state phase difference and (c) the critical current as functions of Δ_{L2}/Δ_{L1} and γ_L are shown in the contour plots. $T = 0.5T_c$, $\Delta_{L1} = \Delta_{R1} = 10^{-3} \mu$, $\Delta_{R2} = 0$, $\gamma_R = \pi/4$.

Junction between a $d + id$ superconductor and a s -wave superconductor. The situation is similar for the junction between a $d + id$ superconductor and a s -wave superconductor. The s -wave pairing in the right superconductor is described by setting $\Delta_R(\theta) = \Delta_R$ where the additional phase $\delta_R(\theta) = 0$. When $\Delta_{L2}/\Delta_{L1} = 1$, the additional phase in $\Delta_L(\theta)$ is $\delta_L(\theta) = 2(\theta - \gamma_L)$. The angle-resolved Josephson current in the first harmonic approximation is $I(\theta) \propto |\Delta_L| \Delta_R \sin(\varphi + 2\theta - 2\gamma_L)$. The total Josephson current is

$$\begin{aligned}
 J(\varphi) &\propto \int_{-\pi/2}^{\pi/2} I(\varphi, \theta) \cos \theta d\theta \\
 &\propto |\Delta_L| \Delta_R \sin(\varphi - 2\gamma_L).
 \end{aligned}
 \tag{7}$$

This conclusion of $\varphi_0 = 2\gamma_L$ is consistent with the numerical result shown in Fig. 4(b). When $\Delta_{L2}/\Delta_{L1} = 0$, the junction reduces to a d -wave/normal metal/ s -wave junction. Only the conventional $0-\pi$ transition is possible due to the sign change in the d -wave pairing with increasing γ_L . The evolution from the tunable φ_0 -state for $\Delta_{L2}/\Delta_{L1} = 1$ to conventional $0-\pi$ transition for $\Delta_{L2}/\Delta_{L1} = 0$ is similar to that discussed in Fig. 3. For the general case between $\Delta_{L2}/\Delta_{L1} = 1$ and 0, the change of φ_0 with increasing γ_L is simultaneously accompanied with the oscillation in J_c . The oscillation in J_c is achieved by second and higher harmonic terms as shown in Fig. 4(a).

Finally, we comment on the experimental feasibility of observing the $d + id$ superconductivity. In view of the fact that there are various possible pairing mechanisms^{32,33} in graphene and graphene-like materials, we suggest that SrPtAs should be the most promising candidate to observe the $d + id$ pairing. The experimental challenge lies in the observing the ground-state phase difference, and can be met by employing the superconducting quantum




Figure 4. Josephson current for the Junction between a $d + id$ superconductors and a s -wave superconductor. (a) CPR for various γ_L with fixed $\Delta_{L2}/\Delta_{L1} = 0.3$. (b) The ground-state phase difference and (c) the critical current as functions of Δ_{L2}/Δ_{L1} and γ_L are shown in the contour plots. $T = 0.5T_c$, $\Delta_{L1} = \Delta_R = 10^{-3}\mu$.

interference device²⁷. We argue that the slight nonmagnetic impurities do not affect the ground-state phase difference qualitatively if only the $d + id$ pairing is not destroyed. But the interplay between the magnetic impurities and the superconductivity is complicated and beyond the topic of this paper, and may be the next aim in our further study. Otherwise, we employ a quadratic dispersion relation to describe the quasiparticles in the normal state for the $d + id$ paired superconducting leads. This quadratic dispersion is obviously invalid for graphene and graphene-like materials. But the result for the ground-state phase difference will not be changed because it is only affected by the phase of the pair potential. The critical current will be qualitatively unchanged but quantitatively modified if the linear dispersion is adopted.

Conclusion

In conclusion, we study the anomalous Josephson effect in junctions with chiral $d + id$ superconductor induced by the time-reversal breaking of superconducting order. The ground-state phase difference ϕ_0 other than 0 and π is predicted and should be the definite evidence of the $d + id$ pairing. The ground-state phase difference and the critical current are shown to depend on the ratio between two types of d -wave components and the direction of the α -axis of the $d + id$ superconductor. The demonstration of a smooth evolution from conventional $0-\pi$ transitions to tunable ϕ_0 -states advances the understanding of Josephson effect. And the simple ϕ_0 -junction consisting of the $d + id$ superconductor is important to applications in superconducting electronics and superconducting quantum computation.

References

- Vučičević, J., Goerbig, M. O. & Milovanović, M. V. d-wave superconductivity on the honeycomb bilayer. *Phys. Rev. B* **86**, 214505 (2012).
- Liu, F., Liu, C.-C., Wu, K., Yang, F. & Yao, Y. d+id' Chiral Superconductivity in Bilayer Silicene. *Phys. Rev. Lett.* **111**, 066804 (2013).
- Vafeek, O., Murray, J. M. & Cvetkovic, V. Superconductivity on the Brink of Spin-Charge Order in a Doped Honeycomb Bilayer. *Phys. Rev. Lett.* **112**, 147002 (2014).
- Ye, J. T. *et al.* Superconducting dome in a gate-tuned band insulator. *Science* **338**, 1193 (2012).
- Taniguchi, K., Matsumoto, A., Shimotani, H. & Takagi, H. Electric-field-induced superconductivity at 9.4 K in a layered transition metal disulphide MoS₂. *Appl. Phys. Lett.* **101**, 042603 (2012).
- Yuan, N. F. Q., Mak, K. F. & Law, K. T. Possible topological superconducting phases of MoS₂. *Phys. Rev. Lett.* **113**, 097001 (2014).
- Wu, W., Scherer, M. M., Honerkamp, C. & Hur, K. Le. Correlated Dirac particles and superconductivity on the honeycomb lattice. *Phys. Rev. B* **87**, 094521 (2013).
- Biswas, P. K. *et al.* Evidence for superconductivity with broken time-reversal symmetry in locally noncentrosymmetric SrPtAs. *Phys. Rev. B* **87**, 180503 (2013).
- Fischer, M. H. *et al.* Chiral d-wave superconductivity in SrPtAs. *Phys. Rev. B* **89**, 020509 (2014).
- Okamoto, S. Doped mott insulators in (111) bilayers of perovskite transition-metal oxides with a strong spin-orbit coupling. *Phys. Rev. Lett.* **110**, 066403 (2013).
- Okamoto, S. Global phase diagram of a doped Kitaev-Heisenberg model. *Phys. Rev. B* **87**, 064508 (2013).
- Huang, W., Taylor, E. & Kallin, C. Vanishing edge currents in non-p-wave topological chiral superconductors. *Phys. Rev. B* **90**, 224519 (2014).
- Brydon, P. M. R., Kastening, B., Morr, D. K. & Manske, D. Interplay of ferromagnetism and triplet superconductivity in a Josephson junction. *Phys. Rev. B* **77**, 104504 (2008).
- Reynoso, A. A., Usaj, G., Balseiro, C. A., Feinberg, D. & Avignon, M. Anomalous Josephson current in junctions with spin polarizing quantum point contacts. *Phys. Rev. Lett.* **101**, 107001 (2008).
- Buzdin, A. Direct coupling between magnetism and superconducting current in the Josephson φ_0 junction. *Phys. Rev. Lett.* **101**, 107005 (2008).
- Zazunov, A., Egger, R., Jonckheere, T. & Martin, T. Anomalous Josephson current through a Spin-Orbit coupled quantum dot. *Phys. Rev. Lett.* **103**, 147004 (2009).
- Liu, J.-F. & Chan, K. S. Relation between symmetry breaking and the anomalous Josephson effect. *Phys. Rev. B* **82**, 125305 (2010).
- Liu, J.-F. & Chan, K. S. Anomalous Josephson current through a ferromagnetic trilayer junction. *Phys. Rev. B* **82**, 184533 (2010).
- Yokoyama, T., Eto, M. & Nazarov, Y. V. Anomalous Josephson effect induced by spin-orbit interaction and Zeeman effect in semiconductor nanowires. *Phys. Rev. B* **89**, 195407 (2014).
- Kulagina, I. & Linder, J. Spin supercurrent, magnetization dynamics, and φ -state in spin-textured Josephson junctions. *Phys. Rev. B* **90**, 054504 (2014).
- Zhang, H., Wang, J. & Liu, J.-F. Anomalous Josephson effect in noncentrosymmetric superconductors. *Appl. Phys. Lett.* **108**, 102601 (2016).
- Klam, L., Epp, A., Chen, W., Sigrist, M. & Manske, D. Josephson effect and triplet-singlet ratio of noncentrosymmetric superconductors. *Phys. Rev. B* **89**, 174505 (2014).
- Rahnavard, Y., Manske, D. & Annunziata, G. Magnetic Josephson junctions with noncentrosymmetric superconductors. *Phys. Rev. B* **89**, 214501 (2014).
- Tanaka, Y., Yokoyama, T. & Nagaosa, N. Manipulation of the Majorana fermion, Andreev reflection, and Josephson current on topological insulators. *Phys. Rev. Lett.* **103**, 107002 (2009).
- Dolcini, F., Houzet, M. & Meyer, J. S. Topological Josephson φ_0 junctions. *Phys. Rev. B* **92**, 035428 (2015).
- Wang, J., Hao, L. & Liu, J.-F. Electric control of the Josephson current-phase relation in a topological circuit. *Phys. Rev. B* **93**, 155405 (2016).
- Szombati, D. B. *et al.* Josephson φ_0 -junction in nanowire quantum dots. *Nat. Phys.* **12**, 568 (2016).
- Gingrich, E. C. *et al.* Controllable $0-\pi$ Josephson junctions containing a ferromagnetic spin valve. *Nat. Phys.* **12**, 564 (2016).
- Reynoso, A. A., Usaj, G., Balseiro, C. A., Feinberg, D. & Avignon, M. Spin-orbit-induced chirality of Andreev states in Josephson junctions. *Phys. Rev. B* **86**, 214519 (2012).
- Padurariu, C. & Nazarov, Y. V. Theoretical proposal for superconducting spin qubits. *Phys. Rev. B* **81**, 144519 (2010).
- Kalenkov, M. S., Galaktionov, A. V. & Zaikin, A. D. Josephson current in ballistic heterostructures with spin-active interfaces. *Phys. Rev. B* **79**, 014521 (2009).
- Uchoa, B. & Neto, A. H. C. Superconducting States of Pure and Doped Graphene. *Phys. Rev. Lett.* **98**, 146801 (2007).
- Pellegrino, F. M. D., Angilella, G. G. N. & Pucci R. Pairing symmetry of superconducting graphene. *Eur. Phys. J. B* **76**, 469 (2010).

Acknowledgements

The work described in this paper is supported by the National Natural Science Foundation of China (NSFC, Grant Nos 11204187, and 11274059).

Author Contributions

J.F.L. and J.W. conceived the study. J.F.L. and Y.X. performed the numerical calculations. J.F.L. wrote the main manuscript text. All authors contributed to discussion and reviewed the manuscript.

Additional Information

Competing Interests: The authors declare no competing financial interests.

How to cite this article: Liu, J.-F. *et al.* Identifying the chiral *d*-wave superconductivity by Josephson φ_0 -states. *Sci. Rep.* **7**, 43899; doi: 10.1038/srep43899 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017