

SCIENTIFIC REPORTS



OPEN

Experimental contextuality in classical light

Tao Li^{1,2}, Qiang Zeng¹, Xinbing Song¹ & Xiangdong Zhang¹

Received: 20 December 2016

Accepted: 07 February 2017

Published: 14 March 2017

The Klyachko, Can, Binicioglu, and Shumovsky (KCBS) inequality is an important contextuality inequality in three-level system, which has been demonstrated experimentally by using quantum states. Using the path and polarization degrees of freedom of classical optics fields, we have constructed the classical trit (cetrit), tested the KCBS inequality and its geometrical form (Wright's inequality) in this work. The projection measurement has been implemented, the clear violations of the KCBS inequality and its geometrical form have been observed. This means that the contextuality inequality, which is commonly used in test of the conflict between quantum theory and noncontextual realism, may be used as a quantitative tool in classical optical coherence to describe correlation characteristics of the classical fields.

Contextuality is a curious property in quantum world. In classical world, an observable is predefined and has a joint probability distribution when it is measured with the other observables. This observable is independent of other compatible observables that are measured simultaneously with it. This independency is called noncontextuality. Kochen and Specker proposed the original contextual theory in 1967¹. The theory gives the conflict between quantum mechanics and noncontextuality reality. The original theory needs 117 tests in dimension $d = 3$, and it is complex and nearly impossible to demonstrate experimentally. Afterwards the theory is simplified by many researchers²⁻⁷. These simplified contextuality theories have been tested experimentally, for instance in photon⁸⁻¹³, neutron^{14,15}, trapped ion¹⁶ and nuclear magnetic resonance^{17,18} systems. For the three-level system, Yu and Oh proposed a contextuality inequality, which needs only 13 variables and 24 pair correlations¹⁹. The scenario is state independent and only requires a single qutrit. Klyachko, Can, Binicioglu, and Shumovsky also found a state dependent contextuality inequality in the three-level system²⁰. The three-level system is the simplest case to show quantum contextuality. The Klyachko-Can-Binicioglu-Shumovsky (KCBS) inequality is fundamentally important, because it is behind the violation of other noncontextualities²¹.

The state independent inequality (13 variables and 24 pair correlations as described in ref. 19) has been demonstrated by Zu *et al.* using the qutrit encoded in three paths of the single photons²². In a single trapped ion system, the inequality has also been tested by Zhang *et al.*²³. Using the polarization and path of single photons, the demonstration of the inequality has been implemented by Huang *et al.*²⁴. These three-level systems are indivisible. For the KCBS inequality as shown in ref. 20, several experiments have also been performed with the single photons, trapped ions and biphotons. Lapkiewicz *et al.* firstly proved the violation of the KCBS inequality with the single photon qutrit²⁵. Subsequently, the experimental demonstrations of the KCBS inequality and the Wright inconsistency have been reported by Ahrens *et al.*²⁶. The demonstration of violation of the KCBS inequality and the random-number production have been implemented by a single trapped ion²⁷. Using nitrogen-vacancy centers, Kong *et al.* have tested the violation of the geometrical form of the KCBS inequality²⁸. An experiment for the KCBS contextuality and nonlocality has also been implemented by Shaham *et al.*²⁹. In addition, the no-disturbance monogamy relation about the KCBS contextuality and the nonlocality has also been discussed^{30,31}. The contextuality is important for applications³² and theoretical investigations, but it is difficult to implement experimentally. Thus, it is necessary to develop new simplified methods.

On the other hand, recent investigation has shown that the quantum bound is not exclusive to quantum theory³³. It has been demonstrated that the quantum bound exists similarly in classical microwaves. The local and nonlocal correlations in the classical optical beams, which violate the Bell inequality, have been demonstrated in a series of works³⁴⁻⁴¹. However, the KCBS contextuality has not been studied in classical wave systems so far.

In this paper we study the KCBS contextuality in classical light systems. The polarization and path of the classical light beams are used to build the classical trit, which is called the "cetrit" like the cebit in ref. 42. Based on the

¹Beijing Key Laboratory of Nanophotonics & Ultrafine Optoelectronic Systems, School of Physics, Beijing Institute of Technology, 100081, Beijing, China. ²Beijing Key Laboratory of Nanophotonics & Ultrafine Optoelectronic Systems, Kunming Institute of Physics, Kunming, 650223, China. Correspondence and requests for materials should be addressed to X.Z. (email: zhangxd@bit.edu.cn)

bases of the cetrit, the input state is constructed, and then the observables are tested by projective measurement. Here the typical KCBS inequality and its geometrical form are studied, and the results are compared with the noncontextuality cases.

Results and Discussion

KCBS inequality and its expression in classical optical systems. According to ref. 20, if we consider the five observables A'_i ($i = 1, 2, \dots, 5$), which have the values $+1$ or -1 , the algebraic inequality for any choice of them can be expressed as

$$A'_1 A'_2 + A'_2 A'_3 + A'_3 A'_4 + A'_4 A'_5 + A'_5 A'_1 \geq -3. \quad (1)$$

For the contextuality inequality, the observables A'_i correspond to the five unit vectors, which are expressed with a'_i . The eigenvalues of the operators $|a'_i\rangle\langle a'_i|$ are 0 or 1, as well as $A'_i = \mathbb{I} - 2|a'_i\rangle\langle a'_i|$ (where \mathbb{I} is a 3×3 identity matrix), thus the eigenvalues of A'_i are $+1$ or -1 . The unit vectors a'_i are presented at the five vertexes of pentagram, where a'_i and a'_{i+1} ($i + 1$ modulo 5) are orthogonal, and they are compatible. When a'_i and a'_{i+1} project to the pentagram plane, the angle between the projected vectors of a'_i and a'_{i+1} is $4\pi/5$, and the angle between the direction vector and the symmetry axis is approximately $4\pi/15$. It is easy to obtain:

$$|a'_i\rangle = N \left[\cos\left(\frac{4\pi i}{5}\right)|0\rangle + \sin\left(\frac{4\pi i}{5}\right)|1\rangle + \sqrt{\cos\left(\frac{\pi}{5}\right)}|2\rangle \right], \quad (2)$$

where $N = 1/\sqrt{1 + \cos(\pi/5)}$ is a normalization constant. For the KCBS inequality, the input state is at the symmetry axis of the pentagram²⁰. The input state $|\varphi\rangle$ projects to the five couples of observables $A'_i A'_{i+1}$, and the inequality has the maximum violation,

$$\langle A'_1 A'_2 \rangle + \langle A'_2 A'_3 \rangle + \langle A'_3 A'_4 \rangle + \langle A'_4 A'_5 \rangle + \langle A'_5 A'_1 \rangle \geq 5 - 4\sqrt{5} \approx -3.944, \quad (3)$$

where $\langle A'_i A'_{i+1} \rangle$ is used to denote the average of the measurement outcome of $A'_i A'_{i+1}$. The KCBS inequality is state dependent, here the input state is taken as the specific input state $|\varphi\rangle = |2\rangle$. The geometric form of the KCBS inequality or Wright's inequality is expressed as²⁰

$$\sum_{i=1}^5 |\langle a'_i | \varphi \rangle|^2 \leq 2 \quad (4)$$

Under the Kochen-Specker rules, the value at every direction vector is predetermined to 1 or 0, but the maximum is only one for the two orthogonal directions. Thus, the maximum is two for the five directions of pentagram. In quantum mechanics we select the same input state like that in the KCBS inequality, and $|\langle a'_i | \varphi \rangle|^2 = \cos^2 \frac{4\pi}{15} = \frac{1}{\sqrt{5}}$ (the angle between a'_i and $|\varphi\rangle$ is approximately $4\pi/15$) can be obtained. Thus, we have

$$\sum_{i=1}^5 |\langle a'_i | \varphi \rangle|^2 = \sqrt{5} \approx 2.236 > 2. \quad (5)$$

The above inequalities (Equations (3) and (5)) have been demonstrated experimentally using quantum states^{25–29}.

Now we use classical optical fields to express the input state and the direction unit vectors for the KCBS inequality and its geometric form as showed in Fig. 1. The input state is written as $|\chi\rangle = E_0|\vec{e}_0\rangle + E_1|\vec{e}_1\rangle + E_2|\vec{e}_2\rangle$, where E_0, E_1 and E_2 represent amplitudes of the classical optical fields, and $|\vec{e}_0\rangle, |\vec{e}_1\rangle$ and $|\vec{e}_2\rangle$ are the cetrit's bases corresponding to quantum bases $|0\rangle, |1\rangle$ and $|2\rangle$ in three-dimensional space⁴³. Here a slightly modified version of the familiar bra-ket notation of quantum mechanics is taken to express these vectors in the classical optical fields. Meanwhile the direction unit vector can be denoted as $|a_i\rangle = \alpha_i|\vec{e}_0\rangle + \beta_i|\vec{e}_1\rangle + \gamma_i|\vec{e}_2\rangle$, where $\alpha_i = N \cos \frac{4\pi i}{5}$, $\beta_i = N \sin \frac{4\pi i}{5}$ and $\gamma_i = N \sqrt{\cos \frac{\pi}{5}}$ are the coefficients corresponding to those in Eq. (2).

In the following we calculate the correlations $A_i A_{i+1}$ of classical optical fields. In order to describe the KCBS inequality, $|\chi\rangle = E_2|\vec{e}_2\rangle$ is taken as the input state, which is at the symmetry axis of the pentagram as shown in Fig. 1. $A_i = \mathbb{I} - 2|a_i\rangle\langle a_i|$ is used to express the observable in the classical optical systems, where $|a_i\rangle$ is the direction unit vector. For the input state $|\chi\rangle$, from the correlation $A_i A_{i+1} = [\mathbb{I} - 2|a_i\rangle\langle a_i|][\mathbb{I} - 2|a_{i+1}\rangle\langle a_{i+1}|] = \mathbb{I} - 2|a_i\rangle\langle a_i| - 2|a_{i+1}\rangle\langle a_{i+1}|$, we can obtain the average value $\overline{(A_i A_{i+1})}$ of $A_i A_{i+1}$ as $\overline{(A_i A_{i+1})} = \langle \chi | A_i A_{i+1} | \chi \rangle = E_2^* E_2 \langle \vec{e}_2 | A_i A_{i+1} | \vec{e}_2 \rangle = |E_2|^2 (1 - 2\gamma_i^2 - 2\gamma_{i+1}^2) = |E_2|^2 \left(1 - \frac{2}{\sqrt{5}} - \frac{2}{\sqrt{5}}\right) = |E_2|^2 \left(1 - \frac{4}{\sqrt{5}}\right)$. The KCBS inequality requires the specific input state, thus the amplitude E_2 need to be normalized. When it is normalized, the value $\overline{(A_i A_{i+1})} = 1 - \frac{4}{\sqrt{5}}$, therefore

$$\overline{A_1 A_2} + \overline{A_2 A_3} + \overline{A_3 A_4} + \overline{A_4 A_5} + \overline{A_5 A_1} = 5 - 4\sqrt{5} \approx -3.944 < -3. \quad (6)$$

Equation (6) corresponds to Eq. (3), which is the corresponding form of the KCBS inequality in the classical optical systems. For the geometric form of the KCBS inequality (or called Wright's inequality), the input state is $|\chi\rangle = E_2|\vec{e}_2\rangle$, similarly we can obtain the equation $|\langle a_i | \chi \rangle|^2 = E_2^2 \gamma_i^2 = E_2^2 \cos^2 \frac{4\pi}{15} = E_2^2 \frac{1}{\sqrt{5}}$. The amplitude of the optical field needs to be normalized, $|\langle a_i | \chi \rangle|^2 = \frac{1}{\sqrt{5}}$, thus




Figure 1. Representation of the measurement for the KCBS inequality in three-dimensional space. A_1, A_2, A_3, A_4, A_5 are observables at five directions, and the direction vectors are pairwise orthogonal. $|\chi\rangle$ is used to express the input state in the classical light systems.




Figure 2. The experimental setup for the test of the KCBS inequality. The laser pulses from the He-Ne laser transmit through the GRIN lens (GL), and we obtain the horizontal polarization optical beam. Then the beam is modulated by HWPs and PBSs to achieve our construction. At last the light intensities are detected by the three photoelectric detectors (PD1, PD2 and PD3) at the three output ports. The HWP3, 4, and 6 are used for path-length compensation. HWP, half-wave plate; PBS, polarizing beam splitter; PD, photoelectric detector.

$$\sum_{i=1}^5 |a_i|\chi\rangle|^2 = \sqrt{5} \approx 2.236 > 2. \tag{7}$$

The geometric form of the KCBS inequality is also violated in the classical optical systems. In the following, we test experimentally the KCBS inequality and its geometrical form in the classical light systems.

Experimental demonstration of the KCBS inequality in classical light systems. The test of the KCBS inequality needs a three-level system. Using classical optical beams, we can construct the corresponding system. The scheme is shown in Fig. 2. It consists of two parts: state preparation and measurement. In the state preparation stage, the laser beam from He-Ne laser transmits through Glan prism and then the laser beam possessing the certain polarization appears. Here the center wavelength of the laser beam is 633 nm. The beam is split through polarizing beam splitters (PBSs) and is modulated by half-wave plates (HWPs) to perform the construction of classical trit (cetrir). The different polarizations in three paths of classical optical fields are used to carry out the process as showed in Fig. 2. The horizontal polarization for the first path, the vertical polarization for the second path, and the horizontal polarization for the third path are encoded to $E_0|\vec{e}_0\rangle, E_1|\vec{e}_1\rangle$ and $E_2|\vec{e}_2\rangle$, respectively, that is

$$|H_1\rangle \rightarrow E_0|\vec{e}_0\rangle, |V_2\rangle \rightarrow E_1|\vec{e}_1\rangle, |H_3\rangle \rightarrow E_2|\vec{e}_2\rangle. \quad (8)$$

where $|\vec{e}_0\rangle, |\vec{e}_1\rangle$ and $|\vec{e}_2\rangle$ are the basis vectors, E_0, E_1 and E_2 represent the amplitudes of classical optical fields for the different paths. With tuning the HWP1 and HWP2, the different amplitudes of classical optical fields in three paths can be obtained, thus the desired input state can be represented with these bases, namely, $|\chi\rangle = E_0|\vec{e}_0\rangle + E_1|\vec{e}_1\rangle + E_2|\vec{e}_2\rangle$. In our experiment, the HWP1 and HWP2 all are set to 45° , and the specific input state $|\chi\rangle = E_2|\vec{e}_2\rangle$ for the KCBS inequality is obtained.

In the KCBS experiment, these observables correspond to their respective operators. After the input state is projected onto the eigenstates of each operator, the measurement for the observable can be implemented. In the three-level system, the operator owns three different eigenstates and the corresponding eigenvalues. Here the input state is projected onto the eigenstate and then the probability of the eigenvalue is obtained. When the eigenvalue is multiplied by its probability, we can obtain the observable by summing the product. Nevertheless, the probability can be obtained by detecting the light intensity in our classical optical experiments (see Methods section for details).

For the KCBS inequality, we need to test the correlation $\overline{A_i A_{i+1}}$, and the observables A_i and A_{i+1} need to be tested simultaneously. Here the operator $A_i = \mathbb{I} - 2|a_i\rangle\langle a_i|$, and its eigenvalues are $+1, +1$ and -1 , which correspond to the values of A'_i in Eq.(1). The eigenvalues $+1$ and -1 for the operator $\mathbb{I} - 2|a_i\rangle\langle a_i|$ correspond to the eigenvalues 0 and 1 for $|a_i\rangle\langle a_i|$, and their eigenstates are also one-to-one correspondence. We adopt the scheme of joint measurement to test the contextuality inequality²². In order to measure the correlation $A_i A_{i+1}$, we need to know their respective eigenstates corresponding to eigenvalues $+1, +1$ and -1 . As A_i and A_{i+1} are orthogonal, the eigenstate of A_i corresponding to the eigenvalue -1 is just the eigenstate of A_{i+1} corresponding to the eigenvalue $+1$. In contrast, for the eigenstate corresponding to $A_{i+1} = -1$, it is just the eigenstate corresponding to $A_i = +1$. When the input state is projected onto the respective eigenstates, we can obtain the joint probabilities $P(A_i = -1, A_{i+1} = +1)$ and $P(A_i = +1, A_{i+1} = -1)$. At the same time, there is the third eigenstate that the eigenvalues all are $+1$ for A_i and A_{i+1} , and the probability is $P(A_i = +1, A_{i+1} = +1)$. However, the eigenstate is nonexistent when the corresponding eigenvalues for both A_i and A_{i+1} are taken as -1 , there is no the joint probability in such a case. With these probabilities, we can obtain $\overline{A_i A_{i+1}} = -P(A_i = -1, A_{i+1} = +1) - P(A_i = +1, A_{i+1} = -1) + P(A_i = +1, A_{i+1} = +1)$. Then, the contextuality can be calculated from Eq. (6).

The above measurement process can be completed by the experimental setup of measurement part in Fig. 2. For the measurement of the observable $A_i A_{i+1}$, because A_i and A_{i+1} possess a joint probability, they are required to measure simultaneously in the experiment. Due to the complexity of expressions for a_i , it is difficult to establish the eigenstates of A_i and A_{i+1} simultaneously. The experiment setting needs to be designed skillfully and the observable $A_i A_{i+1}$ needs to be measured carefully. In order to obtain the eigenstates of A_i and A_{i+1} simultaneously, some designs are done and several interferometers are cascaded to achieve the above purpose. Then the cetrit bases can map to these eigenstates at the three output ports as showed in Fig. 2. Using the HWPs (5, 7 and 8) and PBSs (4, 5 and 6), the polarization and path are combined to accomplish the construction. The expressions of the state vectors at the three output ports are showed in Eq. (11) (see Methods section).

With suitably setting up the angles of HWP5, HWP7 and HWP8, the eigenstates of these observables can be obtained at three output ports. They meet the requirements of joint measurement. Taking A_1 and A_2 for examples, the setting angles of HWP5, HWP7 and HWP8 are $117^\circ, 24^\circ$, and 32° , respectively. Hence the state vectors at output ports 1, 2 and 3 are $-0.6012|\vec{e}_0\rangle + 0.43682|\vec{e}_1\rangle + 0.66912|\vec{e}_2\rangle, 0.2288|\vec{e}_0\rangle - 0.70822|\vec{e}_1\rangle + 0.66792|\vec{e}_2\rangle, -0.7656|\vec{e}_0\rangle - 0.5547|\vec{e}_1\rangle - 0.3258|\vec{e}_2\rangle$, respectively. The output states at the port 1 describe the eigenstates with eigenvalues $A_1 = -1$ and $A_2 = +1$. The output states at the port 2 correspond to the eigenstates with $A_1 = +1$ and $A_2 = -1$. But the output states at the port 3 are the eigenstates with $A_1 = +1$ and $A_2 = +1$. The state vectors of output ports 1 and 2 correspond to the expressions in Eq. (9) (see Methods section), and they are the interrelated eigenstates of A_1 and A_2 . Because of the imperfect setting angles of HWPs, the expressions have a little deviation. With the input state being projected onto these eigenstates, the joint probabilities are obtained when we measure the light intensities at the three output ports. The light intensities are detected by three photodetectors (PD1, PD2 and PD3). The light intensity of each output port is normalized (divided by the total light intensities), and the probability of eigenvalue is obtained. The measurement probabilities for the output ports 1, 2 and 3 correspond to $P(A_1 = -1, A_2 = +1), P(A_1 = +1, A_2 = -1)$ and $P(A_1 = +1, A_2 = +1)$, respectively, thus the observable $A_1 A_2$ can be obtained.

The measurement methods for all five pairs of observables are the same. We tune the angles of the HWPs and obtain the desired eigenstates of the pairs of observables. The setting angles of the HWPs for the measurements of the different correlation pairs are given in Methods section. Then the input state is projected onto these eigenstates and the light intensities of the output ports are measured for every setting. The light intensity of each output port is normalized, therefore the joint probabilities of A_i and A_{i+1} are acquired, $P_{PD1} = I_1/(I_1 + I_2 + I_3)$, $P_{PD2} = I_2/(I_1 + I_2 + I_3)$ and $P_{PD3} = I_3/(I_1 + I_2 + I_3)$. And then using the relation: $\overline{A_i A_{i+1}} = -P(A_i = -1, A_{i+1} = +1) - P(A_i = +1, A_{i+1} = -1) + P(A_i = +1, A_{i+1} = +1)$, the five pairs of observables can be calculated. The probabilities and measurement values of $\overline{A_i A_{i+1}}$ are given in the following Table 1.

With summing the five sets of results, we can calculate the contextuality for Eq. (6). The result is -3.4196 ± 0.0057 and less than the minimum value -3 . Due to inaccuracy of the experiment measurements, the experiment results have some deviations from the theoretical values. However, they still show strong violation of the noncontextuality inequality. This means that the KCBS contextuality can be realized in the classical optical systems similar to the cases in the single photon systems.

Terms	PD1	PD2	PD3	Calculated result
$\overline{A_1 A_2}$	$P(A_1 = -1, A_2 = +1)$ 0.4308 ± 0.0046	$P(A_1 = +1, A_2 = -1)$ 0.4204 ± 0.0049	$P(A_1 = +1, A_2 = +1)$ 0.1488 ± 0.0012	-0.7024 ± 0.0024
$\overline{A_3 A_2}$	$P(A_3 = -1, A_2 = +1)$ 0.4313 ± 0.0039	$P(A_3 = +1, A_2 = -1)$ 0.3974 ± 0.0053	$P(A_3 = +1, A_2 = +1)$ 0.1713 ± 0.0023	-0.6574 ± 0.0046
$\overline{A_3 A_4}$	$P(A_3 = -1, A_4 = +1)$ 0.4313 ± 0.0026	$P(A_3 = +1, A_4 = -1)$ 0.4154 ± 0.0027	$P(A_3 = +1, A_4 = +1)$ 0.1533 ± 0.0036	-0.6935 ± 0.0071
$\overline{A_5 A_4}$	$P(A_5 = -1, A_4 = +1)$ 0.4297 ± 0.0029	$P(A_5 = +1, A_4 = -1)$ 0.4011 ± 0.0068	$P(A_5 = +1, A_4 = +1)$ 0.1691 ± 0.0042	-0.6618 ± 0.0084
$\overline{A_5 A_1}$	$P(A_5 = -1, A_1 = +1)$ 0.4313 ± 0.0045	$P(A_5 = +1, A_1 = -1)$ 0.4210 ± 0.0035	$P(A_5 = +1, A_1 = +1)$ 0.1477 ± 0.0012	-0.7045 ± 0.0024
				-3.4196 ± 0.0057

Table 1. The experimental probabilities and calculated results for the KCBS inequality.




Figure 3. Experimental setup for the test of the geometric form of the KCBS inequality. The establishment of the input state and the eigenstates, and the detection method are similar to the test of the KCBS inequality. The HWP3 and HWP4 are used for the path-length compensation similarly.

Experimental demonstration for the geometric form of the KCBS inequality in classical light systems.

For the geometric form of the KCBS inequality, the input state needs to be projected onto the eigenstates of the operator $|a_i\rangle\langle a_i|$. The operator owns eigenvalues 0, 0, +1 and the corresponding eigenstates. When the input state is projected onto the relevant eigenstates, the probability of the eigenvalue is obtained. Each eigenvalue is multiplied by its respective probability. Then with summing the three computation results, we got these observables. In such a case, the experimental setting shown in Fig. 2 can be simplified to Fig. 3 to measure the geometric form of the KCBS inequality.

In the state preparation stage as showed in Fig. 3, the different polarizations of the classical light fields in three paths indicate three-dimensional basis vectors. Just like the case in the Fig. 2, these bases construct the input state $|\chi\rangle = E_0|\vec{e}_0\rangle + E_1|\vec{e}_1\rangle + E_2|\vec{e}_2\rangle$. Likewise, the angles of the HWP1 and HWP2 all are set to 45° to obtain the particular input state $|\chi\rangle = E_2|\vec{e}_2\rangle$ for the geometric form of the KCBS inequality.

Similarly we need to construct these eigenstates of the operator $|a_i\rangle\langle a_i|$ and implement the projective measurements in the measurement stage. As showed in Fig. 3, with utilizing the appropriate settings for HWP5, HWP6 and the PBSs, the three eigenstates of $|a_i\rangle\langle a_i|$ can be constructed using the classical optical fields at the three output ports. The state vectors at the three output ports are expressed in Methods section. The output ports 1, 2 and 3 correspond to the eigenstates with eigenvalues 0, 0 and +1, respectively. Take $|a_1\rangle\langle a_1|$ as an example, the setting angles of HWP5 and HWP6 are 117° and 24° , respectively. We assume that the amplitudes of the three input bases all are unit amplitude, the light fields can be expressed as $-0.5878|\vec{e}_0\rangle - 0.8090|\vec{e}_1\rangle$, $0.5413|\vec{e}_0\rangle - 0.3933|\vec{e}_1\rangle + 0.7431|\vec{e}_2\rangle$, and $-0.6012|\vec{e}_0\rangle + 0.4368|\vec{e}_1\rangle + 0.6691|\vec{e}_2\rangle$ at the three output ports 1, 2 and 3, respectively. They correspond to the three eigenstates with eigenvalues 0, 0 and +1 of the operator $|a_1\rangle\langle a_1|$. In order to obtain the probabilities corresponding to eigenvalues 0, 0 and +1, we measure the light intensities at the respective output ports. After the normalization, we obtain the probabilities of eigenvalues, namely $P_{PD1} = I_1/(I_1 + I_2 + I_3)$, $P_{PD2} = I_2/(I_1 + I_2 + I_3)$ and $P_{PD3} = I_3/(I_1 + I_2 + I_3)$. With the probabilities we can calculate the value $|a_1|\chi|^2 = P_{PD3}$ and further obtain $\sum_{i=1}^5 |a_i|\chi|^2$. Of course, the light intensity may also be regarded as the modular square of the input state $|\chi\rangle$ being projected directly onto the direction vector a_i , which corresponds to the expression in Eq. (7). For the other terms, the methods are the same. The Eq. (7) can be calculated using these measurement results and then we can detect the violation of the geometric form of the KCBS inequality. The experimental results are shown in Table 2.

The upper bound of the inequality for Eq. (4) is 2 and the maximum theoretical prediction is 2.236. Our experiment result is 2.1880 ± 0.0064 , which shows obvious violation of the inequality. This means that the violation of the geometric form of the KCBS inequality can also be observed in classical light systems. The above experimental results of the KCBS inequality and its geometrical form have always some deviations with the theoretical

Terms	PD1 $P(a_i=0)$	PD2 $P(a_i=0)$	PD3 $P(a_i=1)$	Theoretical value
a_1	0.0381 ± 0.0004	0.5214 ± 0.0019	0.4405 ± 0.0021	0.4472
a_2	0.0402 ± 0.0003	0.5232 ± 0.0020	0.4366 ± 0.0019	0.4472
a_3	0.0388 ± 0.0003	0.5239 ± 0.0014	0.4373 ± 0.0014	0.4472
a_4	0.0401 ± 0.0007	0.5214 ± 0.0031	0.4386 ± 0.0034	0.4472
a_5	0.0395 ± 0.0003	0.5255 ± 0.0030	0.4350 ± 0.0029	0.4472
			2.1880 ± 0.0064	2.236

Table 2. The experimental results and the theoretical predictions of the observables for the geometric form of the KCBS inequality. Because the eigenvalues for the output port 1 and 2 all are 0, the result of the output port 3 is the value of the observable.

maximum violation. This is because there are some imprecise measurements and imperfect operations on optical elements in the experiments. For example, the PBS cannot transmit the horizontal polarization light and reflect the vertical polarization light absolutely. The mixed polarization lights always enter into the light path because of the inaccurate PBS. Although there are some imprecise measurements, the presented experiments still yield strong violation of the KCBS inequality and its geometrical form.

Conclusions

We have performed the experimental test for the KCBS inequality and its geometry form (or Wright's inequality) in the classical light systems. Using the polarization and path, the cetrir has been constructed, the projection measurement has been implemented, the clear violations of the KCBS inequality and its geometrical form have been observed. This indicates that the contextuality inequality, which is commonly used in test of the conflict between quantum theory and noncontextual realism, can be used in the classical optical coherence to describe correlation characteristics of the classical fields. In addition, recent investigations have shown that there is a remarkable equivalence between the onset of contextuality and the possibility of universal quantum computation³². Thus, our results also imply that the strong quantum computing power is likely to be simulated in the classical light systems.

Methods

The probability of eigenvalue is tested by measuring light intensity in classical optics systems. In the classical optical systems, the direction unit vector can be expressed as $|a_i\rangle = \alpha_i|\vec{e}_0\rangle + \beta_i|\vec{e}_1\rangle + \gamma_i|\vec{e}_2\rangle$, and in the general case the input state is $|\chi\rangle = E_0|\vec{e}_0\rangle + E_1|\vec{e}_1\rangle + E_2|\vec{e}_2\rangle$. In Eq. (2) the expression of the state vector is clear for the geometric comprehension of pentagram direction unit vector, but it is not direct for the algebra calculation. Thus, we transform it into the decimal forms:

$$\begin{aligned}
 |a_1\rangle &= -0.6014|\vec{e}_0\rangle + 0.4370|\vec{e}_1\rangle + 0.6687|\vec{e}_2\rangle \\
 |a_2\rangle &= 0.2297|\vec{e}_0\rangle - 0.7071|\vec{e}_1\rangle + 0.6687|\vec{e}_2\rangle \\
 |a_3\rangle &= 0.2297|\vec{e}_0\rangle + 0.7071|\vec{e}_1\rangle + 0.6687|\vec{e}_2\rangle \\
 |a_4\rangle &= -0.6014|\vec{e}_0\rangle - 0.4370|\vec{e}_1\rangle + 0.6687|\vec{e}_2\rangle \\
 a_5 &= |0.7435|\vec{e}_0\rangle + 0.6687|\vec{e}_2\rangle
 \end{aligned} \tag{9}$$

In the measurement stage, the input state needs to be projected onto the different eigenstates of the operator A_i , and then we can calculate the observables. Here the eigenstates are also expressed with the classical optical fields, where $|a_i\rangle$ is exactly the eigenstate corresponding to the eigenvalue -1 for the operator A_i ($A_i = \mathbb{I} - 2|a_i\rangle\langle a_i|$). The probability for the eigenvalue -1 is $|(a_i|\chi)|^2$ when the input state $|\chi\rangle$ is projected onto the eigenstate $|a_i\rangle$. The probability $|(a_i|\chi)|^2$ is the modular square of the amplitude of the optical field in the experiment, and the modular square of the amplitude is the light intensity I

$$|(a_i|\chi)|^2 = |\alpha_i^*E_0 + \beta_i^*E_1 + \gamma_i^*E_2|^2 = I. \tag{10}$$

Thus the probability of the eigenvalue can be expressed with the light intensity and we can detect the light intensity to obtain the probability of the eigenvalue in our experiment. The probabilities of two other eigenvalues can be measured similarly.

The expressions of state vectors at the three output ports and the setting angles of HWPs for the two experiments. For the KCBS inequality, we need to construct the interrelated eigenstates that satisfy the joint measurement scheme at the three output ports. The transformation matrix of the HWP is $\begin{pmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{pmatrix}$, where θ is the rotation angle of the HWP. We assume that all amplitudes for the three input bases are unit amplitudes, and at the three output ports the output expressions may be written as

	HWP5	HWP7	HWP8
$\overline{A_1A_2}$	117°	24°	32°
$\overline{A_3A_2}$	81°	24°	58°
$\overline{A_3A_4}$	81°	24°	122°
$\overline{A_5A_4}$	45°	24°	148°
$\overline{A_5A_1}$	45°	24°	32°

Table 3. The setting angles of the HWPs for the test of the KCBS inequality.

	a_1	a_2	a_3	a_4	a_5
HWP5	117°	9°	81°	−27°	45°
HWP6	24°	24°	24°	24°	24°

Table 4. The setting angles of HWP5 and HWP6 for the test of the geometric form of the KCBS inequality.

$$\begin{aligned}
 & \sin 2\theta_5 \sin 2\theta_7 |\vec{e}_0\rangle - \cos 2\theta_5 \sin 2\theta_7 |\vec{e}_1\rangle + \cos 2\theta_7 |\vec{e}_2\rangle \\
 & (\cos 2\theta_5 \cos 2\theta_8 - \sin 2\theta_5 \cos 2\theta_7 \sin 2\theta_8) |\vec{e}_0\rangle + \\
 & (\sin 2\theta_5 \cos 2\theta_8 + \cos 2\theta_5 \cos 2\theta_7 \sin 2\theta_8) |\vec{e}_1\rangle + \sin 2\theta_7 \sin 2\theta_8 |\vec{e}_2\rangle \\
 & (\cos 2\theta_5 \sin 2\theta_8 + \sin 2\theta_5 \cos 2\theta_7 \cos 2\theta_8) |\vec{e}_0\rangle + \\
 & (\sin 2\theta_5 \sin 2\theta_8 - \cos 2\theta_5 \cos 2\theta_7 \cos 2\theta_8) |\vec{e}_1\rangle - \sin 2\theta_7 \cos 2\theta_8 |\vec{e}_2\rangle
 \end{aligned} \quad (11)$$

In the Table 3, we display the rotation angles of HWPs for the measurements of five pairs of observables for the KCBS inequality. For the geometric form of the KCBS inequality, we need to construct the eigenstates of the operator $|a_i\rangle\langle a_i|$ with the combination of HWPs and PBSs. As showed in Fig. 3, at the three output ports 1, 2 and 3, the state vectors can be expressed as $\cos 2\theta_5 |\vec{e}_0\rangle + \sin 2\theta_5 |\vec{e}_1\rangle$, $-\sin 2\theta_5 \cos 2\theta_6 |\vec{e}_0\rangle + \cos 2\theta_5 \cos 2\theta_6 |\vec{e}_1\rangle + \sin 2\theta_6 |\vec{e}_2\rangle$ and $\sin 2\theta_5 \sin 2\theta_6 |\vec{e}_0\rangle - \cos 2\theta_5 \sin 2\theta_6 |\vec{e}_1\rangle + \cos 2\theta_6 |\vec{e}_2\rangle$, respectively. With suitably choosing the angles of HWP5 and HWP6, the desired eigenstates are obtained at the three output ports. In the following Table 4, it is the setting angles of HWP5 and HWP6 for the five measurements.

References

- Kochen, S. & Specker, E. P. The Problem of Hidden Variables in Quantum Mechanics. *J. Math. Mech.* **17**, 59 (1967).
- Peres, A. Two simple proofs of the Kochen–Specker theorem. *J. Phys. A: Math. Gen.* **24**, L175–L178 (1991).
- Cabello, A., Estebaran, J. M. & Garcia-Alcaine, G. Bell–Kochen–Specker theorem: A proof with 18 vectors. *Phys. Lett. A* **212**, 183–187 (1996).
- Pavicic, M., Merlet, J. P., McKay, B. & Megill, N. D. Kochen–Specker vectors. *J. Phys. A: Math. Gen.* **38**, 1577–1592 (2005).
- Cabello, A. How many questions do you need to prove that unasked questions have no answers? *Int. J. Quantum. Inform.* **4**, 55–61 (2006).
- Toh, S. P. & Zainuddin, H. Kochen–Specker theorem for a three-qubit system: A state-dependent proof with seventeen rays. *Phys. Lett. A* **374**, 4834–4837 (2010).
- Bengtsson, I., Blanchfield, K. & Cabello, A. A Kochen–Specker inequality from a SIC. *Phys. Lett. A* **376**, 374–376 (2012).
- Huang, Y.-F., Li, C.-F.; Zhang, Y.-S., Pan, J.-W. & Guo, G.-C. Experimental Test of the Kochen–Specker Theorem with Single Photons. *Phys. Rev. Lett.* **90**, 250401 (2003).
- Amselem, E., Radmark, M., Bourennane, M. & Cabello, A. State-Independent Quantum Contextuality with Single Photons. *Phys. Rev. Lett.* **103**, 160405 (2009).
- Amselem, E. *et al.* Experimental Fully Contextual Correlations. *Phys. Rev. Lett.* **108**, 200405 (2012).
- D'Ambrosio, V. *et al.* Experimental Implementation of a Kochen–Specker Set of Quantum Tests. *Phys. Rev. X* **3**, 011012 (2013).
- Hu, X.-M. *et al.* Experimental Test of Compatibility–Loop-hole-Free Contextuality with Spatially Separated Entangled Qutrits. *Phys. Rev. Lett.* **117**, 170403 (2016).
- Mazurek, M. D., Pusey, M. F., Kunjwal, R., Resch, K. J. & Spekkens, R. W. An experimental test of noncontextuality without unphysical idealizations. *Nat. Commun.* **7**, 11780 (2016).
- Bartosik, H. *et al.* Experimental Test of Quantum Contextuality in Neutron Interferometry. *Phys. Rev. Lett.* **103**, 040403 (2009).
- Hasegawa, Y., Loidl, R., Badurek, G., Baron, M. & Rauch, H. Quantum Contextuality in a Single-Neutron Optical Experiment. *Phys. Rev. Lett.* **97**, 230401 (2006).
- Kirchmair, G. *et al.* State-independent experimental test of quantum contextuality. *Nature* **460**, 494–497 (2009).
- Moussa, O., Ryan, C. A., Cory, D. G. & Laflamme, R. Testing Contextuality on Quantum Ensembles with One Clean Qubit. *Phys. Rev. Lett.* **104**, 160501 (2010).
- Dogra, S. & Dorai, K. Arvind. Experimental demonstration of quantum contextuality on an NMR qutrit. *Phys. Lett. A* **380**, 1941–1946 (2006).
- Yu S. & Oh, C. H. State-Independent Proof of Kochen–Specker Theorem with 13 Rays. *Phys. Rev. Lett.* **108**, 030402 (2012).
- Klyachko, A. A., Can, M. A., Binicioglu, S. & Shumovsky, A. S. Simple Test for Hidden Variables in Spin-1 Systems. *Phys. Rev. Lett.* **101**, 020403 (2008).
- Cabello, A. Simple Explanation of the Quantum Violation of a Fundamental Inequality. *Phys. Rev. Lett.* **110**, 060402 (2013).
- Zu, C. *et al.* State-Independent Experimental Test of Quantum Contextuality in an Indivisible System. *Phys. Rev. Lett.* **109**, 150401 (2012).
- Zhang, X. *et al.* State-Independent Experimental Test of Quantum Contextuality with a Single Trapped Ion. *Phys. Rev. Lett.* **110**, 070401 (2013).

24. Huang, Y.-F. *et al.* Experimental test of state-independent quantum contextuality of an indivisible quantum system. *Phys. Rev. A* **87**, 052133 (2013).
25. Lapkiewicz, R. *et al.* Experimental non-classicality of an indivisible quantum system. *Nature* **474**, 490 (2011).
26. Ahrens, J.; Amselem, E.; Cabello, A. & Bourennane, M. Two fundamental experimental tests of nonclassicality with qutrits. *Sci. Rep.* **3**, 2170 (2013).
27. Um, M. *et al.* Experimental Certification of Random Numbers via Quantum Contextuality. *Sci. Rep.* **3**, 1627 (2013).
28. Kong, X. *et al.* An experimental test of the non-classicality of quantum mechanics using an unmovable and indivisible system. *arXiv*: 1210.0961.
29. Shaham A. & Eisenberg, H. S. Effect of decoherence on the contextual and nonlocal properties of a biphoton. *Phys. Rev. A* **91**, 022123 (2015).
30. Kurzyński, P., Cabello, A. & Kaszlikowski, D. Fundamental Monogamy Relation between Contextuality and Nonlocality. *Phys. Rev. Lett.* **112**, 100401 (2014).
31. Zhan, X. *et al.* Realization of the Contextuality-Nonlocality Tradeoff with a Qubit-Qutrit Photon Pair. *Phys. Rev. Lett.* **116**, 090401 (2016).
32. Howard, M., Wallman, J., Veitch, V. & Emerson, J. Contextuality supplies the ‘magic’ for quantum computation. *Nature* **510**, 351–355 (2014).
33. Frustaglia, D. *et al.* Classical Physics and the Bounds of Quantum Correlations. *Phys. Rev. Lett.* **116**, 250404 (2016).
34. Töppel, F., Aiello, A., Marquardt, C., Giacobino, E. & Leuchs, G. Classical entanglement in polarization metrology. *New J. Phys.* **16**, 073019 (2014).
35. Ghose, P. & Mukherjee, A. Entanglement in classical optics. *Rev. Theor. Sci.* **2**, 274–288 (2014).
36. Aiello, A., Töppel, F., Marquardt, C., Giacobino, E. & Leuchs, G. Quantum-like nonseparable structures in optical beams. *New J. Phys.* **17**, 043024 (2015).
37. Goldin, M. A., Francisco, D. & Ledesma, S. Simulating Bell inequality violations with classical optics encoded qubits. *J. Opt. Soc. Am. B* **27**, 779 (2010).
38. Lee K. F. & Thomas, J. E. Experimental Simulation of Two-Particle Quantum Entanglement using Classical Fields. *Phys. Rev. Lett.* **88**, 097902 (2002).
39. Qian, X.-F., Little, B., Howell, J. C. & Eberly, J. H. Shifting the quantum-classical boundary: theory and experiment for statistically classical optical fields. *Optica* **2**, 611–615 (2015).
40. Sun, Y. *et al.* Non-local classical optical correlation and implementing analogy of quantum teleportation. *Sci. Rep.* **5**, 9175 (2015).
41. Song, X., Sun, Y., Li, P., Qin H. & Zhang, X. Bell’s measure and implementing quantum Fourier transform with orbital angular momentum of classical light. *Sci. Rep.* **5**, 14113 (2015).
42. Spreuw, R. J. C. A Classical Analogy of Entanglement. *Foundations of Physics* **28**, 361–374 (1998).
43. Spreuw, R. J. C. Classical wave-optics analogy of quantum-information processing. *Phys. Rev. A* **63**, 062302 (2001).

Acknowledgements

This work is supported by the National Natural Science Foundation of China (Grant No. 11574031 and 61421001).

Author Contributions

The experiments are performed by T.L., the corresponding theoretical method is presented by T.L. In doing the experiments, T.L. get the help of Q.Z. and X.S., the idea and physical analysis are given by X.Z. All authors reviewed the manuscript.

Additional Information

Competing Interests: The authors declare no competing financial interests.

How to cite this article: Li, T. *et al.* Experimental contextuality in classical light. *Sci. Rep.* **7**, 44467; doi: 10.1038/srep44467 (2017).

Publisher’s note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017