## SCIENTIFIC REPORTS

## **OPEN** Erratum: Antibodies from multiple sclerosis patients preferentially recognize hyperglucosylated adhesin of non-typeable Haemophilus influenzae

Marthe T. C. Walvoort, Chiara Testa, Raya Eilam, Rina Aharoni, Francesca Nuti, Giada Rossi, Feliciana Real-Fernandez, Roberta Lanzillo, Vincenzo Brescia Morra, Francesco Lolli, Paolo Rovero, Barbara Imperiali & Anna Maria Papini

Scientific Reports 6:39430; doi: 10.1038/srep39430; published online 23 December 2016; updated 22 March 2017

The original version of this Article contained typographical errors in the Abstract.

In autoimmune diseases, there have been proposals that exogenous "molecular triggers", i.e., specific this should be 'non-self antigens' accompanying infectious agents, might disrupt control of the adaptive immune system resulting in serious pathologies'.

now reads:

'In autoimmune diseases, there have been proposals that exogenous "molecular triggers", i.e., specific 'non-self antigens' accompanying infectious agents, might disrupt control of the adaptive immune system resulting in serious pathologies'.

'The etiology of the multiple sclerosis (MS) remains unclear'.

now reads:

'The etiology of multiple sclerosis (MS) remains unclear'.

These errors have been corrected in the HTML and PDF versions of this Article.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/

© The Author(s) 2017