

# SCIENTIFIC REPORTS

OPEN

## Linoleic acid isomerase gene *FgLAI12* affects sensitivity to salicylic acid, mycelial growth and virulence of *Fusarium graminearum*

Received: 14 July 2016

Accepted: 13 March 2017

Published: 07 April 2017

Ya-Zhou Zhang<sup>1,\*</sup>, Zhen-Zhen Wei<sup>1,\*</sup>, Cai-Hong Liu<sup>1,\*</sup>, Qing Chen<sup>1,\*</sup>, Bin-Jie Xu<sup>1,\*</sup>, Zhen-Ru Guo<sup>1</sup>, Yong-Li Cao<sup>1</sup>, Yan Wang<sup>1</sup>, Ya-Nan Han<sup>1</sup>, Chen Chen<sup>1</sup>, Xiang Feng<sup>1</sup>, Yuan-Yuan Qiao<sup>1</sup>, Lu-Juan Zong<sup>1</sup>, Ting Zheng<sup>1</sup>, Mei Deng<sup>1</sup>, Qian-Tao Jiang<sup>1</sup>, Wei Li<sup>2</sup>, You-Liang Zheng<sup>1</sup>, Yu-Ming Wei<sup>1</sup> & Peng-Fei Qi<sup>1</sup>


*Fusarium graminearum* is the major causal agent of fusarium head blight in wheat, a serious disease worldwide. Linoleic acid isomerase (LAI) catalyses the transformation of linoleic acid (LA) to conjugated linoleic acid (CLA), which is beneficial for human health. We characterised a *cis*-12 LAI gene of *F. graminearum* (FGSG\_02668; *FgLAI12*), which was downregulated by salicylic acid (SA), a plant defence hormone. Disruption of *FgLAI12* in *F. graminearum* resulted in decreased accumulation of *cis*-9, *trans*-11 CLA, enhanced sensitivity to SA, and increased accumulation of LA and SA in wheat spikes during infection. In addition, mycelial growth, accumulation of deoxynivalenol, and pathogenicity in wheat spikes were reduced. Re-introduction of a functional *FgLAI12* gene into  $\Delta FgLAI12$  recovered the wild-type phenotype. Fluorescent microscopic analysis showed that *FgLAI12* protein was usually expressed in the septa zone of conidia and the vacuole of hyphae, but was expressed in the cell membrane of hyphae in response to exogenous LA, which may be an element of LA metabolism during infection by *F. graminearum*. The *cis*-12 LAI enzyme encoded by *FgLAI12* is critical for fungal response to SA, mycelial growth and virulence in wheat. The gene *FgLAI12* is potentially valuable for biotechnological synthesis of *cis*-9, *trans*-11 CLA.

*Fusarium graminearum* (teleomorph *Gibberella zeae* [Schwein.] Petch) is an ascomycete fungus that causes fusarium head blight (FHB) in wheat. FHB can lead to yield loss and accumulation of trichothecene mycotoxins (predominantly deoxynivalenol [DON]) in wheat seeds, which threatens human and animal health<sup>1–3</sup>. Control of FHB by using natural resistance and traditional breeding in wheat is difficult, and thus FHB remains a serious disease worldwide<sup>4,5</sup>.

Linoleic acid (LA; *cis*-9, *cis*-12 C18:2) plays an important role in wheat resistance to *F. graminearum* infection. Accumulation of LA is greater in FHB-resistant wheat lines than in susceptible lines<sup>6</sup>, suggesting that LA may contribute to FHB resistance. Enhanced accumulation of LA and other free fatty acids can reinforce the cuticle, which acts as a barrier to pathogen entry<sup>7</sup>. LA is metabolised by linoleate diol synthase (LDS)<sup>8,9</sup> and linoleic acid isomerase (LAI) in fungi (Fig. 1). LDS has been well studied in fungi, and is involved in fungal development, reproduction, synthesis of trichothecenes and cytochrome metabolism<sup>8,9,12,13</sup>. However, limited knowledge on LAI in fungi is available.

Activity of LAI was first analysed in the bacterium *Butyrivibrio fibrisolvens*<sup>14</sup> and subsequently in the red alga *Ptilota flicina*<sup>15</sup>. To date, the enzyme has been detected mostly frequently in bacteria<sup>16,17</sup>. LAI catalyses the transformation of LA to conjugated linoleic acid (CLA) in microbes. Two categories of LAI (i.e., *cis*-9 and *cis*-12) have been identified: *cis*-9 LAI catalyses conversion of LA to *trans*-10, *cis*-12 CLA (*trans*-10, *cis*-12 C18:2); and *cis*-12 LAI transforms LA to *cis*-9, *trans*-11 CLA (*cis*-9, *trans*-11 C18:2). *Trans*-10, *cis*-12 CLA and *cis*-9, *trans*-11 CLA are

<sup>1</sup>Triticaceae Research Institute, Sichuan Agricultural University, Chengdu, Sichuan 611130, China. <sup>2</sup>Agronomy College, Sichuan Agricultural University, Chengdu, Sichuan 611130, China. \*These authors contributed equally to this work. Correspondence and requests for materials should be addressed to Y.-M.W. (email: ymwei@sicau.edu.cn) or P.-F.Q. (email: pengfeiqi@hotmail.com)



**Figure 1.** Schematic illustration of LA isomers. FgLAI12 is marked by a black box. PAI, *Propionibacterium acnes* cytosolic LA isomerase<sup>10,11</sup>.

then transformed into *trans-10* C18:1 and *trans-11* C18:1, respectively, by unknown enzymes (Fig. 1). In fungi, the gene encoding *cis-9* LAI (PAI) was identified in *Propionibacterium acnes*<sup>10,11,18</sup>. Activity of *cis-12* LAI was measured in *Clostridium sporogenes*; however, no *cis-12* LAI gene has yet been reported<sup>19</sup>.

In this study, we characterised a *cis-12* LAI gene of *F. graminearum* (FGSG\_02668; FgLAI12; *cis-12* LAI of *F. graminearum*), which was downregulated by salicylic acid (SA), a plant defence hormone<sup>20</sup>. The objectives were to investigate the function of FgLAI12 in LA metabolism and to clarify the gene's effects on fungal response to SA, fungal development, synthesis of DON and pathogenicity in wheat.

## Results

**Deletion and complementation of FgLAI12.** The FgLAI12 gene contains three exons and two introns (Fig. 2A). To determine the function of FgLAI12 in *F. graminearum*, knockout mutants ( $\Delta$ FgLAI12) and complementation mutants (C-FgLAI12) were created (Fig. 2A,B). The mutants were verified by PCR (Fig. 2C,D) and sequencing (data not shown).


The copy number of integrated T-DNA inserts was determined by targeting hygromycin B phosphotransferase genes (HPH). As expected, HPH was not detected in the wild type (WT). One and two normalised copies of HPH were detected in the  $\Delta$ FgLAI12 and C-FgLAI12 mutants, respectively, confirming the single insertion of the T-DNA construct for gene knockout and complementation (Fig. 2E).

Expression of FgLAI12 under treatment with 3 mmol L<sup>-1</sup> SA or 10 mg mL<sup>-1</sup> LA was determined by reverse transcription-PCR (RT-PCR). As expected, FgLAI12 expression was downregulated by SA and upregulated by LA (Fig. 2F).


**FgLAI12 catalyses conversion of LA to *cis-9,trans-11* CLA.** Three peaks were detected in gas chromatography-mass spectrometry (GC-MS) analysis of compounds derived from LA (Fig. 3A), namely LA (the first peak), stearic acid (the second peak) and *cis-9,trans-11* CLA (the third peak), by matching with GC-MS spectrometry gallery. The peaks for LA, stearic acid and *cis-9,trans-11* CLA in GC-MS were confirmed by using of corresponding standard chemicals (Fig. 3B). In the presence of exogenous LA, enzyme extracts from mycelia of WT and C-FgLAI12 produced stearic acid and *cis-9,trans-11* CLA, compared with no *cis-9,trans-11* CLA and a small quantity of stearic acid for the extract from  $\Delta$ FgLAI12. After addition of *cis-9,trans-11* CLA, the enzyme extract from  $\Delta$ FgLAI12 produced stearic acid as for the WT. FgLAI12 contains no known conserved domain of transcription factors, further suggesting that FgLAI12 is directly responsible for conversion of LA to *cis-9,trans-11* CLA. These results showed that FgLAI12 encoded a LAI enzyme that catalysed the transformation of LA to *cis-9,trans-11* CLA (Fig. 1), and that FgLAI12 played a major role in the isomerism of LA in *F. graminearum* under the experimental conditions.

SA inhibited the expression of FgLAI12<sup>20</sup> (Fig. 2F). To further confirm the function of FgLAI12 and clarify the relationship between SA and FgLAI12, 3 mL mSNA-2 liquid medium was supplemented with 10 mg SA and cultures of WT,  $\Delta$ FgLAI12 and C-FgLAI12 were incubated for 24 h before collection of mycelia. Compared with cultures that lacked SA, the enzyme extracts from WT and C-FgLAI12 metabolised a smaller quantity of LA in the presence of SA (Fig. 3C). As expected, following addition of 10 mg SA to the enzyme extract, no significant difference in LA concentration was observed in the presence and absence of SA (Fig. 3D). These results indicated that FgLAI12 was a target for manipulation of LA metabolism by SA in *F. graminearum*.

**Influence of FgLAI12 on mycelial growth and conidial morphology.** To observe the changes in the growth phenotype caused by disruption of FgLAI12 in *F. graminearum*, the WT,  $\Delta$ FgLAI12 and C-FgLAI12



**Figure 2. Deletion and complementation of *FgLAI12* in *F. graminearum*.** (A) The left border (LB) and right border (RB) were amplified from the wild type (a).  $\Delta FgLAI12$  mutant (d) was created by recombination between the knockout vector (b) and *FgLAI12* (c). (B) The nucleotide sequence of *FgLAI12* was cloned (a), and ligated into the complementation vector (b). The T-DNA region of the complementation vector was recombined into  $\Delta FgLAI12$  to create C-*FgLAI12* (c). *SacI*, *Apal*, *SpeI*, *HindIII*, *NcoI* and *SpeI* indicate the restriction enzymes used. (C) PCR verification of  $\Delta FgLAI12$  using the primer pairs P5 forward + P5 reverse and P7 forward + P7 reverse. (D) Verification of expression of *FgLAI12* using the primer pair RJ-*FgLAI12* forward + RJ-*FgLAI12* reverse. *FgGAPDH* were used as a control. (E) Copy numbers of *HPH* determined by qPCR. Different letters (a and b) above each column indicate a significant difference ( $P < 0.05$ ;  $n = 3$ ). (F) Effect of SA and LA on expression of *FgLAI12* in hyphae, as determined using the primer pair RJ-*FgLAI12* (Table 1). Hyphae were collected on the 4<sup>th</sup> day after adding SA and LA. *FgGAPDH* was used as a reference. “F”, forward; “R”, reverse.



**Figure 3.** *FgLA112* encodes a *cis*-12 LA isomerase. (A) GC-MS analysis of chemicals derived from LA. (B) The peaks for LA, stearic acid and *cis*-9,*trans*-11 CLA when using standard chemicals. (C) Quantitative analysis of LA added to liquid medium containing growing mycelia. (D) Quantitative analysis of LA added to enzyme extract from mycelia. The same amount of mycelia was used. Different letters above each column indicate a significant difference ( $P < 0.05$ ;  $n = 3$ ). “a<sub>1</sub>, b<sub>1</sub>” and “a<sub>2</sub>, b<sub>2</sub>” are used to show the significance within WT and C-FgLA112 treatments, respectively, since there are more than one treatments in the same chart.

strains were inoculated on modified Synthetischer Nährstoffarmer Agar-1 (mSNA-1) (Fig. 4A) and potato dextrose agar (PDA) plates (Fig. 4B). On mSNA-1 plates,  $\Delta FgLA112$  formed much less aerial mycelium compared with WT and C-FgLA112. On the 6<sup>th</sup> day after inoculation, aerial mycelium of  $\Delta FgLA112$  was clearly observed (control in Fig. 4A). Similarly,  $\Delta FgLA112$  mycelia grew more slowly than those of WT and C-FgLA112 on PDA plates, as indicated on the 4<sup>th</sup> day after inoculation (control in Fig. 4B). Compared with the control, the spread of mycelia was inhibited by LA (LA treatment in Fig. 4A,B). Consistent with the role of *FgLA112* in LA metabolism, the spread and density of  $\Delta FgLA112$  mycelia were more strongly inhibited by LA. Mycelial growth of  $\Delta FgLA112$  was more sensitive to SA compared with that of WT and C-FgLA112 (SA treatment in Fig. 4A,B). These results indicated that *FgLA112* plays important roles in mycelial growth and fungal response to SA.


To clarify the influence of *FgLA112* on fungal growth within agar, a 0.5-cm<sup>2</sup> agar plug that excluded aerial mycelia was cut from the edge of 6-d-old mSNA-1 plates, and was observed using a stereomicroscope at 100 $\times$  (Fig. 4C). The hyphae of  $\Delta FgLA112$  grew more deeply on mSNA-1 plates than hyphae of WT and C-FgLA112. In the presence of LA or SA, the hyphae of WT,  $\Delta FgLA112$  and C-FgLA112 grew more shallowly than those of the controls. Interestingly, the hyphae of  $\Delta FgLA112$  grew more shallowly under LA treatment than those of WT and C-FgLA112. No notable difference between the strains under SA treatment.

To clarify whether *FgLA112* affected conidial morphology, we compared the length, width and number of septa of  $1 \times 10^3$  randomly selected conidia. No significant difference among WT,  $\Delta FgLA112$  and C-FgLA112 was observed for each parameter (Fig. 4D).

**Subcellular localisation of FgLA112 protein.** The C-FgLA112 strain harbouring a green fluorescent protein gene (GFP) tag was used to examine the subcellular localisation of FgLA112 protein. Conidia and hyphae at different vegetative growth stages were used to detect GFP fluorescence. As expected, no morphological differences among conidia of the WT,  $\Delta FgLA112$  and C-FgLA112 strains were observed (Fig. 5A). FgLA112 protein was predominantly localised on septa of conidia (Fig. 5A), and in vacuoles within hyphae (Fig. 5B). In response to LA treatment, FgLA112 protein was mainly localised on the cell membrane (Fig. 5C). In the presence of exogenous SA, FgLA112 protein remained localised in vacuoles but the fluorescence intensity was reduced, reflecting the inhibitory effect of SA on *FgLA112* expression (Fig. 5C).


***FgLA112* affects pathogenicity in wheat.** To clarify whether *FgLA112* was involved in the pathogenicity of *F. graminearum*, we point-inoculated two fully developed florets of a central spikelet within a spike with






**Figure 4. Effect of *FgLAII2* on fungal biology.** Mycelial growth of the WT,  $\Delta FgLAII2$  and C-*FgLAII2* on mSNA-1 (A) and PDA plates (B) under treatment with LA and SA on the 4<sup>th</sup>, 6<sup>th</sup> and 8<sup>th</sup> days after inoculation. (C) Microscopic observation of mycelia growth within mSNA-1 plates supplemented with LA and SA. The depth of hyphae is marked with green lines. Agar plugs (0.5 cm<sup>2</sup>) cut from the edge of growing mycelia (Fig. 4A) were photographed with an Olympus SZ51 (Japan). (D) Comparison of conidial length, width and the number of septa of the WT,  $\Delta FgLAII2$  and C-*FgLAII2*. Different letters above each column indicate a significant difference ( $P < 0.05$ ;  $n = 1000$ ). The same rule as mentioned in the legend of Fig. 3 is used for indicating the significance within treatment, when there are more than one treatments in the same chart.

conidial suspensions of WT,  $\Delta FgLAII2$  and C-*FgLAII2*.  $\Delta FgLAII2$  spikes showed much lower severity of disease compared with those of WT and C-*FgLAII2* (Fig. 6A,B). *FgLAII2* was associated with variation in fungal biomass in wheat spikes, which was indicated by the relative expression level of the *F. graminearum* *GAPDH* gene. The spikes inoculated with  $\Delta FgLAII2$  conidia showed a significantly lower *GAPDH* expression level than spikes



**Figure 5. Subcellular localisation of FgLAI12 protein.** Conidia (A) and mycelia (B and C) of the WT,  $\Delta FgLAI12$  and C-FgLAI12 strains photographed with a Nikon-80i fluorescence microscope (Japan). The fluorescent protein signal is marked with red arrows. (E), Optical microscope; (F), fluorescence microscope. Scale bar, 10  $\mu\text{m}$ .





**Figure 6.** Effect of *FgLAI12* on pathogenicity of *F. graminearum* in wheat. (A) Symptoms of Fusarium head blight in spikes of the WT,  $\Delta FgLAI12$  and C- $FgLAI12$  strains on the 12<sup>th</sup> day after inoculation. (B) Numbers of infected and bleached spikelets on the 4<sup>th</sup>, 8<sup>th</sup> and 12<sup>th</sup> days after inoculation. (C) Relative expression level of *FgGAPDH* in wheat spikes. (D) Concentrations of salicylic acid (SA), jasmonic acid (JA) and linoleic acid (LA) in spikes inoculated with WT and  $\Delta FgLAI12$ . (E) Concentration of deoxynivalenol (DON) in wheat spikes. (F) Measurement of DON in liquid medium (the same amount of mycelia was used). Values are the mean  $\pm$  standard deviation of three biological replicates per treatment. Different letters above each column indicate a significant difference ( $P < 0.05$ ;  $n = 3$ ). FW, fresh weight. The same rule as mentioned in the legend of Fig. 3 is used for indicating the significance within treatment, when there are more than one treatments in the same chart.

inoculated with WT conidia (Fig. 6C). In addition, the concentration of DON in the liquid culture medium and in wheat spikes was compared.  $\Delta FgLAI12$  showed much lower DON production than that of WT (Fig. 6E,F).

Both SA, jasmonic acid (JA) and LA are important components in the wheat defence response against *F. graminearum*, and infection of spikes by *F. graminearum* leads to a significant increase in their accumulation<sup>6,21</sup>. To further understand the reduced disease severity of  $\Delta FgLAI12$  spikes compared with WT spikes, we compared the amount of SA, JA and LA in spikes inoculated with WT and  $\Delta FgLAI12$  conidia. Spikes infected with  $\Delta FgLAI12$  accumulated higher concentrations of SA and LA, and a lower concentration of JA (Fig. 6D).

## Discussion

*Fusarium graminearum* is a fungal pathogen that causes severe damage to wheat crops worldwide. The genetics and biology of *F. graminearum* have been well studied. However, the metabolism of LA and its role in fungal biology and pathogenicity has attracted little attention. The *FgLAI12* gene of *F. graminearum* encodes a *cis*-12 LAI. Our results demonstrated that this *cis*-12 LAI was responsible for transformation of LA to *cis*-9,*trans*-11 CLA. In addition, *FgLAI12* was necessary for fungal growth, fungal response to SA, aerial mycelium development, synthesis of a mycotoxin and pathogenicity in wheat.

Both LDS<sup>8,9</sup> and LAI metabolise LA in fungi (Fig. 1). We previously reported that SA downregulates expression of *FgLAI12*, which encodes a predicted LDS enzyme<sup>20</sup>. Surprisingly, GC-MS analysis demonstrated that *FgLAI12* functioned as a LAI *in vivo*. Deletion of *FgLAI12* had no significant effect on conidial production (data

not shown) and morphology (Fig. 4D), even though *FgLAI12* played a major role in the isomerism of LA in *F. graminearum* (Fig. 3A). Disruption of LDS expression in *Emerella nidulans* and *Neosartorya fumigata* results in a decrease in ascospore production<sup>8</sup>, because one product of LDS activity, (8R)-hydroperoxylinoleate, is a sexual sporulation hormone<sup>22</sup>. These results indicate the functional difference of LDS and LAI in fungal biology. Additional studies of LDS genes would help to understand the role of LA in *F. graminearum*.

LA plays an important role in wheat and barley in defence against *F. graminearum* and has been established to be a FHB resistance-related metabolite<sup>6,7,23</sup>. Higher concentrations of LA accumulate in FHB-resistant wheat lines than in susceptible lines<sup>7</sup>, which suggests that LA contributes to FHB resistance. LA is a precursor in the biosynthesis of the phytohormone JA, which is a critical signalling molecule for wheat resistance against FHB<sup>21,24</sup>. Spikes inoculated with  $\Delta FgLAI12$  conidia accumulated greater quantities of LA than those inoculated with WT conidia (Fig. 6D), which was consistent with the hypothesised function of *FgLAI12*. Unexpectedly, spikes inoculated with  $\Delta FgLAI12$  accumulated lower concentrations of JA, possibly because of the lower fungal biomass (Fig. 6C). Besides being a chemical inhibitor, LA can reinforce the cuticle, which acts as a barrier to pathogen entry<sup>7</sup>. In this regard, the lower disease severity of  $\Delta FgLAI12$ -inoculated spikes compared with spikes inoculated with WT is understandable.

Fluorescent microscopic observation showed that *FgLAI12* protein was usually localised in the septa of conidia and the vacuoles of hyphae. However, exogenous LA treatment resulted in localisation of *FgLAI12* in the cell membrane of hyphae, which might reflect the mechanism by which *FgLAI12* protein metabolises LA synthesised by the host plant.

SA and JA are important defence phytohormones, for which both antagonistic and synergistic interactions are reported<sup>25–28</sup>. Both SA and JA accumulate in wheat spikes infected by *F. graminearum*, which indicates that synergistic interactions between the two phytohormones may contribute to FHB resistance<sup>21</sup>. Expression of *FgLAI12* was downregulated by SA<sup>20</sup>, and degradation of LA was reduced in response to exogenous SA (Fig. 3C). Spikes inoculated with  $\Delta FgLAI12$  accumulated higher quantities of SA and reduced amounts of JA compared with spikes inoculated with the WT strain (Fig. 6D), indicating that *FgLAI12* plays a role in manipulating biosynthesis of SA and JA *in planta*. Therefore, we speculate that *FgLAI12* represents a link between the SA and JA defence-response pathways during the host–pathogen interaction, which warrants future investigation.

Mycelial growth of  $\Delta FgLAI12$  was reduced on mSNA-1 and PDA plates compared with that of WT (Fig. 4A). This finding suggests that *FgLAI12* is essential for normal growth of *F. graminearum*. Polyunsaturated fatty acids (PUFAs), such as  $\gamma$ -linolenic, arachidonic and eicosapentaenoic acids, can modify membrane-bound proteins, ATPase and the histocompatibility complex, and control the reaction of fatty acid binding proteins<sup>29,30</sup>. As a PUFA, LA may perform a similar function to other PUFAs. Disruption of *FgLAI12* is likely to have disturbed the balance of PUFAs and negatively affected growth of fungal mycelia.

The mycotoxin DON is a threat to human and animal health<sup>1</sup>. Many countries have adopted a maximum allowable level for DON in grains for human consumption and animal feedstuffs. We previously showed that SA significantly reduced DON production<sup>20</sup> and SA also significantly inhibited expression of *FgLAI12*. In the present study, we demonstrated that disruption of *FgLAI12* reduced accumulation of DON in wheat spikes and in the culture medium. Therefore, *FgLAI12* may be a regulatory target for SA to downregulate DON production in *F. graminearum*.

*FgLAI12* catalyses the transformation of LA to *cis*-9,*trans*-11 CLA. CLA has been shown to lower cancer risk, enhance immunity and reduce body fat while increasing lean body mass in animals<sup>31,32</sup>. The *cis*-9,*trans*-11 CLA and *trans*-10,*cis*-12 CLA isomers show the highest biological activities and are transformed by *cis*-12 and *cis*-9 isomerases, respectively (Fig. 1). CLA is currently marketed as a dietary supplement, which is produced by alkaline isomerization of LA<sup>33</sup> or vegetable oils containing triglyceride esters of LA<sup>34</sup>. However, chemically synthesised CLA contains a variety of other isomers and is unsuitable for nutritional purposes<sup>35</sup>. Industrial-scale production of CLA by biotransformation from LA is increasingly attractive. Therefore, identification of the genes encoding LAI enzymes is warranted. The *cis*-12 LAI gene characterised in this study is potentially valuable for commercial use.

## Methods

**Experimental materials and growth conditions.** Plants of *Triticum aestivum* ‘Roblin’ were grown in greenhouses under 16/8 h (day/night) cycles at 23/18 °C. Plants were watered as needed and fertilised before planting with 15-15-15 (N-P-K) fertiliser. ‘Roblin’ is highly susceptible to *F. graminearum* infection.

The *F. graminearum* isolate DAOM180378 (Canadian Fungal Culture Collection, AAFC, Ottawa, ON, Canada), which is highly virulent on wheat, was used in all experiments. This isolate was also used for measurement of the expression level of FGSG\_02668 (*FgLAI12*) under SA treatment<sup>20</sup>. The isolate was cultured on modified mSNA-1 (1 g KH<sub>2</sub>PO<sub>4</sub>, 1 g KNO<sub>3</sub>, 0.5 g MgSO<sub>4</sub>, 0.5 g KCl, 1 g glucose, 1 g sucrose and 20 g agar per litre) plates at 25 °C. Conidia were produced in carboxymethyl cellulose liquid medium at 28 °C, with shaking (180 rpm) for 5 d<sup>36</sup>. For mycelial growth studies, PDA (BD Difco, Sparks, MD, USA) and mSNA-1 plates were used. LA and SA were added to media after autoclaving. Each plate was inoculated with  $1 \times 10^3$  conidia of *F. graminearum* and all inoculated plates were incubated in a dark cabinet at 28 °C. There were 10 replicates per treatment and the growth experiments were repeated three times. For deletion and complementation of the FGSG\_02668 gene, *Agrobacterium tumefaciens* strain AGL-1, which was used for transformation of *F. graminearum*, was grown at 28 °C in yeast extract broth (5 g nutrient broth, 1 g yeast extract, 5 g peptone, 5 g sucrose and 0.5 g MgSO<sub>4</sub> per litre; pH 7.4). *Escherichia coli* strain DH5 $\alpha$  (Tiagen, Beijing, China) was cultured in Luria–Bertani broth at 37 °C<sup>37</sup>. Unless specifically noted, all chemicals were purchased from Sigma-Aldrich (St Louis, MO, USA).



Primer	Forward (5'–3')	Reverse (5'–3')
FgLAI12-LB	GGAAGCTTCAAAGCGCAACAC	GGACTAGTTTCCCATCAACAACATAT
FgLAI12-RB	GCGGGCCCGGCTTATCTCAACGACA	GCGAGCTCTACCTGAATCCAACAT
P1 <sup>45</sup>	CTTTTCTCTTAGGTTTACCCG	TAATGCAGGAGTCGCATAAG
P3 <sup>45</sup>	CCCAAAAAGTGCTCCTTCAA	TGTGCTGCAAGGCGATTAA
P5 <sup>45</sup>	TTGTCACCGCCAAAGC	ACCTACTACTGGGCTGCTT
P7 <sup>45</sup>	CTGACCAGTTGCCTAA	ACAGTTCATTCCGAGAC
R-FgLAI12	GGACTAGTTGCTCAACTTACGCCACTG	GGCAGCTGTCGACGCAGAAGCT
RJ-FgLAI12	ATCCCAAGCCACCAG	GCCAAGTTCTCCGTC
Fg-GAPDH <sup>20</sup>	TGACTTGACTGTTCGCCTCGAGAA	ATGGAGGAGTTGGTGTTCGCGTTA
w-GAPDH <sup>20</sup>	AACTGTTTCAATGCCATCACTGCCAC	AGGACATACCAGTGAGCTTGCCAT
Aox <sup>20</sup>	GACTTGTCATGGTAGATGCCTG	CAGGACGAGCATAACCATTCTC
hn-RNP-Q <sup>20</sup>	TCACCTTCGCCAAGCTCAGAACTA	AGTTGAACCTTGCCCGAAACATGCC
Fg- $\beta$ -tubulin <sup>20</sup>	GTTGATCTCCAAGATCCGTG	CATGCAATATGCTAGAGGG
Fg-Factor1 <sup>20</sup>	CCTCCAGGATGTCTACAAGA	CTCAACGGACTTGACTTCAG
HPH	CGATCTTAGCCAGACGAGCG	TTGCCCTCGGACGAGTGCTG

**Table 1. Primers used in this study.** Horizontal lines under nucleotides indicate the cutting sites for restriction enzymes.

**Sequence analysis and primer design.** The nucleotide sequence of *FGSG\_02668* was downloaded from the Ensembl Fungi database (<http://fungi.ensembl.org/index.html>). Primer Premier 5.0 software (Premier Biosoft, Palo Alto, Canada) was used to design PCR primers (Table 1).

**Disruption and complementation of *FGSG\_02668*.** Genomic DNA of *F. graminearum* was extracted from mycelia, which were cultured on mSNA-1 plates for 5 d at 25 °C, by the CTAB method<sup>38</sup>. Disruption of *FGSG\_02668* in *F. graminearum* is schematically shown in Fig. 2A. The pRF-HU2 vector was used for targeted gene replacement in *F. graminearum* through *A. tumefaciens*-mediated transformation<sup>39,40</sup>. The left border (LB) and right border (RB) of *FGSG\_02668* targeted for homologous recombination were amplified with the primer pairs FgLAI12-LB forward + FgLAI12-LB reverse and FgLAI12-RB forward + FgLAI12-RB reverse, respectively. The resulting recombinant plasmids were verified by PCR and sequencing (Invitrogen, Shanghai, China) with the primer pairs P1 forward + P1 reverse and P3 forward + P3 reverse (Fig. 2A). *Agrobacterium*-mediated transformation of *F. graminearum* was performed as described previously<sup>41</sup>. The resulting mutants were verified by PCR and sequencing (Invitrogen) (Fig. 2A,C). The full open reading frame of *FGSG\_02668* was inserted into the pCambia1302 vector. The 3' terminus of *FGSG\_02668* was fused with a GFP tag under the control of the 35S promoter. The recombinant pCambia1302 vector was transformed into  $\Delta$ FgLAI12 to create complementation mutants. The complementation mutants were verified by PCR and sequencing (Fig. 2B,D). Five knockout mutants and four complementation mutants were used throughout the research.

To confirm the single integration of the *FGSG\_02668* knockout construct into the genome, quantitative real-time PCR (qPCR) was performed to determine the copy number of T-DNA inserts in fungal transformants as described previously<sup>42</sup>, with some modification. Genomic DNA was used as the qPCR template. The *HPH* gene was used to determine the copy number of the integrated T-DNA construct for gene knockout. The primer pairs Fg-GAPDH forward + Fg-GAPDH reverse, Fg- $\beta$ -tubulin forward + Fg- $\beta$ -tubulin reverse, and Fg-Factor1 forward + Fg-Factor1 forward, which target the single-copy house-keeping gene *FgGAPDH* (*FGSG\_06257*),  $\beta$ -tubulin (*FGSG\_09530*) and elongation factor 1 (*FGSG\_08811*) of *F. graminearum*, respectively, were used as references for normalisation of data. The primer pair HPH-F + HPH-R was used to amplify the *HPH* gene in qPCR reactions. The qPCR analyses were carried out as described previously<sup>43</sup> using a MyiQ Real-Time PCR Detection System (Bio-Rad, Hercules, CA, USA).

**RT-PCR analyses.** Total RNA was extracted from 100 mg (fresh weight) of mycelia grown on mSNA-1 plates using the E.Z.N.A.® Total RNA Kit I (Omega Bio-Tek, Norcross, GA, USA) in accordance with the manufacturer's instructions. RNA was reverse transcribed using the PrimeScript™ RT Reagent Kit with genomic DNA Eraser (Takara, Dalian, China) following the protocol of the manufacturer. The primer pair RJ-FgLAI12 forward + RJ-FgLAI12 reverse was used to measure the expression level of *FGSG\_02668* by RT-PCR. The *FgGAPDH* gene was used as a reference.

**LAI enzyme activity assay.** The crude enzyme was extracted as described previously<sup>13</sup>, which was utilised for extracting recombinant LDS enzyme. Mycelia of the WT,  $\Delta$ FgLAI12 and C-FgLAI12 strains were cultured and collected on liquid mSNA-2 (1 g KH<sub>2</sub>PO<sub>4</sub>, 1 g KNO<sub>3</sub>, 0.5 g MgSO<sub>4</sub>, 0.5 g KCl, 6 g glucose and 6 g sucrose per litre) medium, and incubated at 28 °C on an orbital shaker for 6 d in darkness at 120 rpm. The collected mycelia (100 mg fresh weight) were supplemented with 0.1 mol L<sup>-1</sup> sodium borate and 10 mg LA, then incubated with shaking (90 rpm) for 4 h at 28 °C. Subsequently, mycelia were collected by filtration and dried with filter paper, before being ground in liquid nitrogen. For each strain, three biological replicates were performed.

One millilitre of enzyme extraction solution (0.2 mol L<sup>-1</sup> sucrose, 0.1 mol L<sup>-1</sup> Tris, 10 mmol L<sup>-1</sup> ethylene diamine tetraacetic acid, 0.1 mol L<sup>-1</sup> sodium borate and 20% [v/v] glycerol; pH 8.0) was added to each 50 mg of

mycelia powder. The supernatant was collected by centrifugation at  $1 \times 10^4$  rpm (Eppendorf centrifuge 5417 R, Hamburg, Germany) for 10 min at 4 °C, and then immediately mixed with 10 mg LA for 1 h at 28 °C. Finally, the targeted chemicals were enriched with SepPak/C<sub>18</sub>, eluted with 0.5 mL methyl alcohol, and dried over Na<sub>2</sub>SO<sub>4</sub>. The enriched samples were analysed by preparative GC-MS. A GC system (6890 N Network, Agilent, China) with a non-polar column (30 m, DB-5; J&W Scientific, Folsom, CA, USA; film, 0.25 µm; diameter, 0.25 mm; carrier gas He, 15 psi; flow rate, 1 mL min<sup>-1</sup>) and MS detector (5973 Network; Agilent, USA) were used. After splitless injections of samples in heptane, the GC was programmed from 80 °C for 4 min, to 200 °C at a rate of 30 °C min<sup>-1</sup>, to 233 °C at 3 °C min<sup>-1</sup> and then to 285 °C at 10 °C min<sup>-1</sup>. The split ratio was 1:25 and the injected sample size was 1 µL. The standard chemicals for GC-MS (LA, stearic acid and *cis*-9,*trans*-11 CLA) were purchased from Sigma-Aldrich (St Louis, MO, USA).

**Virulence assay.** To determine whether *FGSG\_02668* affected the severity of FHB disease symptoms in wheat spikes, two florets of a single central spikelet per spike were each point-inoculated with a micropipette at the mid-anthesis stage with  $1 \times 10^3$  conidial suspension of *F. graminearum*. The spikes were enclosed in plastic wrap, and placed in a room for 48 h at 25 °C under 16/8 h (day/night) cycles. Blight symptoms were assessed on the 4<sup>th</sup>, 8<sup>th</sup> and 12<sup>th</sup> days after inoculation. Five to ten plants were used per treatment.

To examine whether disruption of *FGSG\_02668* impacted on DON synthesis, the accumulation of DON in liquid media and wheat spikes were tested. Measurement of DON content in liquid media was carried out following a two-stage protocol as described previously<sup>20,44</sup>. Two fully developed florets of each spikelet of one head were each point inoculated with a micropipette at the mid-anthesis stage with  $1 \times 10^3$  conidia of *F. graminearum*. The inoculated spikes were treated as above to induce FHB disease. On the 4<sup>th</sup> day after inoculation, each spikelet was harvested and ground in liquid nitrogen. Three biological replicates were conducted with at least six heads per treatment. Measurement of DON accumulation in wheat spikes was performed as described previously<sup>45</sup>. The DON concentrations were measured using the DON ELISA kit (Beacon, Hebei, China) and Multiskan Spectrum (Thermo Scientific, Finland).

**Measurement of SA, JA, LA and fungal biomass in wheat spikes.** To prepare spike samples, two florets of each fully developed spikelet in an entire spike at mid-anthesis were inoculated with  $1 \times 10^3$  conidia. After inoculation, the wheat plants were placed in a room as described above. At 48 h after initial inoculation, the spikes were harvested and ground to fine powder in liquid nitrogen for chemical measurement and RNA extraction. Three biological replicates were conducted with at least six heads per treatment. Quantification of SA, JA and LA was performed as described previously<sup>46</sup>.

RNA was extracted as described above. The relative amount of *F. graminearum* was estimated by measuring the expression level of *FgGAPDH* in the RNA samples by qPCR, with normalisation to the three wheat reference genes (*w*-GAPDH, NCBI UniGene Ta.66461; *Aox*, Ta.6172; *hn*-RNP-Q, Ta.10105)<sup>20</sup>.

**Microscopic assay.** For optical microscopic and fluorescence microscopic assays, conidia were produced in carboxymethyl cellulose liquid medium at 28 °C with shaking (180 rpm) for 5 d. Conidia ( $1 \times 10^3$ ) were inoculated into 3 mL mSNA-2 liquid media, incubated at 28 °C on an orbital shaker for 6 d in darkness at 120 rpm, and then 20 mg LA was added 1 d before microscopic observation. The conidia and hyphae were observed with a Nikon-80i fluorescence microscope (Nikon, Japan).

**Statistical analysis.** Student's *t*-test (implemented in DPS version 12.01 software) was used to test the significance of differences in the means for LA metabolism, conidial morphology, DON production and disease severity between treatments.

## References

1. Fernando, W. G., Paulitz, T. C., Seaman, W. L., Dutilleul, P. & Miller, J. D. Head blight gradients caused by *Gibberella zeae* from area sources of inoculum in wheat field plots. *Phytopathology* **87**, 414–421 (1997).
2. Chandler, E. A. & Simpson, D. R. Development of PCR assays to *TRI7* and *TRI13* trichothecene biosynthetic genes, and characterization of chemotypes of *Fusarium graminearum*, *Fusarium culmorum* and *Fusarium cerealis*. *Physiol. Mol. Plant P.* **62**, 355–367 (2003).
3. Goswami, R. S. & Kistler, H. C. Heading for disaster: *Fusarium graminearum* on cereal crops. *Mol. Plant Pathol.* **5**, 515–525 (2004).
4. Anand, A. *et al.* Greenhouse and field testing of transgenic wheat plants stably expressing genes for thaumatin-like protein, chitinase and glucanase against *Fusarium graminearum*. *J. Exp. Bot.* **54**, 1101–1111 (2003).
5. Mesterhazy, A. Types and components of resistance to *Fusarium* head blight. *Plant Breeding* **114**, 377–386 (2006).
6. Ellinger, D., Sode, B., Falter, C. & Voigt, C. A. Resistance of callose synthase activity to free fatty acid inhibition as an indicator of *Fusarium* head blight resistance in wheat. *Plant Signal Behav.* **9**, e28982 (2014).
7. Blümke, A. *et al.* Secreted fungal effector lipase releases free fatty acids to inhibit innate immunity-related callose formation during wheat head infection. *Plant Physiol.* **165**, 346–358 (2014).
8. Garscha, U. *et al.* Identification of dioxygenases required for *Aspergillus* development. *J. Biol. Chem.* **282**, 34707–34718 (2007).
9. Hoffmann, I. & Oliu, E. H. 7,8- and 5,8-Linoleate diol synthases support the heterolytic scission of oxygen-oxygen bonds by different amide residues. *Arch. Biochem. Biophys.* **539**, 87–91 (2013).
10. Hao, G. F. *et al.* Role of malic enzyme during fatty acid synthesis in the oleaginous fungus *Mortierella alpina*. *Appl. Environ. Microb.* **80**, 2672–2678 (2014).
11. Hao, D. *et al.* Production of conjugated linoleic acid by heterologous expression of linoleic acid isomerase in oleaginous fungus *Mortierella alpina*. *Biotechnol. Lett.* **37**, 1983–1992 (2015).
12. Tsitsigiannis, D. I., Zarnowski, R. & Keller, N. P. The lipid body protein, *PpoA*, coordinates sexual and asexual sporulation in *Aspergillus nidulans*. *J. Biol. Chem.* **279**, 11344–11353 (2004).
13. Sooman, L. & Oliu, E. H. Discovery of a novel linoleate dioxygenase of *Fusarium oxysporum* and linoleate diol synthase of *Colletotrichum graminicola*. *Lipids* **50**, 1243–1252 (2015).
14. Kepler, C. R., Hirons, K. P., McNeill, J. J. & Tove, S. B. Intermediates and products of the biohydrogenation of linoleic acid by *Butyrivibrio fibrilvolvens*. *J. Biol. Chem.* **241**, 1350–1354 (1966).

15. Wise, M. L., Hamberg, M. & Gerwick, W. H. Biosynthesis of conjugated triene-containing fatty acids by a novel isomerase from the red marine alga *Ptilota filicina*. *Biochemistry* **33**, 15223–15232 (1994).
16. Destailhats, F., Trottier, J. P., Galvez, J. M. & Angers, P. Analysis of alpha-linolenic acid biohydrogenation intermediates in milk fat with emphasis on conjugated linoleic acids. *J. Dairy Sci.* **88**, 3231–3239 (2005).
17. Churrua, I., Fernández-Quintela, A. & Portillo, M. P. Conjugated linoleic acid isomers: differences in metabolism and biological effects. *Biofactors* **35**, 105–111 (2009).
18. Rosberg-Cody, E., Johnson, M. C., Fitzgerald, G. F., Ross, P. R. & Stanton, C. Heterologous expression of linoleic acid isomerase from *Propionibacterium acnes* and anti-proliferative activity of recombinant trans-10, cis-12 conjugated linoleic acid. *Microbiology* **153**, 2483–2490 (2007).
19. Peng, S. S., Deng, M. D., Grund, A. D. & Rosson, R. A. Purification and characterization of a membrane-bound linoleic acid isomerase from *Clostridium sporogenes*. *Enzyme Microb. Tech.* **40**, 831–839 (2007).
20. Qi, P. F. *et al.* Effect of salicylic acid on *Fusarium graminearum*, the major causal agent of fusarium head blight in wheat. *Fungal Biol.* **116**, 413–426 (2012).
21. Qi, P. F. *et al.* Jasmonic acid and abscisic acid play important roles in host-pathogen interaction between *Fusarium graminearum* and wheat during the early stages of fusarium head blight. *Physiol. Mol. Plant P.* **93**, 39–48 (2016).
22. Champe, S. P. & El-zayat, A. A. E. Isolation of a sexual sporulation hormone from *Aspergillus nidulans*. *J. Bacteriol.* **171**, 3982–3988 (1989).
23. Kumaraswamy, K. G., Kushalappa, A. C., Choo, T. M., Dion, Y. & Rioux, S. Mass spectrometry based metabolomics to identify potential biomarkers for resistance in barley against fusarium head blight (*Fusarium graminearum*). *J. Chem. Ecol.* **37**, 846–856 (2011).
24. Li, G. & Yen, Y. Jasmonate and ethylene signaling pathway may mediate fusarium head blight resistance in wheat, *Crop Sci.* **48**, 1888–1896 (2008).
25. Robert-Seilant, A., Navarro, L., Bari, R. & Jones, J. D. G. Pathological hormone imbalances. *Curr. Opin. Plant Biol.* **10**, 372–379 (2007).
26. Koornneef, A. & Pieterse, C. M. J. Cross talk in defense signalling. *Plant Physiol.* **146**, 839–844 (2008).
27. Leon-Reyes, A. *et al.* Ethylene signaling renders the jasmonate response of *Arabidopsis* insensitive to future suppression by salicylic acid. *Mol. Plant Microbe Interact.* **23**, 187–197 (2010).
28. Mika, A., Boenisch, M. J., Hopff, D. & Luthje, S. Membrane-bound guaiacol peroxidases from maize (*Zea mays* L.) roots are regulated by methyl jasmonate, salicylic acid, and pathogen elicitors. *J. Exp. Bot.* **61**, 831–841 (2010).
29. Needleman, P., Turk, J., Jakschik, B. A., Morrison, A. R. & Lefkowitz, J. B. Arachidonic acid metabolism. *Annu. Rev. Biochem.* **55**, 69–102 (1986).
30. Zhang, X. *et al.* Disruption of the fatty acid Delta6-desaturase gene in the oil-producing fungus *Mortierella isabellina* by homologous recombination. *Curr. Microbiol.* **55**, 128–134 (2007).
31. Pariza, M., Park, Y. & Cook, M. The biologically active isomers of conjugated linoleic acid. *Prog. Lipid Res.* **40**, 293–298 (2001).
32. Pariza, M. W. Perspective on the safety and effectiveness of conjugated linoleic acid. *Am. J. Clin. Nutr.* **79**, 1132S–1136S (2004).
33. Zheng, G., Gengwang, Z. & Yan, S. Preparation of conjugated linoleic acid and identification of its isomers. *Chinese J. Chem. Eng.* **11**, 130–135 (2003).
34. Deng, M. D. *et al.* Linoleic acid isomerase from *Propionibacterium acnes*: purification, characterization, molecular cloning, and heterologous expression. *Appl. Biochem. Biotechnol.* **143**, 199–211 (2007).
35. Sehat, N. *et al.* Silver-ion high-performance liquid chromatographic separation and identification of conjugated linoleic acid isomers. *Lipids* **33**, 217–221 (1998).
36. Capellini, R. A. & Peterson, J. L. Macroconidium formation in submerged cultures by a non-sporulating strain of *Gibberella zeae*. *Mycologia* **57**, 962–966 (1965).
37. Bertani, G. Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia coli*. *J. Bacteriol.* **62**, 293–300 (1951).
38. Lodhi, M. A., Ye, G. N., Weeden, N. F. & Reisch, B. I. A simple and efficient method for DNA extraction from grapevine cultivars and Vitis species. *Plant Mol. Biol. Rep.* **12**, 6–13 (1994).
39. Frandsen, R. J., Andersson, J. A., Kristensen, M. B. & Giese, H. Efficient four fragment cloning for the construction of vectors for targeted gene replacement in filamentous fungi. *BMC Mol. Biol.* **9**, 1–11 (2008).
40. Frandsen, R. J., Frandsen, M. & Giese, H. Targeted gene replacement in fungal pathogens via *Agrobacterium tumefaciens*-mediated transformation. *Methods Mol. Biol.* **835**, 17–45 (2012).
41. Maier, F. J., Malz, S., Lösch, A. P., Lacour, T. & Schäfer, W. Development of a highly efficient gene targeting system for *Fusarium graminearum* using the disruption of a polyketide synthase gene as a visible marker. *FEMS Yeast Res.* **5**, 653–662 (2005).
42. Solomon, P. S., Ipcho, S. V. S., Hane, J. K., Tan, K. C. & Oliver, R. P. A quantitative PCR approach to determine gene copy number. *Fungal Genet. Rep.* **55**, 5–8 (2008).
43. Wang, J. R. *et al.* RNA profiling of fusarium head blight-resistant wheat addition lines containing the *Thinopyrum elongatum* chromosome 7E. *Can. J. Plant Pathol.* **32**, 188–214 (2010).
44. Miller, D. & Blackwell, B. A. Biosynthesis of 3-acetyldeoxynivalenol and other metabolites by *Fusarium culmorum* HLK 1503 in a stirred jar fermentor. *Can. J. Bot.* **64** (1986).
45. Zhang, Y. Z. *et al.* Chitin synthase gene *FgCHS8* affects virulence and fungal cell wall sensitivity to environmental stress in *Fusarium graminearum*. *Fungal Biol.* **120**, 764–774 (2016).
46. Siciliano, I., Amaral Carneiro, G., Spadaro, D., Garibaldi, A. & Gullino, M. L. Jasmonic acid, abscisic acid, and salicylic acid are involved in the phytoalexin responses of rice to *Fusarium fujikuroi*, a high gibberellin producer pathogen. *J. Agric. Food Chem.* **63**, 8134–8142 (2015).

## Acknowledgements

This research was supported by the National Natural Science Foundation of China (31230053) and International S&T Cooperation program of China (2015DFA30600). We thank Prof. Wei Wu of Sichuan Agricultural University and Dr Kai Zhong of Sichuan University for technical assistance. We thank Dr Ji-Rui Wang of the Triticeae Research Institute for kind advice. The authors offer special thanks to Dr Therese Ouellet of Agriculture and Agri-Food Canada for providing research materials.

## Author Contributions

P.F.Q. and Y.M.W. designed the experiments. P.F.Q., Y.Z.Z. and Q.C. wrote the manuscript and analysed the data. P.F.Q. and Y.Z.Z. prepared the figures. P.F.Q., Y.Z.Z., Z.Z.W., Q.C., C.H.L., Z.R.G., Y.L.C., L.J.Z., Y.N.H., C.C., Y.W., Y.Y.Q., X.F., T.Z., D.M. and B.J.X. performed the experiments. W.L., Q.T.J. and Y.L.Z. provided key reagents and advice. All authors reviewed the results and approved the final version of the manuscript.

## Additional Information

**Competing Interests:** The authors declare no competing financial interests.



**How to cite this article:** Zhang, Y.-Z. *et al.* Linoleic acid isomerase gene *FgLAI12* affects sensitivity to salicylic acid, mycelial growth and virulence of *Fusarium graminearum*. *Sci. Rep.* 7, 46129; doi: 10.1038/srep46129 (2017).

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017