

# Transgenerational impact of intimate partner violence on methylation in the promoter of the glucocorticoid receptor

KM Radtke<sup>1,2,4</sup>, M Ruf<sup>1,4</sup>, HM Gunter<sup>2,3,4</sup>, K Dohrmann<sup>1</sup>, M Schauer<sup>1</sup>, A Meyer<sup>2</sup> and T Elbert<sup>1</sup>

Prenatal exposure to maternal stress can have lifelong implications for psychological function, such as behavioral problems and even the development of mental illness. Previous research suggests that this is due to transgenerational epigenetic programming of genes operating in the hypothalamic–pituitary–adrenal axis, such as the glucocorticoid receptor (GR). However, it is not known whether intrauterine exposure to maternal stress affects the epigenetic state of these genes beyond infancy. Here, we analyze the methylation status of the *GR* gene in mothers and their children, at 10–19 years after birth. We combine these data with a retrospective evaluation of maternal exposure to intimate partner violence (IPV). Methylation of the mother's *GR* gene was not affected by IPV. For the first time, we show that methylation status of the *GR* gene of adolescent children is influenced by their mother's experience of IPV during pregnancy. As these sustained epigenetic modifications are established *in utero*, we consider this to be a plausible mechanism by which prenatal stress may program adult psychosocial function.

*Translational Psychiatry* (2011) 1, e21; doi:10.1038/tp.2011.21; published online 19 July 2011

## Introduction

Prenatal stress can have a lasting detrimental impact on psychological health; however, the molecular mechanisms that transmit this experience to adult behavior are not fully characterized.<sup>1–5</sup> The hypothalamic–pituitary–adrenal (HPA) axis is critical for homeostasis—it controls growth, reproduction, metabolism and behavior; it is also the primary line of the 'defence cascade' that helps humans to deal with crises. Hyperactivity of the HPA-axis can cause anything from a long-lasting head cold to depression. Hypoactivity of the HPA-axis can cause undesirable consequences such as abdominal fat, loss of muscle mass and mental ill-health.<sup>6,7</sup> Tuning of the HPA-axis and its physiological pathways is highly susceptible to the influence of early life events. Prenatal stress such as antenatal exposure to maternal anxiety has sustained effects on HPA-axis function<sup>8</sup> and is associated with behavioral and emotional problems arising during development.<sup>9,10</sup>

DNA methylation could be a mechanism by which prenatal stress is translated into changes in gene expression and physiology,<sup>11</sup> and ultimately psychologically vulnerable phenotypes.<sup>12,13</sup> The glucocorticoid receptor (GR), a major regulator of the HPA-axis,<sup>14</sup> could be involved in such a transmission, controlling many aspects of development, metabolism and immune function. Landmark experiments in rodents show that both the hippocampal expression of this gene and the behavioral responses to stress are modulated by the amount of care mothers invest into their offspring in the first days of postnatal life.<sup>15,16</sup> This is likely to be the result of epigenetic

modifications, specifically, through methylation of exon 1F of the *GR* promoter.<sup>17</sup> This exon contains a response element for the nerve-growth-factor-inducible protein A (NGFI-A),<sup>18</sup> and binding of NGFI-A to its response element increases *GR* expression. Methylation of the NGFI-A response element inhibits the association with its 'ligand' thereby decreasing *GR* expression.<sup>19,20</sup> Also, the human *GR* gene is affected by aversive social environments, as childhood abuse leads to increased methylation in exon 1F of the *GR* promoter.<sup>21</sup>

The vast majority of studies analyzing the influence of stress on the epigenetic regulation of the HPA-axis have focussed on postnatal stressors. However, recent research suggests that the intrauterine environment can also impact the epigenetic state of HPA-axis genes—prenatal exposure to maternal depressive mood was shown to correlate with *GR* promoter methylation in newborns.<sup>22</sup> Furthermore, increased *GR* promoter methylation was associated with higher cortisol responses to stress in this study and could therefore represent fetal programming of the HPA-axis. However, this study focused on methylation status in umbilical cord blood. Therefore, it is not known whether epigenetic marks that are established *in utero* due to the psychosocial situation of the mother are maintained beyond infancy.

Here, we investigate whether gestational maternal aversive experiences can have a prolonged effect, 10–19 years after birth, on DNA methylation of the offspring based on bisulfite sequencing of DNA from whole blood. Both intrauterine exposure to marital discord and *GR* promoter methylation

<sup>1</sup>Department of Psychology, University of Konstanz and Center for Psychiatry Reichenau, Konstanz, Germany; <sup>2</sup>Lehrstuhl für Zoologie und Evolutionsbiologie, Department of Biology, University of Konstanz, Konstanz, Germany and <sup>3</sup>Zukunftskolleg, University of Konstanz, Konstanz, Germany

Correspondence: Professor T Elbert, Department of Psychology, University of Konstanz and Center for Psychiatry Reichenau, 78457 Konstanz, Germany or Professor A Meyer, Lehrstuhl für Zoologie und Evolutionsbiologie, Department of Biology, University of Konstanz, 78457 Konstanz, Germany.  
E-mail: thomas.elbert@uni-konstanz.de or axel.meyer@uni-konstanz.de

<sup>4</sup>These authors contributed equally to this work.

**Keywords:** epigenetics; glucocorticoid receptor; hypothalamic–pituitary–adrenal axis; intimate partner violence; methylation; prenatal stress

Received 6 June 2011; accepted 7 June 2011

can predict psychopathology.<sup>21,23,24</sup> We aimed to determine whether prenatal exposure to intimate partner violence (IPV) leads to increased *GR* promoter methylation later in life, which might mediate increased susceptibility to psychopathology.

## Materials and methods

**Participants.** The study cohort represents a convenience sample, as there were no specific criteria that the participants had to fulfil. Mother–child pairs were either recruited via advertisements or taken from another study.<sup>25</sup>

**Psychological parameters.** We studied the impact of exposure to IPV in women from a variety of ethnic backgrounds (for example, Kosovo, Russia, Turkey etc.), who reside in Germany (Table 1). IPV was evaluated using the composite abuse scale (CAS)<sup>26</sup> applied as structured interview by experienced clinical psychologists. The CAS is a validated tool,<sup>27</sup> which measures the degree of domestic violence experienced by an individual in the four dimensions of severe combined abuse, physical abuse, emotional abuse and harassment.<sup>26</sup> It consists of 30 items, which are scaled from 0 (never) to 5 (daily). Missing items, due to problems concerning the translation, were replaced by the mean of the existing items (1 of 25 interviews). According to previously described methods,<sup>28,29</sup> a resulting ‘sumscore’ of 7 or higher was used as the criterion for exposure to IPV. In order to

match the design of our study, the CAS was conducted three times separately focusing on the periods before, during and after the pregnancy with the particular child whose blood was analyzed in this study. For the periods before and after pregnancy, participants were asked to report acts of domestic violence whenever these happened before or after pregnancy with the relevant child. These periods were not limited to an absolute time span.

**Sodium bisulfite sequencing.** Sodium bisulfite conversion was performed, according to previously described methods,<sup>30,31</sup> on DNA extracted from whole blood (DNeasy Blood and Tissue Kit, Qiagen, Hilden, Germany). Briefly, 5 µg of DNA was denatured by incubating in 0.4 M NaOH (Riedel-deHaën, Seelze, Germany) for 30 min at 42 °C. After the addition of hydroquinone (Sigma-Aldrich, Steinheim, Germany) and sodium bisulfite (Sigma-Aldrich) to a final concentration of 0.5 mM and 2.6 mM, respectively, the reaction mixture was incubated at 55 °C for 18 h. To remove free sodium bisulfite, the reaction mixture was cleaned up using a silica-based method,<sup>32</sup> and was eluted in 100 µl of double-distilled H<sub>2</sub>O. To remove the bisulfite adduct, 11 µl of 4.0 M NaOH was added, and the reaction mixture was incubated at 37 °C for 15 min. The reaction mixture was then neutralized by adding 111 µl of 7.5 M ammonium acetate, pH 7.0, (Riedel-deHaën). DNA was precipitated with ethanol<sup>33</sup> and resuspended in 30 µl of double-distilled H<sub>2</sub>O.

**Table 1** Maternal IPV and sociodemographics

Label	Maternal exposure to IPV				Children's gender	Children's age (years)	Mode of birth	Maternal age at birth (years)	Maternal country of origin
	Before pregnancy	During pregnancy	After pregnancy	Total					
1	No	No	No	No	Male	19	Spontaneous birth	19	Rus.
2	No	No	No	No	Male	17	Spontaneous birth	31	Pl.
3	No	No	No	No	Female	13	Emergency caesarean section	25	Tr.
4	No	No	No	No	Male	10	Spontaneous birth	29	Ger.
5	No	No	Yes	Yes	Female	16	Spontaneous birth	19	Irq.
6	No	No	No	No	Female	14	Elective caesarean delivery	28	Ger.
7	No	No	No	No	Male	13	Spontaneous birth	28	Rus.
8	No	No	No	No	Female	14	Elective caesarean delivery	27	Rus.
10	No	Yes	Yes	Yes	Female	15	Other	29	Ger.
11	No	No	Yes	Yes	Female	12	Spontaneous birth	31	Tw.
12	No	No	Yes	Yes	Male	17	Emergency caesarean section	21	Ir.
13	No	No	No	No	Female	17	Spontaneous birth	26	Cs.
14	No	Yes	No	Yes	Female	13	Spontaneous birth	38	Rus.
17	No data	No data	Yes	Yes	Female	12	Emergency caesarean section	18	Irq.
18	No	No	No	No	Female	13	Spontaneous birth	29	Sos.
20	No	No	No	No	Female	12	Spontaneous birth	26	Sos.
21	Yes	Yes	Yes	Yes	Male	11	Elective caesarean delivery	23	Sos.
22	No	No	No	No	Male	12	Spontaneous birth	20	Scg.
23	No	Yes	No	Yes	Female	10	Spontaneous birth	19	Irq.
24	No	No	No	No	Male	13	Spontaneous birth	29	Sos.
25	Yes	Yes	Yes	Yes	Female	14	Spontaneous birth	24	Sos.
26	No	No	Yes	Yes	Female	17	Spontaneous birth	16	Sos.
27	No data	Yes	No data	Yes	Female	14	Spontaneous birth	22	Tr.
28	Yes	Yes	Yes	Yes	Female	18	Spontaneous birth	22	Tr.
29	No data	Yes	No data	Yes	Female	17	Spontaneous birth	30	Tr.
<i>n</i>	3 (22)	8 (24)	9 (23)	13 (25)	25	25	25	25	25
Range						10–19		16–38	
Mean ± s.e.m.						14.1 ± 0.5		25.2 ± 1.0	

Abbreviations: Cs., Czechoslovakia; Ger., Germany; IPV, intimate partner violence; Ir., Iran; Irq., Iraq; Pl., Poland; Rus., Russia; Scg., Serbia and Montenegro; Sos., Kosovo; Tw., Taiwan; Tr., Turkey.

‘Total’ indicates exposure to IPV regardless of timing of exposure. Whenever IPV was experienced before, during or after pregnancy ‘total’ was computed as ‘yes’. In three cases we could not obtain CAS-scores across all periods as indicated by ‘no data’.

The converted DNA was amplified with PCR, using Fast-Start Taq DNA Polymerase (Roche, Mannheim, Germany), with the previously published PCR-primers:<sup>34</sup> forward, -142763905-5'-GTTGTTATTYGTAGGGGTATTGG-3'-142763883 and reverse, 142763770-5'-AAACCACCRAATTTCT CCAA-3'-142763789 (sequence numbering is according to the published nucleotide position on chromosome 5; GenBank accession number: AJ877168). PCR conditions: 94 °C for 5 min, then 34 cycles of denaturation (30 s, 95 °C), annealing (2 min, 56 °C) and elongation (30 s, 72 °C), plus a final elongation step (72 °C, 7 min). The PCR was designed to amplify a 93 bp fragment in the *GR* promoter spanning 10 CpG sites. Products were purified (QIAquick, Qiagen) and then cloned into pCR4-TOPO-vector using the TOPO TA Cloning kit (Invitrogen, Darmstadt, Germany). Positive clones were identified with colony PCR using the primers designed to the plasmid: M13-Fwd and M13-Rev primers (Invitrogen). Between 14 and 25 clones were sequenced using the same forward and reverse primers used for colony PCR (BigDye Terminator 3.1 Cycle Sequencing kit, Applied Biosystems, Darmstadt, Germany). The sequences for each individual clone were aligned and analyzed in Sequencher (version 4.2.2., Gene Codes Corporation, Ann Arbor, MI, USA).

**Statistical analysis.** Statistical analysis was conducted using PASW statistics (version 18.0, SPSS incorporation, Munich, Germany). To test for a potential association between maternal exposure to IPV and the presence of methylation in either the mothers or the children, we conducted Fisher's exact tests thereby treating methylation as a binary variable. The *GR* promoter was referred to as being methylated, if a methylated CpG site was detected in at least one of up to 25 clones that were sequenced. For the mothers, we did not distinguish between different periods of exposure to IPV. Therefore, we computed an overall variable for exposure to IPV, assigning exposure to IPV whenever it happened before, during or after pregnancy. For the children, as we were interested in the relationship between methylation and maternal exposure to IPV before, during and after pregnancy, we performed tests independently for each interval.

To test for an association between the methylation status of the *GR* promoter in the mothers and in the children, we used Fisher's exact test.

To further analyze the direction of the relationship between methylation and maternal exposure to IPV during pregnancy, we conducted a Man-Whitney U-test with the exposure to IPV during pregnancy as the between group factor, and the percentage of methylated clones in the children as the dependent variable. The percentage of methylated clones was calculated, according to previously described methods,<sup>21</sup> as the number of clones containing at least one methylated CpG site divided by the total number of clones. Additionally, we analyzed the relationship between maternal exposure to IPV before and after gestation and the percentage of methylated clones in the children in the same way.

## Results

**Maternal stress profiling.** A total of 25 women of various ethnic backgrounds (Table 1) participated in the study. They


were aged between 29 and 51 years. All of them had children aged between 10 and 19 years (Table 1). We used the CAS to screen the mothers, but not the children, for exposure to IPV before, during and after their pregnancies. We identified three women, who were exposed to IPV in the period before pregnancy, eight women, who were exposed during and nine after pregnancy. It should be noted that these groups were overlapping, that is, for some women exposure to IPV was apparent in more than one period. In three cases we could not obtain CAS-values for each period, therefore, sample sizes varied slightly in the statistical analyses (Table 1).

***GR* promoter methylation.** We examined the methylation status of 10 CpG sites in the *GR* promoter, to determine whether there is a link between exposure to IPV and methylation in a gene that is involved in the HPA-axis. This region was selected because it was previously shown to contain transcription factor binding sites, whose methylation statuses are influenced by early life stress.<sup>21</sup> The *GR* promoter was referred to as being methylated, if a methylated CpG site was detected in at least one of up to 25 clones that were sequenced. We characterized *GR* promoter methylation of 25 women and of 24 of their children (for one mother, we could not obtain a blood sample of any of her children). We detected methylation in 7 of 10 CpG sites (sites 01, 02, 03, 07, 08, 09 and 10) in the *GR* promoter in our samples. The degree of methylation in these CpG sites ranged from 0 to 20% of all analyzed clones (Figure 1).


We detected methylation in the *GR* promoter of 10 mothers and seven children in at least one clone. There was no association between maternal methylation status and the methylation status of the children ( $n=24$ ;  $P=1.0$ ).

**Impact of IPV on maternal methylation pattern.** Using fisher's exact test, we did not find a significant relationship between exposure to IPV and maternal methylation status ( $n=25$ ,  $P=0.7$ ), irrespective of when IPV was experienced.

**Impact of IPV on methylation patterns in offspring.** The presence of methylated residues was significantly associated with maternal exposure to IPV during pregnancy ( $n=23$ ,  $P<0.05$ ). Remarkably, there was no association between maternal exposure to IPV and the presence of methylation in the offspring, when IPV was experienced either before or after pregnancy (before:  $n=21$ ,  $P=1.0$ ; after:  $n=22$ ,  $P=1.0$ ). Furthermore, we tested the influence of a variety of third variables (country of origin; maternal age; age of children; marital status; graduation from high school, completion of vocational training or academic degree; pregnancy problems; smoking, alcohol or drug consumption during pregnancy; use of painkillers during birth; anxiety, panic or helplessness during birth; skin contact after birth; severe diseases of children; birth weight; gestational age at birth; mode of birth) on the methylation status, but did not find a significant relationship (Supplementary Table S1 and S2). According to the guidelines of the world health organization (WHO) none of the analyzed children were of extreme immaturity (<28 completed weeks of gestation, WHO) or extreme low birth weight (birth weight of 999 g or less, WHO).



**Figure 1** Degree of methylation per individual and CpG site. Each line represents the analyzed fraction of the glucocorticoid receptor gene of a woman (Mu) and her child (Ki). The rows correspond to individual CpG sites. The degree of methylation for each individual CpG site, that is, the number of clones containing methylation in the particular site divided by the total number of sequenced clones, is colour-coded. In the case of one child we could not obtain a blood sample as indicated by 'no data'. (i) Putative NGFI-A-binding site; (ii) Known NGFI-A-binding site.



**Figure 2** Gestational effects of IPV on methylation of the *GR* promoter in the children. Mean  $\pm$  s.e.m. of percentage of methylated clones for the children of women exposed to IPV. IPV only associates with increased methylation, if maternal exposure occurred during pregnancy. The percentage of methylated clones was calculated as the number of clones containing at least one methylated CpG site divided by the total number of clones. \* $P < 0.05$ ; IPV, intimate partner violence.

However, one child was born prematurely (28 completed weeks or more but less than 37 completed weeks of gestation, WHO) and four children were born with low birth weights (1000–2499 g, WHO). None of these children showed methylation at the *GR* promoter. Furthermore, the vast majority of the children ( $n = 17$ ) were delivered spontaneously, indicating that most mothers did not take any measures to delay or induce labor. However, six were delivered via caesarean section and one via a different procedure (Supplementary Table S2). Neither the birth weight, the gestational age nor the mode of birth affected the methylation status of the *GR* gene (Supplementary Table S1). This provides evidence that methylation in the

offspring is directly affected by adverse experiences of the mother during gestation.

We then sought to determine whether gestational exposure to IPV influenced the degree of methylation in the *GR* promoter in the offspring. To achieve this we evaluated the percentage of methylated clones (that is, the number of clones containing at least one methylated CpG site divided by the total number of clones). This value ranged from 0 to 20% across all of our experimental subjects. In line with our previous analysis, we found a significant relationship between the percentage of methylated clones and IPV during pregnancy ( $n = 23$ ,  $U = 30.5$ ,  $P = 0.015$ ; Figure 2; Supplementary Figure S3). Additionally, we did not find an association between the percentage of methylated residues and maternal exposure to IPV before or after pregnancy (before:  $n = 21$ ,  $U = 25.0$ ,  $P = 0.9$ ; after:  $n = 22$ ,  $U = 55.0$ ,  $P = 0.7$ ; Figure 2; Supplementary Figure S4-5).

## Discussion

We examined the methylation pattern of the *GR* promoter in 25 women and their children and analyzed its association with maternal stress during gestation. We find a positive relationship between methylation of the children's *GR* promoter and maternal exposure to IPV during pregnancy, defined as 'any behavior within an intimate relationship that causes physical, psychological or sexual harm to those in the relationship.'<sup>35</sup> We present evidence that *GR* methylation is related to maternal stress during pregnancy, as when IPV is experienced before or after pregnancy, it has no impact on the children's methylation. To the best of our knowledge, this is the first study to demonstrate that prenatal psychological stress can result in sustained, rather than just short-term alteration of methylation in a regulator gene of the HPA-axis.

We found that maternal gestational IPV is associated with methylation of exon 1F in the *GR* promoter of the offspring, which is hence a transgenerational effect that may exert a lifelong influence on HPA-axis regulation in these individuals. This is consistent with previous observations that prenatal anxiety is associated with a sustained elevation of basal HPA activity<sup>8</sup> as well as with behavioral/emotional problems that may continue throughout the lifetime.<sup>9,36</sup> It seems likely that the intrauterine environment that the fetus experiences can differentially affect the methylation pattern. Maternal stress may lead to changes in HPA regulatory circuits with an alteration of catecholamines and glucocorticoids among other factors and effects.<sup>14</sup> Prenatal exposure to either of these hormones is known to influence the development of the HPA-axis.<sup>37</sup> Therefore in the fetus, the *GR* gene, which is a key regulator of the HPA-axis, constitutes a likely target upon which maternal IPV could be acting. Our data are consistent with observations in rat, where maternal effects evoke increased HPA-axis activity and fearfulness in response to stress, which are associated with methylation of the *GR* promoter in the hippocampus.<sup>17</sup> In these experiments, the first days of postnatal life constitute a critical period for the manifestation of maternal effects, which corresponds to late gestation in humans.<sup>38</sup> Although the observed increase in methylation was only very subtle, it might be sufficient to alter HPA-axis function. Subtle increases in methylation in a single CpG site of the *GR* promoter in cord blood were reported to correlate positively with the cortisol stress response in neonates.<sup>22</sup> This suggests that IPV elicits fetal programming of the HPA-axis through methylation of the *GR* gene. However, although a link between *GR* promoter methylation in blood and cortisol stress responses has already been provided,<sup>22</sup> it remains inconclusive whether methylation in blood cells reflects methylation in the hippocampus or whether it impacts HPA-axis function. This emphasizes the need for follow-up studies, which evaluate both HPA-axis function and psychological function. This is of particular interest, as blood—as opposed to brain tissue—represents an easily sampled tissue.

The results of our study suggest that maternal stress-induced changes in intrauterine environment can have an impact on the methylation pattern of the children's *GR* gene, rather than a direct maternal transmission of methylation pattern via the germ line. The CAS assesses exclusively environmental factors, episodes of domestic violence in this case, thereby minimizing the influence of genetic components. However, an individual's genotype might affect others' behavior towards them, thus their own social environment, or predispose them to seek risky or unfavorable social environments.<sup>39</sup> As this type of indirect effect is difficult to avoid in studies with humans, we believe that our chosen methods go a long way to minimizing the influence of underlying genetic background. Hence, we consider gestational IPV to be the main source of variation in our study.

The methodologies employed by our study present some limitations: we have adapted a commonly used psychological survey—the CAS, to examine the influence of past events on current methylation patterns of mothers and their children. The effectiveness of retrospective surveys relies on the accuracy of our participants' memories. We believe that our

use of an event-based analysis tool is particularly robust because of its reliance on emotionally arousing events, which are known to create long-lasting memories.<sup>40</sup> Our data represent correlative findings and thus cannot prove a causal relationship between changes in methylation and adverse experiences. For instance, it might be possible that IPV correlates with aspects of maternal diet that in turn might have affected the observed methylation patterns. Such a scenario, however, seems unlikely, as we would then predict an association between IPV and methylation in the mothers. This, however, was not observed.

## Conclusions

Our findings show that prenatal exposure to IPV is associated with a sustained increase in methylation of the human *GR* promoter in the blood. Prenatal stress is known to alter HPA-axis regulatory function later in life.<sup>8</sup> Specifically gestational marital discord is associated with psychopathology of the offspring.<sup>23,24</sup> This is the first demonstration that gestational exposure to psychological stressors can have a lasting impact on methylation status in human offspring. Our results provide a potential mechanism—methylation of the *GR* promoter—upon which prenatal stress could act, to influence psychological function. This emphasizes the importance of IPV interventions to assure the well-being not only of the mother but also of the unborn child. This mechanism opens up many new avenues for research on the transgenerational epigenetic effects of stress and aggression on human behavior.

## Conflict of interest

The authors declare no conflict of interest.

**Acknowledgements.** We thank IT Kolassa, A Ashe, Ö Hatimaz and WA Schulz for helpful suggestions on the methods, M Roth and H Riedke for help during sample collection, and PM Hull and F Henning for helpful suggestions on the manuscript. HMG is supported by the Zukunftscolleg of University of Konstanz. Research was supported by the DFG.

1. St Clair D, Xu M, Wang P, Yu Y, Fang Y, Zhang F *et al*. Rates of adult schizophrenia following prenatal exposure to the Chinese famine of 1959–1961. *JAMA* 2005; **294**: 557–562.
2. Susser E, Neugebauer R, Hoek HW, Brown AS, Lin S, Labovitz D *et al*. Schizophrenia after prenatal famine. Further evidence. *Arch Gen Psychiatry* 1996; **53**: 25–31.
3. Susser ES, Lin SP. Schizophrenia after prenatal exposure to the Dutch Hunger Winter of 1944–1945. *Arch Gen Psychiatry* 1992; **49**: 983–988.
4. Watson JB, Mednick SA, Huttunen M, Wang X. Prenatal teratogens and the development of adult mental illness. *Dev Psychopathol* 1999; **11**: 457–466.
5. Talge NM, Neal C, Glover V. Antenatal maternal stress and long-term effects on child neurodevelopment: how and why? *J Child Psychol Psychiatry* 2007; **48**: 245–261.
6. Elbert T, Rockstroh B. Stress factors. The science of our flexible responses to an unpredictable world. *Nature* 2003; **421**: 477–478.
7. McEwen BS, Lasley EN. *The End of Stress as We Know It*. Joseph Henry Press/ Dana Press: Washington DC, USA, 2002.
8. O'Connor TG, Ben-Shlomo Y, Heron J, Golding J, Adams D, Glover V. Prenatal anxiety predicts individual differences in cortisol in pre-adolescent children. *Biol Psychiatry* 2005; **58**: 211–217.
9. O'Connor TG, Heron J, Golding J, Beveridge M, Glover V. Maternal antenatal anxiety and children's behavioural/emotional problems at 4 years. Report from the Avon Longitudinal Study of Parents and Children. *Br J Psychiatry* 2002; **180**: 502–508.
10. O'Connor TG, Heron J, Golding J, Glover V. Maternal antenatal anxiety and behavioural/emotional problems in children: a test of a programming hypothesis. *J Child Psychol Psychiatry* 2003; **44**: 1025–1036.

11. Murgatroyd C, Patchev AV, Wu Y, Micale V, Bockmuhl Y, Fischer D *et al*. Dynamic DNA methylation programs persistent adverse effects of early-life stress. *Nat Neurosci* 2009; **12**: 1559–1566.
12. Gregory SG, Connelly JJ, Towers AJ, Johnson J, Biscocho D, Markunas CA *et al*. Genomic and epigenetic evidence for oxytocin receptor deficiency in autism. *BMC Med* 2009; **7**: 62.
13. Uddin M, Aiello AE, Wildman DE, Koenen KC, Pawelec G, de Los Santos R *et al*. Epigenetic and immune function profiles associated with posttraumatic stress disorder. *Proc Natl Acad Sci USA* 2010; **107**: 9470–9475.
14. de Kloet ER, Joels M, Holsboer F. Stress and the brain: from adaptation to disease. *Nat Rev Neurosci* 2005; **6**: 463–475.
15. Liu D, Diorio J, Tannenbaum B, Caldji C, Francis D, Freedman A *et al*. Maternal care, hippocampal glucocorticoid receptors, and hypothalamic-pituitary-adrenal responses to stress. *Science* 1997; **277**: 1659–1662.
16. Francis D, Diorio J, Liu D, Meaney MJ. Nongenomic transmission across generations of maternal behavior and stress responses in the rat. *Science* 1999; **286**: 1155–1158.
17. Weaver ICG, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR *et al*. Epigenetic programming by maternal behavior. *Nat Neurosci* 2004; **7**: 847–854.
18. McCormick JA, Lyons V, Jacobson MD, Noble J, Diorio J, Nyirenda M *et al*. 5'-heterogeneity of glucocorticoid receptor messenger RNA is tissue specific: differential regulation of variant transcripts by early-life events. *Mol Endocrinol* 2000; **14**: 506–517.
19. Szyf M, Weaver ICG, Champagne FA, Diorio J, Meaney MJ. Maternal programming of steroid receptor expression and phenotype through DNA methylation in the rat. *Front Neuroendocrinol* 2005; **26**: 139–162.
20. Weaver ICG, D'Alessio AC, Brown SE, Hellstrom IC, Dymov S, Sharma S *et al*. The transcription factor nerve growth factor-inducible protein a mediates epigenetic programming: altering epigenetic marks by immediate-early genes. *J Neurosci* 2007; **27**: 1756–1768.
21. McGowan PO, Sasaki A, D'Alessio AC, Dymov S, Labonté B, Szyf M *et al*. Epigenetic regulation of the glucocorticoid receptor in human brain associates with childhood abuse. *Nat Neurosci* 2009; **12**: 342–348.
22. Oberlander TF, Weinberg J, Papsdorf M, Grunau R, Misri S, Devlin AM. Prenatal exposure to maternal depression, neonatal methylation of human glucocorticoid receptor gene (NR3C1) and infant cortisol stress responses. *Epigenetics* 2008; **3**: 97–106.
23. Stott DH. Follow-up study from birth of the effects of prenatal stresses. *Dev Med Child Neurol* 1973; **15**: 770–787.
24. Ward AJ. Prenatal stress and childhood psychopathology. *Child Psychiatry Hum Dev* 1991; **22**: 97–110.
25. Roth M, Schauer M, Ruf M. Generationsübergreifende Folgen von Trauma und Posttraumatischer Belastungsstörung bei Flüchtlingen in Deutschland. *Z Klin Psychol Psychother*, submitted.
26. Hegarty K, Sheehan M, Schonfeld C. A multidimensional definition of partner abuse: development and preliminary validation of the composite abuse scale. *J Fam Violence* 1999; **14**: 399–415.
27. Hegarty K, Fracgp, Bush R, Sheehan M. The composite abuse scale: further development and assessment of reliability and validity of a multidimensional partner abuse measure in clinical settings. *Violence Vict* 2005; **20**: 529–547.
28. Taft A, Small R, Hegarty K, Lumley J, Watson L, Gold L. MOSAIC (MOthers' Advocates In the Community): protocol and sample description of a cluster randomised trial of mentor mother support to reduce intimate partner violence among pregnant or recent mothers. *BMC Public Health* 2009; **9**: 159–159.
29. MacMillan HL, Wathen CN, Jamieson E, Boyle MH, Shannon HS, Ford-Gilboe M *et al*. Screening for intimate partner violence in health care settings: a randomized trial. *JAMA* 2009; **302**: 493–501.
30. Frommer M, McDonald LE, Millar DS, Collis CM, Watt F, Grigg GW *et al*. A genomic sequencing protocol that yields a positive display of 5-methylcytosine residues in individual. *Proc Natl Acad Sci USA* 1992; **89**: 1827–1831.
31. Clark SJ, Harrison J, Paul CL, Frommer M. High-sensitivity mapping of methylated cytosines. *Nucleic Acids Res* 1994; **22**: 2990–2997.
32. Boyle JS, Lew AM. An inexpensive alternative to glassmilk for DNA purification. *Trends Genet* 1995; **11**: 8.
33. Sambrook J, Russell DW. *Molecular Cloning*. CSHL Press, 2001.
34. Moser D, Molitor A, Kumsta R, Tatschner T, Riederer P, Meyer J. The glucocorticoid receptor gene exon 1-F promoter is not methylated at the NGFI-A binding site in human hippocampus. *World J Biol Psychiatry* 2007; **8**: 262–268.
35. Krug EG, Mercy JA, Dahlberg LL, Zwi AB, Lozano R. *World report on violence and health*. World Health Organization: Geneva, 2002.
36. Seckl JR, Meaney MJ. Glucocorticoid 'programming' and PTSD risk. *Ann NY Acad Sci* 2006; **1071**: 351–378.
37. Kapoor A, Dunn E, Kostaki A, Andrews MH, Matthews SG. Fetal programming of hypothalamo-pituitary-adrenal function: prenatal stress and glucocorticoids. *J Physiol* 2006; **572**: 31–44.
38. Dobbing J, Sands J. Comparative aspects of the brain growth spurt. *Early Hum Dev* 1979; **3**: 79–83.
39. Scarr S, McCartney K. How people make their own environments: a theory of genotype greater than environment effects. *Child Dev* 1983; **54**: 424–435.
40. McGaugh JL. *Memory and Emotion: the Making of lasting Memories*. Columbia University Press: New York, USA, 2003.



**Translational Psychiatry** is an open-access journal published by Nature Publishing Group. This work is licensed under the Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies the paper on the Translational Psychiatry website (<http://www.nature.com/tp>)